



STAT6 and PARP Family Members in the Development of T Cell-dependent Allergic Inflammation

Purna Krishnamurthy^{1,2} and Mark H. Kaplan^{1,2*}

¹Department of Pediatrics, Wells Center for Pediatric Research, ²Department of Microbiology and Immunology, Indiana University School of Medicine, Indianapolis, IN 46202, USA

Allergic inflammation requires the orchestration of altered gene expression in the target tissue and in the infiltrating immune cells. The transcription factor STAT6 is critical in activating cytokine gene expression and cytokine signaling both in the immune cells and in target tissue cells including airway epithelia, keratinocytes and esophageal epithelial cells. STAT6 is activated by the cytokines IL-4 and IL-13 to mediate the pathogenesis of allergic disorders such as asthma, atopic dermatitis, food allergy and eosinophilic esophagitis (EoE). In this review, we summarize the role of STAT6 in allergic diseases, its interaction with the co-factor PARP14 and the molecular mechanisms by which STAT6 and PARP14 regulate gene transcription.

[Immune Network 2016;16(4):201-210]

Keywords: STAT, PARP, Allergic inflammation, T cell

INTRODUCTION

STATs (Signal Transducer and Activator of Transcription) are latent cytoplasmic transcription factors that activate gene transcription in response to cytokines. In response to extracellular cytokine signals and growth factors, STAT proteins undergo tyrosine phosphorylation and are involved in many different regulatory events such as hematopoiesis, immunomodulation and development (1). Phosphorylated STAT6 is transported to the nucleus, where along with co-factors, STAT6 regulates the expression of genes that are involved in allergic inflammatory responses. Upon exposure to an environmental allergen, epithelial cells in the airways, esophagus and the keratinocytes in the skin secrete cytokines that initiate the development of an immune response (2). Chronic challenge with allergen results in organ-specific diseases such as asthma or food allergy that are characterized by pulmonary and

gastrointestinal inflammation respectively.

In this review, we describe the role of STAT6 in the development of allergic disease with a focus on PARP14, a STAT6 transcriptional co-factor, and the related factor PARP1. Although neither PARP protein is required for T cell development, deficiency in either factor has effects on T helper cell differentiation and allergic inflammation. We review the role of STAT6 in allergic airway disease (AAD), atopic dermatitis (AD), food allergy and eosinophilic esophagitis (EoE) (Fig. 1). We then summarize research on the STAT6/PARP14 pathway and the cells where this pathway is important.

STAT6

Signal transducer and activator of transcription 6 (STAT6) is a member of the STAT family of proteins, which

Received on May 4, 2016. Revised on July 17, 2016. Accepted on July 20, 2016.

© This is an open access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Corresponding Author. Mark H. Kaplan, Department of Pediatrics, Indiana University School of Medicine, 1044 West Walnut St., Room 202, Indianapolis, IN 46202, USA. Tel: 1-317-278-3696; Fax: 1-317-274-5378; E-mail: mkaplan2@iu.edu


Abbreviations: AAD, allergic airway disease; AD, atopic dermatitis; AHR, airway hyperreactivity; EoE, eosinophilic esophagitis; PARP, poly-ADP ribosyl-transferase

ASTHMA

T-cells – Th2 differentiation
 Airway Epithelium – Mucous production, Chemokine release
 Airway smooth muscle - Contractility, chemokine release

EOSINOPHILIC ESOPHAGITIS

Esophageal epithelium - Chemokine release

**ATOPIC DERMATITIS**

T-cells – Th2 differentiation
 Keratinocytes – Chemokine release, affects EDC gene expression, Anti-Microbial peptides

FOOD ALLERGY

T-cells – Th2 differentiation
 Intestinal epithelium - Mucous secretion, anti-microbial peptides

Figure 1. Function of PARP14 and STAT6 in allergic inflammation. Although the studies on PARP-14 are still emerging, this figure illustrates the function of PARP14 and STAT6 in asthma, atopic dermatitis, eosinophilic esophagitis and food allergy.

consists of STAT1, 2, 3, 4, 5A, 5B and 6. Human STAT6 is located on chromosome 12q13.3. The cytokines IL-4 and IL-13 activate STAT6 (3). The type I IL-4 receptor is composed of the common gamma chain (γ_c) and the IL4R α chain and binds only IL-4. The type-II IL-4 receptor is composed of IL4R α chain and IL13R α 1 chain and binds to and transmits signals of both IL-4 and IL-13. Three central tyrosine residues in IL4R α chain (Y575, Y603 and Y631) become phosphorylated by stimulation of the receptor and are important for the activation of STAT6. These phosphotyrosines on the receptor chain become docking sites for SH2 domains of STAT6 monomers. The SH2 domains then get tyrosine phosphorylated themselves by receptor associated Jak kinases (Jak1 & 3). This allows for dimerization of STAT proteins via their SH2 domains. The resulting STAT6 dimer can now migrate to the nucleus where it can bind DNA directly (4-6).

Early studies in Stat6^{-/-} mice showed that STAT6 is involved in IL-4 mediated functions including Th2 cell differentiation, class switching to IgE and expression of cell surface markers such as MHC-II and CD23 (7-9). STAT6 regulates the expression of Gata-3 and Gata-3 reconstitutes Th2 development in STAT6-deficient T cells, making it a critical factor for Th2 cell differentiation and function (10).

Asthma

Asthma is a chronic lung disease with intermittent re-

curring periods of coughing and wheezing caused by airway obstruction, abnormal lung physiology and airway hyperresponsiveness (AHR). It affects people of all ages but predominantly starts at childhood. Pathologic manifestations of asthma include airway inflammation, airway remodeling and increased secretion of mucus. Disease initiation and progression depends on Th2 cytokines IL-4, IL-5 and IL-13 and STAT6 activation is involved in IL-4/IL-13 mediated responses and development of inflammation in allergic airway disease (AAD). In an Ovalbumin-induced model of AAD, STAT6 deficient mice have greatly attenuated eosinophilia in BAL, airway hyperresponsiveness (AHR), mucus production and lung inflammation caused by the antigen (11-15). Increased expression of IL-13 and IL-4 in mouse airway epithelial cells are able to reproduce the pathological conditions characteristic of asthma (16,17). As STAT6 is responsible for the clinical manifestations of allergic asthma, many groups are studying it as a therapeutic target. STAT6 inhibitors such as STAT6-IP (Stat6 inhibitory protein) inhibit the Ova-induced development and secretion of Th2 cytokines, mucus production, lung inflammation and AHR (18,19). These findings suggest that STAT6 plays a significant role in the development and progression of AAD.

Atopic dermatitis

Atopic dermatitis (AD) is a chronic, relapsing inflammatory condition of the skin. The key characteristics of

AD include pruritus, scratching and chronic relapsing lesions. AD is characterized by increases in total IgE levels, infiltration of mast cells and eosinophils, decreases in anti-microbial peptides and increased expression of Th2 cytokines (20). The cause of AD is still unknown though genetic factors play an important role. Mutations in the epidermal differentiation complex (EDC, important during terminal differentiation of keratinocytes) genes such as filaggrin (*FLG*) are among those that can cause skin barrier impairment (21,22). STAT6 and Th2 cytokines, IL-4 and IL-13 decrease the expression of EDC genes filaggrin (*FLG*), loricrin (*LOR*) and involucrin (*IVL*) in human skin biopsies and in differentiated primary human keratinocytes (23,24). IL-4 and IL-13 are elevated in the skin of patients with AD and either cytokine can lead to decreased expression of antimicrobial genes like human β defensins (HBD)-2 and 3 (20,25). Atopy is generally associated with increased activation of STAT6, hence a transgenic mouse model with constitutively phosphorylated STAT6 in T cells was generated to define a direct role for STAT6 in allergic inflammation (26). STAT6 transgenic mice expressing a constitutively active form of STAT6 in T cells have a hyper Th2 environment and are predisposed to develop allergic inflammation in the lung, skin and ocular tissue (27-29). The absence of endogenous IL-4 protects the STAT6VT transgenic mice from allergic inflammation (27,28). Transgenic mice expressing IL-4 or IL-13 in the epidermis of the skin developed chronic inflammatory pruritic skin disease, which is very similar to AD in humans (30,31). Thus, STAT6 and Th2 cytokines along with other factors mediate skin barrier disruption and play a crucial role in atopic dermatitis.

Food allergy and eosinophilic esophagitis

As defined by the National Institutes of Allergy and Infectious Diseases (NIAID), food allergy is an abnormal immune response to food. The prevalence of food allergy appears to have increased over the last 2 decades, occurring more frequently in children than adults (32). Food allergy is more common in children with other allergic diseases and is most often IgE mediated. Reaction to food can result in skin manifestations, oral allergy syndrome, and can even result in anaphylaxis. Nut allergy is a common food allergy that has an early onset and is life threatening due to risk of anaphylactic shock. A single nucleotide polymorphism in the 3'UTR of the STAT6 gene is associated with susceptibility and severity to nut allergy in a UK population (33) and has been associated with food sensitization in a Mexican

population (34). Anaphylactic response is usually the result from the activation of intestinal mast cells. IL-4 has an important role in IgE production and *IL4*-deficient mice do not develop experimental food allergy (35,36). Studies have shown that IL-4 signaling has a direct effect on survival and proliferation of intestinal mast cells in food allergy (37). Along with IL-4, IL-9 also has an essential role in promoting intestinal mastocytosis and driving experimental food allergy (38). Studies in mice have shown the presence of multifunctional IL-9 producing mucosal mast cells that induce intestinal mastocytosis and increases in symptoms of food allergy (39). Increased IL-4R α signaling confers increased intestinal permeability and sensitivity to food allergens (40). Oral tolerance to foods requires the development of allergen specific Treg cells and failure of Treg mediated oral tolerance is a major reason for food allergy. Allergen-specific Treg cells undergo Th2 cell reprogramming as a result of enhanced IL4R-STAT6 signaling which is sufficient to impart disease susceptibility (41).

Eosinophilic esophagitis (EoE) is a chronic inflammation of the esophagus and an emerging disease worldwide with a high rate of associated atopic disease (42). It is currently defined as a chronic, immune-mediated or antigen-mediated esophageal disease that is characterized by esophageal dysfunction and eosinophil-predominant inflammation. Children present with food impaction, dysphagia and have esophageal mucosal eosinophilia of >15 eosinophils per high power field (43). STAT6 mediates the expression of eotaxin-3/CCL26 (42,44,45) and hence eosinophil accumulation in EoE (46-48). Esophageal biopsies from children with EoE have an increased expression of CCL26 (49). Th2 cytokines IL-5 and IL-13 downregulate the expression of desmoglein-1 (DSG1) and genes of the epidermal differentiation complex (EDC) like filaggrin (*FLG*) (50,51) and impair barrier function. Altered esophageal barrier could thus lead to enhanced antigen presentation and greater eosinophil recruitment. Taken together, these studies suggest an important role and function of STAT6 and Th2 cytokines in food antigen susceptibility and thus causing gastrointestinal inflammation such as food allergy and eosinophilic esophagitis.

MECHANISMS OF STAT6-DEPENDENT GENE REGULATION

The protein structure and domain organization of STAT6 includes the N-terminal domain that is involved

in dimerization, the coiled-coil domain required for interaction with other proteins, the DNA binding domain, the SH2 domain (src homology 2 domain) that interacts with phosphorylated tyrosine residues in the receptor and the transactivation domain at the C-terminal end. STAT6 preferentially binds to a TTCN₄GAA sequence and activates gene transcription. STAT6 recruits specific sets of co-activators to provide specificity for promoter activation (5). STAT6 recruits co-activators such as CBP/p300 and NCoAs to activate gene expression by providing histone acetyltransferase (HAT) activity. PARP-14 has been identified to be one such co-factor that associates with STAT6 and regulates gene expression.

PARPs

Poly (ADP-ribose) polymerases (PARPs) family of proteins contains 17 homologues, of which PARP1 is the founding member. PARPs are enzymes that catalyze the transfer of ADP ribose moieties from donor NAD⁺ molecules to target proteins although members of the family vary in their enzymatic abilities. Mono- and poly-ADP ribosylation is a protein modification that impacts the regulation of transcription, cell proliferation, apoptosis, signaling cascades and cell differentiation. Since the catalytic product of many of the members is unclear and some don't have a catalytic activity, a new nomenclature was proposed. Based on the type of enzymatic reaction and on structural features, PARP family members are now referred to as ADP-ribosyl transferase Diphtheria-toxin like (ARTD) family of proteins (52).

Of the 17 PARP family members, PARP1 is the most extensively studied. Reports using PARP inhibitors and knockout mice have shown that, in addition to its role in surveillance and maintaining genome integrity, PARP1 also functions in inflammatory diseases by triggering cell death in injured tissue (53). PARP family members including PARP1 and PARP2 are involved in T cell development in the thymus and in the periphery (54,55). PARP1 regulates T cell activation and absence of PARP1 alters expression of genes and chemokines that are involved in maintaining Th1-Th2 balance (56). PARP1 and PARP14 have been studied to have a role in allergic inflammation.

PARP1 (ARTD1) and allergic inflammation: PARP1 was originally characterized for its role as a DNA damage sensor and its function in DNA repair. More recently, PARP1 has been identified for its role in transcription, chromatin structure modification, cell death pathways and in the mitotic apparatus among many others, thus

having outcomes in carcinogenesis, genome maintenance, inflammation, neuronal function and aging (57). PARP1 participates in inflammation by directly or indirectly regulating the expression of inflammatory factors including cytokines, adhesion molecules and iNOS (58). Inflammation in the lung or any other organ causes an increase in PARP expression. PARP1 has emerged as a critical player in the development and the progression of allergic airway disease. Animal models of allergen (Ovalbumin) induced airway inflammation elucidates the positive involvement of PARP1 in airway inflammation, airway hyper responsiveness (AHR) (59,60).

In an Ova triggered model of asthma, mice treated using pharmacological PARP inhibitors to target PARP1 and *Parp1*^{-/-} mice had diminished AAD characterized by decreasing infiltration of inflammatory cells into the lung, reducing AHR and mucus secretion (59-62). This protective effect of absence of PARP1 is attributed to the inhibition of NF- κ B activation and regulating the expression of molecules like TNF- α , iNOS and NFATc (59,63). Decreases in the amount of IL-5 was linked to reduction in the number of eosinophils infiltrating the lung of *Parp1*^{-/-} mice, suggesting that PARP1 regulates IL-5 production (58). The absence of PARP1 or its enzymatic activity severely decreases STAT6 protein without having an effect on STAT6 mRNA levels. STAT6 protein degradation was allergen stimulated and mediated by calpain as observed in an asthma model in *Parp1*^{-/-} mice. This in turn results in reduced IL-5 levels, decreased eosinophilia and lung inflammation (64). Together, studies of PARP1 in models of asthma suggest that PARP1 directly regulates STAT6 and its absence can mitigate all signs of inflammation.

PARP14 (ARTD8) and allergic inflammation: PARP family members are classified into sub-families based on their domain architecture. One such sub-family is macro-PARPs that include PARP9, PARP14 and PARP15. Members of this family contain repeats of a domain found in the non-histone-like region of variant histone macroH2A (65,66). PARP14 is enzymatically active and catalyzes mono-ADP ribosylation. Studies from many labs have shown that PARP14 functions in transcription regulation, signal transduction pathways and the actin cytoskeleton (67). PARP14 lacks the conventional transcriptional co-regulator domains such as HAT, Bromo, Chromo, SET or ATPase domains.

PARP14 associates with STAT6 in the presence or the absence of IL-4, but not with STAT1. The ability of PARP14 to promote transcriptional enhancement of IL-4 dependent and not IFN γ dependent gene activation lead to

this factor initially being named CoaSt6 (for collaborator of STAT6) (Fig. 2) (68). Although the macro domains of histone macro H2A are known to participate in gene silencing, the macro domains of PARP14 interact with STAT6 and increase IL-4 induced gene expression (68). PARP14 and its associated PARP catalytic activity can mono-ADP-ribosylate itself and a STAT6 co-activator, p100 (Fig. 2). The enzymatic activity of PARP14 is required to enhance gene transcription and blocking PARP activity using pharmacological inhibitors blocked IL-4 dependent gene transcription *in vivo* (69). Studies show that PARP14 regulates STAT6-dependent gene transcription by functioning as a transcriptional switch. Under non-stimulating conditions, PARP14 binds to the STAT6-activated promoter and recruits HDAC2 and 3 to repress gene transcription. Upon IL-4 stimulation and STAT6 activation, the catalytic activity of PARP14 is increased resulting in ADP-ribosylation of itself and HDACs. The HDAC repressor complex leaves the promoter relieving gene repression and making the promoter more accessible to histone acetyl transferases (CBP/p300, NCoA-1 and NCoA-3) and increase gene transcription (Fig. 2) (70). As PARP14 functions as a transcriptional switch for STAT6 dependent gene transcription, naïve T cells from *Parp14*^{-/-} mice cultured under Th2 conditions secrete reduced amounts of Th2 cytokines IL-4, IL-5 and IL-13 (71). Inhibiting PARP

enzymatic activity using a pharmacological inhibitor confirmed a dose dependent reduction in Th2 cytokines, with similar amounts of IFN γ produced from Th1 skewed cells (71). The mechanism for promoting Th2 differentiation is based on the ability of PARP14 or its enzymatic activity to regulate the binding of STAT6 to the *Gata3* promoter. In a model of AAD, *Parp14*^{-/-} mice have reduced numbers of inflammatory cells in BAL and reduced AHR.

Until recently, the role of PARP14 in other T helper cell subset differentiation and development was not known. Th17 cells are defined by their ability to produce IL-17A, IL-17F and IL-22 and to a lesser degree TNF and IL-6. They play a pivotal role in host defense and in induction and propagation of autoimmunity (72). Increases in IL-17A is correlated with increases in neutrophil recruitment and airway resistance in patients with asthma (73). T follicular helper (Tfh) cells are responsible for providing cognate help to B cells during formation of germinal centers. CD4 Tfh cells secrete IL-4, IL-10 and IL-21. IL-21 is required for germinal center formation for T-cell dependent antigens (74). Tfh cells are essential for IgE production and play an important role in allergic responses (75). ChIP-Seq analysis of WT and *Parp14*^{-/-} Th2 cells identified that PARP14 along with its enzymatic activity regulated the expression of *Ii21*, and IL-21 promotes Th17 and Tfh development and function (76). It was also




Figure 2. PARP14 (ARTD8) and its enzymatic activity functions as an activator of STAT6. In the presence of IL-4, PARP14 mono-ADP ribosylates histone deacetylase-2 and -3 (HDAC-2 and HDAC-3), leading to their dissociation from the IL-4 responsive promoter. This allows STAT6 and transcription co-factors such as p100 to bind to the promoter, resulting in active gene transcription.

identified from the ChIP-Seq data that the expression of a number of genes were dependent on the enzymatic activity of PARP-14 but were not regulated by STAT6, suggesting that PARP14 has STAT6 independent functions (76). Upon further studies, the absence of PARP14 or its enzymatic activity resulted in decreases in Th17 cells in the allergic airway inflammation model. PARP14 also promotes Tfh development by regulating the activation of STAT3, which is crucial for the differentiation of both Th17 and Tfh cell development (77). Similarly, Th17 and Tfh differentiation is diminished in PARP14 deficient T cells coincident with decreases in the phosphorylation of STAT3 (77).

PARP14 and its enzymatic activity also have functions in other cell types. In B-cells, PARP14 is required to maintain balance of B-cell subsets in the spleen and in generating IgA responses to antigen. PARP14 induces the expression of B-cell survival factors such as Pim-1 and Mcl-1 and protects IL-4 treated cells from apoptosis (78). PARP-14 deficient mice have significantly reduced numbers and frequency of Tfh cells. Tfh cells promote GC B cell development and a 45% reduction in the numbers and frequency of GC B cells was observed along with reductions in serum IgG1 and IgG2a/c titres in PARP14 deficient mice in response to SRBC immunizations. These could either be B-cell intrinsic effects or extrinsic mechanisms due to reduced Tfh numbers (77). Epithelial cells express PARP14 and this pathway has a significant role in airway epithelial cells where PARP14 deficient mice subjected to OVA induced AAD had diminished AHR (71). In esophageal epithelial cells, STAT6 regulates the expression of eotaxin-3 and it is found to be dramatically increased in patients with EoE (79,80). PARP14 expression is increased in EoE biopsy samples with a positive correlation to eotaxin-3 expression, which was entirely dependent on STAT6 binding to eotaxin-3 promoter (49). Inhibiting the enzymatic activity of PARP14 in human esophageal epithelial cells cultured with IL4 and/or IL13, decreased the expression of eosinophil chemoattractant – eotaxin-3 (*CCL26*) (49). Keratinocytes, which are highly specialized epithelial cells also express PARP14. Our preliminary studies suggest that unlike its function in airway epithelial cells and esophageal epithelial cells, PARP14 does not impact expression of *CCL26* or other epidermal differentiation genes (unpublished observations). These data together propose that PARP14 is expressed by a wide array of cells and STAT6/PARP14 pathway functions differently in specific cell types.

Contrasting the roles of PARP1 and PARP14 in allergic

inflammation: As described in the previous sections, PARP1 and PARP14 work in distinct but overlapping pathways. In T cells, PARP1 has also been shown to modulate transcription factors such as STAT6, NFAT and NF- κ B (59,63), while PARP14 primarily regulates STAT6-dependent gene induction (70) and the activation of STAT3 (77). Although deficiency in PARP1 or PARP14 has similar outcomes in STAT6-mediated allergic inflammation, they regulate *GATA3* gene expression and hence Th2 differentiation in very different ways. In the absence of PARP1, calpain mediates the degradation of STAT6 to reduce protein concentration within the cell (64), whereas in the absence of PARP14, HDACs remain bound to the *Gata3* promoter, inhibiting gene expression (70). The absence of PARP14 and its enzymatic activity resulted in a reduced expression of IL-17A, IL-17F and IL-21 (77), whereas the absence of PARP1 has no effect on IL17 production (81). Lung cells of ovalbumin challenged *Parp1*^{-/-} mice fail to synthesize GM-CSF and IL-5 resulting in impaired eosinophil recruitment (58). In comparison, the significant reduction in lung eosinophilia in *Parp14*^{-/-} mice is attributed to the decrease in expression of *CCL24* (71). Production of IL-4 was inhibited to a lesser extent than IL-5 and IL-13 following PARP-1 inhibition (58). Highlighting differences in mechanism, in clinical samples, there was a positive correlation between PARP14 and *CCL26* expression in EoE patient biopsies, the same was not observed between PARP1 and *CCL26* (49). Together, these studies suggest that PARP1 and PARP14 interact with different partners and regulate inflammation through independent functions.

CONCLUDING REMARKS

STAT6 plays varied functions in allergic inflammation both in immune cells and in resident tissue cells such as airway, esophageal, and intestinal epithelial cells, and keratinocytes (Fig. 1). Current therapies to treat allergic disease are targeted towards disease symptoms rather than the causative factor. Since STAT6 plays a pivotal role in allergic diseases, targeting the function of STAT6 has been an appealing therapeutic strategy. Along with STAT6, inhibiting the transcriptional co-activator PARP14 would be helpful. PARP inhibition might have beneficial effects in inflammatory conditions such as asthma (58,62) and contact hypersensitivity (82,83). In a mouse model of AAD, *Parp14*^{-/-} mice have decreased lung inflammation, airway hyper responsiveness and serum IgE (71). Administering a pharmacological PARP

inhibitor –PJ34 (that inhibits the enzymatic activity of PARPs) to wild-type mice that were subjected to AAD, reduced the severity of airway hyperresponsiveness and lung inflammation along with a reduction in the Th2 responses. Administering PJ34 either during challenge or sensitization and challenge or multiple times during the sensitization phase were effective (71). Recent studies show that PARP14 is required for STAT3 activation (77), suggesting that it could be a potential therapeutic target not only for allergic inflammation but also in other immune responses where this signaling pathway is important.

ACKNOWLEDGEMENTS

The authors dedicate this article to the memory of Shreevrat Goenka who began the PARP14/ARTD8/Coast6 field a decade ago.

REFERENCES

1. Levy, D. E., and J. E. Darnell. 2002. Stats: transcriptional control and biological impact. *Nat. Rev. Mol. Cell Biol.* 3: 651-662.
2. Galli, S. J., M. Tsai, and A. M. Piliponsky. 2008. The development of allergic inflammation. *Nature* 454: 445-454.
3. Quelle, F. W., K. Shimoda, W. Thierfelder, C. Fischer, A. Kim, S. M. Ruben, J. L. Cleveland, J. H. Pierce, A. D. Keegan, K. Nelms, W. E. Paul, and J. N. Ihle. 1995. Cloning of murine Stat6 and human Stat6, Stat proteins that are tyrosine phosphorylated in responses to IL-4 and IL-3 but are not required for mitogenesis. *Mol. Cell. Biol.* 15: 3336-3343.
4. Mikita, T., D. Campbell, P. Wu, K. Williamson, and U. Schindler. 1996. Requirements for interleukin-4-induced gene expression and functional characterization of Stat6. *Mol. Cell. Biol.* 16: 5811-5820.
5. Wurster, A. L., T. Tanaka, and M. J. Grusby. 2000. The biology of Stat4 and Stat6. *Oncogene* 19: 2577-2584.
6. Hebenstreit, D., G. Wirnsberger, J. Horejs-hoeck, and A. Duschl. 2006. Signaling mechanisms, interaction partners, and target genes of STAT6. *Cytokine Growth Factor Rev.* 17: 173-188.
7. Takeda, K., T. Tanaka, W. Shi, M. Matsumoto, M. Minami, S. Kashiwamura, K. Nakanishi, N. Yoshida, T. Kishimoto, and S. Akira. 1996. Essential role of Stat6 in IL-4 signalling. *Nature* 380: 627-630.
8. Shimoda, K., J. van Deursen, M. Y. Sangster, S. R. Sarawar, R. T. Carson, R. A. Tripp, C. Chu, F. W. Quelle, T. Nosaka, D. A. Vignali, P. C. Doherty, G. Grosveld, W. E. Paul, and J. N. Ihle. 1996. Lack of IL-4-induced Th2 response and IgE class switching in mice with disrupted Stat6 gene. *Nature* 380: 630-633.
9. Kaplan, M. H., U. Schindler, S. T. Smiley, and M. J. Grusby. 1996. Stat6 is required for mediating responses to IL-4 and for the development of Th2 cells. *Immunity* 4: 313-319.
10. Ouyang, W., M. Löhning, Z. Gao, M. Assenmacher, S. Ranganath, a Radbruch, and K. M. Murphy. 2000. Stat6-independent GATA-3 autoactivation directs IL-4-independent Th2 development and commitment. *Immunity* 12: 27-37.
11. Kuperman, D., B. Schofield, M. Wills-Karp, and M. J. Grusby. 1998. Signal transducer and activator of transcription factor 6 (Stat6)-deficient mice are protected from antigen-induced airway hyperresponsiveness and mucus production. *J. Exp. Med.* 187: 939-948.
12. Akimoto, T., F. Numata, M. Tamura, Y. Takata, N. Higashida, T. Takashi, K. Takeda, and S. Akira. 1998. Abrogation of bronchial eosinophilic inflammation and airway hyperreactivity in signal transducers and activators of transcription (STAT)6-deficient mice. *J. Exp. Med.* 187: 1537-1542.
13. Tomkinson, A., A. Kanehiro, N. Rabinovitch, A. Joetham, G. Cieslewicz, and E. W. Gelfand. 1999. The failure of STAT6-deficient mice to develop airway eosinophilia and airway hyperresponsiveness is overcome by interleukin-5. *Am. J. Respir. Crit. Care Med.* 160: 1283-1291.
14. Webb, D. C., A. N. J. McKenzie, A. M. L. Koskinen, M. Yang, J. Mattes, and P. S. Foster. 2000. Integrated signals between IL-13, IL-4, and IL-5 regulate airways hyperreactivity. *J. Immunol.* 165: 108-113.
15. Hoshino, A., T. Tsuji, J. Matsuzaki, T. Jinushi, S. Ashino, T. Teramura, K. Chamoto, Y. Tanaka, Y. Asakura, T. Sakurai, Y. Mita, A. Takaoka, S. Nakaike, T. Takeshima, H. Ikeda, and T. Nishimura. 2004. STAT6-mediated signaling in Th2-dependent allergic asthma: Critical role for the development of eosinophilia, airway hyper-responsiveness and mucus hypersecretion, distinct from its role in Th2 differentiation. *Int. Immunol.* 16: 1497-1505.
16. Elias, J. A., Z. Zhu, G. Chupp, and R. J. Homer. 1999. Airway remodeling in asthma. *J. Clin. Invest.* 104: 1001-1006.
17. Kuperman, D. A., X. Huang, L. L. Koth, G. H. Chang, G. M. Dolganov, Z. Zhu, J. A. Elias, D. Sheppard, and D. J. Erle. 2002. Direct effects of interleukin-13 on epithelial cells cause airway hyperreactivity and mucus overproduction in asthma. *Nat. Med.* 8: 885-889.
18. McCusker, C. T., Y. Wang, J. Shan, M. W. Kinyanjui, A. Villeneuve, H. Michael, and E. D. Fixman. 2007. Inhibition of experimental allergic airways disease by local application of a cell-penetrating dominant-negative STAT-6 peptide. *J. Immunol.* 179: 2556-2564.

19. Chiba, Y., M. Todoroki, Y. Nishida, M. Tanabe, and M. Misawa. 2009. A novel STAT6 inhibitor AS1517499 ameliorates antigen-induced bronchial hypercontractility in mice. *Am. J. Respir. Cell Mol. Biol.* 41: 516-524.
20. Hamid, Q., M. Boguniewicz, and D. Y. M. Leung. 1994. Differential in situ cytokine gene expression in acute versus chronic atopic dermatitis. *J. Clin. Invest.* 94: 870-6.
21. Palmer, C. N. A., A. D. Irvine, A. Terron-Kwiatkowski, Y. Zhao, H. Liao, S. P. Lee, D. R. Goudie, A. Sandilands, L. E. Campbell, F. J. D. Smith, G. M. O'Regan, R. M. Watson, J. E. Cecil, S. J. Bale, J. G. Compton, J. J. DiGiovanna, P. Fleckman, S. Lewis-Jones, G. Arseculeratne, A. Sergeant, C. S. Munro, B. El Houate, K. McElreavey, L. B. Halkjaer, H. Bisgaard, S. Mukhopadhyay, and W. H. I. McLean. 2006. Common loss-of-function variants of the epidermal barrier protein filaggrin are a major predisposing factor for atopic dermatitis. *Nat. Genet.* 38: 441-446.
22. Baurecht, H., A. D. Irvine, N. Novak, T. Illig, B. Bühler, J. Ring, S. Wagenpfeil, and S. Weidinger. 2007. Toward a major risk factor for atopic eczema: meta-analysis of filaggrin polymorphism data. *J. Allergy Clin. Immunol.* 120: 1406-1412.
23. Howell, M. D., B. E. Kim, P. Gao, A. V. Grant, M. Boguniewicz, A. DeBenedetto, L. Schneider, L. A. Beck, K. C. Barnes, and D. Y. M. Leung. 2007. Cytokine modulation of atopic dermatitis filaggrin skin expression. *J. Allergy Clin. Immunol.* 120: 150-155.
24. Kim, B. E., D. Y. M. Leung, M. Boguniewicz, and M. D. Howell. 2008. Loricrin and involucrin expression is down-regulated by Th2 cytokines through STAT-6. *Clin. Immunol.* 126: 332-337.
25. Nomura, I., E. Goleva, M. D. Howell, Q. A. Hamid, P. Y. Ong, C. F. Hall, M. A. Darst, B. Gao, M. Boguniewicz, J. B. Travers, and D. Y. M. Leung. 2003. Cytokine milieu of atopic dermatitis, as compared to psoriasis, skin prevents induction of innate immune response genes. *J. Immunol.* 171: 3262-3269.
26. Bruns, H. A., U. Schindler, and M. H. Kaplan. 2003. Expression of a constitutively active Stat6 *in vivo* alters lymphocyte homeostasis with distinct effects in T and B cells. *J. Immunol.* 170: 3478-3487.
27. Sehra, S., H. A. Bruns, A.-N. N. Ahyi, E. T. Nguyen, N. W. Schmidt, E. G. Michels, G.-U. von Bülow, and M. H. Kaplan. 2008. IL-4 is a critical determinant in the generation of allergic inflammation initiated by a constitutively active Stat6. *J. Immunol.* 180: 3551-3559.
28. Sehra, S., Y. Yao, M. D. Howell, E. T. Nguyen, G. S. Kansas, D. Y. M. Leung, J. B. Travers, and M. H. Kaplan. 2010. IL-4 regulates skin homeostasis and the predisposition toward allergic skin inflammation. *J. Immunol.* 184: 3186-3190.
29. Turner, M. J., S. Dasilva-arnold, N. Luo, X. Hu, C. C. West, L. Sun, C. Hall, J. Bradish, M. H. Kaplan, J. B. Travers, and Y. Sun. 2014. STAT6-mediated keratitis and blepharitis : a novel murine model of ocular atopic dermatitis. *Invest. Ophthalmol. Vis. Sci.* 55: 3803-3808.
30. Chan, L. S., N. Robinson, and L. Xu. 2001. Expression of interleukin-4 in the epidermis of transgenic mice results in a pruritic inflammatory skin disease: an experimental animal model to study atopic dermatitis. *J. Invest. Dermatol.* 117: 977-983.
31. Zheng, T., M. H. Oh, S. Y. Oh, J. T. Schroeder, A. B. Glick, and Z. Zhu. 2009. Transgenic expression of interleukin-13 in the skin induces a pruritic dermatitis and skin remodeling. *J. Invest. Dermatol.* 129: 742-751.
32. Sicherer, S. H., and D. Y. M. Leung. 2015. Advances in allergic skin disease, anaphylaxis, and hypersensitivity reactions to foods, drugs, and insects in 2014. *J. Allergy Clin. Immunol.* 135: 357-367.
33. Amoli, M., S. Hand, A. Hajeer, K. Jones, S. Rolf, C. Sting, B. Davies, and W. Ollier. 2002. Polymorphism in the Stat6 gene encodes risk for nut allergy. *Genes Immun.* 3: 220-224.
34. Hancock, D. B., I. Romieu, G. Y. Chiu, J.-J. Sienra-Monge, H. Li, B. E. del Rio-Navarro, and S. J. London. 2012. STAT6 and LRP1 polymorphisms are associated with food allergen sensitization in Mexican children. *J. Allergy Clin. Immunol.* 129: 1673-1676.
35. Kweon, M., M. Yamamoto, M. Kajiki, I. Takahashi, and H. Kiyono. 2000. Systemically derived large intestinal CD4⁺ Th2 cells play a central role in STAT6-mediated allergic diarrhea. *J. Clin. Invest.* 106: 199-206.
36. Brandt, E. B., A. Munitz, T. Orekov, M. K. Mingler, M. McBride, F. D. Finkelman, and M. E. Rothenberg. 2009. Targeting IL-4 / IL-13 signaling to alleviate oral allergen - induced diarrhea. *J. Allergy Clin. Immunol.* 123: 53-58.
37. Burton, O. T., A. R. Darling, J. S. Zhou, M. Noval-Rivas, T. G. Jones, M. F. Gurish, T. A. Chatila, and H. C. Oettgen. 2013. Direct effects of IL-4 on mast cells drive their intestinal expansion and increase susceptibility to anaphylaxis in a murine model of food allergy. *Mucosal Immunol.* 6: 740-750.
38. Forbes, E. E., K. Groschwitz, J. P. Abonia, E. B. Brandt, E. Cohen, C. Blanchard, R. Ahrens, L. Seidu, A. Mckenzie, R. Strait, F. D. Finkelman, P. S. Foster, K. I. Matthaei, M. E. Rothenberg, and S. P. Hogan. 2008. IL-9 - and mast cell - mediated intestinal permeability predisposes to oral antigen hypersensitivity. *J. Exp. Med.* 205: 897-913.
39. Chen, C., J. Lee, B. Liu, S. Ohta, P. Wang, A. V Kartashov, L. Mugge, J. P. Abonia, A. Barski, K. Izuhara, M. E. Rothenberg, F. D. Finkelman, and S. P. Hogan. 2015. Induction of interleukin-9-producing mucosal mast cells promotes susceptibility to IgE-mediated article induction of interleukin-9-producing mucosal mast cells promotes susceptibility to IgE-

- mediated experimental food allergy. *Immunity* 43: 1-15.
40. Mathias, C. B., S. A. Hobson, M. Garcia-Iloret, G. Lawson, D. Poddighe, E.-J. Freyschmidt, W. Xing, M. F. Gurish, T. A. Chatila, and H. C. Oettgen. 2011. Mechanisms of allergy and clinical immunology IgE-mediated systemic anaphylaxis and impaired tolerance to food antigens in mice with enhanced IL-4 receptor signaling. *J. Allergy Clin. Immunol.* 127: 795-805.
 41. Rivas, M. N., O. T. Burton, P. Wise, L. Charbonnier, P. Georgiev, H. C. Oettgen, R. Rachid, and T. A. Chatila. 2015. Regulatory T cell reprogramming toward a Th2-cell-like lineage impairs oral tolerance and promotes regulatory T cell reprogramming toward a Th2-cell-like lineage impairs oral tolerance and promotes food allergy. *Immunity* 42: 512-523.
 42. Blanchard, C., S. Durual, M. Estienne, S. Emami, S. Vasseur, and J. C. Cuber. 2005. Eotaxin-3/CCL26 gene expression in intestinal epithelial cells is up-regulated by interleukin-4 and interleukin-13 via the signal transducer and activator of transcription 6. *Int. J. Biochem. Cell Biol.* 37: 2559-2573.
 43. Furuta, G. T., and D. A. Katzka. 2015. Eosinophilic Esophagitis. *N. Engl. J. Med.* 373: 1640-1648.
 44. Hoeck, J., and M. Woiseschläger. 2001. Activation of eotaxin-3/CCL26 gene expression in human dermal fibroblasts is mediated by STAT6. *J. Immunol.* 167: 3216-3222.
 45. Kagami, S., H. Saeki, M. Komine, T. Kakinuma, Y. Tsunemi, K. Nakamura, K. Sasaki, A. Asahina, and K. Tamaki. 2005. Interleukin-4 and interleukin-13 enhance CCL26 production in a human keratinocyte cell line, HaCaT cells. *Clin. Exp. Immunol.* 141: 459-466.
 46. Mishra, A., and M. E. Rothenberg. 2003. Intratracheal IL-13 induces eosinophilic esophagitis by an IL-5, eotaxin-1, and STAT6-dependent mechanism. *Gastroenterology* 125: 1419-1427.
 47. Zhang, X., E. Cheng, X. Huo, C. Yu, Q. Zhang, T. H. Pham, D. H. Wang, S. J. Spechler, and R. F. Souza. 2012. Omeprazole blocks STAT6 binding to the eotaxin-3 promoter in eosinophilic esophagitis cells. *PLoS One* 7: e50037.
 48. Niranjana, R., M. Rayapudi, A. Mishra, P. Dutt, S. Dynda, and A. Mishra. 2013. Pathogenesis of allergen-induced eosinophilic esophagitis is independent of interleukin (IL)-13. *Immunol. Cell Biol.* 91: 408-415.
 49. Krishnamurthy, P., J. D. Sherrill, K. Parashette, S. Goenka, M. E. Rothenberg, S. Gupta, and M. H. Kaplan. 2014. Correlation of increased PARP14 and CCL26 expression in biopsies from children with eosinophilic esophagitis. *J. Allergy Clin. Immunol.* 133: 577-580.
 50. Blanchard, C., E. M. Stucke, K. Burwinkel, J. M. Caldwell, M. H. Collins, A. Ahrens, B. K. Buckmeier, S. C. Jameson, A. Greenberg, A. Kaul, J. P. Franciosi, J. P. Kushner, L. J. Martin, P. E. Putnam, J. P. Abonia, S. I. Wells, and M. E. Rothenberg. 2010. Coordinate interaction between IL-13 and epithelial differentiation cluster genes in eosinophilic esophagitis. *J. Immunol.* 184: 4033-4041.
 51. Sherrill, J. D., K. Kc, D. Wu, Z. Djukic, J. M. Caldwell, E. M. Stucke, K. A. Kemme, M. S. Costello, M. K. Mingler, C. Blanchard, M. H. Collins, J. P. Abonia, P. E. Putnam, E. S. Dellon, R. C. Orlando, S. P. Hogan, and M. E. Rothenberg. 2014. Desmoglein-1 regulates esophageal epithelial barrier function and immune responses in eosinophilic esophagitis. *Mucosal Immunol.* 7: 718-729.
 52. Hottiger, M. O., P. O. Hassa, B. Lüscher, H. Schüler, and F. Koch-Nolte. 2010. Toward a unified nomenclature for mammalian ADP-ribosyltransferases. *Trends Biochem. Sci.* 35: 208-219.
 53. Schreiber, V., F. Dantzer, J.-C. Ame, and G. de Murcia. 2006. Poly(ADP-ribose): novel functions for an old molecule. *Nat. Rev. Mol. Cell Biol.* 7: 517-528.
 54. Yélamos, J., Y. Monreal, L. Saenz, E. Aguado, V. Schreiber, R. Mota, T. Fuente, A. Minguela, P. Parrilla, G. de Murcia, E. Almarza, P. Aparicio, and J. Ménissier-de Murcia. 2006. PARP-2 deficiency affects the survival of CD4⁺CD8⁺ double-positive thymocytes. *EMBO J.* 25: 4350-4360.
 55. Rosado, M. M., E. Bennici, F. Novelli, and C. Pioli. 2013. Beyond DNA repair, the immunological role of PARP-1 and its siblings. *Immunology* 139: 428-437.
 56. Saenz, L., J. J. Lozano, R. Valdor, A. Baroja-Mazo, P. Ramirez, P. Parrilla, P. Aparicio, L. Sumoy, and J. Yélamos. 2008. Transcriptional regulation by poly(ADP-ribose) polymerase-1 during T cell activation. *BMC Genomics* 9: 171.
 57. Kim, M. Y., T. Zhang, and W. L. Kraus. 2005. Poly (ADP-ribose) ation by PARP-1 : “ PAR-laying ” NAD⁺ into a nuclear signal. *Genes Dev* 19: 1951-1967.
 58. Oumouna, M., R. Datta, K. Oumouna-Benachour, Y. Suzuki, C. Hans, K. Matthews, K. Fallon, and H. Boulares. 2006. Poly(ADP-ribose) polymerase-1 inhibition prevents eosinophil recruitment by modulating Th2 cytokines in a murine model of allergic airway inflammation: a potential specific effect on IL-5. *J. Immunol.* 177: 6489-6496.
 59. Boulares, A. H., A. J. Zoltoski, Z. A. Sherif, P. Jolly, D. Massaro, and M. E. Smulson. 2003. Gene knockout or pharmacological inhibition of poly(ADP-ribose) polymerase-1 prevents lung inflammation in a murine model of asthma. *Am. J. Respir. Cell Mol. Biol.* 28: 322-329.
 60. Suzuki, Y., E. Masini, C. Mazzocca, S. Cuzzocrea, A. Ciampa, H. Suzuki, and D. Bani. 2004. Inhibition of Poly (ADP-Ribose) polymerase prevents allergen- induced asthma-like reaction in sensitized guinea pigs. *Pharmacology* 311: 1241-1248.
 61. Virág, L., P. Bai, I. Bak, P. Pacher, J. G. Mabley, L. Liaudet, E. Bakondi, P. Gergely, M. Kollai, and C. Szabó. 2004. Effects of poly(ADP-ribose) polymerase inhibition on inflammatory cell

- migration in a murine model of asthma. *Med. Sci. Monit.* 10: BR77-R83.
62. Naura, A. S., C. P. Hans, M. Zerfaoui, D. You, S. a Cormier, M. Oumouna, and A. H. Boulares. 2008. Post-allergen challenge inhibition of poly(ADP-ribose) polymerase harbors therapeutic potential for treatment of allergic airway inflammation. *Clin. Exp. Allergy* 38: 839-846.
 63. Olabisi, O. A., N. Soto-Nieves, E. Nieves, T. T. C. Yang, X. Yang, R. Y. L. Yu, H. Y. Suk, F. Macian, and C.-W. Chow. 2008. Regulation of transcription factor NFAT by ADP-ribosylation. *Mol. Cell. Biol.* 28: 2860-2871.
 64. Datta, R., A. S. Naura, M. Zerfaoui, Y. Errami, M. Oumouna, H. Kim, J. Ju, V. P. Ronchi, A. L. Haas, and A. H. Boulares. 2011. PARP-1 deficiency blocks IL-5 expression through calpain-dependent degradation of STAT-6 in a murine asthma model. *Allergy Eur. J. Allergy Clin. Immunol.* 66: 853-861.
 65. Pehrson, J.R., V. A. Fried. 1992. MacroH2A, a core histone containing a large nonhistone region. *Science* 257:1398-400
 66. Ladurner, A. G. 2003. Inactivating chromosomes: A macro domain that minimizes transcription. *Mol. Cell* 12: 1-4.
 67. Vyas, S., I. Matic, L. Uchima, J. Rood, R. Zaja, R. T. Hay, I. Ahel, and P. Chang. 2014. Family-wide analysis of poly(ADP-ribose) polymerase activity. *Nat. Commun.* 5: 4426.
 68. Goenka, S., and M. Boothby. 2006. Selective potentiation of Stat-dependent gene expression by collaborator of Stat6 (CoaSt6), a transcriptional cofactor. *Proc. Natl. Acad. Sci. U. S. A.* 103: 4210-4215.
 69. Goenka, S., H. C. Sung, and M. Boothby. 2007. Collaborator of Stat6 (CoaSt6)-associated poly(ADP-ribose) polymerase activity modulates Stat6-dependent gene transcription. *J. Biol. Chem.* 282: 18732-18739.
 70. Mehrotra, P., J. P. Riley, R. Patel, F. Li, L. Voss, and S. Goenka. 2011. PARP-14 functions as a transcriptional switch for Stat6-dependent gene activation. *J. Biol. Chem.* 286: 1767-1776.
 71. Mehrotra, P., A. Hollenbeck, J. P. Riley, F. Li, R. J. Patel, N. Akhtar, and S. Goenka. 2013. Poly (ADP-ribose) polymerase 14 and its enzyme activity regulates TH2 differentiation and allergic airway disease. *J. Allergy Clin. Immunol.* 131: 521-531.
 72. Bettelli, E., T. Korn, and V. K. Kuchroo. 2007. Th17 : the third member of the effector T cell trilogy. *Curr. Opin. Immunol.* 19: 652-657.
 73. Alcorn, J. F., C. R. Crowe, and J. K. Kolls. 2010. TH17 cells in asthma and COPD. *Annu. Rev. Physiol.* 72: 495-516.
 74. Kemeny, D. M. 2012. The role of the T follicular helper cells in allergic disease. *Cell. Mol. Immunol.* 9: 386-389.
 75. Crotty, S. 2014. Review T follicular helper cell differentiation , function , and roles in disease. *Immunity* 41: 529-542.
 76. Riley, J. P., A. Kulkarni, P. Mehrotra, B. Koh, N. B. Perumal, M. H. Kaplan, and S. Goenka. 2013. PARP-14 binds specific DNA sequences to promote Th2 cell gene expression. *PLoS One* 8: e83127.
 77. Mehrotra, P., P. Krishnamurthy, J. Sun, S. Goenka, and M. H. Kaplan. 2015. Poly-ADP-ribosyl polymerase-14 promotes T helper 17 and follicular T helper development. *Immunology* 146: 537-546.
 78. Cho, S. H., S. Goenka, T. Henttinen, P. Gudapati, A. Reini-kainen, C. M. Eischen, R. Lahesmaa, and M. Boothby. 2009. PARP-14, a member of the B aggressive lymphoma family, transduces survival signals in primary B cells. *Blood* 113: 2416-2425.
 79. Blanchard, C., N. Wang, K. F. Stringer, A. Mishra, P. C. Fulkerson, J. P. Abonia, S. C. Jameson, C. Kirby, M. R. Konikoff, M. H. Collins, M. B. Cohen, R. Akers, S. P. Hogan, A. H. Assa`ad, P. E. Putnam, B. J. Aronow, and M. E. Rothenberg. 2006. Eotaxin-3 and a uniquely conserved gene expression profile in eosinophilic esophagitis. *J. Clin. Invest.* 116: 536-547.
 80. Gupta, S. K., J. F. Fitzgerald, T. Kondratyuk, and H. HogenEsch. 2006. Cytokine expression in normal and inflamed esophageal mucosa: a study into the pathogenesis of allergic eosinophilic esophagitis. *J. Pediatr. Gastroenterol. Nutr.* 42: 22-26.
 81. Nasta, F., F. Laudisi, M. Sambucci, M. M. Rosado and C. Pioli. 2010. Increased Foxp3+ regulatory T cells in poly (ADP-ribose) polymerase -1 deficiency. *J. Immunol.* 184: 3470-3477.
 82. Bai, P., C. Hegedus, E. Szabo, L. Gyure, E. Bakondi, A. Brunyanszki, S. Gergely, C. Szabo, and L. Virag. 2009. Poly (ADP-Ribose) polymerase mediates inflammation in a mouse model of contact hypersensitivity. *J. Invest. Dermatol.* 129: 234-238.
 83. Brunyanszki, A., C. Hegedus, M. Szanto, K. Erdelyi, K. Kovacs, V. Schreiber, S. Gergely, B. Kiss, E. Szabo, L. Virag, and P. Bai. 2010. Genetic ablation of PARP-1 protects against oxazolone-induced contact hypersensitivity by modulating oxidative stress. *J. Invest. Dermatol.* 130: 2629-2637.