

Inhibition of the *Aspergillus flavus* Growth and Aflatoxin B1 Contamination on Pistachio Nut by Fengycin and Surfactin-Producing *Bacillus subtilis* UTBSP1

Mohsen Farzaneh^{1,2*}, Zhi-Qi Shi², Masoud Ahmadzadeh³, Liang-Bin Hu^{2,4}, and Alireza Ghassempour¹

¹Medicinal Plants and Drugs Research Institute, Shahid Beheshti University, Tehran 19835-389, Iran

²Institute of Food Safety and Quality, Jiangsu Academy of Agricultural Sciences, Nanjing 210014, China

³Department of Plant Protection, University of Tehran, Karaj 4111, Iran

⁴School of Food, Henan Institute of Science and Technology, Xinxiang 453003, China

(Received on November 24, 2015; Revised on January 18, 2016; Accepted on January 20, 2016)

In this study, the treatment of pistachio nuts by *Bacillus subtilis* UTBSP1, a promising isolate to degrade aflatoxin B1 (AFB1), caused to reduce the growth of *Aspergillus flavus* R5 and AFB1 content on pistachio nuts. Fluorescence probes revealed that the cell free supernatant fluid from UTBSP1 affects spore viability considerably. Using high-performance liquid chromatographic (HPLC) method, 10 fractions were separated and collected from methanol extract of cell free supernatant fluid. Two fractions showed inhibition zones against *A. flavus*. Mass spectrometric analysis of the both antifungal fractions revealed a high similarity between these anti-*A. flavus* compounds and cyclic-lipopeptides of surfactin, and fengycin families. Coproduction of surfactin and fengycin acted in a synergistic manner and consequently caused a strong antifungal activity against *A. flavus* R5. There was a positive significant correlation between the reduction of *A. flavus* growth and the reduction of AFB1 contamination on pistachio nut by UTBSP1. The results indicated that fengycin and surfactin-producing *B. subtilis* UTBSP1 can potentially reduce *A. flavus* growth and AFB1 content in pistachio nut.

Keywords : aflatoxin B1, antifungal activity, *Bacillus subtilis*, fengycin, surfactin

*Corresponding author.

Phone) +98-2129904068, FAX) +98-2122431783

E-mail) m_farzaneh@sbu.ac.ir

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Aspergillus flavus is a cosmopolitan organism, able to contaminate a wide variety of natural substrates such as grains, nuts and food products (Cotty et al., 1994; González et al., 2005). *A. flavus* produces aflatoxin which is carcinogenic and toxic to both human and livestock populations (Bennett and Klich, 2003; Shephard, 2005). Furthermore, *A. flavus* is an opportunistic pathogen causing invasive and non-invasive aspergillosis in humans (Perfect, 2001). In almost all cases, fungal contamination, colonization and infection are usually initiated by airborne conidia that invade the host tissue and subsequently by germination (Samson et al., 1995).

Of the many research approaches being used to reduce and eliminate *A. flavus* contamination, antifungal agents produced by microorganisms consider as environmentally friendly method to control of *A. flavus* (Dorner, 2004; Shephard, 2005). Several bacteria particularly *Bacillus subtilis* strains have been reported as biological agents to control *A. flavus* (Reddy et al., 2009; Zhang et al., 2007). It is demonstrated that some *B. subtilis* strains produce antifungal compounds including lipopeptides (LPs) such as surfactin (Hosono and Suzuki, 1983; Vater et al., 2002), fengycin (Vanittanakom et al., 1986), and iturin compounds (Chitarra et al., 2003; Peypoux et al., 1978; Zhang et al., 2007). These antifungal peptides inhibit the growth of large number of fungi, such as *Aspergillus*, *Rhizopus*, *Botrytis* and *Penicillium* (Chitarra et al., 2003; Munimbazi and Bullerman, 1998; Sun et al., 2006). In some studies, combination of antifungal agents synergistically acted against different plant pathogens (Koumoutsis et al., 2004; Kuroda et al., 2000; Maget-Dana and Ptak, 1990).

In previous study, *B. subtilis* UTBSP1 considerably remedied aflatoxin B1 in both liquid culture and pistachio

nut (Farzaneh et al., 2012). In addition, this strain showed an antifungal activity against *A. flavus* in dual culture assay (unpublished data). Therefore, the purposes of this work were as follows: (1) Determine the potential of *B. subtilis* UTBSP1 to inhibit *A. flavus* and AFB1 on pistachio. (2) Evaluate the destructive effect of cell-free supernatant fluid (SF) of *B. subtilis* UTBSP1 against the *A. flavus* spore. (3) Separate, purify and identify the anti-*A. flavus* compounds produced by *B. subtilis* UTBSP1.

Materials and Methods

Microorganisms. *A. flavus* R5, with strong ability to produce the high amount of AFB1, and *B. subtilis* UTBSP1, with destructive effect on AFB1 (Farzaneh et al., 2012), were obtained from University of Tehran.

Inhibition of *A. flavus* and Aflatoxin B1 in pistachio.

Mature pistachios with undamaged and non-adhering hulls were collected from Rafsanjan orchard. The hulls were aseptically removed from the pistachios and a total of 100 pistachios with opened shells were selected and injured by using a scribe to make a puncture wound approximately 1 mm wide by 2 mm deep in each kernel. Ten wounded nuts were placed in 100-ml flasks containing 20 ml of different concentration of bacterial suspension in saline solution, sterile 0.85% NaCl (10^5 , 10^6 , and 10^7 cfu/ml) for 10 minutes and then were placed on sterile paper. After 15 minutes, approximately 200 spores of *A. flavus* R5 in 0.05% Tween 80 were inoculated on the wound of kernel and then were placed on 100-ml flasks and incubated for 7 days at 28°C in darkness. The following controls were (1), treated nuts with saline solution and *A. flavus* spore, and (2), treated nuts with saline solution. The growth of *A. flavus* (population of spores) on gram nut was evaluated by adding 50 ml saline solution, shaking vigorously at 250 rpm for 30 minutes and counting the spores with hemocytometer and light microscope. In addition, the AFB1 content was extracted and purified by using immune-affinity column cleanup (Stroka et al., 2001) and then determined by high-performance thin-layer chromatography (HPTLC) as well as high-performance liquid chromatographic (HPLC) method (Farzaneh et al., 2012).

Isolation of antifungal compounds produced by *B. subtilis* UTBSP1.

Forty milliliter of 18 hours pre-incubation of UTBSP1 in nutrient broth (NB; 1% peptone, 0.3% beef extract, 0.5% NaCl, pH 7.0) was transferred to 700 ml fresh NB and incubated at 30°C for 48 hours. Bacteria cells were removed by centrifugation at 4,024g for 15 minutes. The pH of cell free SF was adjusted to 2.0 and

the precipitate was allowed to settle at 4°C. Subsequently, the acid precipitate was collected by centrifugation at 1,369g for 20 minutes at 4°C. The antifungal compounds from the precipitate were extracted twice using methanol and then methanol evaporated by rotary-evaporator at 45°C under a vacuum and then stored at 4°C as a HCl precipitate.

Inhibitory effects on spore. Inhibitory effects of SF (at the three concentrations; 10%, 25%, and 50% in potato dextrose broth medium) and its HCl precipitate (100, 250, and 500 $\mu\text{g}\cdot\text{ml}^{-1}$) on spore germination were determined after 8 hours at 30°C by using light microscope according to Chitarra et al. (2003). In addition, the effect of HCl precipitate on spore viability was carried out by using fluorescent probes (Chitarra et al., 2003). Briefly, spores were incubated at 30°C for 3 hours in the absence and presence of HCl precipitate (500 $\mu\text{g}\cdot\text{ml}^{-1}$) and were subsequently labeled with the fluorescent probes carboxy-fluorescein (cF) diacetate (cFDA) and propidium iodide (PI; Molecular Probes, Eugene, OR, USA) at a final concentration of 10 $\mu\text{mol}\cdot\text{l}^{-1}$ for 15 minutes at 30°C. The incubated spores in water:methanol solution (50:50 v/v) for 3 hours were considered as control. The staining of the spores was determined by the Inverted Epi-fluorescence microscope (ECLIPSE TE2000-S; Nikon Instech Co., Ltd., Tokyo, Japan) equipped with a 100 W mercury lamp, a laser beam (488 nm), a fluorescent filter set and a CFI Plan Apo TIRF 60X oil objective lens.

Purification of antifungal compounds. The methanol fraction of SF acid precipitation was purified by reverse phase-HPLC (RP-HPLC) with a C18 column (COSMOSIL 5 μm C18-AR-300, 4.6 \times 250 mm; Nacalai Tesque, Kyoto, Japan) and injection volume of 100 μl on Agilent 1,100 HPLC and monitored by absorbance at 214 and 280 nm. The solvents were (A) methanol and (B) 0.1% trifluoroacetic acid (TFA) in water. Elution (0.9 ml/min) starts with 45% A for 20 minutes, which was continued for 40 minutes with 100% A, holding for 5 minutes and finally continued for 10 minutes with 45% A. This step was repeated several times, and fractions with the same retention time were collected and dried under N_2 at 35°C and finally their antifungal activities were determined against *A. flavus* (Chitarra et al., 2003).

HPLC/mass spectrometric (MS) analysis of antifungal substances.

A coupled liquid chromatography-tandem MS system consisting of an Agilent HP1200 liquid chromatography and a triple-quadrupole mass spectrometer (Agilent 6410 LC/MS; Agilent Technologies, Palo Alto, CA, USA) were employed to separate and determine

the molecular weight of compounds in unpurified methanol fraction of acid precipitation and purified antifungal compounds collected from RP-HPLC step. The symmetry ZORBAX SB-C18 column (rapid resolution, 3.5 μm , 30 \times 2.1 mm; Agilent Technologies) oven was at 23°C, the flow rate was 0.4 ml/min, and the injection volume was 5 μl . Solvents (A) 10% formic acid in water and (B) methanol were used as mobile phases. The electrospray source was operated at a capillary voltage of 30 V, a spray voltage of 5 kV, a capillary temperature of 350°C and full scan spectra were recorded over the range m/z 50 to 2,000.

Statistical analysis. The experiment was repeated two times in a randomized complete design with four replicates. Data analysis was performed using the SAS ver. 9.1 software (SAS Institute Inc., Cary, NC, USA). The analysis of variance was done by the GLM module.

Results

Inhibit the *A. flavus* growth and Aflatoxin B1 contamination in pistachio. There was significant difference among different populations (10^5 , 10^6 , and 10^7 cfu·ml⁻¹) of *B. subtilis* UTBSP1 to inhibit *A. flavus* growth as well as reduce AFB1 accumulation on pistachio nuts (Fig. 1). In control treatment (without UTBSP1 inoculation), the infected pistachio nut was extensively colonized by *A. flavus* R5 (1.3×10^7 spores/g nut) and its AFB1 content was 12,465.1 ng·g⁻¹ nut. Although the dipping




Fig. 1. Reduction of *Aspergillus flavus* growth (spore population) and aflatoxin B1 (AFB1) content in pistachio nuts by using different concentrations of *Bacillus subtilis* UTBSP1, including 10^5 , 10^6 , and 10^7 cfu/ml saline solution. In the infected control (inoculated nut with saline solution and *A. flavus* R5), 1.3×10^7 spores and 12,465.1 ng AFB1 per g nut were detected. The treatments with same letter were not significantly different ($P \leq 0.05$). Error bars indicate standard error of four replications.

of pistachio nuts in 10^5 cfu·ml⁻¹ of UTBSP1 couldn't significantly decrease the *A. flavus* growth but it could significantly reduce AFB1 accumulation in nut. By using of 10^6 cfu·ml⁻¹ of UTBSP1, spore population and AFB1 content in pistachio nut decreased to 54.2% and 82.1%, respectively. However, by increasing the cfu of UTBSP1 to 10^7 per ml, spore population decrease to 94.1% while no AFB1 content was detected.

Antifungal activity against spore. The germination efficiency of *A. flavus* spores in the absence of SF or its HCl precipitate was 95%. After 8 hours of incubation, the percentage of spore germination efficiency decreased to 25, 5% and 0%, in the presence of 10%, 25%, and 50% of SF, respectively. In addition, none spore was germinated in the presence of HCl precipitate (at 500 $\mu\text{g}\cdot\text{ml}^{-1}$).

The membrane permeabilization and viability of the spore were assessed using fluorescent probes. The control spores could considerably be stained with cFDA (99% spores), but not significantly with PI (4% spores). Incubation of the spores with HCl precipitate for 3 hours caused to stain 98% spores with PI whereas they could not significantly be stained with cFDA (results are shown in the Fig. 2).

Isolation and purification of antifungal compounds.

The methanol fraction of HCl precipitate produced several obvious peaks in HPLC (results are shown in the Fig. 3). All the fractions (10 peaks) were collected manually as separated ones. At the concentration of 200 $\mu\text{g}\cdot\text{ml}^{-1}$, two major fractions (A) and (B) showed anti-




Fig. 2. Fluorescence microscopy pictures of *Aspergillus flavus* spores that were incubated for 3 hours in the HCl precipitate of *Bacillus subtilis* UTBSP1 ($500 \mu\text{g}\cdot\text{ml}^{-1}$) and then stained with PI (A) or cFDA (F). The fluorescence microscopy pictures of PI-stained (C) or cFDA-stained (H) spores in control (water:methanol [50:50 v/v]) are presented. In addition, the light microscopy pictures of HCl precipitate-treated spores (B, E) and untreated (control) spores (D, G) are shown. PI, propidium iodide; cFDA, carboxyfluorescein diacetate.

fungal activities against *A. flavus* after 4 days. The inhibition zones caused by the fractions (A) and (B) were 17 and 5 mm, respectively. Mixture of the both fractions showed stronger antifungal activity with 19 mm inhibition zone (Fig. 4). The inhibition zone of the fraction (A) and the combination of both fractions were stable after 10 days whereas the inhibitory zone arising from fraction (B) disappeared after 3 days. Therefore, LC/MS was applied to identify both anti-*Aspergillus* fractions.

Identification using MS analysis. HPLC analysis of methanol fraction of SF acid precipitate and antifungal activity tests, revealed that *B. subtilis* UTBSP1 produces two peaks with antifungal activities against *A. flavus* R5 (results are shown in the Fig. 3, 4). Full MS analysis of fraction A showed a series of molecular ions with 1,022.6; 1,036.6 m/z for $(M + H)^+$, and 1,044.6; 1,058.6 m/z for $(M + Na)^+$.




Fig. 3. Purification chromatogram of the methanol extraction of the HCl precipitate of *Bacillus subtilis* UTBSP1 by using reverse phase high-performance liquid chromatography (HPLC) with C18 column. Mobile phase: water (0.1% trifluoroacetic acid) and methanol; detector: 214 and 280 nm. All the distinct fractions (10 peaks) were collected manually. Peaks 'A' and 'B' showed inhibition zones against *Aspergillus flavus* R5.

Also, the mass spectrum of fraction B showed a series of molecular ions with 1,463.8; 1,477.8; 1,491.8; 1,505.8 m/z, for $(M + H)^+$ (results are shown in the Fig. 5).

With regards to the comparing of mass data with standard and those reported in previous studies, the antifungal fractions, (A) and (B), represented surfactin homologues and fengycin homologues, respectively (Kowall et al., 1998; Stein, 2005; Vanittanakom et al., 1986; Vater et al., 2002).

HPLC/MS spectra of unpurified methanol fraction of SF acid precipitation exhibited several peaks at different m/z values. Two fractions showed m/z values between 1,000–1,060 as well as 1,460–1,510, symbolized surfactin homologues and fengycin homologues, respectively (Table 1).

Discussion

In this study, *B. subtilis* UTBSP1 could reduce *A. flavus* growth and AFB1 contamination in pistachio nuts and there was a positive correlation between the reduction of *A. flavus* colonization and the reduction of AFB1 accumulation in pistachio nut. Similarly, the antifungal activities of *B. subtilis* strains and other biological agents against *A. flavus* and AFB1-production have been re-



Fig. 4. The inhibition zones against *Aspergillus flavus* growth caused by the fraction A (right) and mixture of fractions A + B (left) after 4 days incubation at 30°C in darkness.





Fig. 5. Full mass spectrometric spectra of surfactin (A), and fengycin (B), produced by *Bacillus subtilis* UTBSP1, and obtained from electrospray ionization triple-quadrupole mass spectrometer. The peaks belong to the HCl precipitate of *B. subtilis* UTBSP1.

Table 1. Assignments of all surfactin and fengycin mass peaks obtained by electrospray ionization triple-quadrupole mass spectrometry of the lipopeptide fractions from *Bacillus subtilis* UTBSP1

| Mass peak (m/z) | Assignment | Structure | Amino acid at position 6 (7) |
|-----------------|--------------------------------------|-----------|------------------------------|
| 1,008.5 | C13 surfactin, (M + H) ⁺ | | Val (7) |
| 1,022.6 | C14 surfactin, (M + H) ⁺ | | Val (7) |
| 1,030.6 | C13 surfactin, (M + Na) ⁺ | | Leu/Ile (7) |
| 1,036.6 | C15 surfactin, (M + H) ⁺ | | Leu/Ile (7) |
| 1,044.6 | C14 surfactin, (M + Na) ⁺ | | Leu/Ile (7) |
| 1,058.6 | C15 surfactin, (M + Na) ⁺ | | Leu/Ile (7) |
| 1,463.8 | C16 fengycin, (M + H) ⁺ | | Ala/Val (6) |
| 1,477.8 | C17 fengycin, (M + H) ⁺ | | Ala/Val (6) |
| 1,491.8 | C18 fengycin, (M + H) ⁺ | | Ala/Val (6) |
| 1,505.8 | C19 fengycin, (M + H) ⁺ | | Ala/Val (6) |

Val, valine; Leu, leucine; Ile, isoleucine; Ala, alanine.

ported in several studies (Dorner et al., 1998; Dorner, 2004; Mohammadipour et al., 2009; Reddy et al., 2009; Zhang et al., 2007). However, contrary to our results, several studies indicated that although *Bacillus* spp., exhibit considerable antifungal activities against some phytopathogenic fungi but their inhibitory activities against *A. flavus* are no significant (Hu et al., 2009; Sun et al., 2006).

In our study, cell free SF from *B. subtilis* UTBSP1 showed destructive effect against spore of *A. flavus*. Therefore, the fluorescent probes including PI and cFDA were applied to perceive the influence of UTBSP1 on fungal spore precisely. The fluorescent probe cFDA is a hydrophobic enzyme activity probe that can pass through the hydrophobic cell membrane and is immediately converted to hydrophilic cF by non-specific esterases and then release the fluorescent light. Therefore, cF slowly accumulates in the cell because of the low permeability of the membrane to cF. In dead cells, there is no cF-accumulation, since the membrane is perforated by many holes (Brul, 1997; Chitarra et al., 2003). In contrast, the fluorescent probe PI is a nucleic acid probe which is supposed to cross only through damaged cell membranes. Therefore, the spore mortality can be correlated with the accumulation of PI. However, fluorescence methods using probes such as cFDA and PI have been used to evaluate conidial viability (Chitarra et al., 2003; Schading et al., 1995). Furthermore, fluorescent labelling in combination with flow-cytometry is a rapid, reliable and sensitive method to assess viability (Brul, 1997; Bunthof et al., 2001). In our study, the incubation of the spores with HCl precipitate for 3 hours, caused to spore staining with PI which indicates that the spores membranes were damaged and permeabilized, similarly to the results

reported in the previous studies (Brul, 1997; Chitarra et al., 2003; Schading et al., 1995). It is obvious that in the presence of SF and its HCl precipitate, the fungal spores lose their abilities to initiate biochemical activities and morphological changes as well.

By HPLC analysis and antifungal activity tests, two fractions arising from HCl precipitation with anti-*A. flavus* activity were isolated in this study. In addition, MS analysis of the two anti-*A. flavus* fractions unveils an exponential similarity between their compounds and cyclic LPs of surfactin and fengycin families. The mass peak at (M + H)⁺ m/z = 1,477.8 (Fig. 5B) is attributed to a fengycin isoform containing hydrogen adduct of a β -hydroxy fatty acid with a chain length of 17 carbon atoms and an alanine at position 6 of the octapeptide ring (Vater et al., 2002). A set of mass peaks with an intervals of 14 Da [(M + H)⁺ ions at m/z 1,463.8, 1,491.8 and 1,505.8] are fengycins with different number of methylene groups (-CH₂-). The mass peak at (M + H)⁺ m/z = 1,022.6 and (M + Na)⁺ m/z = 1,044.6 (Fig. 5A) is attributed to a surfactin isoform containing proton and sodium adducts respectively, with 14 carbon atoms in carbon chain of β -hydroxy fatty acids and a valine at position 7 of the peptide ring (Vater et al., 2002). Molecular ion with 1,008.5 and 1,036.6 m/z for (M + H)⁺, and 1,030.6 and 1,058.6 m/z for (M + Na)⁺ are surfactin isoforms with different length of the carbon chain of β -hydroxy fatty acid (C₁₃ and C₁₅) and a valine or leucine/isoleucine at position 7 of the heptapeptide ring (Morikawa et al., 2000), showed that a set of mass peaks with an intervals of 14 Da are often observed for LPs with different number of methylene groups (-CH₂-) in fatty acyl chains.

This study demonstrates that antifungal activity of *B. subtilis* UTBSP1, producer of both fengycin and sur-

factin, is attributed to both antibiotics. However, some *Bacillus* strains which produce surfactin and/or fengycin couldn't inhibit the growth of *A. flavus* (Hu et al., 2009; Sun et al., 2006), whereas (Mohammadipour et al., 2009) reported that *B. subtilis* BS119m, as a surfactin-producer in a high level, completely inhibit the growth of *A. flavus*. Furthermore, this study indicates that surfactin and fengycin act in a synergistic manner and provide stronger antifungal activity against *A. flavus* R5, similar to the synergistic antifungal manners of surfactin and iturin (Maget-Dana and Ptak, 1990), surfactin and fengycin (Ongena et al., 2007; Sun et al., 2006) and iturin and fengycin (Romero et al., 2007). Iturin-producer *Bacillus* strains also showed antifungal activities against *A. flavus* (Moyné et al., 2001; Zhang et al., 2007).

The surfactin antibiotic has amphiphilic nature and consequently readily associates and tightly anchors into lipid layers. At low concentration it induces only limited perturbation, but at high concentration causes the formation of irreversible pore (Deleu et al., 2005; Heerklotz and Seelig, 2007). Fengycins have strong fungitoxic activity, especially against filamentous fungi (Hu et al., 2007; Sun et al., 2006). Mechanistically, their actions are less well known in comparison with the other LPs, but they readily interact with lipid layers and alter cell membrane structure and permeability in a dose-dependent way (Deleu et al., 2005).

In conclusion, to sum up, *B. subtilis* UTBSP1, co-producer of surfactin and fengycin, is potentially used to decrease *A. flavus* growth and aflatoxin contamination in pistachio nut. In addition, *B. subtilis* UTBSP1 is an AFB1-degrading agent (Farzaneh et al., 2012) and can use for reducing AFB1-content in foods, feeds and agricultural crops. Therefore, the inoculation of agricultural crops by *B. subtilis* UTBSP1 in cultivation, harvest and post-harvest stages might noticeably reduce aflatoxin in nuts, grains and other susceptible crops. However, the significance of *Bacillus* species and, on the other hand, their ability to produce toxic compounds call for careful monitoring of their environmental occurrence (Outtrup and Jorgensen, 2002; Schallmey et al., 2004).

Acknowledgments

We gratefully acknowledge University of Tehran and Institute of Food Safety and Quality, Jiangsu Academy of Agricultural Sciences in China for providing financial supports in this study.

References

Bennett, J. W. and Klich, M. 2003. Mycotoxins. *Clin. Microbiol.*

- Rev.* 16:497-516.
- Brul, S., Nussbaum, J. and Dielbandhoesing, S. K. 1997. Fluorescent probes for wall porosity and membrane integrity in filamentous fungi. *J. Microbiol. Methods* 28:169-178.
- Bunthof, C. J., Bloemen, K., Breeuwer, P., Rombouts, F. M. and Abee, T. 2001. Flow cytometric assessment of viability of lactic acid bacteria. *Appl. Environ. Microbiol.* 67:2326-2335.
- Chitarra, G. S., Breeuwer, P., Nout, M. J. R., Van Aelst, A. C., Rombouts, F. M. and Abee, T. 2003. An antifungal compound produced by *Bacillus subtilis* YM 10-20 inhibits germination of *Penicillium roqueforti* conidiospores. *J. Appl. Microbiol.* 94:159-166.
- Cotty, P. J., Bayman, P., Egel, D. S. and Elis, K. S. 1994. Agriculture, aflatoxins and *Aspergillus*. In: *The genus Aspergillus: from taxonomy and genetics to industrial applications*, eds. by K. A. Powell, A. Renwick and J. F. Peberdy, pp. 1-27. Plenum Press, New York, NY, USA.
- Deleu, M., Paquot, M. and Nylander, T. 2005. Fengycin interaction with lipid monolayers at the air-aqueous interface--implications for the effect of fengycin on biological membranes. *J. Colloid. Interface Sci.* 283:358-365.
- Dorner, J. W., Cole, R. J. and Blankenship, P. D. 1998. Effect of inoculum rate of biological control agents on preharvest aflatoxin contamination of peanuts. *Biol. Control* 12:171-176.
- Dorner, J. W. 2004. Biological control of aflatoxin contamination of crops. *J. Toxicol.: Toxin Rev.* 23:425-450.
- Farzaneh, M., Shi, Z. Q., Ghassempour, A., Sedaghat, N., Ahmadzadeh, M., Mirabolghathy, M. and Javan-Nikkah, M. 2012. Aflatoxin B1 degradation by *Bacillus subtilis* UTBSP1 isolated from pistachio nuts of Iran. *Food Control* 23:100-106.
- González, G., Hinojo, M. J., Mateo, R., Medina, A. and Jiménez, M. 2005. Occurrence of mycotoxin producing fungi in bee pollen. *Int. J. Food Microbiol.* 105:1-9.
- Heerklotz, H. and Seelig, J. 2007. Leakage and lysis of lipid membranes induced by the lipopeptide surfactin. *Eur. Biophys. J.* 36:305-314.
- Hosono, K. and Suzuki, H. 1983. Acylpeptides, the inhibitors of cyclic adenosine 3, 5-monophosphate phosphodiesterase I, Purification, physicochemical properties and structures of fatty acid residues. *J. Antibiot.* 36:667-673.
- Hu, L. B., Shi, Z. Q., Zhang, T. and Yang, Z. M. 2007. Fengycin antibiotics isolated from B-FS01 culture inhibit the growth of *Fusarium moniliforme* Sheldon ATCC 38932. *FEMS Microbiol. Lett.* 272:91-98.
- Hu, L. B., Zhang, T., Yang, Z. M. and Shi, Z. Q. 2009. Inhibition of fengycins on the production of fumonisin B1 from *Fusarium verticillioides*. *Lett. Appl. Microbiol.* 48:84-89.
- Koumoutsis, A., Chen, X. H., Henne, A., Liesegang, H., Hitzeroth, G., Franke, P., Vater, J. and Borriss, R. 2004. Structural and functional characterization of gene clusters directing nonribosomal synthesis of bioactive cyclic lipopeptides in *Bacillus amyloliquefaciens* strain FZB42. *J. Bacteriol.* 186:1084-1096.
- Kowall, M., Vater, J., Kluge, B., Stein, T., Franke, P. and Ziessow, D. 1998. Separation and characterization of

- surfactin isoforms produced by *Bacillus subtilis* OKB 105. *J. Colloid Interface Sci.* 204:1-8.
- Kuroda, J., Fukai, T., Konishi, M., Uno, J., Kurusu, K. and Nomura, T. 2000. LI-F antibiotics, a family of antifungal cyclic depsipeptides produced by *Bacillus polymyxa* L-1129. *Heterocycles* 53:1533-1549.
- Maget-Dana, R. and Ptak, M. 1990. Iturin lipopeptides: interactions of mycosubtilin with lipids in planar membranes and mixed monolayers. *Biochim. Biophys. Acta* 1023:34-40.
- Mohammadipour, M., Mousivand, M., Salehi-Jouzani, G. and Abbasalizadeh, S. 2009. Molecular and biochemical characterization of Iranian surfactin-producing *Bacillus subtilis* isolates and evaluation of their biocontrol potential against *Aspergillus flavus* and *Colletotrichum gloeosporioides*. *Can. J. Microbiol.* 55:395-404.
- Morikawa, M., Hirata, Y. and Imanaka, T. 2000. A study on the structure-function relationship of lipopeptide biosurfactants. *Biochim. Biophys. Acta* 1488:211-218.
- Moyne, A. L., Shelby, R., Cleveland, T. E. and Tuzun, S. 2001. Bacillomycin D: an iturin with antifungal activity against *Aspergillus flavus*. *J. Appl. Microbiol.* 90:622-629.
- Munimbazi, C. and Bullerman, L. B. 1998. Isolation and partial characterization of antifungal metabolites of *Bacillus pumilus*. *J. Appl. Microbiol.* 84:959-968.
- Ongena, M., Jourdan, E., Adam, A., Paquot, M., Brans, A., Joris, B., Arpigny, J. L. and Thonart, P. 2007. Surfactin and fengycin lipopeptides of *Bacillus subtilis* as elicitors of induced systemic resistance in plants. *Environ. Microbiol.* 9: 1084-1090.
- Outtrup, H. and Jorgensen, S. T. 2002. The importance of *Bacillus* species in the production of industrial enzymes. In: *Applications and systematics of Bacillus and relatives*, eds. by R. Berkeley, M. Heyndrickx, N. Logan and P. De Vos, pp. 206-218. Wiley-Blackwell, Malden, MA, USA.
- Perfect, J. R., Cox, G. M., Lee, J. Y., Kauffman, C. A., De Repentigny, L., Chapman, S. W., Morrison, V. A., Pappas, P., Hiemenz, J. W. and Stevens, D. A. 2001. The impact of culture isolation of *Aspergillus* species: a hospital-based survey of aspergillosis. *Clin. Infect. Dis.* 33:1824-1833.
- Peypoux, F., Guinand, M., Michel, G., Delcambe, L., Das, B. C. and Lederer, E. 1978. Structure of iturine A, a peptidolipid antibiotic from *Bacillus subtilis*. *Biochemistry* 17:3992-3996.
- Reddy, K. R. N., Reddy, C. S. and Muralidharan, K. 2009. Potential of botanicals and biocontrol agents on growth and aflatoxin production by *Aspergillus flavus* infecting rice grains. *Food Control* 20:173-178.
- Romero, D., De Vicente, A., Rakotoaly, R. H., Dufour, S. E., Veening, J. W., Arrebola, E., Cazorla, F. M., Kuipers, O. P., Paquot, M. and Perez-Garcia, A. 2007. The iturin and fengycin families of lipopeptides are key factors in antagonism of *Bacillus subtilis* toward *Podosphaera fusca*. *Mol. Plant-Microbe Interact.* 20:430-440.
- Samson, R. A., Hoekstra, E. S., Frisvad, J. C. and Filtenborg, O. 1995. Introduction to food and airborne fungi. 4th ed. Centraalbureau Voor Schimmelcultures, Baarn, the Netherlands. 389 pp.
- Schading, R. L., Carruthers, R. I. and Mullin-Schading, B. A. 1995. Rapid determination of conidial viability for entomopathogenic hyphomycetes using fluorescence microscopy techniques. *Biocontrol Sci. Technol.* 5:201-208.
- Schallmeyer, M., Singh, A. and Ward, O. P. 2004. Developments in the use of *Bacillus* species for industrial production. *Can. J. Microbiol.* 50:1-17.
- Shephard, G. S. 2005. Aflatoxin and food safety: recent African perspectives. CRC Taylor & Francis, Boca Raton, FL, USA. pp. 13-28.
- Stein, T. 2005. *Bacillus subtilis* antibiotics: structures, syntheses and specific functions. *Mol. Microbiol.* 56:845-857.
- Stroka, J., Anklam, E., Joerissen, U., Gilbert, J. 2001. Determination of aflatoxin B1 in baby food (infant formula) by immunoaffinity column cleanup liquid chromatography with postcolumn bromination: collaborative study. *J. AOAC Int.* 84:1116-1123.
- Sun, L., Lu, Z., Bie, X., Lu, F. and Yang, F. 2006. Isolation and characterization of a co-producer of fengycins and surfactins, endophytic *Bacillus amyloliquefaciens* ES-2, from *Scutellaria baicalensis* Georgi. *World J. Microbiol. Biotechnol.* 22:1259-1266.
- Vanittanakom, N., Loeffler, W., Koch, U. and Jung, G. 1986. Fengycin-a novel antifungal lipopeptide antibiotic produced by *Bacillus subtilis* F-29-3. *J. Antibiot.* 39:888-901.
- Vater, J., Kablitz, B., Wilde, C., Franke, P., Mehta, N. and Cameotra, S. S. 2002. Matrix-assisted laser desorption ionization-time of flight mass spectrometry of lipopeptide biosurfactants in whole cells and culture filtrates of *Bacillus subtilis* C-1 isolated from petroleum sludge. *Appl. Environ. Microbiol.* 68:6210-6219.
- Zhang, T., Hu, L. B., Wang, F., Cheng, L. G. and Shi, Z. Q. 2007. Identification of B-FS06 and the antagonistic activity of its cultural productions against *Aspergillus flavus*. *Chin. J. Biol. Control* 23:160-165.