

Antioxidant Activities of Various Extracts from Korean Yam (*Dioscorea batatas* DECNE.)

Yishan Duan¹ · Gyeong-Hwui Kim² · Han-Soo Kim^{1†}

¹Department of Food Science and Technology, Pusan National University, Miryang 50463, Korea

²Department of Biological Science and Technology, Yonsei University, Seoul 03722, Korea

(Received May 1, 2016; Revised May 23, 2016; Accepted June 20, 2016)

Abstract : The objective of this study was to investigate the antioxidant activity of Korean yam (*Dioscorea batatas* DECNE.) by different extraction solvents including 70% methanol, 70% ethanol and chloroform-methanol mixture (CM, 2:1, v/v). Raw yam was analyzed for its color property, total phenol content and antioxidant activity. Yam possessed high L^* value and H^* value, which were 81.64 ± 2.59 and 83.36 ± 0.15 , respectively. Raw yam was found to have great antioxidant activity evaluated through ABTS [2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt] radical scavenging activity, DPPH (2,2-diphenyl-1-picrylhydrazyl) radical scavenging activity, reducing power, and ferric reducing antioxidant power. Total phenol contents of various extracts from raw yam increased in the following order: 70% methanol extract (37.62 ± 0.88 mg CAE/g), 70% ethanol extract (43.38 ± 0.66 mg CAE/g) and CM extract (67.17 ± 0.12 mg CAE/g), respectively. The same trend was also could be found in the ABTS radical scavenging activity, DPPH radical scavenging activity and ferric reducing antioxidant power assays. These results implied that Korean yam might play an important role in antioxidation and serve as the bio-health functional food to take a good part in prevention of human diseases and aging.

Keywords : yam (*Dioscorea batatas* DECNE.), extraction solvents, color values, total phenol, antioxidant activities

1. Introduction

Yam (*Dioscorea batatas* DECNE.) is widely produced throughout East Asia such as Korea, China and Japan, which is the perennial trailing herb and belongs to the *Dioscoreaceae* family [1]. Due to its component characteristics, yam is usually served as the

crucial staple food [2] as well as traditional medicine ingredient in many parts of world. Yam is mainly composed of starch (75.6-84.3%) with small amounts of crude protein, crude fat, crude fiber and crude ash, whose contents are in the range of 6.7-7.9%, 1.0-1.2%, 1.2-1.8%, 2.8-3.8%, respectively. Furthermore, yam tubers also contain vitamin C (13.0-24.7 mg/100 g dry weight), minerals (K, Na, P, Ca, Mg, Cu, Fe, Mn, Zn) [3], organic acids (succinic acid, citric acid, malic

[†]Corresponding author
(E-mail: kimhs777@pusan.ac.kr)

acid, oxalic acid) [4], steroidal saponins [1], musin (glycoprotein) (2.11 g/100 g dry weight) [5] and dioscorins [6]. Numerous studies have conclusively conducted that dioscorins are the storage proteins of yam tubers, which exhibit dehydroascorbate reductase and monodehydroascorbate reductase activities as well as antioxidant activities [6,7]. And glycoprotein not only has been demonstrated to have an antioxidative potential as one of natural antioxidants, but also can serve as a potent anti-inflammatory agent [8,9]. Dioscin is a steroidal saponin of yam which can be hydrolyzed to form diosgenin [10,11]. And diosgenin of yam can be changed into dehydroepiandrosterone on some degree, which showed antioxidative activity against lipid peroxidation and lowered serum cholesterol and phospholipids levels as well as increased high density lipoprotein level in elderly people [12,13]. The overproduction of oxidative radicals such as reactive oxygen species (ROS) or reactive nitrogen species (RNS) results in oxidative stress, which can cause many human diseases, including cancer, aging, atherosclerosis, coronary heart diseases and neurodegenerative diseases [14,15]. The using of antioxidant system such as glutathione (GSH), Se, vitamin C, vitamin E, superoxide dismutase (SOD) and glutathione peroxidase (GSH-Px) can help to eliminate the excessive oxidative radicals [16]. It was reported that people who have diets rich in natural water-soluble antioxidants such as fresh fruits and vegetables can improve the antioxidant activity and reduce the risk of these mentioned diseases [17].

In the present study, the total phenol contents and antioxidant activities of raw yam extracts by 70% methanol, 70% ethanol and chloroform-methanol mixture (CM, 2:1, v/v) were determined and compared *in vitro*.

2. Materials and Methods

2.1. Materials and chemicals

Yam (*Dioscorea batatas* DECNE.) was purchased from Andong (Korea), which was seeded in March or April and harvested in the end of October or December. The fresh yam was washed, sliced (thickness, 0.4–0.6 cm) and dried in the hot air for 18–24 h at 60–70°C. Then the dried chips were smashed (150-mesh) into raw yam meals (RY, commonly called white yam).

2.2. Preparation of yam extracts

Yam meals and extraction solvents including 70% methanol, 70% ethanol and chloroform-methanol mixture (CM, 2:1, v/v) were mixed in a ratio of 1:10 and kept in the dark about 3 h, and then used the Advantec No. 1 filter paper (Tokyo, Japan) to filter. The process of extraction was repeated 3 times. The filtrate was evaporated by rotary vacuum evaporator (EYELA, N-N series, Tokyo, Japan) until the solvents were completely removed. The yam extracts were collected and sealed in brown reagent bottles and frozen at -80°C until required for further analyses.

2.3. Measurement of total phenol contents (TPC)

The phenolic compounds were determined by using Folin-Ciocalteu's phenol reagent [18] and caffeic acid was used as a standard for the calibration curve. In brief, samples (0.5 mL) were mixed with 2.4 mL of distilled water, 2 mL of 2% sodium carbonate (w/v) and 0.1 mL Folin-Ciocalteu's phenol reagent in the test tubes. Then the mixture was incubated in the temperature for 60 min. The absorbance of the reaction mixture was measured at 700 nm using uv/vis-spectrophotometer (Specord 200, Analytikjena, Jena, Germany). Total phenol contents were expressed as mg of caffeic acid equivalents per g of extracts (mg CAE/ g extract).

2.4. ABTS radical scavenging assay

ABTS radical scavenging activity was evaluated according to the method of Sun, Hayakawa, Ogawa, and Izumori [19]. The mixture of 30 mL of 7 mM ABTS and 528 μ L of 140 mM potassium persulfate was stored at room temperature in the dark for 16 h to get the green-blue free radical ABTS^{•+}. Then the solution was diluted with ethanol until the absorbance was 0.7 ± 0.02 at 734 nm. Samples (0.1 mL) were mixed with 2.9 mL of ABTS working solution. After 6 min of reaction, the absorbance was taken at 734 nm. BHA was used as positive control. The percentage of ABTS radical scavenging effect was calculated as follow:

$$\text{ABTS radical scavenging effect (\%)} = [1 - (A_s/A_c)] 100$$

where A_s is the absorbance in the presence of sample or BHA, and A_c is the absorbance of control reaction.

2.5. DPPH radical scavenging assay

DPPH radical scavenging activity was measured according to the method of Blois and Duan [20, 21]. Samples (1 mL) of yam extraction were mixed with 0.2 mM DPPH (1.5 mL) and then vigorously shaken. The mixture solution was stood in the dark for 30 min at 37°C water bath. BHA was used as positive control. Then the absorbance of the reaction mixture was read with spectrophotometer at 517 nm. The percentage inhibition of DPPH radical scavenging activity was calculated based on the control reading using the following calculation:

$$\text{DPPH radical scavenging activity (\%)} = [1 - (A_s/A_c)] 100$$

where A_s is the absorbance in the presence of sample or BHA, and A_c is the absorbance of control reaction.

2.6. Reducing power assay

The reducing power of yam was determined according to the method of Barros et al. [22]. 1.5 mL of phosphate buffer (0.2 M, pH 6.6),

1.5 mL of sample and 1.5 mL of potassium ferricyanide (1%, w/v) were mixed in test tubes, incubated at 50°C water bath for 20 min. An aliquot of 1.5 mL trichloroacetic acid (10%, w/v) was added to the mixture, which was centrifuged (MF-550, Hanil Science Industrial Co., Ltd, Incheon, Korea) at 3,000 rpm for 10 min. And then the supernatant (1 mL) was mixed with 3 mL of distilled water and 0.3 mL of ferric chloride (0.1%, w/v). BHA was used for the positive control. The absorbance was measured at 700 nm after 10 min of reaction at room temperature.

2.7. Ferric reducing antioxidant power (FRAP) assay

The working FRAP reagent was prepared by mixing 10 mL of 0.3 M sodium acetate buffer (pH 3.6), 1 mL of 10 mM TPTZ in 40 mM hydrochloride acid and 1 mL of 20 mM ferric chloride. The freshly prepared FRAP reagent (1.5 mL) was incubated at 37°C water bath for 10 min, at the same time, $A_{\text{reagent blank}}$ was read. Then sample (150 μ L) was added to the FRAP reagent. The reaction mixture was incubated at 37°C water bath for 4 min; the absorbance was read at 593 nm (A_{sample}). The mixture of sodium acetate buffer (1.5 mL) and sample (150 μ L) was used as sample blank. BHT was used as the positive control. And the difference between A_{sample} , $A_{\text{sample blank}}$ and $A_{\text{reagent blank}}$ was used to calculate the FRAP values. Aqueous solution of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ were used for calibration curve and final results were expressed as $\mu\text{M Fe}^{\text{II}}$ [23].

2.8. Color assessment

Color determination was performed by using colorimeter (CR-400, Minolta Co., Osaka, Japan) and described the color values in CIE $L^*a^*b^*$ color system. The instrument was standardized each time with a white ($L^* = 97.79$, $a^* = -0.38$, $b^* = 2.05$) tile. The L^* value is the lightness (dark-light) and correspond to black ($L^* = 0$) and white ($L^* = 100$). A positive a^* value represent redness and

negative one is greenness. A positive b^* value corresponds to yellowness and negative one is blueness. C^* for the metric chroma and H^* for the hue angle were calculated by the transformation of a^* and b^* the following equations:

$$C^* = (a^{*2} + b^{*2})^{1/2}$$

$$H^* = \tan^{-1} (b^*/a^*) \quad (a^* > 0, b^* > 0)$$

$$H^* = 180^\circ + \tan^{-1} (b^*/a^*) \quad (a^* < 0, b^* > 0)$$

2.9. Statistical analysis

The experimental data in triplicate were subjected to analysis of variance (ANOVA) and expressed as mean \pm standard deviation ($n=3$). Analyses of variance were performed by using the one-way analysis of variance procedures. Duncan's multiple-range test was used to analysis the significant difference of means, and $p < 0.05$ was considered to be statistically significant for all statistic procedures. IBM SPSS statistic 21 program was used for data analysis.

3. Results and discussion

3.1. Yields

The various extraction yields of raw yam (RY) by 70% methanol, 70% ethanol and chloroform-methanol (CM, 2:1, v/v) were shown in Table 1. The extraction yield of RY

by 70% methanol exhibited the maximum value (25.32%), CM was found to be the lowest (2.31%). Additionally, the extraction yield of RY by 70% ethanol was found to be 19.24%.

3.2. Total phenol contents (TPC)

Due to the phenolic structure of hydroxyl substituent on the aromatic ring, phenolics can behave as antioxidants [24]. Furthermore, a positively and highly significant relationship between total phenolics and antioxidant activity was documented by Velioglu, Mazza, Gao, and Oomah [25], which implied a compound with higher content of phenol possessed higher antioxidant activity. Total phenol contents were determined by according to the colorimetric Folin-Ciocalteu method with caffeic acid as a stand compound ($y = 2.4819x + 0.0133$, $R^2 = 0.9997$). The total phenol contents of RY by different extraction solvents (70% methanol, 70% ethanol and CM) were showed in Table 1. Total phenol contents of various extracts from RY increased in the order: 70% methanol extract (37.62 ± 0.88 mg CAE/g), 70% ethanol extract (43.38 ± 0.66 mg CAE/g) and CM extract (67.17 ± 0.12 mg CAE/g), respectively. According to Li et al. [26], there were several parameters include condition of sample, temperature of extraction, solvent concentration and solvent type can influence the extraction of phenolics presented in yam.

Table 1. Extraction yields, total phenol contents and IC_{50} values in the antioxidant activity evaluation assays of raw yam (*Dioscorea batatas* DECNE.)

	70% methanol	70% ethanol	CM
Extraction yields (%) δ	25.32	19.24	2.31
Total phenol content (mg GAE/g)	37.62 ± 0.88^a	43.38 ± 0.66^b	67.17 ± 0.12^c
ABTS (IC_{50} , mg/mL)	2.24 ± 0.10^c	2.08 ± 0.20^b	1.15 ± 0.05^a
DPPH (IC_{50} , mg/mL)	1.34 ± 0.02^c	1.22 ± 0.03^b	0.71 ± 0.00^a

The values are means \pm standard deviation ($n=3$). Values with the different letters in the same row are significantly different ($p < 0.05$) by Duncan's multiple range tests.

CM: chloroform-methanol mixture (2:1, v/v).

3.3. ABTS radical scavenging activity

The radical-cation ABTS^{•+} is produced by the oxidation of ABTS. In the absence of antioxidants, ABTS is rather stable, but it reacts actively with an H-atom donor (i.e. phenolics). Therefore, the blue/green chromophore would discolor gradually or be converted into a non-colored form of ABTS up to the antioxidant capacity of antioxidants [27]. Fig. 1 showed the inhibitory effect of all kinds of extract (70% methanol, 70% ethanol and CM extracts) on ABTS radical. ABTS radical scavenging activity was marked and concentration-related (0.4 mg/mL, 0.7 mg/mL and 1.0 mg/mL). The IC₅₀ values of different fractions of RY decreased in the following order: 70% methanol extract (IC₅₀=2.24±0.10 mg/mL), 70% ethanol extract (IC₅₀=2.08±0.20 mg/mL), respectively. The CM extract exhibited the strongest scavenging activity




Fig. 1. ABTS radical scavenging activities of various extracts from raw yam (*Dioscorea batatas* DECNE.).

*The values are means±standard deviation ($n=3$). Bars with the different letters are significantly different ($p<0.05$) by Duncan's multiple range tests.

**CM: chloroform-methanol mixture (2:1, v/v).

scavenging activity. In our anticipation, the extracts with higher total phenol content showed higher ABTS radical scavenging activity. There was a positive correlation between scavenging effects of various extracts against ABTS radical and their amounts of total phenol contents. And our results were consistent with the finding of Velioglu, Mazza, Gao, and Oomah [25].

3.4. DPPH radical scavenging activity

DPPH is a stable free radical and can be scavenged by antioxidants through donating hydrogen. The discoloration from purple to yellow induces the absorbance of reaction mixture decreases at 517 nm [28]. Various fractions obtained by using different extraction solvents indicated different DPPH scavenging capacity (Fig. 2 and Table 1). The DPPH radical scavenging abilities of various extracts appeared to be associated with increasing concentrations. All extracts exhibited excellent DPPH radical scavenging ability even if their effects were lower than that of BHA. Results showed that CM extract (IC₅₀=0.71±0.00 mg/mL) and 70% methanol extract (IC₅₀=1.34±0.02 mg/mL) possessed the highest and lowest activity upon the elimination of DPPH radical, respectively. From the results showed in Table 1 and Fig. 2, remarkable correlation between total phenol contents and DPPH radical scavenging activities can be found clearly, which were in keeping with the results of Hsu et al. [29]. The strong scavenging capacities of the extracts on DPPH radical were most likely on account of the hydrogen donating ability of the phenolic compounds presented in the extracts.




Fig. 2. DPPH radical scavenging activities of various extracts from raw yam (*Dioscorea batatas* DECNE.).

*The values are means \pm standard deviation ($n=3$). Bars with the different letters are significantly different ($p<0.05$) by Duncan's multiple range tests.

**CM: chloroform-methanol mixture (2:1, v/v).

3.5. Reducing power

In the reducing power assay, ferric chloride and potassium ferricyanide combined to form Fe^{3+} /ferricyanide complex. Because of the presence of reductants (i.e. antioxidants), the Fe^{3+} /ferricyanide complex was reduced to its ferrous form. As a consequence, the color of the test solution changed from yellow to different shades of green and blue, up to the antioxidant ability. Hence, the reducing power can be indicated by the number of the Fe^{2+} complex, which was monitored through measuring the formation of Perl's Prussian blue at 700 nm [22]. The reducing power of various extracts at varying concentrations was measured and the results were depicted in Fig. 3. The reducing power of various extracts and reference compound BHA steadily increased with the increasing concentrations up to 1 mg/mL. BHA showed significantly higher reducing power than that of RY. As can be seen from the data, the reducing power values of various extracts did not have significant

difference at a concentration of 1.0 mg/mL. In contrast to our anticipation, the reducing power and total phenol content didn't exit a significant correlation. And our results were confirmed by the report of Bhandari and Kawabata [4], who found total phenol content had no significant correlation with antioxidant activity. We guessed not only total phenol content but also some nonphenolic compounds might affect the antioxidant activity of yam.




Fig. 3. Reducing power of various extracts from raw yam (*Dioscorea batatas* DECNE.).

*The values are means \pm standard deviation ($n=3$). Bars with the different letters are significantly different ($p<0.05$) by Duncan's multiple range tests.

**CM: chloroform-methanol mixture (2:1, v/v).

3.6. Ferric reducing antioxidant power (FRAP)

Antioxidant potential of different fractions from RY was estimated from their ability to reduce the ferric-tripyridyltriazine (Fe^{III} -TPTZ) complex to ferrous-tripyridyltriazine (Fe^{II} -TPTZ) at low pH, forming an intense blue color with an absorption maximum at 593 nm develops [23]. The antioxidant activities through the ferric reducing antioxidant power model system of RY extracts at 0.4 to 1.0 mg/mL concentrations compared with BHT were presented in the Fig. 4. The results




Fig. 4. Ferric reducing antioxidant power of various extracts from raw yam (*Dioscorea batatas* DECNE.).

*The values are means \pm standard deviation ($n=3$). Bars with the different letters are significantly different ($p<0.05$) by Duncan's multiple range tests.

**CM: chloroform-methanol mixture (2:1, v/v).

exhibited concentration-dependent ferric reducing antioxidant activities in all the tested concentrations of various extracts. At a concentration of 1.0 mg/mL, the FRAP values of various extracts increased in the following order: 70% methanol extract ($91.67 \pm 0.48 \mu\text{M Fe}^{\text{II}}$), 70% ethanol extract ($108.33 \pm 0.00 \mu\text{M Fe}^{\text{II}}$) and CM extract ($220.56 \pm 0.73 \mu\text{M Fe}^{\text{II}}$), respectively. The CM extract showed pronounced ferric reducing antioxidant power values compared with other extracts although lower than that of BHT. Significant differences in ferric reducing antioxidant power values between various extracts could be found. Similar to the results obtained from DPPH

radical and ABTS radical scavenging activity, correlation between FRAP values and total phenol contents could also be observed from our research. This significant correlation was in accordance with the reports of the Bertoneclet et al. [30], who found total phenol content strongly correlated with antioxidant activity evaluated by FRAP assay in the honey.

3.7. Color property

The color values was measured as L^* , a^* , b^* values and found 81.64 ± 2.59 , 1.73 ± 0.01 and 14.87 ± 0.18 , respectively (Table 2). C^* value and H^* value were calculated as 14.97 ± 0.17 and 83.36 ± 0.15 , respectively. The H^* value of RY (83.36 ± 0.15) was corresponding to yellow. The L^* and H^* values were significantly higher than the other values. And the yellowness-blueness (b^*) value was higher than that of redness-greenness (a^*) value.

4. Conclusions

Korean yam (*Dioscorea batatas* DECNE.) was analyzed for its color property, total phenol content and antioxidant activity. Yam possessed high L^* value and H^* value, which were 81.64 ± 2.59 and 83.36 ± 0.15 , respectively. Raw yam was found to have great antioxidant activity evaluated through ABTS [2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt] radical scavenging activity, DPPH (2,2-diphenyl-1-picrylhydrazyl) radical scavenging activity, reducing power, and ferric reducing antioxidant power. Total phenol contents of various extracts from raw yam

Table 2. Color properties (L^* , a^* , b^* , C^* and H^*) of raw yam (*Dioscorea batatas* DECNE.)

	L^*	a^*	b^*	C^*	H^*
RY	81.64 ± 2.59^b	1.73 ± 0.01^a	14.87 ± 0.18^a	14.97 ± 0.17^a	83.36 ± 0.15^b

The values are means \pm standard deviation ($n=3$). Values with the different letters in the same row are significantly different ($p<0.05$) by Duncan's multiple range tests.

increased in the following order: 70% methanol extract (37.62 ± 0.88 mg CAE/g), 70% ethanol extract (43.38 ± 0.66 mg CAE/g) and CM extract (67.17 ± 0.12 mg CAE/g), respectively. The same trend was also found in the ABTS radical scavenging activity, DPPH radical scavenging activity and ferric reducing antioxidant power assays. These results implied that Korean yam might play an important role in antioxidation and serve as the bio-health functional food to take a good part in prevention of lifestyle related diseases and aging.

References

1. J. T. Lin, D. J. Yang, Determination of steroidal saponins in different organs of yam (*Dioscorea pseudojaponica* Yamamoto), *Food Chem.*, **108**, 1068 (2008).
2. C. S. Hariprakash, B. Nambisan, Carbohydrate metabolism during dormancy and sprouting in yam (*Dioscorea*) tubers: changes in carbohydrate constituents in yam (*Dioscorea*) tubers during dormancy and sprouting, *J. Agric. Food Chem.*, **44**, 3066 (1996).
3. J. P. D. Wanasundera, G. Ravindran, Nutritional assessment of yam (*Dioscorea alata*) tubers, *Plant Foods Human Nutr.*, **46**, 33 (1994).
4. M. R. Bhandari, J. Kawabata, Organic acid, phenolic content and antioxidant activity of wild yam (*Dioscorea* spp.) tubers of Nepal, *Food Chem.*, **88**, 163 (2004).
5. J. P. D. Wanasundera, G. Ravindran, Effects of cooking on the nutrient and antinutrient contents of yam tubers (*Dioscorea alata* and *Dioscorea esculenta*), *Food Chem.*, **45**, 247 (1992).
6. W. C. Hou, H. J. Chen, Y. H. Lin, Dioscorins, the major tuber storage proteins of yam (*Dioscorea batatas* Decne), with dehydroascorbate reductase and monodehydroascorbate reductase activities, *Plant Sci.*, **149**, 151 (1999).
7. W. C. Hou, M. H. Lee, H. J. Chen, W. L. Liang, C. H. Han, Y. W. Liu, Y. H. Lin, Antioxidant activities of dioscorin, the storage protein of yam (*Dioscorea batatas* Decne) tuber, *Food Chem.*, **49**, 4956 (2001).
8. P. S. Oh, K. T. Lim, Plant glycoprotein modulates the expression of interleukin-1 β via inhibition of MAP kinase in HMC-1 cells, *Biosc. Biotechnol. Biochem.*, **72**, 2133 (2008).
9. P. S. Oh, K. T. Lim, Antioxidant activity of *Dioscorea batatas* Decne glycoprotein, *European Food Res. Technol.*, **226**, 507 (2008).
10. R. F. Dawson, Diosgenin production North America, *Hort. Technol.*, **1**, 22-27 (1991).
11. W. G. Taylor, J. L. Elder, P. R. Chang, K. W. Richards, Microdetermination of diosgenin from fenugreek (*Trigonella foenum-graecum*) seeds, *J. Agric. Food Chem.*, **48**, 5206 (2000).
12. M. Araghiniknam, S. Chung, T. Nelson-White, C. Eskelson, R. R. Waston, Antioxidant activity of dioscorea and dehydroepiandrosterone (DHEA) in older humans, *Life Sci.*, **59**, 147 (1996).
13. Y. M. Liu, K. W. Lin, Antioxidative ability, dioscorin stability, and the quality of yam chips from various yam species as affected by processing method, *Food Chem.*, **74**, 118 (2009).
14. D. Harrison, K. K. Griendling, U. Landmesser, B. Hornig, H. Drexler, Role of oxidative stress in atherosclerosis, *Am. J. Cardiology*, **91**, 7 (2003).
15. L. Migliore, F. Coppedè, Genetic and environmental factors in cancer and neurodegenerative diseases, *Mutation Research*, **512**, 135 (2002).
16. E. C. Cadenas, K. J. A. Davies, Mitochondrial free radical generation,

- oxidative stress, and aging, *Free Radic. Biol. Med.*, **29**, 222 (2000).
17. J. Chen, W. M. Da, D. W. Zhang, Q. Liu, J. H. Kang, Water-soluble antioxidants improve the antioxidant and anticancer activity of low concentrations of curcumin in human leukemia cells, *Pharmazie*, **60**, 57 (2005).
 18. M. Shaghghi, J. L. Manzoori, A. Jouyban, Determinations of total phenols in tea infusions, tomato and apple juice by terbium sensitized fluorescence method as an alternative approach to the Folin-Ciocalteu spectrophotometric method, *Food Chem.*, **108**, 695 (2007).
 19. Y. X. Sun, S. Hayakawa, M. Ogawa, K. Izumori, Antioxidant properties of custard pudding dessert containing rare hexose, D-psicose, *Food Control*, **18**, 220 (2005).
 20. M. S. Blois, Antioxidant determinations by the use of a stable free radical, *Nature*, **181**, 1199 (1958).
 21. Y. S. Duan, M. A. Kim, J. H. Seong, H. S. Chung, H. S. Kim, Antioxidative activities of various solvent extracts from haw (*Crataegus pinnatifida* Bunge), *Kor. J. Food Preserv.*, **21**, 246 (2014).
 22. L. Barros, M. J. Ferreira, B. Queirós, C. F. R. Ferreira, Isabel, P. Baptista, Total phenols, ascorbic acid, β -carotene and lycopene in Portuguese wild edible mushrooms and their antioxidant activities, *Food Chem.*, **103**, 413 (2007).
 23. F. F. Benzie, Iris, J. J. Strain, The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the FRAP assay, *Anal. Biochem.*, **239**, 70 (1996).
 24. R. J. Robbins, Phenolic acids in foods: an overview of analytical methodology, *J. Agric. Food Chem.*, **51**, 2866 (2003).
 25. Y. S. Velioglu, G. Mazza, L. Gao, B. D. Oomah, Antioxidant activity and total phenolics in selected fruits, vegetables, and grain products, *J. Agric. Food Chem.*, **46**, 4113 (1998).
 26. B. B. Li, B. Smith, M. M. Hossain, Extraction of phenolics from citrus peels I. solvent extraction method, *Separ. Purifi. Technol.*, **48**, 182 (2006).
 27. V. Roginsky, E. A. Lissi, Review of methods to determine chain-breaking antioxidant activity in food, *Food Chem.*, **92**, 235 (2005).
 28. L. S. Lai, S. T. Chou, W. W. Chao, Studies on the antioxidative activities of hsian-tsao (*Mesona procumbens* Hemsl) leaf gum, *J. Agric. Food Chem.*, **49**, 963 (2001).
 29. C. L. Hsu, W. Chen, Y. M. Weng, C. Y. Tseng, Chemical composition, physical properties, and antioxidant activities of yam flours as affected by different drying methods, *Food Chem.*, **83**, 85 (2003).
 30. J. Bertoneclj, U. Doberšek, M. Jamnik, T. Golob, Evaluation of the phenolic content, antioxidant activity and colour of Slovenian honey, *Food Chem.*, **105**, 822 (2007).