

**NONLINEAR ALGORITHMS FOR A COMMON  
SOLUTION OF A SYSTEM OF VARIATIONAL  
INEQUALITIES, A SPLIT EQUILIBRIUM PROBLEM  
AND FIXED POINT PROBLEMS**

JAE UG JEONG

ABSTRACT. In this paper, we propose an iterative algorithm for finding a common solution of a system of generalized equilibrium problems, a split equilibrium problem and a hierarchical fixed point problem over the common fixed points set of a finite family of nonexpansive mappings in Hilbert spaces. Furthermore, we prove that the proposed iterative method has strong convergence under some mild conditions imposed on algorithm parameters. The results presented in this paper improve and extend the corresponding results reported by some authors recently.

## 1. Introduction

Let  $H$  be a real Hilbert space with inner product  $\langle \cdot, \cdot \rangle$  and norm  $\| \cdot \|$ . Let  $C$  be a nonempty closed convex subset of  $H$ . Let  $T : C \rightarrow H$  be a nonlinear mapping. We use  $\text{Fix}(T)$  to denote the set of fixed points of  $T$ , *i.e.*,  $\text{Fix}(T) = \{x \in C : Tx = x\}$ . A mapping  $T$  is called nonexpansive if the following inequality holds:

$$\|Tx - Ty\| \leq \|x - y\|$$

---

Received May 23, 2016. Revised August 25, 2016. Accepted September 9, 2016.  
2010 Mathematics Subject Classification: 47H09, 47H10, 49J30, 49J40.

Key words and phrases: Nonexpansive mapping; mixed equilibrium problem; fixed point; variational inequality.

© The Kangwon-Kyungki Mathematical Society, 2016.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

for all  $x, y \in C$ . Moreover, we also denote by  $\mathbb{R}$  the set of all real numbers.

Recently, Ceng and Yao [10] considered the following system of generalized equilibrium problems, which involves finding  $(x^*, y^*) \in C \times C$  such that

$$\begin{cases} F_1(x^*, x) + \langle B_1 y^*, x - x^* \rangle + \frac{1}{\mu_1} \langle x^* - y^*, x - x^* \rangle \geq 0, \forall x \in C, \\ F_2(y^*, y) + \langle B_2 x^*, y - y^* \rangle + \frac{1}{\mu_2} \langle y^* - x^*, y - y^* \rangle \geq 0, \forall y \in C, \end{cases} \quad (1.1)$$

where  $F_1, F_2 : C \times C \rightarrow \mathbb{R}$  are two bifunctions,  $B_1, B_2 : C \rightarrow H$  are two nonlinear mappings and  $\mu_1, \mu_2 > 0$  are two constants. The solution set of (1.1) is denoted by  $\Omega$ .

If  $F_1 = F_2 = 0$ , then problem (1.1) reduces to the following general system of variational inequalities: Find  $(x^*, y^*) \in C \times C$  such that

$$\begin{cases} \langle \mu_1 B_1 y^* + x^* - y^*, x - x^* \rangle \geq 0, & \forall x \in C, \\ \langle \mu_2 B_2 x^* + y^* - x^*, y - y^* \rangle \geq 0, & \forall y \in C, \end{cases} \quad (1.2)$$

which is introduced and considered by Ceng et al. [9].

If  $B_1 = B_2 = B$  in (1.2), then problem (1.2) reduces to finding  $(x^*, y^*) \in C \times C$  such that

$$\begin{cases} \langle \mu_1 B y^* + x^* - y^*, x - x^* \rangle \geq 0, & \forall x \in C, \\ \langle \mu_2 B x^* + y^* - x^*, x - y^* \rangle \geq 0, & \forall x \in C, \end{cases} \quad (1.3)$$

which has been introduced and studied by Verma [23,24].

If  $x^* = y^*$  and  $\mu_1 = \mu_2$ , then problem (1.3) collapses to the classical variational inequality: Find  $x^* \in C$  such that

$$\langle Bx^*, x - x^* \rangle \geq 0, \quad \forall x \in C.$$

The theory of variational inequality emerged as rapidly growing area of research because of its applications in nonlinear analysis, optimization, economics and game theory ( see [1,2,4,5] and the references cited therein).

The equilibrium problem is to find  $x \in C$  such that

$$F(x, y) \geq 0, \quad \forall y \in C. \quad (1.4)$$

The solution set of (1.4) is denoted by  $EP(F)$ . Numerous problems in physics, optimization and economics reduce to finding a solution of (1.4) ( see [12,19]).

In 1994, Censor and Elfving [11] introduced and studied the following split feasibility problem:

Let  $C$  and  $K$  be nonempty closed convex subsets of the infinite-dimensional real Hilbert spaces  $H_1$  and  $H_2$ , respectively, and let  $A : H_1 \rightarrow H_2$  be a bounded linear operator. Then the split feasibility problem is to find  $x^* \in C$  such that  $Ax^* \in K$ .

In this paper, we study the following split equilibrium problem:

Let  $\Theta : K \times K \rightarrow \mathbb{R}$  be a nonlinear bifunction and  $A : H_1 \rightarrow H_2$  be a bounded linear operator. Then the split equilibrium problem (SEP) is to find  $x^* \in C$  such that

$$y^* = Ax^* \in K \quad \text{solves} \quad \Theta(y^*, y) \geq 0, \quad \forall y \in K. \quad (1.5)$$

The solution set of SEP (1.5) is denoted by  $\Lambda = \{p \in C : Ap \in EP(\Theta)\}$ .

Let  $S : C \rightarrow H$  be a nonexpansive mapping. The following problem is called a hierarchical fixed point problem: find  $x \in F(T)$  such that

$$\langle x - Sx, y - x \rangle \geq 0, \quad \forall y \in F(T). \quad (1.6)$$

It is well known that the iterative methods for finding hierarchical fixed points of nonexpansive mappings can be used to solve a convex minimization problem ( see [25,26] and the references therein).

In 2001, Yamada [26] considered the following hybrid steepest-descent iterative method:

$$x_{n+1} = Tx_n - \mu\lambda_n F(Tx_n),$$

where  $F$  is  $\kappa$ -Lipschitzian continuous and  $\eta$ -strongly monotone operator with  $\kappa > 0$ ,  $\eta > 0$  and  $0 < \mu < \frac{2\eta}{\kappa^2}$ . Under some appropriate conditions, the sequence  $\{x_n\}$  converges strongly to the unique solution of the variational inequality

$$\langle F(x^*), x - x^* \rangle \geq 0, \quad \forall x \in \text{Fix}(T). \quad (1.7)$$

Zhou and Wang [28] proposed a simple explicit iterative algorithm for finding a solution of variational inequality over the set of common fixed points of a finite family nonexpansive mappings. They introduced an explicit scheme as follows:

**THEOREM 1.1.** *Let  $H$  be a real Hilbert space and  $F : H \rightarrow H$  be an  $\kappa$ -Lipschitzian continuous and  $\eta$ -strongly monotone mapping with  $\kappa > 0$  and  $\eta > 0$ . Let  $\{T_i\}_{i=1}^N$  be  $N$  nonexpansive self-mappings of  $H$  such that*

$C = \bigcap_{i=1}^N \text{Fix}(T_i) \neq \phi$ . For any point  $x_0 \in H$ , define a sequence  $\{x_n\}$  as follows:

$$x_{n+1} = (1 - \alpha_n \mu F) T_N^n T_{N-1}^n \cdots T_1^n x_n, \quad n \geq 0,$$

where  $\mu \in (0, \frac{2\eta}{\kappa^2})$  and  $T_i^n = (1 - \sigma_n^i)I + \sigma_n^i T_i$  for  $i = 1, 2, \dots, N$ . When the parameters satisfy appropriate conditions, the sequence  $\{x_n\}$  converges strongly to the unique solution of the variational inequality (1.7).

Recently, Zhang and Yang [27] proposed an explicit iterative algorithm based on the viscosity method for finding a solution for a class of variational inequalities over the common fixed points set of a finite family of nonexpansive mappings as follows:

**THEOREM 1.2.** *Let  $H$  be a real Hilbert space and  $F : H \rightarrow H$  be an  $\kappa$ -Lipschitzian continuous and  $\eta$ -strongly monotone mapping with  $\kappa > 0$  and  $\eta > 0$ . Let  $\{T_i\}_{i=1}^N$  be  $N$  nonexpansive mappings of  $H$  such that  $C = \bigcap_{i=1}^N \text{Fix}(T_i) \neq \phi$  and  $V$  be an  $\rho$ -Lipschitzian continuous on  $H$  with  $\rho > 0$ . For any point  $x_0 \in H$ , define a sequence  $\{x_n\}$  as follows:*

$$x_{n+1} = \alpha_n \gamma V(x_n) + (I - \alpha_n \mu F) T_N^n T_{N-1}^n \cdots T_1^n x_n, \quad n \geq 0,$$

where  $0 < \gamma \rho < \tau$  with  $\tau = \mu(2\eta - \mu\kappa^2)$ ,  $0 < \mu < \frac{2\eta}{\kappa^2}$ ,  $T_i^n = (1 - \sigma_n^i)I + \sigma_n^i T_i$  for  $i = 1, 2, \dots, N$  and  $\sigma_n^i \in (\zeta_1, \zeta_2)$  for some  $\zeta_1, \zeta_2 \in (0, 1)$ . When the parameters satisfy appropriate conditions, the sequence  $\{x_n\}$  converges strongly to the unique solution  $x^* \in C$  of the variational inequality:

$$\langle (\mu F - \gamma V)x^*, x - x^* \rangle \geq 0, \quad \forall x \in \bigcap_{i=1}^N \text{Fix}(T_i). \quad (1.8)$$

In this paper, motivated by the above works, we introduce a new iterative algorithm for finding the approximate element of the common set of solutions of (1.1), (1.5) and (1.8) in real Hilbert spaces. Strong convergence theorems for common elements are established. Our result improves and extends many known results for solving a system of variational inequality problems, split equilibrium problems and hierarchical fixed point theorems (see [6,8,15,18,22,27,28] and the references cited therein).

## 2. Preliminaries

Let  $C$  and  $K$  be nonempty closed convex subsets of real Hilbert spaces  $H_1$  and  $H_2$ , respectively. We denote the strong convergence and the weak convergence of  $\{x_n\}$  to  $x \in H_1$  by  $x_n \rightarrow x$  and  $x_n \rightharpoonup x$ , respectively. For every point  $x \in H_1$ , there exists a unique nearest point in  $C$ , denoted by  $P_C(x)$ , such that

$$\|x - P_C(x)\| \leq \|x - y\|, \quad \forall y \in C.$$

Then  $P_C$  is called the metric projection of  $H_1$  onto  $C$ . It is well known that  $P_C$  is nonexpansive and satisfies the following property:

$$\langle x - P_C(x), P_C(x) - y \rangle \geq 0, \quad \forall x \in H_1, y \in C. \quad (2.1)$$

DEFINITION 2.1. A mapping  $T : H_1 \rightarrow H_1$  is said to be

(1)  $\eta$ -strongly monotone if there exists  $\eta > 0$  such that

$$\langle Tx - Ty, x - y \rangle \geq \eta \|x - y\|^2, \quad \forall x, y \in H_1;$$

(2)  $\delta$ -inverse strongly monotone if there exists  $\delta > 0$  such that

$$\langle Tx - Ty, x - y \rangle \geq \delta \|Tx - Ty\|^2, \quad \forall x, y \in H_1;$$

(3)  $\kappa$ -Lipschitzian continuous if there exists  $\kappa > 0$  such that

$$\|Tx - Ty\| \leq \kappa \|x - y\|, \quad \forall x, y \in H_1;$$

(4)  $\sigma$ -averaged if there exists  $\sigma \in (0, 1)$  such that  $T = (1 - \sigma)I + \sigma S$ , where  $I : H_1 \rightarrow H_1$  is the identity mapping and  $S : H_1 \rightarrow H_1$  is nonexpansive.

In order to prove our main results in the next section, we need the following lemmas.

LEMMA 2.1. For all  $x, y \in H_1$ , there holds the inequality

$$\|x + y\|^2 \leq \|x\|^2 + 2\langle y, x + y \rangle.$$

ASSUMPTION 2.1 [3]. Let  $F : C \times C \rightarrow \mathbb{R}$  be a bifunction satisfying the following assumptions:

(A1)  $F(x, x) = 0, \forall x \in C$ ;

(A2)  $F$  is monotone, i.e.,  $F(x, y) + F(y, x) \leq 0, \forall x, y \in C$ ;

(A3) for each  $x, y, z \in C, \lim_{t \rightarrow 0} F(tz + (1 - t)x, y) \leq F(x, y)$ ;

(A4) for each  $x \in C, y \mapsto F(x, y)$  is convex and lower semicontinuous.

LEMMA 2.2 [14]. Let  $C$  be a nonempty closed convex subset of  $H_1$ . Let  $F_1 : C \times C \rightarrow \mathbb{R}$  satisfy (A1)-(A4). Assume that for  $r > 0$ , define a mapping  $T_r^{F_1} : H_1 \rightarrow C$  as follows:

$$T_r^{F_1}(x) = \{z \in C : F_1(z, y) + \frac{1}{r} \langle y - z, z - x \rangle \geq 0, \forall y \in C\}, \forall x \in H_1.$$

Then the following hold:

(i)  $T_r^{F_1}$  is nonempty and single-valued;

(ii)  $T_r^{F_1}$  is firmly nonexpansive, i.e.,

$$\|T_r^{F_1}(x) - T_r^{F_1}(y)\|^2 \leq \langle T_r^{F_1}(x) - T_r^{F_1}(y), x - y \rangle, \forall x, y \in H_1;$$

(iii)  $\text{Fix}(T_r^{F_1}) = EP(F_1)$ ;

(iv)  $EP(F_1)$  is closed and convex.

LEMMA 2.3. [13] Assume that  $F_1 : C \times C \rightarrow \mathbb{R}$  satisfies Assumption 2.1 and let  $T_r^{F_1}$  be defined as in Lemma 2.2. Let  $x, y \in H_1$  and  $r_1, r_2 > 0$ . Then

$$\|T_{r_2}^{F_1}(y) - T_{r_1}^{F_1}(x)\| \leq \|y - x\| + \left| \frac{r_2 - r_1}{r_2} \right| \|T_{r_2}^{F_1}(y) - y\|.$$

LEMMA 2.4. [7] Let  $F_1, F_2 : C \times C \rightarrow \mathbb{R}$  be two bifunctions satisfying (A1)-(A4). For any  $(x^*, y^*) \in C \times C$ ,  $(x^*, y^*)$  is a solution of (1.1) if and only if  $x^*$  is a fixed point of the mapping  $Q : C \rightarrow C$  defined by

$$Q(x) = T_{\mu_1}^{F_1}[T_{\mu_2}^{F_2}(x - \mu_2 B_2 x) - \mu_1 B_1 T_{\mu_2}^{F_2}(x - \mu_2 B_2 x)], \forall x \in C,$$

where  $y^* = T_{\mu_2}^{F_2}(x^* - \mu_2 B_2 x^*)$ ,  $\mu_i \in (0, 2\theta_i)$  and  $B_i : C \rightarrow C$  is a  $\theta_i$ -inverse strongly monotone mapping for each  $i = 1, 2$ .

LEMMA 2.5. [21] Suppose that  $\lambda \in (0, 1)$  and  $\mu > 0$ . Let  $F : C \rightarrow C$  be a  $\kappa$ -Lipschitzian continuous and  $\eta$ -strongly monotone mapping with  $\kappa > 0$  and  $\eta > 0$ . In association with a nonexpansive mapping  $T : C \rightarrow C$ , define the mapping  $T^\lambda : C \rightarrow H_1$  by

$$T^\lambda x = Tx - \lambda \mu F(Tx), \quad \forall x \in C.$$

Then  $T^\lambda$  is a contraction provided  $\mu < \frac{2\eta}{\kappa^2}$ , i.e.,

$$\|T^\lambda x - T^\lambda y\| \leq (1 - \lambda\tau)\|x - y\|, \quad \forall x, y \in C,$$

where  $\tau = 1 - \sqrt{1 - \mu(2\eta - \mu\kappa^2)}$ .

LEMMA 2.6. [17]

(i) The composite of finitely many averaged mappings is averaged. That is, if each of the mappings  $\{T_i\}_{i=1}^N$  is averaged, then so is the

composite  $T_1 \cdots T_N$ . In particular, if  $T_1$  is  $\alpha_1$ -averaged and  $T_2$  is  $\alpha_2$ -averaged, where  $\alpha_1, \alpha_2 \in (0, 1)$ , then both  $T_1T_2$  and  $T_2T_1$  are  $\alpha$ -averaged, where  $\alpha = \alpha_1 + \alpha_2 - \alpha_1\alpha_2$ .

(ii) If the mappings  $\{T_i\}_{i=1}^N$  are averaged and have a common fixed point, then  $\bigcap_{i=1}^N \text{Fix}(T_i) = \text{Fix}(T_1 \cdots T_N)$ . In particular, if  $N = 2$ , then we have  $\text{Fix}(T_1) \cap \text{Fix}(T_2) = \text{Fix}(T_1T_2) = \text{Fix}(T_2T_1)$ .

LEMMA 2.7. [16] Let  $H_1$  be a Hilbert space,  $C$  be a closed convex subset of  $H_1$  and  $T : C \rightarrow C$  be a nonexpansive mapping with  $\text{Fix}(T) \neq \emptyset$ . If  $\{x_n\}$  is a sequence in  $C$  weakly converging to  $x \in C$  and  $\{(I - T)x_n\}$  converges strongly to  $y \in C$ , then  $(I - T)x = y$ . In particular, if  $y = 0$ , then  $x \in \text{Fix}(T)$ .

LEMMA 2.8. [20] Let  $\{x_n\}$  and  $\{y_n\}$  be bounded sequences in a Banach space  $E$  and let  $\{\beta_n\}$  be a sequence in  $[0, 1]$  with  $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$ . Suppose  $x_{n+1} = \beta_n x_n + (1 - \beta_n)y_n$  for all integers  $n \geq 0$  and  $\limsup_{n \rightarrow \infty} (\|y_{n+1} - y_n\| - \|x_{n+1} - x_n\|) \leq 0$ . Then  $\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0$ .

LEMMA 2.9. [25]. Assume  $\{a_n\}$  is a sequence of nonnegative real numbers such that

$$a_{n+1} \leq (1 - \gamma_n)a_n + \delta_n,$$

where  $\{\gamma_n\}$  is a sequence in  $(0, 1)$  and  $\delta_n$  is a sequence such that

- (1)  $\sum_{n=1}^{\infty} \gamma_n = \infty$ ;
  - (2)  $\limsup_{n \rightarrow \infty} \frac{\delta_n}{\gamma_n} \leq 0$  or  $\sum_{n=1}^{\infty} |\delta_n| < \infty$ .
- Then  $\lim_{n \rightarrow \infty} a_n = 0$ .

### 3. Main results

THEOREM 3.1. Let  $H_1$  and  $H_2$  be two real spaces, and  $C \subseteq H_1$  and  $K \subseteq H_2$  be two nonempty closed convex subsets. Let  $A : H_1 \rightarrow H_2$  be a bounded linear operator with its adjoint  $A^*$ . Assume that  $F_1, F_2 : C \times C \rightarrow \mathbb{R}$  and  $\Theta : K \times K \rightarrow \mathbb{R}$  are the bifunctions satisfying Assumption 2.1 and  $\Theta$  is upper semicontinuous in the first argument. Let  $B_i : C \rightarrow H_1$  be a  $\gamma_i$ -inverse strongly monotone mapping for each  $i = 1, 2$  and  $T_i : C \rightarrow C$  be a nonexpansive mapping for each  $i = 1, 2, \dots, N$  such that  $\Gamma = \bigcap_{i=1}^N \text{Fix}(T_i) \cap \Omega \cap \Lambda \neq \emptyset$ . Let  $F : C \rightarrow C$  be a  $\kappa$ -Lipschitzian continuous and  $\eta$ -strongly monotone mapping with  $\kappa > 0$  and  $\eta > 0$ , and  $V : C \rightarrow C$  be a  $\sigma$ -Lipschitzian continuous mapping with  $\sigma > 0$ . Let

$0 < \mu < \frac{2\eta}{\kappa^2}$  and  $0 < \rho\sigma < \tau$ , where  $\tau = 1 - \sqrt{1 - \mu(2\eta - \mu\kappa^2)}$ . Suppose  $\{\alpha_n\}$  and  $\{\beta_n\}$  are two sequences in  $(0, 1)$ . Given  $x_1 \in C$ , let  $\{x_n\}$  be defined by

$$\begin{cases} z_n = P_C(x_n + \delta A^*(T_{r_n}^\Theta - I)Ax_n), \\ y_n = T_{\mu_1}^{F_1}[T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - \mu_1 B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n)], \\ x_{n+1} = \alpha_n \rho V(x_n) + \beta_n x_n + ((1 - \beta_n)I - \alpha_n \mu F)T_N^n T_{N-1}^n \cdots T_1^n y_n, \forall n \geq 1, \end{cases} \tag{3.1}$$

where  $T_i^n = (1 - \sigma_n^i)I + \sigma_n^i T_i$  for  $i = 1, 2, \dots, N$ ,  $\{r_n\} \subset (0, 2\zeta)$ ,  $\zeta > 0$ ,  $\mu_i \in (0, 2\gamma_i)$  for each  $i = 1, 2$ ,  $\delta \in (0, \frac{1}{L})$ ,  $L$  is the spectral radius of the operator  $A^*A$ ,  $A^*$  is the adjoint of  $A$  and  $\sigma_n^i \in (\zeta_1, \zeta_2)$  for some  $\zeta_1, \zeta_2 \in (0, 1)$ . If the following conditions are satisfied:

- (i)  $\lim_{n \rightarrow \infty} \alpha_n = 0$  and  $\sum_{n=1}^\infty \alpha_n = \infty$ ;
- (ii)  $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$ ;
- (iii)  $0 < \liminf_{n \rightarrow \infty} r_n \leq \limsup_{n \rightarrow \infty} r_n < 2\zeta$  and  $\lim_{n \rightarrow \infty} |r_{n+1} - r_n| = 0$ ;
- (iv)  $\lim_{n \rightarrow \infty} |\sigma_{n+1}^i - \sigma_n^i| = 0$  for  $i = 1, 2, \dots, N$ .

Then the sequence  $\{x_n\}$  converges strongly to  $x^* \in \Gamma = \cap_{i=1}^N \text{Fix}(T_i) \cap \Omega \cap \Lambda$ , where  $x^* = P_\Gamma(I - \mu F + \rho V)x^*$  is the unique solution of the variational inequality:

$$\langle (\mu F - \rho V)x^*, x - x^* \rangle \geq 0, \quad \forall x \in \Gamma. \tag{3.2}$$

*Proof.* Since our methods easily deduce the general case, we prove Theorem 3.1 for  $N = 2$ . Now we divide the proof into several steps.

**Step 1.**  $\{x_n\}$  is bounded.

Indeed, take  $x^* \in \Gamma = \cap_{i=1}^N \text{Fix}(T_i) \cap \Omega \cap \Lambda$  arbitrarily. Since  $Ax^* = T_{r_n}^\Theta(Ax^*)$ , we have

$$\begin{aligned} \|Ax_n - Ax^*\|^2 &= \|(T_{r_n}^\Theta - I)Ax_n - (T_{r_n}^\Theta Ax_n - Ax^*)\|^2 \\ &= \|(T_{r_n}^\Theta - I)Ax_n\|^2 - 2\langle (T_{r_n}^\Theta - I)Ax_n, T_{r_n}^\Theta Ax_n - Ax^* \rangle \\ &\quad + \|T_{r_n}^\Theta Ax_n - Ax^*\|^2 \\ &\leq \|(T_{r_n}^\Theta - I)Ax_n\|^2 - 2\langle (T_{r_n}^\Theta - I)Ax_n, T_{r_n}^\Theta Ax_n - Ax^* \rangle \\ &\quad + \|Ax_n - Ax^*\|^2. \end{aligned}$$



It follows that

$$\langle (T_{r_n}^\Theta - I)Ax_n, T_{r_n}^\Theta Ax_n - Ax^* \rangle \leq \frac{1}{2} \|(T_{r_n}^\Theta - I)Ax_n\|^2. \quad (3.3)$$

Note that  $x^* = P_C(x^*)$ . From (3.1) and (3.3), it follows that

$$\begin{aligned} & \|z_n - x^*\|^2 \\ &= \|P_C(x_n + \delta A^*(T_{r_n}^\Theta - I)Ax_n) - P_C(x^*)\|^2 \\ &\leq \|x_n + \delta A^*(T_{r_n}^\Theta - I)Ax_n - x^*\|^2 \\ &= \|x_n - x^*\|^2 + 2\delta \langle x_n - x^*, A^*(T_{r_n}^\Theta - I)Ax_n \rangle \\ &\quad + \delta^2 \|A^*(T_{r_n}^\Theta - I)Ax_n\|^2 \\ &= \|x_n - x^*\|^2 + 2\delta \langle A(x_n - x^*), (T_{r_n}^\Theta - I)Ax_n \rangle \\ &\quad + \delta^2 \langle (T_{r_n}^\Theta - I)Ax_n, AA^*(T_{r_n}^\Theta - I)Ax_n \rangle \\ &= \|x_n - x^*\|^2 + 2\delta (\langle T_{r_n}^\Theta(Ax_n) - Ax^*, (T_{r_n}^\Theta - I)Ax_n \rangle - \|(T_{r_n}^\Theta - I)Ax_n\|^2) \\ &\quad + \delta^2 \langle (T_{r_n}^\Theta - I)Ax_n, AA^*(T_{r_n}^\Theta - I)Ax_n \rangle \\ &\leq \|x_n - x^*\|^2 + 2\delta (\frac{1}{2} \|(T_{r_n}^\Theta - I)Ax_n\|^2 - \|(T_{r_n}^\Theta - I)Ax_n\|^2) \\ &\quad + \delta^2 \|AA^*\| \|(T_{r_n}^\Theta - I)Ax_n\|^2 \\ &= \|x_n - x^*\| - \delta(1 - L\delta) \|(T_{r_n}^\Theta - I)Ax_n\|^2 \\ &\leq \|x_n - x^*\|^2. \end{aligned} \quad (3.4)$$

Since  $B_i$  is a  $\gamma_i$ -inverse strongly monotone mapping for each  $i = 1, 2$ ,  $x^* = T_{\mu_1}^{F_1}(y^* - \mu_1 B_1 y^*)$  and  $y^* = T_{\mu_2}^{F_2}(x^* - \mu_2 B_2 x^*)$ , we obtain from (3.4) that

$$\begin{aligned} & \|y_n - y^*\|^2 \\ &= \|T_{\mu_1}^{F_1}[T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - \mu_1 B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n)] \\ &\quad - T_{\mu_1}^{F_1}[T_{\mu_2}^{F_2}(x^* - \mu_2 B_2 x^*) - \mu_1 B_1 T_{\mu_2}^{F_2}(x^* - \mu_2 B_2 x^*)]\|^2 \\ &\leq \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - T_{\mu_2}^{F_2}(x^* - \mu_2 B_2 x^*) \\ &\quad - \mu_1 (B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 T_{\mu_2}^{F_2}(x^* - \mu_2 B_2 x^*))\|^2 \end{aligned}$$

$$\begin{aligned}
&\leq \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - T_{\mu_2}^{F_2}(x^* - \mu_2 B_2 x^*)\|^2 \\
&\quad - 2\mu_1 \langle B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 T_{\mu_2}^{F_2}(x^* - \mu_2 B_2 x^*), \\
&\quad \quad T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - T_{\mu_2}^{F_2}(x^* - \mu_2 B_2 x^*) \rangle \\
&\quad + \mu_1^2 \|B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 T_{\mu_2}^{F_2}(x^* - \mu_2 B_2 x^*)\|^2 \\
&\leq \|T_{\mu_1}^{F_2}(z_n - \mu_2 B_2 z_n) - T_{\mu_2}^{F_2}(x^* - \mu_2 B_2 x^*)\|^2 \\
&\quad - \mu_1(2\gamma_1 - \mu_1) \|B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 T_{\mu_2}^{F_2}(x^* - \mu_2 B_2 x^*)\|^2 \\
&\leq \|(z_n - x^*) - \mu_2(B_2 z_n - B_2 x^*)\|^2 \\
&\quad - \mu_1(2\gamma_1 - \mu_1) \|B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 T_{\mu_2}^{F_2}(x^* - \mu_2 B_2 x^*)\|^2 \\
&= \|z_n - x^*\|^2 - 2\mu_2 \langle B_2 z_n - B_2 x^*, z_n - x^* \rangle + \mu_2^2 \|B_2 z_n - B_2 x^*\|^2 \\
&\quad - \mu_1(2\gamma_1 - \mu_1) \|B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 T_{\mu_2}^{F_2}(x^* - \mu_2 B_2 x^*)\|^2 \\
&\leq \|z_n - x^*\|^2 - \mu_2(2\gamma_2 - \mu_2) \|B_2 z_n - B_2 x^*\|^2 \\
&\quad - \mu_1(2\gamma_1 - \mu_1) \|B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 T_{\mu_2}^{F_2}(x^* - \mu_2 B_2 x^*)\|^2 \\
&\leq \|z_n - x^*\|^2 \\
&\leq \|x_n - x^*\|^2. \tag{3.5}
\end{aligned}$$

From (3.1), (3.5) and Lemma 2.5, we have

$$\begin{aligned}
&\|x_{n+1} - x^*\| \\
&= \|\alpha_n \rho V(x_n) + \beta_n x_n + ((1 - \beta_n)I - \alpha_n \mu F) T_2^n T_1^n y_n - x^*\| \\
&\leq \alpha_n \|\rho V(x_n) - \mu F(x^*)\| + \beta_n \|x_n - x^*\| \\
&\quad + \|((1 - \beta_n)I - \alpha_n \mu F) T_2^n T_1^n y_n - ((1 - \beta_n)I - \alpha_n \mu F) T_2^n T_1^n x^*\| \\
&= \alpha_n \|\rho V(x_n) - \mu F(x^*)\| + \beta_n \|x_n - x^*\| \\
&\quad + (1 - \beta_n) \|(I - \frac{\alpha_n}{1 - \beta_n} \mu F) T_2^n T_1^n y_n - (I - \frac{\alpha_n}{1 - \beta_n} \mu F) T_2^n T_1^n x^*\| \\
&\leq (1 - \beta_n) (1 - \frac{\alpha_n \tau}{1 - \beta_n}) \|y_n - x^*\| + \beta_n \|x_n - x^*\| + \alpha_n \|\rho V(x_n) - \mu F(x^*)\| \\
&\leq (1 - \alpha_n \tau) \|x_n - x^*\| + \alpha_n \rho \|V(x_n) - V(x^*)\| + \alpha_n \|\rho V(x^*) - \mu F(x^*)\| \\
&\leq (1 - \alpha_n(\tau - \rho\sigma)) \|x_n - x^*\| + \alpha_n(\tau - \rho\sigma) \frac{\|\rho V(x^*) - \mu F(x^*)\|}{\tau - \rho\sigma} \\
&\leq \max \left\{ \|x_n - x^*\|, \frac{\|\rho V(x^*) - \mu F(x^*)\|}{\tau - \rho\sigma} \right\}. \tag{3.6}
\end{aligned}$$

It follows from (3.6) and induction that

$$\|x_n - x^*\| \leq \max \left\{ \|x_0 - x^*\|, \frac{\|\rho V(x^*) - \mu F(x^*)\|}{\tau - \rho\sigma} \right\}, \quad n = 1, 2, \dots .$$

Therefore  $\{x_n\}$  is bounded. We also obtain that  $\{y_n\}$  and  $\{z_n\}$  are all bounded.

**Step 2.**  $\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0$ .

Indeed, set  $x_{n+1} = \beta_n x_n + (1 - \beta_n)w_n$  for all  $n \geq 1$ . Then we obtain

$$\begin{aligned} w_{n+1} - w_n &= \frac{x_{n+2} - \beta_{n+1}x_{n+1}}{1 - \beta_{n+1}} - \frac{x_{n+1} - \beta_n x_n}{1 - \beta_n} \\ &= \frac{\alpha_{n+1}\rho V(x_{n+1}) + ((1 - \beta_{n+1})I - \alpha_{n+1}\mu F)T_2^{n+1}T_1^{n+1}y_{n+1}}{1 - \beta_{n+1}} \\ &\quad - \frac{\alpha_n \rho V(x_n) + ((1 - \beta_n)I - \alpha_n \mu F)T_2^n T_1^n y_n}{1 - \beta_n} \\ &= \frac{\alpha_{n+1}}{1 - \beta_{n+1}} (\rho V(x_{n+1}) - \mu F(T_2^{n+1}T_1^{n+1}y_{n+1})) \\ &\quad + \frac{\alpha_n}{1 - \beta_n} (\mu F(T_2^n T_1^n y_n) - \rho V(x_n)) \\ &\quad + T_2^{n+1}T_1^{n+1}y_{n+1} - T_2^{n+1}T_1^{n+1}y_n + T_2^{n+1}T_1^{n+1}y_n - T_2^n T_1^n y_n. \end{aligned}$$

It follows that

$$\begin{aligned} &\|w_{n+1} - w_n\| - \|x_{n+1} - x_n\| \\ &\leq \frac{\alpha_{n+1}}{1 - \beta_{n+1}} (\rho \|V(x_{n+1})\| + \mu \|F(T_2^{n+1}T_1^{n+1}y_{n+1})\|) \\ &\quad + \frac{\alpha_n}{1 - \beta_n} (\mu \|F(T_2^n T_1^n y_n)\| + \rho \|V(x_n)\|) \\ &\quad + \|T_2^{n+1}T_1^{n+1}y_n - T_2^n T_1^n y_n\| + \|y_{n+1} - y_n\| - \|x_{n+1} - x_n\|. \end{aligned} \tag{3.7}$$

Utilizing the nonexpansivity of  $T_2^{n+1}$ , we obtain

$$\begin{aligned} &\|T_2^{n+1}T_1^{n+1}y_n - T_2^n T_1^n y_n\| \\ &= \|T_2^{n+1}T_1^{n+1}y_n - T_2^{n+1}T_1^n y_n\| + \|T_2^{n+1}T_1^n y_n - T_2^n T_1^n y_n\| \\ &\leq \|T_1^{n+1}y_n - T_1^n y_n\| + \|T_2^{n+1}T_1^n y_n - T_2^n T_1^n y_n\|. \end{aligned} \tag{3.8}$$

Note that

$$\begin{aligned} \|T_1^{n+1}y_n - T_1^n y_n\| &= \|(1 - \sigma_{n+1}^1)y_n + \sigma_{n+1}^1 T_1 y_n - (1 - \sigma_n^1)y_n - \sigma_n^1 T_1 y_n\| \\ &\leq |\sigma_{n+1}^1 - \sigma_n^1| (\|y_n\| + \|T_1 y_n\|). \end{aligned}$$

Since  $\lim_{n \rightarrow \infty} |\sigma_{n+1}^i - \sigma_n^i| = 0$  for  $i = 1, 2$  and  $\{y_n\}, \{T_1 y_n\}$  are bounded, we easily obtain

$$\lim_{n \rightarrow \infty} \|T_1^{n+1} y_n - T_1^n y_n\| = 0. \quad (3.9)$$

Similarly, we get

$$\|T_2^{n+1} T_1^n y_n - T_2^n T_1^n y_n\| \leq |\sigma_{n+1}^2 - \sigma_n^2| (\|T_1^n y_n\| + \|T_2 T_1^n y_n\|),$$

which implies that

$$\lim_{n \rightarrow \infty} \|T_2^{n+1} T_1^n y_n - T_2^n T_1^n y_n\| = 0. \quad (3.10)$$

Since  $z_n = P_C(x_n + \delta A^*(T_{r_n}^\ominus - I)Ax_n)$ , it follows from Lemma 2.3 that

$$\begin{aligned} & \|z_{n+1} - z_n\| \\ &= \|P_C(x_{n+1} + \delta A^*(T_{r_{n+1}}^\ominus - I)Ax_{n+1}) - P_C(x_n + \delta A^*(T_{r_n}^\ominus - I)Ax_n)\| \\ &\leq \|x_{n+1} - x_n - \delta A^*A(x_{n+1} - x_n)\| + \delta \|A\| \|T_{r_{n+1}}^\ominus(Ax_{n+1}) - T_{r_n}^\ominus(Ax_n)\| \\ &\quad + |1 - \frac{r_n}{r_{n+1}}| \|P_C(x_{n+1} + \delta A^*(T_{r_{n+1}}^\ominus - I)Ax_{n+1}) \\ &\quad\quad - (x_{n+1} + \delta A^*(T_{r_{n+1}}^\ominus - I)Ax_{n+1})\| \\ &\leq (\|x_{n+1} - x_n\|^2 - 2\delta \|A(x_{n+1} - x_n)\|^2 + \delta^2 \|A\|^4 \|x_{n+1} - x_n\|^2)^{\frac{1}{2}} \\ &\quad + \delta \|A\| (\|A(x_{n+1} - x_n)\| + |1 - \frac{r_n}{r_{n+1}}| \|T_{r_{n+1}}^\ominus(Ax_{n+1}) - Ax_{n+1}\|) \\ &\quad + |1 - \frac{r_n}{r_{n+1}}| \|P_C(x_{n+1} + \delta A^*(T_{r_{n+1}}^\ominus - I)Ax_{n+1}) \\ &\quad\quad - (x_{n+1} + \delta A^*(T_{r_{n+1}}^\ominus - I)Ax_{n+1})\| \\ &= (1 - \delta \|A\|^2) \|x_{n+1} - x_n\| + \delta \|A\|^2 \|x_{n+1} - x_n\| \\ &\quad + \delta \|A\| |1 - \frac{r_n}{r_{n+1}}| \|T_{r_{n+1}}^\ominus(Ax_{n+1}) - Ax_{n+1}\| \\ &\quad + |1 - \frac{r_n}{r_{n+1}}| \|P_C(x_{n+1} + \delta A^*(T_{r_{n+1}}^\ominus - I)Ax_{n+1}) \\ &\quad\quad - (x_{n+1} + \delta A^*(T_{r_{n+1}}^\ominus - I)Ax_{n+1})\| \\ &= \|x_{n+1} - x_n\| + |\frac{r_{n+1} - r_n}{r_{n+1}}| (\delta \|A\| \psi_{n+1} + \chi_{n+1}), \end{aligned}$$

where  $\psi_n = \|T_{r_n}^\ominus(Ax_n) - Ax_n\|$  and  $\chi_n = \|P_C(x_n + \delta A^*(T_{r_n}^\ominus - I)Ax_n) - (x_n + \delta A^*(T_{r_n}^\ominus - I)Ax_n)\|$ . Without loss of generality, let us assume that

there exists a real number  $\mu > 0$  such that  $r_n > \mu > 0$  for all positive integers  $n$ . Then we get

$$\|z_{n+1} - z_n\| \leq \|x_{n+1} - x_n\| + \frac{1}{\mu}|r_{n+1} - r_n|(\delta\|A\|\psi_{n+1} + \chi_{n+1}). \quad (3.11)$$

Next, we estimate that

$$\begin{aligned} & \|y_{n+1} - y_n\|^2 \\ &= \|T_{\mu_1}^{F_1}[T_{\mu_2}^{F_2}(z_{n+1} - \mu_2 B_2 z_{n+1}) - \mu_1 B_1 T_{\mu_2}^{F_2}(z_{n+1} - \mu_2 B_2 z_{n+1})] \\ &\quad - T_{\mu_1}^{F_1}[T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - \mu_1 B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n)]\|^2 \\ &\leq \|T_{\mu_2}^{F_2}(z_{n+1} - \mu_2 B_2 z_{n+1}) - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) \\ &\quad - \mu_1(B_1 T_{\mu_2}^{F_2}(z_{n+1} - \mu_2 B_2 z_{n+1}) - B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n))\|^2 \\ &\leq \|T_{\mu_2}^{F_2}(z_{n+1} - \mu_2 B_2 z_{n+1}) - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n)\|^2 \\ &\quad - \mu_1(2\gamma_1 - \mu_1)\|B_1 T_{\mu_2}^{F_2}(z_{n+1} - \mu_2 B_2 z_{n+1}) - B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n)\|^2 \\ &\leq \|z_{n+1} - z_n - \mu_2(B_2 z_{n+1} - B_2 z_n)\|^2 \\ &\leq \|z_{n+1} - z_n\|^2 - \mu_2(2\gamma_2 - \mu_2)\|B_2 z_{n+1} - B_2 z_n\|^2 \\ &\leq \|z_{n+1} - z_n\|^2. \end{aligned} \quad (3.12)$$

It follows from (3.11) and (3.12) that

$$\begin{aligned} \|y_{n+1} - y_n\| &\leq \|z_{n+1} - z_n\| \\ &\leq \|x_{n+1} - x_n\| + \frac{1}{\mu}|r_{n+1} - r_n|(\delta\|A\|\psi_{n+1} + \chi_{n+1}). \end{aligned} \quad (3.13)$$

Using (3.8), (3.13) in (3.7), we get

$$\begin{aligned} & \|w_{n+1} - w_n\| - \|x_{n+1} - x_n\| \\ &\leq \frac{\alpha_{n+1}}{1 - \beta_{n+1}}(\rho\|V(x_{n+1})\| + \mu\|F(T_2^{n+1}T_1^{n+1}y_{n+1})\|) \\ &\quad + \frac{\alpha_n}{1 - \beta_n}(\mu\|F(T_2^n T_1^n y_n)\| + \rho\|V(x_n)\|) \\ &\quad + \|T_1^{n+1}y_n - T_1^n y_n\| + \|T_2^{n+1}T_1^n y_n - T_2^n T_1^n y_n\| \\ &\quad + \frac{1}{\mu}|r_{n+1} - r_n|(\delta\|A\|\psi_{n+1} + \chi_{n+1}). \end{aligned}$$

Consequently, it follows from (3.9), (3.10) and conditions (i)-(iii) that

$$\limsup_{n \rightarrow \infty} (\|w_{n+1} - w_n\| - \|x_{n+1} - x_n\|) \leq 0.$$

Hence by Lemma 2.8, we have

$$\lim_{n \rightarrow \infty} \|w_n - x_n\| = 0.$$

Consequently,

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = \lim_{n \rightarrow \infty} (1 - \beta_n) \|w_n - x_n\| = 0. \quad (3.14)$$

**Step 3.**  $\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0$  and  $\lim_{n \rightarrow \infty} \|z_n - x_n\| = 0$ .

Indeed, from (3.13), (3.14) and condition (iii), we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \|y_{n+1} - y_n\| &\leq \lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| + \frac{1}{\mu} \lim_{n \rightarrow \infty} |r_{n+1} - r_n| (\delta \|A\| \psi_{n+1} + \chi_{n+1}) \\ &= 0. \end{aligned}$$

Since  $x_{n+1} = \alpha_n \rho V(x_n) + \beta_n x_n + ((1 - \beta_n)I - \alpha_n \mu F) T_2^n T_1^n y_n$ , we obtain

$$\begin{aligned} \|x_n - T_2^n T_1^n y_n\| &\leq \|x_n - x_{n+1}\| + \|x_{n+1} - T_2^n T_1^n y_n\| \\ &\leq \|x_n - x_{n+1}\| + \alpha_n \|\rho V(x_n) - \mu F(T_2^n T_1^n y_n)\| \\ &\quad + \beta_n \|x_n - T_2^n T_1^n y_n\|, \end{aligned}$$

that is,

$$\|x_n - T_2^n T_1^n y_n\| \leq \frac{1}{1 - \beta_n} \|x_n - x_{n+1}\| + \frac{\alpha_n}{1 - \beta_n} \|\rho V(x_n) - \mu F(T_2^n T_1^n y_n)\|.$$

It follows from (3.14) and condition (i) that

$$\lim_{n \rightarrow \infty} \|x_n - T_2^n T_1^n y_n\| = 0. \quad (3.15)$$

From (3.4), (3.5) and Lemma 2.5, we get

$$\begin{aligned} &\|x_{n+1} - x^*\|^2 \\ &= \langle \alpha_n \rho V(x_n) + \beta_n x_n + ((1 - \beta_n)I - \alpha_n \mu F) T_2^n T_1^n y_n - x^*, x_{n+1} - x^* \rangle \\ &= \langle \alpha_n \rho (V(x_n) - V(x^*)), x_{n+1} - x^* \rangle + \alpha_n \langle \rho V(x^*) - \mu F(x^*), x_{n+1} - x^* \rangle \\ &\quad + \beta_n \langle x_n - T_2^n T_1^n y_n, x_{n+1} - x^* \rangle \\ &\quad + \langle (I - \alpha_n \mu F) T_2^n T_1^n y_n - (I - \alpha_n \mu F) T_2^n T_1^n x^*, x_{n+1} - x^* \rangle \\ &\leq \alpha_n \rho \sigma \|x_n - x^*\| \|x_{n+1} - x^*\| + \alpha_n \langle \rho V(x^*) - \mu F(x^*), x_{n+1} - x^* \rangle \\ &\quad + \beta_n \|x_n - T_2^n T_1^n y_n\| \|x_{n+1} - x^*\| + (1 - \alpha_n \tau) \|y_n - x^*\| \|x_{n+1} - x^*\| \end{aligned}$$

$$\begin{aligned}
 &\leq \frac{\alpha_n \rho \sigma}{2} (\|x_n - x^*\|^2 + \|x_{n+1} - x^*\|^2) + \alpha_n \langle \rho V(x^*) - \mu F(x^*), x_{n+1} - x^* \rangle \\
 &\quad + \beta_n \|x_n - T_2^n T_1^n y_n\| \|x_{n+1} - x^*\| + \frac{1 - \alpha_n \tau}{2} (\|y_n - x^*\|^2 + \|x_{n+1} - x^*\|^2) \\
 &\leq \frac{1 - \alpha_n (\tau - \rho \sigma)}{2} \|x_{n+1} - x^*\|^2 + \frac{\alpha_n \rho \sigma}{2} \|x_n - x^*\|^2 \\
 &\quad + \alpha_n \langle \rho V(x^*) - \mu F(x^*), x_{n+1} - x^* \rangle + \beta_n \|x_n - T_2^n T_1^n y_n\| \|x_{n+1} - x^*\| \\
 &\quad + \frac{1 - \alpha_n \tau}{2} \|y_n - x^*\|^2 \\
 &\leq \frac{1 - \alpha_n (\tau - \rho \sigma)}{2} \|x_{n+1} - x^*\|^2 + \frac{\alpha_n \rho \sigma}{2} \|x_n - x^*\|^2 \\
 &\quad + \alpha_n \langle \rho V(x^*) - \mu F(x^*), x_{n+1} - x^* \rangle + \beta_n \|x_n - T_2^n T_1^n y_n\| \|x_{n+1} - x^*\| \\
 &\quad + \frac{1 - \alpha_n \tau}{2} (\|z_n - x^*\|^2 - \mu_2 (2\gamma_2 - \mu_2) \|B_2 z_n - B_2 x^*\|^2 \\
 &\quad - \mu_1 (2\gamma_1 - \mu_1) \|B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 T_{\mu_2}^{F_2}(x^* - \mu_2 B_2 x^*)\|^2) \\
 &\leq \frac{1 - \alpha_n (\tau - \rho \sigma)}{2} \|x_{n+1} - x^*\|^2 + \frac{\alpha_n \rho \sigma}{2} \|x_n - x^*\|^2 \\
 &\quad + \alpha_n \langle \rho V(x^*) - \mu F(x^*), x_{n+1} - x^* \rangle + \beta_n \|x_n - T_2^n T_1^n y_n\| \|x_{n+1} - x^*\| \\
 &\quad + \frac{1 - \alpha_n \tau}{2} (\|x_n - x^*\| - \delta(1 - L\delta)) \|(T_{r_n}^\ominus - I)Ax_n\|^2 \\
 &\quad - \mu_2 (2\gamma_2 - \mu_2) \|B_2 z_n - B_2 x^*\|^2 \\
 &\quad - \mu_1 (2\gamma_1 - \mu_1) \|B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 T_{\mu_2}^{F_2}(x^* - \mu_2 B_2 x^*)\|^2,
 \end{aligned} \tag{3.16}$$

which implies that

$$\begin{aligned}
 \|x_{n+1} - x^*\| &\leq \frac{\alpha_n \rho \sigma}{1 + \alpha_n (\tau - \rho \sigma)} \|x_n - x^*\|^2 \\
 &\quad + \frac{2\alpha_n}{1 + \alpha_n (\tau - \rho \sigma)} \langle \rho V(x^*) - \mu F(x^*), x_{n+1} - x^* \rangle \\
 &\quad + \frac{2\beta_n}{1 + \alpha_n (\tau - \rho \sigma)} \|x_n - T_2^n T_1^n y_n\| \|x_{n+1} - x^*\| \\
 &\quad + \|x_n - x^*\|^2 - \frac{1 - \alpha_n \tau}{1 + \alpha_n (\tau - \rho \sigma)} \{ \delta(1 - L\delta) \|(T_{r_n}^\ominus - I)Ax_n\|^2 \\
 &\quad + \mu_2 (2\gamma_2 - \mu_2) \|B_2 z_n - B_2 x^*\|^2 \\
 &\quad + \mu_1 (2\gamma_1 - \mu_1) \|B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 T_{\mu_2}^{F_2}(x^* - \mu_2 B_2 x^*)\|^2 \}.
 \end{aligned}$$

Then from the above inequality, we get

$$\begin{aligned}
& \frac{1 - \alpha_n \tau}{1 + \alpha_n(\tau - \rho\sigma)} \{ \delta(1 - L\delta) \|(T_{r_n}^\ominus - I)Ax_n\|^2 + \mu_2(2\gamma_2 - \mu_2)\|B_2z_n - B_2x^*\|^2 \\
& \quad + \mu_1(2\gamma_1 - \mu_1)\|B_1T_{\mu_2}^{F_2}(z_n - \mu_2B_2z_n) - B_1T_{\mu_2}^{F_2}(x^* - \mu_2B_2x^*)\|^2 \} \\
& \leq \frac{\alpha_n\rho\sigma}{1 + \alpha_n(\tau - \rho\sigma)} \|x_n - x^*\|^2 + \frac{2\alpha_n}{1 + \alpha_n(\tau - \rho\sigma)} \langle \rho V(x^*) - \mu F(x^*), x_{n+1} - x^* \rangle \\
& \quad + \frac{2\beta_n}{1 + \alpha_n(\tau - \rho\sigma)} \|x_n - T_2^n T_1^n y_n\| \|x_{n+1} - x^*\| + \|x_n - x^*\|^2 - \|x_{n+1} - x^*\|^2 \\
& \leq \frac{\alpha_n\rho\sigma}{1 + \alpha_n(\tau - \rho\sigma)} \|x_n - x^*\|^2 + \frac{2\alpha_n}{1 + \alpha_n(\tau - \rho\sigma)} \langle \rho V(x^*) - \mu F(x^*), x_{n+1} - x^* \rangle \\
& \quad + \frac{2\beta_n}{1 + \alpha_n(\tau - \rho\sigma)} \|x_n - T_2^n T_1^n y_n\| \|x_{n+1} - x^*\| \\
& \quad + (\|x_n - x^*\| + \|x_{n+1} - x^*\|) \|x_{n+1} - x_n\|.
\end{aligned}$$

From  $\delta(1 - L\delta) > 0$ ,  $2\gamma_i - \mu_i > 0$  for  $i = 1, 2$ , (3.14) and (3.15), we obtain

$$\lim_{n \rightarrow \infty} \|(T_{r_n}^\ominus - I)Ax_n\| = 0, \quad \lim_{n \rightarrow \infty} \|B_2z_n - B_2x^*\| = 0 \quad (3.17)$$

and

$$\lim_{n \rightarrow \infty} \|B_1T_{\mu_2}^{F_2}(z_n - \mu_2B_2z_n) - B_1T_{\mu_2}^{F_2}(x^* - \mu_2B_2x^*)\| = 0. \quad (3.18)$$

Since  $P_C$  is firmly nonexpansive, we have

$$\begin{aligned}
& \|z_n - x^*\|^2 \\
& = \|P_C(x_n + \delta A^*(T_{r_n}^\ominus - I)Ax_n) - P_C(x^*)\|^2 \\
& \leq \langle z_n - x^*, x_n + \delta A^*(T_{r_n}^\ominus - I)Ax_n - x^* \rangle \\
& = \frac{1}{2} \{ \|z_n - x^*\|^2 + \|x_n + \delta A^*(T_{r_n}^\ominus - I)Ax_n - x^*\|^2 \\
& \quad - \|z_n - x^* - (x_n + \delta A^*(T_{r_n}^\ominus - I)Ax_n - x^*)\|^2 \} \\
& \leq \frac{1}{2} \{ \|z_n - x^*\|^2 + \|x_n - x^*\|^2 - \|z_n - x_n - \delta A^*(T_{r_n}^\ominus - I)Ax_n\|^2 \} \\
& \leq \frac{1}{2} \{ \|z_n - x^*\|^2 + \|x_n - x^*\|^2 - (\|z_n - x_n\|^2 + \delta^2 \|A^*(T_{r_n}^\ominus - I)Ax_n\|^2 \\
& \quad - 2\delta \langle z_n - x_n, A^*(T_{r_n}^\ominus - I)Ax_n \rangle) \}.
\end{aligned}$$



Hence we get

$$\|z_n - x^*\|^2 \leq \|x_n - x^*\|^2 - \|z_n - x_n\|^2 + 2\delta \|Az_n - Ax_n\| \|(T_{r_n}^\Theta - I)Ax_n\|. \quad (3.19)$$

Utilizing the firm nonexpansivity of  $T_{\mu_2}^{F_2}$ , we have

$$\begin{aligned} & \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y^*\|^2 \\ &= \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - T_{\mu_2}^{F_2}(x^* - \mu_2 B_2 x^*)\|^2 \\ &\leq \langle T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y^*, z_n - \mu_2 B_2 z_n - (x^* - \mu_2 B_2 x^*) \rangle \\ &= \frac{1}{2} \{ \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y^*\|^2 + \|z_n - x^* - \mu_2(B_2 z_n - B_2 x^*)\|^2 \\ &\quad - \|z_n - x^* - \mu_2(B_2 z_n - B_2 x^*) - (T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y^*)\|^2 \} \\ &\leq \frac{1}{2} \{ \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y^*\|^2 + \|z_n - x^*\|^2 - \mu_2(2\gamma_2 - \mu_2) \|B_2 z_n - B_2 x^*\|^2 \\ &\quad - \|z_n - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (x^* - y^*) - \mu_2(B_2 z_n - B_2 x^*)\|^2 \} \\ &\leq \frac{1}{2} \{ \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y^*\|^2 + \|z_n - x^*\|^2 \\ &\quad - \|z_n - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (x^* - y^*)\|^2 \\ &\quad + 2\mu_2 \langle z_n - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (x^* - y^*), B_2 z_n - B_2 x^* \rangle \\ &\quad - \mu_2^2 \|B_2 z_n - B_2 x^*\|^2 \} \\ &\leq \frac{1}{2} \{ \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y^*\|^2 + \|z_n - x^*\|^2 \\ &\quad - \|z_n - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (x^* - y^*)\|^2 \\ &\quad + 2\mu_2 \|z_n - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (x^* - y^*)\| \|B_2 z_n - B_2 x^*\| \}. \end{aligned}$$

It follows from (3.19) that

$$\begin{aligned} & \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y^*\|^2 \\ &\leq \|z_n - x^*\|^2 - \|z_n - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (x^* - y^*)\|^2 \\ &\quad + 2\mu_2 \|z_n - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (x^* - y^*)\| \|B_2 z_n - B_2 x^*\| \\ &\leq \|x_n - x^*\|^2 - \|z_n - x_n\|^2 + 2\delta \|Az_n - Ax_n\| \|(T_{r_n}^\Theta - I)Ax_n\| \\ &\quad - \|z_n - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (x^* - y^*)\|^2 \\ &\quad + 2\mu_2 \|z_n - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (x^* - y^*)\| \|B_2 z_n - B_2 x^*\|. \quad (3.20) \end{aligned}$$

And we have

$$\begin{aligned}
& \|y_n - x^*\|^2 \\
&= \|T_{\mu_1}^{F_1}(T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - \mu_1 B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n)) - T_{\mu_1}^{F_1}(y^* - \mu_1 B_1 y^*)\|^2 \\
&\leq \langle y_n - x^*, T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - \mu_1 B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (y^* - \mu_1 B_1 y^*) \rangle \\
&= \frac{1}{2} \{ \|y_n - x^*\|^2 + \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y^* - \mu_1 (B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 y^*)\|^2 \\
&\quad - \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y^* - \mu_1 (B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 y^*) - (y_n - x^*)\|^2 \} \\
&= \frac{1}{2} \{ \|y_n - x^*\|^2 + \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y^*\|^2 \\
&\quad - 2\mu_1 \langle T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y^*, B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 y^* \rangle \\
&\quad + \mu_1^2 \|B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 y^*\|^2 \\
&\quad - \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y_n - \mu_1 (B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 y^*) + x^* - y^*\|^2 \} \\
&\leq \frac{1}{2} \{ \|y_n - x^*\|^2 + \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y^*\|^2 \\
&\quad - \mu_1 (2\gamma_1 - \mu_1) \|B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 y^*\|^2 \\
&\quad - \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y_n - \mu_1 (B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 y^*) + x^* - y^*\|^2 \} \\
&\leq \frac{1}{2} \{ \|y_n - x^*\|^2 + \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y^*\|^2 \\
&\quad - \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y_n + x^* - y^*\|^2 \\
&\quad + 2\mu_1 \langle T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y_n + x^* - y^*, B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 y^* \rangle \}.
\end{aligned}$$

So, we obtain from (3.20) that

$$\begin{aligned}
& \|y_n - x^*\|^2 \\
&\leq \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y^*\|^2 - \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y_n + x^* - y^*\|^2 \\
&\quad + 2\mu_1 \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y_n + x^* - y^*\| \|B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 y^*\| \\
&\leq \|x_n - x^*\|^2 - \|z_n - x_n\|^2 + 2\delta \|Az_n - Ax_n\| \| (T_{r_n}^\ominus - I) Ax_n \| \\
&\quad - \|z_n - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (x^* - y^*)\|^2 \\
&\quad + 2\mu_2 \|z_n - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (x^* - y^*)\| \|B_2 z_n - B_2 x^*\| \\
&\quad - \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y_n + x^* - y^*\|^2 \\
&\quad + 2\mu_1 \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y_n + x^* - y^*\| \|B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 y^*\|.
\end{aligned} \tag{3.21}$$

From (3.16) and (3.21), we have

$$\begin{aligned}
& \|x_{n+1} - x^*\|^2 \\
& \leq \frac{1 - \alpha_n(\tau - \rho\sigma)}{2} \|x_{n+1} - x^*\|^2 + \frac{\alpha_n\rho\sigma}{2} \|x_n - x^*\|^2 \\
& \quad + \alpha_n \langle \rho V(x^*) - \mu F(x^*), x_{n+1} - x^* \rangle + \beta_n \|x_n - T_2^n T_1^n y_n\| \|x_{n+1} - x^*\| \\
& \quad + \frac{1 - \alpha_n\tau}{2} \|y_n - x^*\|^2 \\
& \leq \frac{1 - \alpha_n(\tau - \rho\sigma)}{2} \|x_{n+1} - x^*\|^2 + \frac{\alpha_n\rho\sigma}{2} \|x_n - x^*\|^2 \\
& \quad + \alpha_n \langle \rho V(x^*) - \mu F(x^*), x_{n+1} - x^* \rangle + \beta_n \|x_n - T_2^n T_1^n y_n\| \|x_{n+1} - x^*\| \\
& \quad + \frac{1 - \alpha_n\tau}{2} \{ \|x_n - x^*\|^2 - \|z_n - x_n\|^2 + 2\delta \|Az_n - Ax_n\| \| (T_{r_n}^\Theta - I) Ax_n \| \\
& \quad - \|z_n - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (x^* - y^*) \|^2 \\
& \quad + 2\mu_2 \|z_n - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (x^* - y^*)\| \|B_2 z_n - B_2 x^*\| \\
& \quad - \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y_n + x^* - y^*\|^2 \\
& \quad + 2\mu_1 \|T_{\mu_1}^{F_2}(z_n - \mu_2 B_2 z_n) - y_n + x^* - y^*\| \|B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 y^*\| \},
\end{aligned}$$

which implies that

$$\begin{aligned}
& \|x_{n+1} - x^*\|^2 \\
& \leq \frac{\alpha_n\rho\sigma}{1 + \alpha_n(\tau - \rho\sigma)} \|x_n - x^*\|^2 + \frac{2\alpha_n}{1 + \alpha_n(\tau - \rho\sigma)} \langle \rho V(x^*) - \mu F(x^*), x_{n+1} - x^* \rangle \\
& \quad + \frac{2\beta_n}{1 + \alpha_n(\tau - \rho\sigma)} \|x_n - T_2^n T_1^n y_n\| \|x_{n+1} - x^*\| \\
& \quad + \frac{1 - \alpha_n\tau}{1 + \alpha_n(\tau - \rho\sigma)} \{ \|x_n - x^*\|^2 - \|z_n - x_n\|^2 + 2\delta \|Az_n \\
& \quad - Ax_n\| \| (T_{r_n}^\Theta - I) Ax_n \| \\
& \quad - \|z_n - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (x^* - y^*) \|^2 \\
& \quad + 2\mu_2 \|z_n - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (x^* - y^*)\| \|B_2 z_n - B_2 x^*\| \} \\
& \quad - \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y_n + x^* - y^*\|^2 \\
& \quad + 2\mu_1 \|T_{\mu_1}^{F_2}(z_n - \mu_2 B_2 z_n) - y_n + x^* - y^*\| \|B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 y^*\| \}
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{\alpha_n \rho \sigma}{1 + \alpha_n(\tau - \rho \sigma)} \|x_n - x^*\|^2 + \frac{2\alpha_n}{1 + \alpha_n(\tau - \rho \sigma)} \langle \rho V(x^*) - \mu F(x^*), x_{n+1} - x^* \rangle \\
&\quad + \frac{2\beta_n}{1 + \alpha_n(\tau - \rho \sigma)} \|x_n - T_2^n T_1^n y_n\| \|x_{n+1} - x^*\| + \|x_n - x^*\|^2 \\
&\quad + 2\delta \|Az_n - Ax_n\| \|(T_{r_n}^\Theta - I)Ax_n\| \\
&\quad + 2\mu_2 \|z_n - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (x^* - y^*)\| \|B_2 z_n - B_2 x^*\| \\
&\quad + 2\mu_1 \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y_n + x^* - y^*\| \|B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 y^*\| \\
&\quad - \frac{1 - \alpha_n \tau}{1 + \alpha_n(\tau - \rho \sigma)} \{ \|z_n - x_n\|^2 + \|z_n - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (x^* - y^*)\|^2 \\
&\quad + \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y_n + x^* - y^*\|^2 \}.
\end{aligned}$$

Therefore we obtain

$$\begin{aligned}
&\frac{1 - \alpha_n \tau}{1 + \alpha_n(\tau - \rho \sigma)} \{ \|z_n - x_n\|^2 + \|z_n - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (x^* - y^*)\|^2 \\
&\quad + \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y_n + x^* - y^*\|^2 \} \\
&\leq \frac{\alpha_n \rho \sigma}{1 + \alpha_n(\tau - \rho \sigma)} \|x_n - x^*\|^2 + \frac{2\alpha_n}{1 + \alpha_n(\tau - \rho \sigma)} \langle \rho V(x^*) - \mu F(x^*), x_{n+1} - x^* \rangle \\
&\quad + \frac{2\beta_n}{1 + \alpha_n(\tau - \rho \sigma)} \|x_n - T_2^n T_1^n y_n\| \|x_{n+1} - x^*\| \\
&\quad + (\|x_n - x^*\| + \|x_{n+1} - x^*\|) \|x_{n+1} - x_n\| + 2\delta \|Az_n - Ax_n\| \|(T_{r_n}^\Theta - I)Ax_n\| \\
&\quad + 2\mu_2 \|z_n - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (x^* - y^*)\| \|B_2 z_n - B_2 x^*\| \\
&\quad + 2\mu_1 \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y_n + x^* - y^*\| \|B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - B_1 y^*\|.
\end{aligned}$$

Thus, from (3.14), (3.15), (3.17) and (3.18), we conclude that

$$\lim_{n \rightarrow \infty} \|z_n - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (x^* - y^*)\| = 0,$$

$$\lim_{n \rightarrow \infty} \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y_n + x^* - y^*\| = 0$$

and

$$\lim_{n \rightarrow \infty} \|z_n - x_n\| = 0. \quad (3.22)$$

Since

$$\begin{aligned}
\|z_n - y_n\| &\leq \|z_n - T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - (x^* - y^*)\| \\
&\quad + \|T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - y_n + x^* - y^*\|,
\end{aligned}$$

we get

$$\lim_{n \rightarrow \infty} \|z_n - y_n\| = 0. \tag{3.23}$$

It follows from (3.22) and (3.23) that

$$\begin{aligned} \lim_{n \rightarrow \infty} \|x_n - y_n\| &\leq \lim_{n \rightarrow \infty} \|x_n - z_n\| + \lim_{n \rightarrow \infty} \|z_n - y_n\| \\ &= 0. \end{aligned} \tag{3.24}$$

**Step 4.**  $\lim_{n \rightarrow \infty} \|x_n - T_2^n T_1^n x_n\| = 0$ .

Indeed, from (3.1), we get

$$\begin{aligned} \|x_n - T_2^n T_1^n x_n\| &\leq \|x_n - x_{n+1}\| + \|x_{n+1} - T_2^n T_1^n x_n\| \\ &\leq \|x_n - x_{n+1}\| + \|\alpha_n(\rho V(x_n) - \mu F(T_2^n T_1^n y_n)) \\ &\quad + \beta_n(x_n - T_2^n T_1^n y_n) + T_2^n T_1^n y_n - T_2^n T_1^n x_n\| \\ &\leq \|x_n - x_{n+1}\| + \alpha_n \|\rho V(x_n) - \mu F(T_2^n T_1^n y_n)\| \\ &\quad + \beta_n \|x_n - T_2^n T_1^n y_n\| + \|y_n - x_n\|. \end{aligned}$$

So, from (3.14), (3.15), (3.24) and condition (i), we have

$$\lim_{n \rightarrow \infty} \|x_n - T_2^n T_1^n x_n\| = 0. \tag{3.25}$$

**Step 5.**  $\limsup_{n \rightarrow \infty} \langle (\mu F - \gamma V)x^*, x^* - x_n \rangle \leq 0$ , where  $x^* = P_\Gamma(I - \mu F + \gamma V)x^*$ .

Indeed, since  $\{\sigma_k^i\}$  is bounded for  $i = 1, 2$ , we can assume that  $\sigma_{k_j}^i \rightarrow \sigma_\infty^i$  as  $j \rightarrow \infty$ , where  $0 < \zeta_1 \leq \sigma_\infty^i \leq \zeta_2 < 1$  for  $i = 1, 2$ . Define  $T_i^\infty = (1 - \sigma_\infty^i)I + \sigma_\infty^i T_i$  for  $i = 1, 2$ . Then we have  $\text{Fix}(T_i^\infty) = \text{Fix}(T_i)$  for  $i = 1, 2$ . Note that

$$\begin{aligned} \|T_i^{k_j} x - T_i^\infty x\| &= \|(1 - \sigma_{k_j}^i)x + \sigma_{k_j}^i T_i x - (1 - \sigma_\infty^i)x - \sigma_\infty^i T_i x\| \\ &\leq |\sigma_{k_j}^i - \sigma_\infty^i| (\|x\| + \|T_i x\|). \end{aligned}$$

Hence we deduce that

$$\limsup_{j \rightarrow \infty} \sup_{x \in D} \|T_i^{k_j} x - T_i^\infty x\| = 0, \tag{3.26}$$

where  $D$  is an arbitrary bounded subset of  $H_1$ . Since  $\text{Fix}(T_1^\infty) \cap \text{Fix}(T_2^\infty) = \text{Fix}(T_1) \cap \text{Fix}(T_2) \neq \emptyset$  and  $T_i^\infty$  is  $\sigma_\infty^i$ -averaged for  $i = 1, 2$ , by Lemma 2.6, we know that  $\text{Fix}(T_2^\infty T_1^\infty) = \text{Fix}(T_2^\infty) \cap \text{Fix}(T_1^\infty)$ . Since  $\{x_n\}$  is

bounded, there exists a subsequence  $\{x_{n_j}\}$  of  $\{x_n\}$  such that  $x_{n_j} \rightharpoonup w$  as  $j \rightarrow \infty$ . Note that

$$\begin{aligned} \|x_{n_j} - T_2^\infty T_1^\infty x_{n_j}\| &\leq \|x_{n_j} - T_2^{n_j} T_1^{n_j} x_{n_j}\| + \|T_2^{n_j} T_1^{n_j} x_{n_j} - T_2^\infty T_1^{n_j} x_{n_j}\| \\ &\quad + \|T_2^\infty T_1^{n_j} x_{n_j} - T_2^\infty T_1^\infty x_{n_j}\| \\ &\leq \|x_{n_j} - T_2^{n_j} T_1^{n_j} x_{n_j}\| + \|T_2^{n_j} T_1^{n_j} x_{n_j} - T_2^\infty T_1^{n_j} x_{n_j}\| \\ &\quad + \|T_1^{n_j} x_{n_j} - T_1^\infty x_{n_j}\| \\ &\leq \|x_{n_j} - T_2^{n_j} T_1^{n_j} x_{n_j}\| + \sup_{x \in D'} \|T_2^{n_j} x - T_2^\infty x\| \\ &\quad + \sup_{x \in D''} \|T_1^{n_j} x - T_1^\infty x\|, \end{aligned}$$

where  $D'$  is a bounded subset including  $\{T_1^{n_j} x_{n_j}\}$  and  $D''$  is a bounded subset including  $\{x_{n_j}\}$ . According to (3.25) and (3.26), we obtain

$$\lim_{n \rightarrow \infty} \|x_{n_j} - T_2^\infty T_1^\infty x_{n_j}\| = 0.$$

First, it is clear from Lemma 2.7 that  $w \in \text{Fix}(T_2^\infty T_1^\infty)$ .

Second, let us show that  $Aw \in EP(\Theta)$ . Since  $A$  is a bounded linear operator,  $Ax_{n_j} \rightarrow Aw$ . Now we set  $\nu_{n_j} = Ax_{n_j} - T_{r_{n_j}}^\Theta Ax_{n_j}$ . It follows from (3.17) that  $\lim_{j \rightarrow \infty} \nu_{n_j} = 0$  and  $Ax_{n_j} - \nu_{n_j} = T_{r_{n_j}}^\Theta Ax_{n_j}$ . Therefore, from the definition of  $T_{r_{n_j}}^\Theta$ , we have

$$\Theta(Ax_{n_j} - \nu_{n_j}, y) + \frac{1}{r_{n_j}} \langle y - (Ax_{n_j} - \nu_{n_j}), (Ax_{n_j} - \nu_{n_j}) - Ax_{n_j} \rangle \geq 0, \forall y \in K.$$

Since  $\Theta$  is upper semicontinuous in the first argument, taking lim sup in the above inequality as  $j \rightarrow \infty$ , we obtain

$$\Theta(Aw, y) \geq 0, \quad \forall y \in K,$$

which implies that  $Aw \in EP(\Theta)$  and  $w \in \Lambda$ .

Next, we show that  $w \in \Omega$ . From (3.22), it is easy to observe that  $z_{n_j} \rightarrow w$ . For any  $x, y \in C$ , we have

$$\begin{aligned}
\|Q(x) - Q(y)\|^2 &= \|T_{\mu_1}^{F_1}[T_{\mu_2}^{F_2}(x - \mu_2 B_2 x) - \mu_1 B_1 T_{\mu_2}^{F_2}(x - \mu_2 B_2 x)] \\
&\quad - T_{\mu_1}^{F_1}[T_{\mu_2}^{F_2}(y - \mu_2 B_2 y) - \mu_1 B_1 T_{\mu_2}^{F_2}(y - \mu_2 B_2 y)]\|^2 \\
&\leq \|(T_{\mu_2}^{F_2}(x - \mu_2 B_2 x) - T_{\mu_2}^{F_2}(y - \mu_2 B_2 y) \\
&\quad - \mu_1(B_1 T_{\mu_2}^{F_2}(x - \mu_2 B_2 x) - B_1 T_{\mu_2}^{F_2}(y - \mu_2 B_2 y))\|^2 \\
&\leq \|T_{\mu_2}^{F_2}(x - \mu_2 B_2 x) - T_{\mu_2}^{F_2}(y - \mu_2 B_2 y)\|^2 \\
&\quad - \mu_1(2\gamma_1 - \mu_1)\|B_1 T_{\mu_2}^{F_2}(x - \mu_2 B_2 x) - B_1 T_{\mu_2}^{F_2}(y - \mu_2 B_2 x)\|^2 \\
&\leq \|T_{\mu_2}^{F_2}(x - \mu_2 B_2 x) - T_{\mu_2}^{F_2}(y - \mu_2 B_2 y)\|^2 \\
&\leq \|(x - \mu_2 B_2 x) - (y - \mu_2 B_2 y)\|^2 \\
&\leq \|x - y\|^2 - \mu_2(2\gamma_2 - \mu_2)\|B_2 x - B_2 y\|^2 \\
&\leq \|x - y\|^2.
\end{aligned}$$

This shows that  $Q : C \rightarrow C$  is nonexpansive. Note that

$$\begin{aligned}
\|y_n - Q(y_n)\| &= \|Q(z_n) - Q(y_n)\| \\
&\leq \|z_n - y_n\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.
\end{aligned}$$

It follows from Lemma 2.7 that  $w = Q(w)$ . According to Lemma 2.4, we obtain  $w \in \Omega$ . Thus we have  $w \in \Gamma = \bigcap_{i=1}^N \text{Fix}(T_i) \cap \Omega \cap \Lambda$ .

Finally, we claim that  $\limsup_{n \rightarrow \infty} \langle (\mu F - \rho V)x^*, x^* - x_n \rangle \leq 0$ . By  $x^* = P_{\Gamma}(I - \mu F + \rho V)x^*$  and (2.1), we have

$$\begin{aligned}
\limsup_{n \rightarrow \infty} \langle (\mu F - \rho V)x^*, x^* - x_n \rangle &= \lim_{n \rightarrow \infty} \langle (\mu F - \rho V)x^*, x^* - x_{n_k} \rangle \\
&= \langle (\mu F - \rho V)x^*, x^* - w \rangle \\
&\leq 0.
\end{aligned}$$

**Step 6.**  $x_n \rightarrow x^*$  as  $n \rightarrow \infty$ .

Indeed, from (3.1), (3.5), Lemma 2.1 and 2.5, we have

$$\begin{aligned}
& \|x_{n+1} - x^*\|^2 \\
&= \|\alpha_n \rho V(x_n) + \beta_n x_n + ((1 - \beta_n)I - \alpha_n \mu F)T_2^n T_1^n y_n - x^*\|^2 \\
&= \|\alpha_n (\rho V(x_n) - \mu F(x^*)) + \beta_n (x_n - x^*) + ((1 - \beta_n)I - \alpha_n \mu F)T_2^n T_1^n y_n \\
&\quad - ((1 - \beta_n)I - \alpha_n \mu F)T_2^n T_1^n x^*\|^2 \\
&\leq \|\beta_n (x_n - x^*) + ((1 - \beta_n)I - \alpha_n \mu F)T_2^n T_1^n y_n - ((1 - \beta_n)I - \alpha_n \mu F)T_2^n T_1^n x^*\|^2 \\
&\quad + 2\alpha_n \langle \rho V(x_n) - \mu F(x^*), x_{n+1} - x^* \rangle \\
&\leq (\beta_n \|x_n - x^*\| + (1 - \beta_n) \|(I - \frac{\alpha_n}{1 - \beta_n} \mu F)T_2^n T_1^n y_n - (I - \frac{\alpha_n}{1 - \beta_n} \mu F)T_2^n T_1^n x^*\|)^2 \\
&\quad + 2\alpha_n \rho \langle V(x_n) - V(x^*), x_{n+1} - x^* \rangle + 2\alpha_n \langle \rho V(x^*) - \mu F(x^*), x_{n+1} - x^* \rangle \\
&\leq (\beta_n \|x_n - x^*\| + (1 - \beta_n)(1 - \frac{\alpha_n}{1 - \beta_n} \tau) \|y_n - x^*\|)^2 \\
&\quad + 2\alpha_n \rho \sigma \|x_n - x^*\| \|x_{n+1} - x^*\| + 2\alpha_n \langle \rho V(x^*) - \mu F(x^*), x_{n+1} - x^* \rangle \\
&\leq (\beta_n \|x_n - x^*\| + (1 - \beta_n - \alpha_n \tau) \|x_n - x^*\|)^2 \\
&\quad + \alpha_n \rho \sigma (\|x_n - x^*\|^2 + \|x_{n+1} - x^*\|^2) + 2\alpha_n \langle \rho V(x^*) - \mu F(x^*), x_{n+1} - x^* \rangle \\
&= [(1 - \alpha_n \tau)^2 + \alpha_n \rho \sigma] \|x_n - x^*\|^2 + \alpha_n \rho \sigma \|x_{n+1} - x^*\|^2 \\
&\quad + 2\alpha_n \langle \rho V(x^*) - \mu F(x^*), x_{n+1} - x^* \rangle.
\end{aligned}$$

This implies that

$$\begin{aligned}
& \|x_{n+1} - x^*\|^2 \\
&\leq \frac{(1 - \alpha_n \tau)^2 + \alpha_n \rho \sigma}{1 - \alpha_n \rho \sigma} \|x_n - x^*\|^2 \\
&\quad + \frac{2\alpha_n}{1 - \alpha_n \rho \sigma} \langle \rho V(x^*) - \mu F(x^*), x_{n+1} - x^* \rangle \\
&= \left(1 - \frac{2(\tau - \rho\sigma)\alpha_n}{1 - \alpha_n \rho \sigma}\right) \|x_n - x^*\|^2 + \frac{\alpha_n^2 \tau^2}{1 - \alpha_n \rho \sigma} \|x_n - x^*\|^2 \\
&\quad + \frac{2\alpha_n}{1 - \alpha_n \rho \sigma} \langle \rho V(x^*) - \mu F(x^*), x_{n+1} - x^* \rangle \\
&\leq \left(1 - \frac{2(\tau - \rho\sigma)\alpha_n}{1 - \alpha_n \rho \sigma}\right) \|x_n - x^*\|^2 + \frac{2(\tau - \rho\sigma)\alpha_n}{1 - \alpha_n \rho \sigma} \left\{ \frac{\alpha_n \tau^2}{2(\tau - \rho\sigma)} M_1 \right. \\
&\quad \left. + \frac{1}{\tau - \rho\sigma} \langle \rho V(x^*) - \mu F(x^*), x_{n+1} - x^* \rangle \right\} \\
&= (1 - \delta_n) \|x_n - x^*\|^2 + \delta_n \sigma_n,
\end{aligned}$$



where  $M_1 = \sup\{\|x_n - x^*\|^2 : n \geq 0\}$ ,  $\delta_n = \frac{2(\tau-\rho\sigma)\alpha_n}{1-\alpha_n\rho\sigma}$  and  $\sigma_n = \frac{\alpha_n\tau^2}{2(\tau-\rho\sigma)}M_1 + \frac{1}{\tau-\rho\sigma}\langle\rho V(x^*) - \mu F(x^*), x_{n+1} - x^*\rangle$ . It is easy to see that  $\delta_n \rightarrow 0$ ,  $\sum_{n=0}^\infty \delta_n = \infty$  and  $\limsup_{n \rightarrow \infty} \sigma_n \leq 0$ . Hence, applying Lemma 2.9 to the last inequality, we immediately obtain that  $x_n \rightarrow x^*$  as  $n \rightarrow \infty$ . This completes the proof.  $\square$

**COROLLARY 3.1.** *Let  $C, K$  be nonempty closed convex subsets of real Hilbert spaces  $H_1, H_2$ , respectively. Let  $A : H_1 \rightarrow H_2$  be a bounded linear operator with its adjoint  $A^*$ . Assume  $F_1, F_2 : C \times C \rightarrow \mathbb{R}$  and  $\Theta : K \times K \rightarrow \mathbb{R}$  are the bifunctions satisfying Assumption 2.1. Let  $B_i : C \rightarrow H_1$  be a  $\gamma_i$ -inverse strongly monotone mapping for each  $i = 1, 2$  such that  $\Gamma = \Omega \cap \Lambda \neq \phi$ . Let  $F : C \rightarrow C$  be a  $\kappa$ -Lipschitzian continuous and  $\eta$ -strongly monotone mapping with  $\kappa > 0$  and  $\eta > 0$ , and  $V : C \rightarrow C$  be a  $\sigma$ -Lipschitzian continuous mapping with  $\sigma > 0$ . Let  $0 < \mu < \frac{2\eta}{\kappa^2}$  and  $0 < \rho\sigma < \tau$ , where  $\tau = 1 - \sqrt{1 - \mu(2\eta - \mu\kappa^2)}$ . Suppose  $\{\alpha_n\}$  and  $\{\beta_n\}$  are two sequences in  $(0, 1)$ . Given  $x_1 \in C$ , let  $\{x_n\}$  be defined by*

$$\begin{cases} z_n = P_C(x_n + \delta A^*(T_{r_n}^\Theta - I)Ax_n), \\ y_n = T_{\mu_1}^{F_1}[T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n) - \mu_1 B_1 T_{\mu_2}^{F_2}(z_n - \mu_2 B_2 z_n)], \\ x_{n+1} = \alpha_n \rho V(x_n) + \beta_n x_n + ((1 - \beta_n)I - \alpha_n \mu F)y_n, \quad \forall n \geq 1, \end{cases}$$

where  $\{r_n\} \subset (0, 2\zeta)$ ,  $\zeta > 0$ ,  $\mu_i \in (0, 2\gamma_i)$  for each  $i = 1, 2$ ,  $\delta \in (0, \frac{1}{L})$ ,  $L$  is the spectral radius of the operator  $A^*A$ ,  $A^*$  is the adjoint of  $A$ . If the following conditions are satisfied:

- (i)  $\lim_{n \rightarrow \infty} \alpha_n = 0$  and  $\sum_{n=1}^\infty \alpha_n = 0$ ;
- (ii)  $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$ ;
- (iii)  $0 < \liminf_{n \rightarrow \infty} r_n \leq \limsup_{n \rightarrow \infty} r_n < 2\zeta$  and  $\lim_{n \rightarrow \infty} |r_{n+1} - r_n| = 0$ .

Then the sequence  $\{x_n\}$  converges strongly to  $x^* \in \Omega \cap \Lambda$ , where  $x^* = P_{\Omega \cap \Lambda}(I - \mu F + \rho V)x^*$  is the unique solution of the variational inequality:

$$\langle (\mu F - \rho V)x^*, x - x^* \rangle \geq 0, \quad \forall x \in \Omega \cap \Lambda.$$

*Proof.* Put  $T_i x = x$  for all  $i = 1, 2, \dots, N$  and  $x \in C$ , and take the finite family of sequences  $\{\sigma_n^i\}_{i=1}^N$  in  $(\zeta_1, \zeta_2)$  for some  $\zeta_1, \zeta_2 \in (0, 1)$  such that  $\lim_{n \rightarrow \infty} |\sigma_{n+1}^i - \sigma_n^i| = 0$  for all  $i = 1, 2, \dots, N$ . In this case,  $T_N^n T_{N-1}^n \dots T_1^n$  is the identity mapping  $I$  of  $C$ . It is easy to see that all

conditions of Theorem 3.1 are satisfied. Thus, the desired result follows from Theorem 3.1.  $\square$

To verify the theoretical assertions, we consider the following example.

**EXAMPLE 3.1.** Let  $H_1 = H_2 = \mathbb{R}$  with the inner product defined by  $\langle x, y \rangle = xy$  for all  $x, y \in \mathbb{R}$  and the standard norm  $|\cdot|$ . Let  $C = [0, +\infty)$ ,  $K = (-\infty, 0]$ ,  $B_1 = B_2 = 0$  and  $\mu_1 = \mu_2 = 1$ . Define the mappings  $A : \mathbb{R} \rightarrow \mathbb{R}$ ,  $F_1, F_2 : C \times C \rightarrow \mathbb{R}$ ,  $\Theta : K \times K \rightarrow \mathbb{R}$ ,  $T_i : C \rightarrow C$  for each  $i = 1, 2, \dots, N$ ,  $F : C \rightarrow C$  and  $V : C \rightarrow C$  as follows:

$$F_1(x, y) = -3x^2 + xy + 2y^2, \quad F_2(x, y) = -5x^2 + xy + 4y^2,$$

$$T_i(x) = 0 \text{ for } i = 1, 2, \dots, N, \quad F(x) = x, \quad V(x) = \frac{1}{2}x, \quad \forall (x, y) \in C \times C,$$

$$\Theta(x, y) = -x(x - y), \quad \forall (x, y) \in K \times K$$

and

$$Ax = -2x, \quad \forall x \in \mathbb{R}.$$

It is easy to see that  $\sigma = \frac{1}{2}$ ,  $\eta = \kappa = 1$  and hence  $0 < \mu < \frac{2\eta}{\kappa^2} = 2$ . Put  $\mu = 1$ . Then  $\tau = 1 - \sqrt{1 - \mu(2\eta - \mu\kappa^2)} = 1$ . From  $0 < \rho\sigma < \tau$ , we have  $\rho \in (0, 2)$ . Without loss of generality, we put  $\rho = 1$ . Let  $\alpha_n = \frac{1}{3n}$ ,  $\beta = \frac{2n-1}{3n}$ ,  $r_n = 1$  and  $\sigma_n^i = \frac{1}{2}$  for each  $i = 1, 2, \dots, N$ . The sequences  $\{\alpha_n\}$ ,  $\{\beta_n\}$ ,  $\{r_n\}$  and  $\{\sigma_n^i\}$  satisfy conditions (i)-(iv). Since  $T_i(x) = 0$  for  $i = 1, 2, \dots, N$  and  $Ax = -2x$  for every  $x \in \mathbb{R}$ , we have  $\bigcap_{i=1}^N \text{Fix}(T_i) = \{0\}$  and  $A$  is a bounded linear operator with  $A^* = A$  and  $\|A\| = 2$ . Obviously,  $EP(\Theta) = \{0\}$ ,  $\Lambda = \{p \in C : Ap \in EP(\Theta)\} = \{0\}$  and  $F_1, F_2, \Theta$  satisfy Assumption 2.1. For  $r_n = \mu_1 = \mu_2 = 1$ ,  $u_n = T_{r_n}^\Theta(Ax_n)$  is equivalent to

$$\Theta(u_n, v) + \langle v - u_n, u_n - Ax_n \rangle \geq 0, \quad \forall v \in K, n \in \mathbb{N}.$$

Hence, we can easily find  $u_n = -x_n \in K$ . It is not hard to compute  $A^*(T_{r_n}^\Theta - I)Ax_n = A^*(u_n - Ax_n) = -2(u_n - Ax_n) = -2x_n$  for all  $n \in \mathbb{N}$ . Hence, for  $\delta = \frac{1}{8}$ , we have

$$x_n + \frac{1}{8}A^*(T_{r_n}^\Theta - I)Ax_n = x_n + \frac{1}{8}(-2x_n) = \frac{3}{4}x_n \in C, \quad \forall n \in \mathbb{N}.$$

So, we have

$$z_n = \frac{3}{4}x_n. \tag{3.27}$$

From  $v_n = T_{\mu_2}^{F_2} z_n$  and  $y_n = T_{\mu_1}^{F_1} v_n$ , we have

$$\begin{aligned} 0 &\leq F_2(v_n, y) + \langle y - v_n, v_n - z_n \rangle \\ &= -5v_n^2 + v_n y + 4y^2 + (y - v_n)(v_n - z_n), \quad \forall y \in C, \end{aligned} \quad (3.28)$$

and

$$\begin{aligned} 0 &\leq F_1(y_n, y) + \langle y - y_n, y_n - v_n \rangle \\ &= -3y_n^2 + y_n y + 2y^2 + (y - y_n)(y_n - v_n), \quad \forall y \in C. \end{aligned} \quad (3.29)$$

From (3.28) and (3.29), we have

$$0 \leq 4y^2 + (2v_n - z_n)y - 6v_n^2 + v_n z_n, \quad \forall y \in C,$$

and

$$0 \leq 2y^2 + (2y_n - v_n)y - 4y_n^2 + y_n v_n, \quad \forall y \in C.$$

Let  $A_1(y) = 4y^2 + (2v_n - z_n)y - 6v_n^2 + v_n z_n$  and  $A_2(y) = 2y^2 + (2y_n - v_n)y - 4y_n^2 + y_n v_n$ . Then we determine the discriminants  $\Delta_1$  of  $A_1$  and  $\Delta_2$  of  $A_2$  as follows:

$$\begin{aligned} \Delta_1 &= (2v_n - z_n)^2 - 16(-6v_n^2 + v_n z_n) \\ &= z_n^2 - 20v_n z_n + 100v_n^2 \\ &= (z_n - 10v_n)^2 \end{aligned}$$

and

$$\begin{aligned} \Delta_2 &= (2y_n - v_n)^2 - 8(-4y_n^2 + y_n v_n) \\ &= v_n^2 - 12y_n v_n + 36y_n^2 \\ &= (v_n - 6y_n)^2. \end{aligned}$$

By  $A_1(y) \geq 0$  and  $A_2(y) \geq 0$ , we have  $\Delta_1 = \Delta_2 = 0$ . So, we get

$$v_n = \frac{1}{10} z_n \quad \text{and} \quad y_n = \frac{1}{6} v_n. \quad (3.30)$$

For every  $n \geq 1$ , from (3.27) and (3.30), we can rewrite (3.1) as follows:

$$\begin{cases} z_n = \frac{3}{4} x_n, \\ y_n = \frac{1}{60} z_n, \\ x_{n+1} = \frac{1}{3n} \frac{1}{2} x_n + \frac{2n-1}{3n} x_n + \left( \left( 1 - \frac{2n-1}{3n} \right) - \frac{1}{3n} \right) \frac{1}{2^N} y_n, \end{cases}$$

that is,

$$\begin{aligned} x_{n+1} &= \frac{4n-1}{6n}x_n + \frac{1}{3 \cdot 2^N}y_n \\ &= \left(\frac{2}{3} + \frac{1}{240 \cdot 2^N} - \frac{1}{6n}\right)x_n. \end{aligned}$$

Observe that for all  $n \geq 1$ ,

$$\begin{aligned} |x_{n+1} - 0| &= \left|\left(\frac{2}{3} + \frac{1}{240 \cdot 2^N} - \frac{1}{6n}\right)x_n - 0\right| \\ &\leq \frac{161}{240}|x_n - 0|. \end{aligned}$$

Hence we have  $|x_{n+1} - 0| \leq \left(\frac{161}{240}\right)^n|x_1 - 0|$  for all  $n \geq 1$ . This implies that  $\{x_n\}$  converges strongly to  $0 \in \Gamma = \bigcap_{i=1}^N \text{Fix}(T_i) \cap \Omega \cap \Lambda$ . Observe that  $\langle (\mu F - \rho V)0, x - 0 \rangle \geq 0$ ,  $x \in \Gamma$ , that is, 0 is the solution of the variational inequality  $\langle (\mu F - \rho V)x^*, x - x^* \rangle \geq 0$ ,  $x \in \Gamma$ .

**REMARK 3.1.** Theorem 3.1 improves and extends many recent corresponding main results of other authors (see [15,21,27,28]) in the following ways:

(a) The explicit iterative methods in [15,27,28] have been extended to the new iterative method (3.1) in Theorem 3.1. So, their iterative methods are some special cases of our iterative method (3.1) and some of their main results have been included in Theorem 3.1.

(b) The iterative approximating point in Theorem 3.1 is also the unique solution of the variational inequality (3.2). In fact, (3.2) is a hierarchical fixed point problem which closely relates to a convex minimization problem. In hierarchical fixed point problem (1.6), if  $S = I - (\rho V - \mu F)$ , then we can get the variational inequality (3.2). In (3.2), if  $V = 0$ , then we get the variational inequality  $\langle Fx^*, x - x^* \rangle \geq 0$ ,  $\forall x \in \Gamma = \bigcap_{i=1}^N \text{Fix}(T_i) \cap \Omega \cap \Lambda$ , which just is the variational inequality studied by Suzuki [21] extending the common set of solutions of a system of variational inequalities (1.1), a split equilibrium problem (1.5), a hierarchical fixed point problem (3.2) and the common fixed points set of a finite family of averaged mappings. So, the result of Theorem 3.1 in this paper have many useful applications.

### Competing interests

The author declares that there is no conflict of interest regarding the publication of this article.

## References

- [1] Q. H. Ansari, C. S. Lalitha and M. Mehta, *Generalized Convexity, Nonsmooth Variational Inequalities and Nonsmooth Optimization*, CRC Press, Boca Raton, 2014.
- [2] Q. H. Ansari, N. C. Wong and J. C. Yao, *The existence of nonlinear inequalities*, Appl. Math Lett. **12** (1999), 89–92.
- [3] E. Blum and W. Cettli, *From optimization and variational inequalities to equilibrium problems*, Math. Stud. **63** (1994), 123–145.
- [4] A. Bnouhachem, *A self-adaptive method for solving general mixed variational inequalities*, J. Math. Anal. Appl. **309** (2005), 136–150.
- [5] A. Bnouhachem, *A new projection and contraction method for linear variational inequalities*, J. Math. Anal. Appl. **314** (2006), 513–525.
- [6] N. Buong and L. T. Duong, *An explicit iterative algorithm for a class of variational inequalities in Hilbert spaces*, J. Optim. Theory Appl. **151** (2011), 513–524.
- [7] L. C. Ceng, Q. H. Ansari, S. Schaible and J. C. Yao, *Iterative methods for generalized equilibrium problems, systems of general generalized equilibrium problems and fixed point problems for nonexpansive mappings in Hilbert spaces*, Fixed Point Theory **12** (2011), 293–308.
- [8] L. C. Ceng, Q. H. Ansari and J. C. Yao, *Some iterative methods for finding fixed points and for solving constrained convex minimization problems*, Nonlinear Anal. **74** (2011), 5286–5302.
- [9] L. C. Ceng, C. Y. Wang and J. C. Yao, *Strong convergence theorems by a relaxed extragradient method for a general system of variational inequalities*, Math Methods Oper. Res. **67** (2008), 375–390.
- [10] L. C. Ceng and J. C. Yao, *A relaxed extragradient-like method for a generalized mixed equilibrium problem, a general system of generalized equilibria and a fixed point problem*, Nonlinear Anal. **72** (2010), 1922–1937.
- [11] Y. Censor and T. Elfving, *A multiprojection algorithm using Bregman projections in a product space*, Numer. Algorithms **8** (1994), 221–239.
- [12] S. S. Chang, H. W. J. Lee and C. K. Chan, *A new method for solving equilibrium problem, fixed point problem and variational inequality problem with application to optimization*, Nonlinear Anal. **70** (2009), 3307–3319.
- [13] F. Cianciaruso, G. Marino, L. Muglia and Y. Yao, *A hybrid projection algorithm for finding solutions of mixed equilibrium problem and variational inequality problem*, Fixed Point Theory Appl. 2010, 2010: 383740.
- [14] P. L. Combettters and S. A. Hirstoaga, *Equilibrium programming using proximal like algorithms*, Math. Program. **78** (1997), 29–41.
- [15] P. Duan and S. He, *Generalized viscosity approximation methods for nonexpansive mappings*, Fixed Point Theory Appl. 2014, 2014: 68.
- [16] K. Goebel and W. A. Kirk, *Topics in Metric Fixed Point Theory*, Stud. Adv. Math. vol.28, Cambridge University Press, Cambridge, 1990.

- [17] G. López, V. Martin and H. K. Xu, *Iterative algorithm for the multiple-sets split feasibility problem*, In: Biomedical Mathematics Promising Directions: In Imaging Therapy Planning and Inverse Problems, pp. 243–279, 2009.
- [18] P. E. Mainge and A. Moudafi, *Strong convergence of an iterative method for hierarchical fixed point problems*, Pac. J. Optim. **3** (2007), 529–538.
- [19] X. Qin, M. Shang and Y. Su, *A general iterative method for equilibrium problem and fixed point in Hilbert spaces*, Nonlinear Anal. **69** (2008), 3897–3909.
- [20] T. Suzuki, *Strong convergence of Krasnoselskii and Mann's type sequences for one parameter nonexpansive semigroups without Bochner integrals*, J. Math. Anal. Appl. **35** (2005), 227–239.
- [21] N. Suzuki, *Moudafi's viscosity approximations with Meir-Keeler contractions*, J. Math. Anal. Appl. **325** (2007), 342–352.
- [22] M. Tian, *A general iterative algorithm for nonexpansive mappings in Hilbert spaces*, Nonlinear Anal. **73** (2010), 689–694.
- [23] R. U. Verma, *Projection methods, algorithms and a new system of nonlinear variational inequalities*, Comput. Math. Appl. **41** (2001), 1025–1031.
- [24] R. U. Verma, *General convergence analysis for two-step projection methods and applications to variational problems*, Appl. Math. Lett. **18** (2005), 1286–1292.
- [25] H. K. Xu, *An iterative approach to quadratic optimization*, J. Optim. Theory and Appl. **116** (2003), 659–678.
- [26] I. Yamada, *The hybrid steepest descent method for the variational inequality problems over the intersection of fixed points sets of nonexpansive mapping*. In: D. Butnariu, Y. Censor, S. Reich (eds). *Inherently Parallel Algorithms in Feasibility and Optimization and Their Application*, pp. 473–504, North-Holland, Amsterdam, 2001.
- [27] C. Zhang and C. Yang, *A new explicit iterative algorithm for solving a class of variational inequalities over the common fixed points set of a finite family of nonexpansive mappings*, Fixed Point Theory Appl. 2014, 2014: 60.
- [28] H. Y. Zhou and P. A. Wang, *A simpler explicit iterative algorithm for a class of variational inequalities in Hilbert spaces*, J. Optim. Theory Appl. **161** (2014), 716–727.

Jae Ug Jeong  
Department of Mathematics  
Donggeui University  
Busan 47340, South Korea  
*E-mail*: jujeong@deu.ac.kr