

## A GENERIC RESEARCH ON NONLINEAR NON-CONVOLUTION TYPE SINGULAR INTEGRAL OPERATORS

GUMRAH UYSAL, VISHNU NARAYAN MISHRA\*,  
OZGE OZALP GULLER, AND ERTAN IBIKLI

ABSTRACT. In this paper, we present some general results on the pointwise convergence of the non-convolution type nonlinear singular integral operators in the following form:

$$T_\lambda(f; x) = \int_{\Omega} K_\lambda(t, x, f(t)) dt, \quad x \in \Psi, \lambda \in \Lambda,$$

where  $\Psi = \langle a, b \rangle$  and  $\Omega = \langle A, B \rangle$  stand for arbitrary closed, semi-closed or open bounded intervals in  $\mathbb{R}$  or these set notations denote  $\mathbb{R}$ , and  $\Lambda$  is a set of non-negative numbers, to the function  $f \in L_{p,w}(\Omega)$ , where  $L_{p,w}(\Omega)$  denotes the space of all measurable functions  $f$  for which  $\left| \frac{f}{w} \right|^p$  ( $1 \leq p < \infty$ ) is integrable on  $\Omega$ , and  $w : \mathbb{R} \rightarrow \mathbb{R}^+$  is a weight function satisfying some conditions.

### 1. Introduction

Approximating functions with simpler functions is one of the major problems of mathematical analysis. The main aim is to make the approximation error smaller, i.e., to make simpler functions closer to the original function. Here, simpler functions refer to the family or sequence

---

Received July 25, 2016. Revised August 29, 2016. Accepted September 11, 2016.  
2010 Mathematics Subject Classification: 41A35, 41A25, 45P05.

Key words and phrases:  $\mu$ -generalized Lebesgue point, Pointwise convergence, Rate of convergence.

\* Corresponding author.

© The Kangwon-Kyungki Mathematical Society, 2016.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

of functions equipped with good properties, such as continuity, differentiability and integrability. There are many approximation methods in the literature. In particular, approximation of integrable functions by a family of integrable functions is a classical method used by many researchers. To make this sentence clear, we outline some of the significant studies as follows:

Taberski [27] analyzed the pointwise convergence of integrable functions and the approximation properties of derivatives of integrable functions in  $L_1 \langle -\pi, \pi \rangle$ , where  $\langle -\pi, \pi \rangle$  is an arbitrary closed, semi-closed or open interval, by the following integral operator

$$(1.1) \quad L_\lambda(f; x) = \int_{-\pi}^{\pi} f(t) K_\lambda(t-x) dt, \quad x \in \langle -\pi, \pi \rangle, \quad \lambda \in \Lambda \subset \mathbb{R}_0^+,$$

where  $K_\lambda(t)$  is the kernel fulfilling appropriate conditions and  $\Lambda$  is a given set of non-negative numbers with accumulation point  $\lambda_0$ . Further, the singularity condition is satisfied by  $K_\lambda(t)$ , i.e.,  $K_\lambda(0)$  tends to infinity as  $\lambda$  tends to  $\lambda_0$ .

Following Taberski [27], Gadjiev [12] and Rydzewska [26] studied pointwise convergence of the operators of type (1.1) at different characteristic points of integrable functions. In addition, Bardaro [2] studied the rate of convergence of the linear singular integral operators in different function spaces. Further, in [3], Bardaro and Vinti obtained some approximation properties of certain non-convolution type integral operators. Later on, Alexits [1], Mamedov [15] and Esen [10] presented necessary conditions satisfied by kernel functions in order to obtain a good approximation, separately.

In the year 1983, Musielak [21] studied the convergence of convolution type nonlinear integral operators in the following form:

$$(1.2) \quad S_w(f; y) = \int_G K_w(x-y, f(x)) dx, \quad y \in G, \quad w \in \Lambda,$$

where  $G$  is a locally compact Abelian group equipped with Haar measure and  $\Lambda \neq \emptyset$  is an index set with any topology, and he extended the scope of the singularity assumption via replacing the linearity property of the integral operators of type (1.1) by an assumption of Lipschitz condition for  $K_w$  with respect to second variable. In [22], Musielak stepped

up his previous analysis by presenting the significant approximation results for the operators of type (1.2) in generalized Orlicz spaces. After these pioneering studies, Swiderski and Wachnicki [23] investigated the pointwise convergence of the operators of type (1.2) at  $p$ -Lebesgue points of functions  $f \in L_p(-\pi, \pi)$ , where  $1 \leq p < \infty$ . The convergence properties of specific types of nonlinear integral operators were investigated, for example, the pointwise convergence of nonlinear Mellin type convolution operators at Lebesgue points was studied by Bardaro and Mantellini [6]. In [14], Karsli investigated the approximation properties of non-convolution type nonlinear integral operators and established the rate of pointwise convergence. Some further approximation results concerning approximation by non-convolution type singular integral operators can be found in [10, 11]. Based on the results given by Esen [10], Guller *et al.* [13] presented some approximation theorems. For some advanced results concerning the convergence of several types of nonlinear singular integral operators in modular function spaces, we refer the reader to the monograph by Bardaro *et al.* [5] and the articles by Bardaro and Vinti [3, 4]. Recently, some weighted pointwise approximation results for the nonlinear counterpart of the operator of type (1.1) were presented in [28] using the approximation method presented by Esen [10]. For some studies concerning the applications of approximation of functions by linear and nonlinear operators, we refer the reader to see the works [8, 9, 16-20, 24].

In this paper, we prove the convergence of non-convolution type nonlinear singular integral operators designated by

$$(1.3) \quad T_\lambda(f; x) = \int_{\Omega} K_\lambda(t, x, f(t)) dt, \quad x \in \Psi, \quad \lambda \in \Lambda,$$

where  $\Psi = \langle a, b \rangle$  and  $\Omega = \langle A, B \rangle$  stand for arbitrary closed, semi-closed or open bounded intervals in  $\mathbb{R}$  or the indicated set notations represent  $\mathbb{R}$ , and  $\Lambda$  is a set of non-negative numbers, to the function  $f \in L_{p,w}(\Omega)$ , where  $L_{p,w}(\Omega)$  is the space of all measurable functions  $f$  for which  $\left|\frac{f}{w}\right|^p$  ( $1 \leq p < \infty$ ) is integrable on  $\Omega$ , and  $w : \mathbb{R} \rightarrow \mathbb{R}^+$  is a weight function, at a common  $\mu$ -generalized Lebesgue point of the functions  $\frac{f}{w}$  and  $w$ . In this work, the weighted approximation method presented by Esen [10] is harnessed.

The paper is organized as follows: In Section 2, we introduce basic definitions. In Section 3, we prove the existence of the operators type

(1.3). In Section 4, we present two theorems concerning the pointwise convergence of  $T_\lambda(f; x)$  to the value  $f(x_0)$  as  $(x, \lambda)$  tends to  $(x_0, \lambda_0)$ , where  $x_0$  is a  $\mu$ -generalized Lebesgue point of  $f \in L_p(\Omega)$ . In Section 5, we present two theorems concerning the pointwise convergence of  $T_\lambda(f; x)$  to the value  $f(x_0)$  as  $(x, \lambda)$  tends to  $(x_0, \lambda_0)$ , where  $x_0$  is a common  $\mu$ -generalized Lebesgue point of  $f \in L_{p,w}(\Omega)$  and  $w \in L_p(\Omega)$ . In Section 6, we give two theorems concerning the rate of pointwise convergence.

## 2. Preliminaries

DEFINITION 2.1. A point  $x_0 \in \Omega$  is called a  $\mu$ -generalized Lebesgue point of the function  $f \in L_p(\Omega)$  if

$$\lim_{h \rightarrow 0} \left( \frac{1}{\mu(h)} \int_0^h |f(t + x_0) - f(x_0)|^p dt \right)^{\frac{1}{p}} = 0, \quad 1 \leq p < \infty,$$

where  $\mu : \mathbb{R} \rightarrow \mathbb{R}$  is an increasing and absolutely continuous function on  $[0, B - A]$  with  $\mu(0) = 0$  [26].

Using the kernel properties given in [11, 14], the following definition is obtained:

DEFINITION 2.2. (*Class  $A_w$* ) Let  $\Lambda$  be a set of non-negative set of numbers with accumulation point  $\lambda_0$ . Suppose that for every fixed  $x \in \Psi$  there exists a point  $x_0 \in \Omega$ . We say that the family of functions  $(K_\lambda)_{\lambda \in \Lambda}$ ,  $K_\lambda : \mathbb{R} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  belongs to *class  $A_w$*  if the following conditions are satisfied:

- $K_\lambda(t, x, 0) = 0$ , for every  $t, x \in \mathbb{R}$  and  $\lambda \in \Lambda$ .
- The function  $L_\lambda : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}_0^+$  which is integrable for every  $t, x \in \mathbb{R}$  and for each  $\lambda \in \Lambda$  exists such that the inequality

$$|K_\lambda(t, x, u) - K_\lambda(t, x, v)| \leq L_\lambda(t, x) |u - v|$$

holds for every  $t, x, u, v \in \mathbb{R}$  and for each  $\lambda \in \Lambda$ .

- $\lim_{(x, \lambda) \rightarrow (x_0, \lambda_0)} \left| \int_{\mathbb{R}} K_\lambda(t, x, u) dt - u \right| = 0$ , for every  $u \in \mathbb{R}$ .
- $\lim_{(x, \lambda) \rightarrow (x_0, \lambda_0)} \left( \sup_{t \in \mathbb{R} \setminus \langle x_0 - \xi, x_0 + \xi \rangle} L_\lambda(t, x) \right) = 0$ , for every  $\xi > 0$ .
- $\lim_{(x, \lambda) \rightarrow (x_0, \lambda_0)} \left( \int_{\mathbb{R} \setminus \langle x_0 - \xi, x_0 + \xi \rangle} L_\lambda(t, x) dt \right) = 0$ , for every  $\xi > 0$ .

- f. There exists  $\delta_0 > 0$  such that  $L_\lambda(t, x)$  is non-decreasing as a function of  $t$  on  $\langle x_0 - \delta_0, x_0 \rangle$  and non-increasing as a function of  $t$  on  $\langle x_0, x_0 + \delta_0 \rangle$ , for each fixed  $\lambda \in \Lambda$ .
- g.  $\left\| \frac{w(\cdot)}{w(x)} L_\lambda(\cdot, x) \right\|_{L_1(\mathbb{R})} \leq M < \infty$ , where  $M$  is a constant whose value does not depend on  $x \in \Psi$  and  $\lambda \in \Lambda$ .
- h.  $\|\alpha L_\lambda\|_{L_q(\mathbb{R} \times \mathbb{R})} \leq N < \infty$ , where  $N$  is a constant whose value does not depend on  $\lambda \in \Lambda$ ,  $1 < q < \infty$  and  $\alpha(t, x) = \frac{w(t)}{w(x)}$ .

Throughout this paper  $K_\lambda$  belongs to class  $A_w$ .

### 3. Existence of the Operators

**THEOREM 3.1.** *Let  $f \in L_{1,w}(\Omega)$ . Then,  $T_\lambda \in L_{1,w}(\Omega)$  and the following inequality*

$$\|T_\lambda(f; x)\|_{L_{1,w}(\Omega)} \leq \left\| \frac{w(\cdot)}{w(x)} L_\lambda(\cdot, x) \right\|_{L_1(\mathbb{R})} \|f\|_{L_{1,w}(\Omega)}$$

holds for every  $x \in \Psi$  and  $\lambda \in \Lambda$ .

*Proof.* Assume that  $\Psi = \langle a, b \rangle$  and  $\Omega = \langle A, B \rangle$  stand for arbitrary closed, semi-closed or open bounded intervals in  $\mathbb{R}$ .

Using conditions (a), (b) of class  $A_w$  and Fubini's Theorem (see, e.g., [7]), we may write

$$\begin{aligned} \|T_\lambda(f; x)\|_{L_{1,w}(\Omega)} &= \int_{\Omega} \frac{1}{w(x)} \left| \int_{\Omega} K_\lambda(t, x, f(t)) dt \right| dx \\ &\leq \int_{\Omega} \frac{1}{w(x)} \int_{\Omega} \left| f(t) \frac{w(t)}{w(t)} L_\lambda(t, x) \right| dt dx \\ &\leq \int_{\Omega} \left| \frac{f(t)}{w(t)} \right| \left[ \int_{\mathbb{R}} \frac{w(t)}{w(x)} L_\lambda(t, x) dx \right] dt \\ &\leq M \|f\|_{L_{1,w}(\Omega)}. \end{aligned}$$

The proof is completed for the indicated case. One may prove the assertion for the case  $\Psi = \Omega = \mathbb{R}$  by following similar steps. Thus the proof is completed. □

**THEOREM 3.2.** *Let  $f \in L_{p,w}(\Omega)$ . Then,  $T_\lambda \in L_{q,w}(\Omega)$  and the following inequality*

$$\|T_\lambda(f; x)\|_{L_{q,w}(\Omega)} \leq \|\alpha L_\lambda\|_{L_q(\mathbb{R} \times \mathbb{R})} \|f\|_{L_{p,w}(\Omega)}$$

*holds for every  $x \in \Psi$  and  $\lambda \in \Lambda$ . Here,  $\frac{1}{q} + \frac{1}{p} = 1$  provided that  $1 < p, q < \infty$ .*

*Proof.* Suppose that  $\Psi = \langle a, b \rangle$  and  $\Omega = \langle A, B \rangle$  stand for arbitrary closed, semi-closed or open bounded intervals in  $\mathbb{R}$ .

Let us define a new function by

$$g(t) := \begin{cases} f(t), & \text{if } t \in \Omega, \\ 0, & \text{if } t \in \mathbb{R} \setminus \Omega. \end{cases}$$

It is easy to see that the following inequality

$$\begin{aligned} \|T_\lambda(f; x)\|_{L_q(\Omega)} &= \|T_\lambda(g; x)\|_{L_q(\Omega)} \\ &= \left( \int_{\Omega} \frac{1}{w^q(x)} \left| \int_{\mathbb{R}} K_\lambda(t, x, g(t)) dt \right|^q dx \right)^{\frac{1}{q}} \\ &\leq \left( \int_{\Omega} \frac{1}{w^q(x)} \left( \int_{\mathbb{R}} |g(t) L_\lambda(t, x)| dt \right)^q dx \right)^{\frac{1}{q}} \end{aligned}$$

holds. Now, applying Hölder's inequality (see, e.g., [25]) to the above inequality, we obtain

$$\begin{aligned} &\|T_\lambda(f; x)\|_{L_q(\Omega)} \\ &\leq \left( \int_{\Omega} \frac{1}{w^q(x)} \left( \left( \int_{\mathbb{R}} \left| \frac{g(t)}{w(t)} \right|^p dt \right)^{\frac{1}{p}} \left( \int_{\mathbb{R}} |w(t) L_\lambda(t, x)|^q dt \right)^{\frac{1}{q}} \right)^q dx \right)^{\frac{1}{q}} \\ &\leq \|f\|_{L_{p,w}(\Omega)} \left( \int_{\Omega} \left( \int_{\mathbb{R}} \left| \frac{w(t)}{w(x)} L_\lambda(t, x) \right|^q dt \right) dx \right)^{\frac{1}{q}} \\ &\leq \|f\|_{L_{p,w}(\Omega)} \left( \int_{\mathbb{R}} \left( \int_{\mathbb{R}} \left| \frac{w(t)}{w(x)} L_\lambda(t, x) \right|^q dt \right) dx \right)^{\frac{1}{q}} \\ &\leq \|f\|_{L_{p,w}(\Omega)} \|\alpha L_\lambda\|_{L_q(\mathbb{R} \times \mathbb{R})} \\ &\leq N \|f\|_{L_{p,w}(\Omega)}. \end{aligned}$$

Hence, the proof is completed for the indicated case. One may prove the assertion for the case  $\Psi = \Omega = \mathbb{R}$  by following similar steps. Thus the proof is completed.  $\square$

#### 4. Convergence at Characteristic Points

Throughout this section we assume that  $w(t) = 1$ . Also, in the following theorem,  $\Psi = \langle a, b \rangle$  and  $\Omega = \langle A, B \rangle$  denote arbitrary closed, semi-closed or open bounded intervals in  $\mathbb{R}$ .

**THEOREM 4.1.** *Let  $x_0 \in \Omega$  be a  $\mu$ -generalized Lebesgue point of function  $f \in L_p(\Omega)$ . Then,*

$$\lim_{(x,\lambda) \rightarrow (x_0,\lambda_0)} |T_\lambda(f; x) - f(x_0)| = 0$$

on any set  $Z$  on which the function

$$\int_{x_0-\delta}^{x_0+\delta} L_\lambda(t, x) |\{\mu(|x_0 - t|)\}'_t| dt$$

is bounded as  $(x, \lambda)$  tends to  $(x_0, \lambda_0)$ .

*Proof.* We prove the theorem for the case  $1 < p < \infty$ . The proof of the case  $p = 1$  is quite similar to the proof of the case  $1 < p < \infty$  and it is omitted.

Let  $\langle x_0 - \delta, x_0 + \delta \rangle \subset \Omega$ , for fixed  $\delta > 0$ , where  $x_0$  is a  $\mu$ -generalized Lebesgue point of function  $f \in L_p(\Omega)$ . Set  $E(x, \lambda) := |T_\lambda(f; x) - f(x_0)|$ . According to condition (c), we shall write

$$\begin{aligned} E(x, \lambda) &= \left| \int_{\Omega} K_\lambda(t, x, f(t)) dt - f(x_0) \right| \\ &\leq \int_{\Omega} |f(t) - f(x_0)| L_\lambda(t, x) dt + \left| \int_{\mathbb{R}} K_\lambda(t, x, f(x_0)) dt - f(x_0) \right| \\ &\quad + |f(x_0)| \int_{\mathbb{R} \setminus \Omega} L_\lambda(t, x) dt \\ &= I_1 + I_2 + I_3. \end{aligned}$$

By condition (e),  $I_3 \rightarrow 0$  as  $(x, \lambda)$  tends to  $(x_0, \lambda_0)$ . Using well known inequality  $(m + n)^p \leq 2^p(m^p + n^p)$ , where  $m$  and  $n$  are positive real numbers (see, e.g., [25]), we have

$$\begin{aligned} & (I_1 + I_2)^p \\ & \leq 2^p(I_1^p + I_2^p) \\ & = 2^p \left( \int_{\Omega} |f(t) - f(x_0)| L_{\lambda}(t, x) dt \right)^p + 2^p \left| \int_{\mathbb{R}} K_{\lambda}(t, x, f(x_0)) dt - f(x_0) \right|^p \\ & = 2^p(I_4 + I_5). \end{aligned}$$

It is clear that  $I_5 \rightarrow 0$  as  $(x, \lambda)$  tends to  $(x_0, \lambda_0)$  in view of condition (c).

Applying Hölder's inequality to the integral  $I_4$  (see, e.g., [25]), we have

$$\begin{aligned} I_4 & = \left( \int_{\Omega} |f(t) - f(x_0)| L_{\lambda}(t, x) dt \right)^p \\ & \leq \int_{\Omega} |f(t) - f(x_0)|^p L_{\lambda}(t, x) dt \left( \int_{\mathbb{R}} L_{\lambda}(t, x) dt \right)^{\frac{p}{q}} \\ & = I_{41} \times I_{42}. \end{aligned}$$

Using condition (g), we see that  $I_{42} \leq M^{\frac{p}{q}} < \infty$ .

Now, let us find an appropriate inequality for the integral  $I_{41}$ . Splitting  $I_{41}$  into two parts, we have that

$$\begin{aligned} & I_{41} \\ & = \int_{\Omega} |f(t) - f(x_0)|^p L_{\lambda}(t, x) dt \\ & = \int_{\langle x_0 - \delta, x_0 + \delta \rangle} |f(t) - f(x_0)|^p L_{\lambda}(t, x) dt + \int_{\Omega \setminus \langle x_0 - \delta, x_0 + \delta \rangle} |f(t) - f(x_0)|^p L_{\lambda}(t, x) dt \\ & = I_{411} + I_{412}. \end{aligned}$$



For the integral  $I_{412}$ , we may write

$$\begin{aligned} I_{412} &= \int_{\Omega \setminus \langle x_0 - \delta, x_0 + \delta \rangle} |f(t) - f(x_0)|^p L_\lambda(t, x) dt \\ &\leq 2^p \left( \sup_{t \in \Omega \setminus \langle x_0 - \delta, x_0 + \delta \rangle} L_\lambda(t, x) \int_{\Omega \setminus \langle x_0 - \delta, x_0 + \delta \rangle} |f(t) - f(x_0)|^p dt \right) \\ &\leq 2^p \left( \sup_{t \in \Omega \setminus \langle x_0 - \delta, x_0 + \delta \rangle} L_\lambda(t, x) \left[ \|f\|_{L^p(\Omega)}^p + |f(x_0)|^p |B - A| \right] \right). \end{aligned}$$

Hence, by condition (d),  $I_{412} \rightarrow 0$  as  $(x, \lambda)$  tends to  $(x_0, \lambda_0)$ .

For the integral  $I_{411}$ , we may write

$$\begin{aligned} I_{411} &= \int_{\langle x_0 - \delta, x_0 + \delta \rangle} |f(t) - f(x_0)|^p L_\lambda(t, x) dt \\ &= \left\{ \int_{x_0 - \delta}^{x_0} + \int_{x_0}^{x_0 + \delta} \right\} |f(t) - f(x_0)|^p L_\lambda(t, x) dt \\ &= I_{4111} + I_{4112}. \end{aligned}$$

Now, by definition of  $\mu$ -generalized Lebesgue point, for every  $\varepsilon > 0$  there exists  $\delta > 0$  such that the inequality

$$\int_{x_0 - h}^{x_0} |f(t) - f(x_0)|^p dt < \varepsilon^p \mu(h)$$

holds for every  $0 < h \leq \delta < \delta_0$ . We define the new function as

$$G(t) = \int_t^{x_0} |f(u) - f(x_0)|^p du.$$

Then, for every  $t$  satisfying  $0 < x_0 - t \leq \delta$ , we have

$$|G(t)| \leq \varepsilon^p \mu(x_0 - t).$$

We can write the integral  $I_{4111}$  as

$$\begin{aligned} |I_{4111}| &= \left| \int_{x_0-\delta}^{x_0} |f(t) - f(x_0)|^p L_\lambda(t, x) dt \right| \\ &= \left| (LS) \int_{x_0-\delta}^{x_0} L_\lambda(t, x) d[-G(t)] \right|, \end{aligned}$$

where  $(LS)$  denotes Lebesgue-Stieltjes integral.

Applying integration by parts method to the Lebesgue-Stieltjes integral, we have

$$\begin{aligned} |I_{4111}| &\leq \int_{x_0-\delta}^{x_0} |G(t)| |d_t L_\lambda(t, x)| + |G(x_0 - \delta)| L_\lambda(x_0 - \delta, x) \\ &\leq \varepsilon^p \int_{x_0-\delta}^{x_0} \mu(x_0 - t) |d_t L_\lambda(t, x)| + \varepsilon^p \mu(\delta) L_\lambda(x_0 - \delta, x) \\ &= \varepsilon^p \int_{x_0-\delta}^{x_0} L_\lambda(t, x) |\{\mu(x_0 - t)\}'_t| dt. \end{aligned}$$

Using similar technic, one may obtain the following inequality for the integral  $I_{4112}$ :

$$|I_{4112}| \leq \varepsilon^p \int_{x_0}^{x_0+\delta} L_\lambda(t, x) |\{\mu(t - x_0)\}'_t| dt.$$

Combining above inequalities, we get

$$|I_{411}| \leq \varepsilon^p \int_{x_0-\delta}^{x_0+\delta} L_\lambda(t, x) |\{\mu(|x_0 - t|)\}'_t| dt.$$

Since the expression on the right hand side of the inequality is bounded by the hypothesis, we get the desired result for the case  $1 < p < \infty$ , i.e., the proof is completed.  $\square$

In the following theorem,  $\Psi = \Omega = \mathbb{R}$ .

**THEOREM 4.2.** *Let  $x_0 \in \mathbb{R}$  be a  $\mu$ -generalized Lebesgue point of function  $f \in L_p(\mathbb{R})$ . Then,*

$$\lim_{(x,\lambda) \rightarrow (x_0,\lambda_0)} |T_\lambda(f; x) - f(x_0)| = 0$$

on any set  $Z$  on which the function

$$\int_{x_0-\delta}^{x_0+\delta} L_\lambda(t, x) |\{\mu(|x_0 - t|)\}'_t| dt$$

is bounded as  $(x, \lambda)$  tends to  $(x_0, \lambda_0)$ .

*Proof.* The proof of this theorem is quite similar to preceding one and we skip it. □

### 5. Weighted Pointwise Approximation

Throughout this section we assume that  $f \notin L_p(\Omega)$ , where  $1 \leq p < \infty$ . Also, in the following theorem,  $\Psi = \langle a, b \rangle$  and  $\Omega = \langle A, B \rangle$  denote arbitrary closed, semi-closed or open bounded intervals in  $\mathbb{R}$ .

**THEOREM 5.1.** *Let  $w$  and  $L_\lambda$  be differentiable functions almost everywhere on  $\mathbb{R}$  with respect to variable  $t$  such that the following inequality*

$$\frac{d}{dt}w(t)\frac{d}{dt}L_\lambda(t, x) > 0, \text{ for any fixed } x \in \mathbb{R}$$

holds. If  $x_0 \in \Omega$  is a common  $\mu$ -generalized Lebesgue point of functions  $f \in L_{p,w}(\Omega)$  and  $w \in L_p(\Omega)$  then

$$\lim_{(x,\lambda) \rightarrow (x_0,\lambda_0)} |T_\lambda(f; x) - f(x_0)| = 0$$

on any set  $Z$  on which the function

$$\int_{x_0-\delta}^{x_0+\delta} w(t)L_\lambda(t, x) |\{\mu(|x_0 - t|)\}'_t| dt$$

is bounded as  $(x, \lambda)$  tends to  $(x_0, \lambda_0)$ .

*Proof.* We prove the theorem for the case  $1 < p < \infty$ . The proof of the case  $p = 1$  is quite similar to the proof of the case  $1 < p < \infty$  and it is omitted.

Let  $\langle x_0 - \delta, x_0 + \delta \rangle \subset \Omega$ , for fixed  $\delta > 0$ , where  $x_0$  is a common  $\mu$ -generalized Lebesgue point of functions  $f \in L_{p,w}(\Omega)$  and  $w \in L_p(\Omega)$ . Using Theorem 4.1, we have

$$\left| \int_{\Omega} K_{\lambda}(t, x, f(t)) dt - f(x_0) \right|$$

$$\leq \int_{\Omega} \left| \frac{f(t)}{w(t)} - \frac{f(x_0)}{w(x_0)} \right| w(t) L_{\lambda}(t, x) dt + \left| \int_{\Omega} K_{\lambda}(t, x, \frac{f(x_0)}{w(x_0)} w(t)) dt - f(x_0) \right|.$$

It is easy to see that the following inequality

$$\left| \int_{\Omega} K_{\lambda}(t, x, f(t)) dt - f(x_0) \right|^p \leq 2^p \left( \int_{\Omega} \left| \frac{f(t)}{w(t)} - \frac{f(x_0)}{w(x_0)} \right| w(t) L_{\lambda}(t, x) dt \right)^p$$

$$+ 2^p \left| \int_{\Omega} K_{\lambda}(t, x, \frac{f(x_0)}{w(x_0)} w(t)) dt - f(x_0) \right|^p$$

$$= 2^p (I_1 + I_2)$$

holds. By Theorem 4.1,  $I_2 \rightarrow 0$  as  $(x, \lambda)$  tends to  $(x_0, \lambda_0)$ . Applying Hölder's inequality to the integral  $I_1$ , we have

$$I_1 \leq \int_{\Omega} \left| \frac{f(t)}{w(t)} - \frac{f(x_0)}{w(x_0)} \right|^p w(t) L_{\lambda}(t, x) dt \left( \int_{\Omega} w(t) L_{\lambda}(t, x) dt \right)^{\frac{p}{q}}$$

$$= I_{11} \times I_{12}.$$

In view of condition (g),  $I_{12} \rightarrow (w(x_0)M)^{\frac{p}{q}} < \infty$  as  $(x, \lambda)$  tends to  $(x_0, \lambda_0)$ . Let us find an appropriate inequality for the integral  $I_{11}$ . Splitting the integral  $I_{11}$  into two parts, we have

$$\begin{aligned} I_{11} &= \int_{\Omega} \left| \frac{f(t)}{w(t)} - \frac{f(x_0)}{w(x_0)} \right|^p w(t) L_{\lambda}(t, x) dt \\ &= \int_{\langle x_0 - \delta, x_0 + \delta \rangle} \left| \frac{f(t)}{w(t)} - \frac{f(x_0)}{w(x_0)} \right|^p w(t) L_{\lambda}(t, x) dt \\ &\quad + \int_{\Omega \setminus \langle x_0 - \delta, x_0 + \delta \rangle} \left| \frac{f(t)}{w(t)} - \frac{f(x_0)}{w(x_0)} \right|^p w(t) L_{\lambda}(t, x) dt \\ &= I_{111} + I_{112}. \end{aligned}$$

For the integral  $I_{112}$ , we may write

$$\begin{aligned} I_{112} &= \int_{\Omega \setminus \langle x_0 - \delta, x_0 + \delta \rangle} \left| \frac{f(t)}{w(t)} - \frac{f(x_0)}{w(x_0)} \right|^p w(t) L_{\lambda}(t, x) dt \\ &\leq 2^p \sup_{t \in \Omega \setminus \langle x_0 - \delta, x_0 + \delta \rangle} L_{\lambda}(t, x) \sup_{t \in \Omega \setminus \langle x_0 - \delta, x_0 + \delta \rangle} w(t) \int_{\Omega \setminus \langle x_0 - \delta, x_0 + \delta \rangle} \left| \frac{f(t)}{w(t)} - \frac{f(x_0)}{w(x_0)} \right|^p dt \\ &\leq 2^p \left( \sup_{t \in \Omega \setminus \langle x_0 - \delta, x_0 + \delta \rangle} L_{\lambda}(t, x) \sup_{t \in \Omega \setminus \langle x_0 - \delta, x_0 + \delta \rangle} w(t) \left[ \|f\|_{L^p, w(\Omega)}^p + \left| \frac{f(x_0)}{w(x_0)} \right|^p |B - A| \right] \right). \end{aligned}$$

By the hypothesis, monotonicity behavior of  $w$  is similar to  $L_{\lambda}$ . Therefore,  $w(t)$  is bounded on  $\mathbb{R} \setminus \Omega$ . Hence by condition (d),  $I_{112} \rightarrow 0$  as  $(x, \lambda)$  tends to  $(x_0, \lambda_0)$ .

Write

$$\begin{aligned} I_{111} &= \int_{\langle x_0 - \delta, x_0 + \delta \rangle} \left| \frac{f(t)}{w(t)} - \frac{f(x_0)}{w(x_0)} \right|^p w(t) K_{\lambda}(t, x) dt \\ &= \left\{ \int_{x_0 - \delta}^{x_0} + \int_{x_0}^{x_0 + \delta} \right\} \left| \frac{f(t)}{w(t)} - \frac{f(x_0)}{w(x_0)} \right|^p w(t) K_{\lambda}(t, x) dt \\ &= I_{1111} + I_{1112}. \end{aligned}$$

Now, by definition of  $\mu$ -generalized Lebesgue point, for every  $\varepsilon > 0$  there exists  $\delta > 0$  such that the inequality

$$\int_{x_0-h}^{x_0} \left| \frac{f(t)}{w(t)} - \frac{f(x_0)}{w(x_0)} \right|^p dt < \varepsilon^p \mu(h)$$

holds for every  $0 < h \leq \delta < \delta_0$ . We define the new function as

$$\tilde{G}(t) = \int_t^{x_0} \left| \frac{f(u)}{w(u)} - \frac{f(x_0)}{w(x_0)} \right|^p du.$$

Then, for every  $t$  satisfying  $0 < x_0 - t \leq \delta$  we have

$$|\tilde{G}(t)| \leq \varepsilon^p \mu(x_0 - t).$$

Hence

$$\begin{aligned} |I_{1111}| &= \left| \int_{x_0-\delta}^{x_0} \left| \frac{f(u)}{w(u)} - \frac{f(x_0)}{w(x_0)} \right|^p w(t) L_\lambda(t, x) dt \right| \\ &= \left| (LS) \int_{x_0-\delta}^{x_0} w(t) L_\lambda(t, x) d[-\tilde{G}(t)] \right|, \end{aligned}$$

where  $(LS)$  denotes Lebesgue-Stieltjes integral.

Applying integration by parts method to the Lebesgue-Stieltjes integral, we have

$$\begin{aligned} I_{1111} &= \int_{x_0-\delta}^{x_0} |\tilde{G}(t)| d_t [w(t) L_\lambda(t, x)] + \tilde{G}(x_0 - \delta) w(x_0 - \delta) L_\lambda(x_0 - \delta, x) \\ &\leq \varepsilon^p \int_{x_0-\delta}^{x_0} \mu(x_0 - t) |d_t [w(t) L_\lambda(t, x)]| + \varepsilon^p \mu(\delta) w(x_0 - \delta) L_\lambda(x_0 - \delta, x) \\ &= \varepsilon^p \int_{x_0-\delta}^{x_0} w(t) L_\lambda(t, x) |\{\mu(x_0 - t)\}'_t| dt. \end{aligned}$$

Using similar technic, one may obtain the following inequality for the integral  $I_{1112}$ :

$$|I_{1112}| \leq \varepsilon^p \int_{x_0}^{x_0+\delta} w(t)L_\lambda(t, x) |\{\mu(t - x_0)\}'_t| dt.$$

Combining above inequalities, we get

$$|I_{1111}| \leq \varepsilon^p \int_{x_0-\delta}^{x_0+\delta} w(t)L_\lambda(t, x) |\{\mu(|x_0 - t|)\}'_t| dt.$$

Since the expression on the right hand side of the inequality is bounded by the hypothesis, we get the desired result for the case  $1 < p < \infty$ , i.e., the proof is completed. □

In the following theorem,  $\Psi = \Omega = \mathbb{R}$ .

**THEOREM 5.2.** *Let  $w$  and  $L_\lambda$  be differentiable functions almost everywhere on  $\mathbb{R}$  with respect to variable  $t$  such that the following inequality*

$$\frac{d}{dt}w(t)\frac{d}{dt}L_\lambda(t, x) > 0, \text{ for any fixed } x \in \mathbb{R}$$

*holds. If  $x_0 \in \mathbb{R}$  is a common  $\mu$ -generalized Lebesgue point of functions  $f \in L_{p,w}(\mathbb{R})$  and  $w \in L_p(\mathbb{R})$  then*

$$\lim_{(x,\lambda) \rightarrow (x_0,\lambda_0)} |T_\lambda(f; x) - f(x_0)| = 0$$

*on any set  $Z$  on which the function*

$$\int_{x_0-\delta}^{x_0+\delta} w(t)L_\lambda(t, x) |\{\mu(|x_0 - t|)\}'_t| dt$$

*is bounded as  $(x, \lambda)$  tends to  $(x_0, \lambda_0)$ .*

*Proof.* Following the same strategy as in Theorem 5.1, we have

$$\left| \int_{\mathbb{R}} K_\lambda(t, x, f(t))dt - f(x_0) \right|^p \leq$$

$$\begin{aligned}
 & 2^{2p} (w(x)M)^{\frac{p}{q}} \sup_{t \in \mathbb{R} \setminus \langle x_0 - \delta, x_0 + \delta \rangle} L_\lambda(t, x) \sup_{t \in \mathbb{R} \setminus \langle x_0 - \delta, x_0 + \delta \rangle} w(t) \|f\|_{L_{p,w}(\mathbb{R})}^p \\
 & + 2^{2p} (w(x)M)^{\frac{p}{q}} \sup_{t \in \mathbb{R} \setminus \langle x_0 - \delta, x_0 + \delta \rangle} w(t) \left| \frac{f(x_0)}{w(x_0)} \right|^p \int_{\mathbb{R} \setminus \langle x_0 - \delta, x_0 + \delta \rangle} L_\lambda(t, x) dt \\
 & + 2^p \varepsilon^p (w(x)M)^{\frac{p}{q}} \int_{x_0 - \delta}^{x_0 + \delta} w(t) L_\lambda(t, x) |\{\mu(|x_0 - t|)\}'_t| dt \\
 & + 2^p \left| \int_{\mathbb{R}} K_\lambda(t, x, \frac{f(x_0)}{w(x_0)} w(t)) dt - f(x_0) \right|^p.
 \end{aligned}$$

The rest of the proof is clear by the hypotheses. □

### 6. Rate of Weighted Pointwise Convergence

In this section, two theorems concerning rate of pointwise convergence will be given.

**THEOREM 6.1.** *Suppose that the hypothesis of Theorem 5.1 is satisfied. Let*

$$\Delta(x, \lambda, \delta) = \int_{x_0 - \delta}^{x_0 + \delta} w(t) L_\lambda(t, x) |\{\mu(|x_0 - t|)\}'_t| dt,$$

where  $0 < \delta < \delta_0$ , for a fixed (and finite!) positive number  $\delta_0$ , and the following conditions are satisfied:

- (i)  $\Delta(x, \lambda, \delta) \rightarrow 0$  as  $(x, \lambda) \rightarrow (x_0, \lambda_0)$ , for some  $\delta > 0$ .
- (ii) For every  $\xi > 0$ ,

$$\sup_{t \in \mathbb{R} \setminus \langle x_0 - \xi, x_0 + \xi \rangle} L_\lambda(t, x) = o(\Delta(x, \lambda, \delta))$$

as  $(x, \lambda) \rightarrow (x_0, \lambda_0)$

- (iii)  $\left| \int_{\Omega} K_\lambda(t, x, \frac{f(x_0)}{\varphi(x_0)} \varphi(t)) dt - f(x_0) \right| = o(\Delta(x, \lambda, \delta)).$

Then, at each common Lebesgue point of functions  $f \in L_{p,\varphi}(\Omega)$  and  $\varphi \in L_p(\Omega)$  we have



$$|T_\lambda(f; x) - f(x_0)|^p = o(\Delta(x, \lambda, \delta))$$

as  $(x, \lambda) \rightarrow (x_0, \lambda_0)$ .

*Proof.* By the hypotheses of Theorem 5.1, we have

$$\begin{aligned} & \left| \int_{\Omega} K_\lambda(t, x, f(t)) dt - f(x_0) \right|^p \leq \\ & 2^{2p} (w(x)M)^{\frac{p}{q}} \sup_{t \in \mathbb{R} \setminus \langle x_0 - \delta, x_0 + \delta \rangle} L_\lambda(t, x) \sup_{t \in \mathbb{R} \setminus \langle x_0 - \delta, x_0 + \delta \rangle} w(t) \|f\|_{L^p, w(\Omega)}^p \\ & + 2^{2p} (w(x)M)^{\frac{p}{q}} \sup_{t \in \mathbb{R} \setminus \langle x_0 - \delta, x_0 + \delta \rangle} w(t) \sup_{t \in \mathbb{R} \setminus \langle x_0 - \delta, x_0 + \delta \rangle} L_\lambda(t, x) \left| \frac{f(x_0)}{w(x_0)} \right|^p |B - A| \\ & + 2^p \varepsilon^p (w(x)M)^{\frac{p}{q}} \int_{x_0 - \delta}^{x_0 + \delta} w(t) L_\lambda(t, x) |\{\mu(|x_0 - t|)\}'_t| dt \\ & + 2^p \left| \int_{\Omega} K_\lambda(t, x, \frac{f(x_0)}{w(x_0)} w(t)) dt - f(x_0) \right|^p. \end{aligned}$$

The remaining part is obvious by the conditions (i) – (iii). The proof of the case  $p = 1$  is quite similar. Thus the proof is completed.  $\square$

**THEOREM 6.2.** *Suppose that the hypothesis of Theorem 5.2 is satisfied. Let*

$$\Delta(x, \lambda, \delta) = \int_{x_0 - \delta}^{x_0 + \delta} w(t) L_\lambda(t, x) |\{\mu(|x_0 - t|)\}'_t| dt,$$

where  $0 < \delta < \delta_0$ , for a fixed (and finite!) positive number  $\delta_0$ , and the following conditions are satisfied:

- (i)  $\Delta(x, \lambda, \delta) \rightarrow 0$  as  $(x, \lambda) \rightarrow (x_0, \lambda_0)$ , for some  $\delta > 0$ .
- (ii) For every  $\xi > 0$ ,

$$\sup_{t \in \mathbb{R} \setminus \langle x_0 - \xi, x_0 + \xi \rangle} L_\lambda(t, x) = o(\Delta(x, \lambda, \delta))$$

as  $(x, \lambda) \rightarrow (x_0, \lambda_0)$ .

(iii) For every  $\xi > 0$ ,

$$\int_{\mathbb{R} \setminus \langle x_0 - \xi, x_0 + \xi \rangle} L_\lambda(t, x) dt = o(\Delta(x, \lambda, \delta))$$

as  $(x, \lambda) \rightarrow (x_0, \lambda_0)$ .

$$(iv) \left| \int_{\mathbb{R}} K_\lambda(t, x, \frac{f(x_0)}{\varphi(x_0)} \varphi(t)) dt - f(x_0) \right| = o(\Delta(x, \lambda, \delta)).$$

Then, at each common Lebesgue point of functions  $f \in L_{p,\varphi}(\mathbb{R})$  and  $\varphi \in L_p(\mathbb{R})$  we have

$$|T_\lambda(f; x) - f(x_0)|^p = o(\Delta(x, \lambda, \delta))$$

as  $(x, \lambda) \rightarrow (x_0, \lambda_0)$ .

*Proof.* Imposing the conditions (i)-(iv) on the resulting inequality of the proof of Theorem 5.2, we obtain the desired result. The proof of the case  $p = 1$  is quite similar. Thus the proof is completed.  $\square$

## 7. Concluding Remark

In this paper, weighted approximation properties of nonlinear non-convolution type singular integral operators are studied. For this aim, the class of kernel functions, called *Class  $A_w$* , is defined. Therefore, main theorems are presented as Theorem 5.1 and Theorem 5.2.

## References

- [1] G. Alexits, *Konvergenzprobleme der Orthogonalreihen*, Verlag der Ungarischen Akademie der Wissenschaften, Budapest, 1960.
- [2] C. Bardaro, *On approximation properties for some classes of linear operators of convolution type*, Atti Sem. Mat. Fis. Univ. Modena, **33** (2) (1984), 329–356.
- [3] C. Bardaro and G. Vinti, *On approximation properties of certain nonconvolution integral operators*, J. Approx. Theory, **62** (3) (1990), 358–371.
- [4] C. Bardaro and G. Vinti, *Approximation by nonlinear integral operators in some modular function spaces*, Ann. Polon. Math., **63** (1996), 173–182.
- [5] C. Bardaro, J. Musielak and G. Vinti, *Nonlinear integral operators and applications: De Gruyter Series in Nonlinear Analysis and Applications, Vol. 9*, Walter de Gruyter, Publ., Berlin, New York, 2003.
- [6] C. Bardaro and I. Mantellini, *Pointwise convergence theorems for nonlinear Mellin convolution operators*, Int. J. Pure Appl. Math., **27** (4) (2006), 431–447.

- [7] P. L. Butzer and R. J. Nessel, *Fourier Analysis and Approximation: Vol. I*, Academic Press, New York, London, 1971.
- [8] Deepmala and H. K. Pathak, *A study on some problems on existence of solutions for nonlinear functional-integral equations*, Acta Mathematica Scientia, **33** B(5) (2013), 1305–1313.
- [9] Deepmala, *A study on fixed point theorems for nonlinear contractions and its applications*, Ph. D. Thesis, Pt. Ravishankar Shukla University, Raipur 492 010, Chhatisgarh, India, 2014.
- [10] S. Esen, *Convergence and the order of convergence of family of non-convolution type integral operators at characteristic points*, Ph. D. Thesis, Ankara University, Ankara, Turkey, 2002.
- [11] S. Esen, *The order of approximation by the family of integral operators with positive kernel*, Proc. Inst. Math. Mech. Natl. Acad. Sci. Azerb., **28** (2008), 117-122.
- [12] A. D. Gadjiev, *The order of convergence of singular integrals which depend on two parameters*, in: Special Problems of Functional Analysis and their Appl. to the Theory of Diff. Eq. and the Theory of Func., Izdat. Akad. Nauk Azerbaidzan. SSR, (1968), 40–44.
- [13] O. O. Guller, G. Uysal and E. Ibikli, *On weighted approximation by singular integral operators depending on two parameters*, 4th International Eurasian Conference on Mathematical Sciences and Applications (IECMSA 2015), Athens, Greece, (2015), 89.
- [14] H. Karsli, *On approximation properties of non-convolution type nonlinear integral operators*, Anal. Theory Appl., **26** (2) (2010), 140–152.
- [15] R. G. Mamedov, *On the order of convergence of  $m$ -singular integrals at generalized Lebesgue points and in the space  $L_p(-\infty, \infty)$* , Izv. Akad. Nauk SSSR Ser. Mat., **27** (2) (1963), 287-304.
- [16] V. N. Mishra and L. N. Mishra, *Trigonometric approximation of signals (Functions) in  $L_p(p \geq 1)$ -norm*, International Journal of Contemporary Mathematical Sciences, **7**(19) (2012), 909–918.
- [17] V. N. Mishra and S. Pandey, *On Chlodowsky variant of  $(p; q)$  Kantorovich-Stancu-Schurer operators*, Int. J. Analysis and Applications, **11** (1) (2016), 28-39.
- [18] L. N. Mishra and M. Sen, *On the concept of existence and local attractivity of solutions for some quadratic Volterra integral equation of fractional order*, Appl. Math. Comput., **285** (2016), 174–183.
- [19] L. N. Mishra, M. Sen and R. N. Mohapatra, *On existence theorems for some generalized nonlinear functional-integral equations with applications*, Filomat, (2016), (in press).
- [20] L. N. Mishra, R. P. Agarwal and M. Sen, *Solvability and asymptotic behavior for some nonlinear quadratic integral equation involving Erdelyi-Kober fractional integrals on the unbounded interval*, Progress in Fractional Differentiation and Applications, **2** (3) (2016), 153-168.
- [21] J. Musielak, *On some approximation problems in modular spaces*, in: Constructive Function Theory, Proceedings of International Conference

- Varna, 1-5 June, 1981, Publication House of Bulgarian Academic of Sciences, Sofia, (1983), 181-189.
- [22] J. Musielak, *Approximation by nonlinear singular integral operators in generalized Orlicz spaces*, Comment. Math. Prace Mat., **31** (1991), 79–88.
- [23] T. Swiderski and E. Wachnicki, *Nonlinear singular integrals depending on two parameters*, Comment. Math., **40** (2000), 181-189.
- [24] H. K. Pathak and Deepmala, *Common fixed point theorems for PD-operator pairs under relaxed conditions with applications*, J. Comp. Appl. Math., **239** (2013), 103-113.
- [25] W. Rudin, *Real and Complex Analysis*, Mc-Graw Hill Book Co., London, 1987.
- [26] B. Rydzewska, *Approximation des fonctions par des intégrales singulières ordinaires*, Fasc. Math., **7** (1973), 71–81.
- [27] R. Taberski, *Singular integrals depending on two parameters*, Prace Mat., **7** (1962), 173-179.
- [28] G. Uysal and E. Ibikli, *A note on nonlinear singular integral operators depending on two parameters*, New Trends Math. Sci. **4** (1) (2016), 104–114.

Gumrah Uysal  
Department of Computer Technologies  
Division of Technology of Information Security  
Karabuk University  
Karabuk 78050, Turkey  
*E-mail:* guysal@karabuk.edu.tr

Vishnu Narayan Mishra  
Sardar Vallabhbhai National Institute of Technology  
Surat 395 007, Gujarat, India  
*E-mail:* vishnunarayanmishra@gmail.com

Ozge Ozalp Guller  
Department of Mathematics  
Ankara University  
Ankara 06100, Turkey  
*E-mail:* osgeosalp@hotmail.com

Ertan Ibikli  
Department of Mathematics  
Ankara University  
Ankara 06100, Turkey  
*E-mail:* Ertan.Ibikli@ankara.edu.tr