

# The Association Among Individual and Contextual Factors and Unmet Healthcare Needs in South Korea: A Multilevel Study Using National Data

Seung Eun Lee<sup>1</sup>, Miyeon Yeon<sup>2</sup>, Chul-Woung Kim<sup>3,4</sup>, Tae-Ho Yoon<sup>5</sup>

<sup>1</sup>Graduate School of Public Health, Seoul National University, Seoul; <sup>2</sup>Department of Statistics, Graduate School, Chungnam National University, Daejeon; <sup>3</sup>Department of Preventive Medicine, Chungnam National University School of Medicine, Daejeon; <sup>4</sup>Research Institute for Medical Sciences, Chungnam National University, Daejeon; <sup>5</sup>Department of Preventive Medicine, Busan National University School of Medicine, Busan, Korea

**Objectives:** The objective of this study is to investigate associations between contextual characteristics and unmet healthcare needs in South Korea after accounting for individual factors.

**Methods:** The present study used data from the 2012 Korean Community Health Survey (KCHS) of 228 902 adults residing within 253 municipal districts in South Korea. A multilevel analysis was conducted to investigate how contextual characteristics, defined by variables that describe the regional deprivation, degree of urbanity, and healthcare supply, are associated with unmet needs after controlling for individual-level variables.

**Results:** Of the surveyed Korean adults, 12.1% reported experiencing unmet healthcare needs in the past. This figure varied with the 253 districts surveyed, ranging from 2.6% to 26.2%. A multilevel analysis found that the association between contextual characteristics and unmet needs varied according to the factors that caused the unmet needs. The degree of urbanity was associated with unmet need due to “financial burden” (odds ratio [OR], 0.53; 95% confidence interval [CI], 0.42 to 0.66 for rural vs. metropolitan), but not unmet need due to “service not available when needed.” There were no significant associations between these unmet need measures and regional deprivation. Among individual-level variables, income level showed the highest association with unmet need due to “financial burden” (OR, 5.63; 95% CI, 4.76 to 6.66), while employment status showed a strong association with unmet need due to “service not available when needed.”

**Conclusions:** Our finding suggests that different policy interventions should be considered for each at-risk population group to address the root cause of unmet healthcare needs.

**Key words:** Health services accessibility, Utilization, Residence characteristics, Multilevel analysis, Health surveys

Received: April 5, 2016 Accepted: September 7, 2016

**Corresponding author:** Chul-Woung Kim, MD, PhD  
266 Munhwa-ro, Jung-gu, Daejeon 35015, Korea

Tel: +82-42-580-8268, Fax: +82-42-580-8261

E-mail: [woung@cnu.ac.kr](mailto:woung@cnu.ac.kr)

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## INTRODUCTION

Although the Korean government contributed to making healthcare more accessible by establishing universal healthcare coverage in 1989, the rate of unmet healthcare needs in South Korea (hereafter Korea) remains high due to high levels of out-of-pocket payment caused by limited benefit coverage

[1]. According to the Korea National Health and Nutrition Examination Survey (KNHANES), in Korea, the rate of unmet healthcare needs has decreased from 22.8% in 2008 to 16.7% in 2012 [2]. However, as of 2012, compared to the average 6.4% rate of unmet healthcare needs in 28 countries in the European Union, the rate in Korea is almost three times higher and the gap based on income levels has not decreased [2,3].

Existing studies pertaining to unmet healthcare needs have focused primarily on identifying the prevalence of self-reported unmet healthcare needs and analyzing individual-level predictors. According to studies conducted in the US, where unmet healthcare need research is the most active, health insurance coverage and income level were the greatest predictors of unmet healthcare needs [4-6]. Income level was a strong predictor of unmet healthcare needs even in Canada and European countries adopting universal healthcare. In addition, gender, self-reported health, educational level, and chronic diseases status were associated with unmet healthcare needs [4-10].

On the other hand, regional contextual characteristics have been gaining attention as a potential determinant of healthcare utilization [11]. In order to more accurately identify the effects of contextual characteristics, it is worth taking a closer look at the multilevel analysis method. Multilevel analysis is considered a very useful method in studying the association between individuals' health and contextual characteristics, in that it allows simultaneous examination of the effects of higher-level and lower-level predictors. Despite the scarcity of studies on contextual-level correlates of variables of unmet need, some studies have confirmed the influence of regional socioeconomic characteristics on unmet healthcare need even when controlling for individual-level characteristics [1,7,12,13]. Nevertheless, the region-level variables examined in these studies were limited to the number of hospital beds and physicians [1], or the samples were drawn from a single state [12,13], making it difficult to generalize the findings. A US study [7] examined the association between unmet healthcare needs and material deprivation measured by poverty rate, unemployment rate, and educational level. However, the study failed to incorporate the concept of social deprivation, which has emerged as a valuable index in recent years. As such, it falls short of reflecting complex regional characteristics influencing residents' health.

In this context, more complex measures, namely deprivation indices, that include various factors such as income, em-

ployment, education, crime, housing environment, and social networks were developed. Prior studies have proven that the deprivation index is associated with mortality and life expectancy [14,15] as well as physical and mental health status, including subjective health status [16-19]. Moreover, a few studies have demonstrated associations between the index and health service use, including medical and dental visits [16,20,21]. These studies reported that the average health status was poor and healthcare use was higher in regions with severe deprivation.

The present study, which concerns the association between unmet healthcare needs and contextual characteristics, is differentiated from existing studies in the following ways. First, a multilevel analysis was used to consider, simultaneously, the effects of both individual and contextual characteristics on unmet healthcare needs. Second, for the regional characteristics variable, the study adopted a composite deprivation index, which integrates a region's material and social deprivation. This study is the first Korean study examining the association between a regional deprivation index and unmet healthcare needs. Third, the reasons for unmet healthcare needs, which vary by individual, include financial burden, time constraints, and mildness of symptoms; despite the fact that the effects on regional variables and policy-based intervention measures are expected to vary depending on these reasons [4], most preceding studies failed to distinguish these issues in their analyses. This study conducted a multilevel analysis for each individual cause of unmet healthcare needs.

## METHODS

### Conceptual Model

The objective of this study is to estimate the associations of contextual characteristics such as deprivation or other indicators of socioeconomic context with individual-level outcomes (unmet healthcare needs) after adjustment for individual-level measures. We also hypothesized that an association between contextual characteristics and unmet healthcare needs would vary for two common reasons ("financial burden" and "service not available when needed").

Despite the myriad of available studies pertaining to region-level characteristics and health outcomes, most studies in their analyses have used widely available yet limited data sources such as the population census data to extract contextual variables in the absence of conceptual frameworks [22]. In

order to gain a comprehensive understanding of the individual-level or region-level variables that influence unmet healthcare needs, the present study uses Andersen and Davidson's healthcare utilization behavior model [11]. This framework is built upon Andersen's initial model [23], which contains three sets of predictive factors in explaining individual characteristics influencing healthcare utilization behavior: predisposing, enabling, and need factors. The present study incorporates communities' contextual characteristics as a determinant of healthcare utilization behavior, which is subsequently categorized into predisposing, enabling, and need factors. Based on this new model, contextual factors known and predicted to be associated with unmet healthcare needs by previous studies are selected for the analysis model.

## Data Sources

This study used data from the 2012 Korean Community Health Survey (KCHS) by the Korea Centers for Disease Control and Prevention (KCDC) [24,25]. The KCHS was selected as a source of national certified statistics (No. 10118). This nationwide health survey covers information on demographic and socioeconomic characteristics, health status, and healthcare utilization. The 2012 KCHS was conducted on an average of 900 subjects selected by the probability proportional to sampling method and the systematic sampling method among adults aged 19 years or older living in each area of Korea from August to October 2012. The KCHS was administered by trained interviewers in face-to-face interviews. Data was drawn from 228 921 individuals from 253 districts nationwide included in the 2012 KCHS; we omitted 19 individuals who did not provide important information including unmet healthcare needs. Finally, a total of 228 902 individuals were analyzed in this study. Eligible contextual-level data were collected from various sources.

## Measures

### Outcome variables

We measured subjective unmet need for healthcare based on the KCHS question: "During the past year, was there ever a time when you wanted medical care but could not get it?" A "yes" response was treated as a definition of an experience of unmet need. Regarding the question, "What are the causes of your unmet healthcare need," "service not available when needed" (35.8%) was the most prevalent response, followed by "financial burden" (22.8%), "mildness of symptoms" (15.9%),

"transportation issues" (8.4%), and "difficult to get an appointment" (2.3%). Of these responses, the two most prevalent were added as outcome variables.

### Individual-level variables

Variables for access-relevant characteristics of individuals consisted of the three domains from our conceptual model. Predisposing factors included age, gender, and educational attainment. Enabling factors included having private health insurance, social support indicated by current cohabitation with spouse, current employment status, and income level. Income was divided into five levels according to respondents' "monthly average household equivalent income," which was calculated by dividing household income by the square root of the number of household members [26]. Need variables were represented by the number of self-reported medical comorbidities including hypertension, diabetes, coronary artery disease, stroke, asthma, and arthritis. Depression history and self-reported health status were considered need factors in predicting healthcare access [27,28]. Additionally, current smoking status was analyzed as a health behavior factor [13].

### Contextual-level variables

For the contextual predisposing condition, we included the regional deprivation index and the degree of urbanity. To characterize regional deprivation, we used the composite regional deprivation index (RDI) for Korea, which was recently calculated by Kim et al. [29] using 10% of the sample survey from the 2010 census, along with district-level data. To summarize how the RDI is calculated, a total of 16 potential deprivation indices predicted to reflect regional socioeconomic statuses in Korea were obtained from the census data. Upon a distribution test, those showing irregular distribution were converted for a factor analysis. The factor analysis found that residential density and unemployment rate had negative values and, as such, were eliminated. The rate of non-homeowner status was also found to be extremely low in rural areas, and, as such, it was eliminated. As for the indices with similar implications, those with the highest factor loading values were adopted for the analysis [29]. Table 1 shows the final subordinate composition measures. The RDI was constructed by summing up the Z-standardized scores of each variable. In this study, the RDI was divided into five categories according to the score quintile. This index was found to have sufficient validity by Kim et al. [29] and previous studies have reported that the RDI

**Table 1.** Description of the regional deprivation index for Korea

Indicator	Definition
Poor residential environments	Percent of households under the minimum housing standard (no separate kitchen, no water supply, no hot water for the bathroom, or no flushing toilet)
No car	Percent of households without a car for commuting
Low education attainment	Percent of individuals with education level below high school graduation among the population of those 30-64 years old
Aging population	Percent of elderly individuals 65 years or older in the total population
Low social class	Percent of all persons in households with a head of the household who is engaged in elementary occupation <sup>1</sup>
Non-apartment	Percent of non-apartment households
Single occupant households	Percent of households living alone
Female heads-of-households	Percent of households with female heads-of-household
The divorced or separated	Percent of divorced or widowed individuals

<sup>1</sup>Elementary occupation is one that belongs in class V and below (unskilled manual labor) as defined by the Korean "social class classification standards by occupation."

index is more suitable in Korea than the Townsend or Carstairs indices, which were developed in England [29,30]. In addition, several studies have suggested that the degree of urbanity, as a region-level socioeconomic variable influencing unmet healthcare need, may influence health status and hence the need for healthcare [7,13]. Therefore, the present study incorporated urbanity, which is designated by the government (metropolitan, urban, and rural), as a contextual predisposing variable, in addition to the regional deprivation index. This study used the governmental administrative district system based on population size, industrial structure, and lifestyle to divide the autonomous areas into "metropolitan," for districts affiliated with major cities; "urban," for districts affiliated with cities other than major cities; and "rural," for counties affiliated with some parts of major cities (e.g., Ongjin County in Incheon City) and provinces [31].

For the contextual enabling characteristics that may impede or facilitate individuals' health service utilization, we selected variables pertaining to healthcare supply. So far, research into the association between supply-side factors and unmet healthcare need is sparse, and the results of available studies do not indicate a clear association [1,7,12]. To measure regional healthcare supplies, we used the number of physicians per 1000 residents in 2012 and the number of hospital beds per 1000 residents in 2011 [1,12,13]. Each variable was divided into quartiles.

### Statistical Analysis

Due to the hierarchical structure of the data with 228 902 individuals (level 1) nested within 253 districts (level 2), we merged district and individual-level data using local address

codes of the municipal districts and conducted a multilevel logistic regression analysis [32]. This study applied a random intercept model instead of a random coefficients regression model because the major explanatory variable, the RDI was not random for each of the 253 districts, and each region had a single fixed value. The two-level random intercept model was fit to the null model. We examined the series of models by adding contextual-level and individual-level variables. First, model 1 examined the effects of contextual-level predisposing variables (regional deprivation and degree of urbanity). Then, model 2 was analyzed, adding contextual-level enabling variables (the number of physicians and hospital beds per 1000 residents) into model 1. Finally, to examine the associations of contextual characteristics with individual-level unmet healthcare needs after adjustment for individual-level confounders, model 3 was analyzed, adding the individual-level variables into model 2. The sample for the community health survey was extracted by complex sample design. Thus, adjusted weights, which were provided by the KCHS, were included in the two-level designs to account for the design effect [32,33].

This study used two steps to determine whether there were any explanatory variables that created a multicollinearity problem. First, correlation coefficients were calculated for the explanatory variables to perform a preliminary evaluation of multicollinearity. Second, the variation inflation factor for each explanatory variable was calculated in the regression model to identify any with a value  $\geq 3$ . The two evaluation steps showed that there were no variables suspected to have multicollinearity; therefore, explanatory variables presented in the study results were included in the model.

Descriptive statistics were conducted in SAS version 9.3 (SAS

**Table 2.** Regional distribution of unmet healthcare needs across districts in Korea (n=253)

Provinces	No. of districts	Unmet healthcare needs (%)		
		Mean (SD)	Min	Max
Seoul	25	13.3 (2.7)	9.0	19.7
Busan	16	10.9 (3.0)	6.5	18.2
Daegu	8	12.6 (2.3)	9.5	15.6
Incheon	10	12.3 (3.6)	6.0	17.4
Gwangju	5	13.5 (2.7)	8.5	16.2
Daejeon	5	9.7 (2.3)	6.1	12.9
Ulsan	5	10.3 (2.3)	6.9	13.9
Gyeonggi	45	11.3 (2.7)	3.1	18.2
Gangwon	18	10.7 (2.9)	5.0	15.1
Chungbuk	13	12.6 (5.0)	3.8	22.7
Chungnam	15	11.9 (3.0)	4.9	16.2
Jeonbuk	14	12.7 (4.0)	7.5	19.6
Jeonnam	22	13.1 (5.6)	4.0	23.1
Gyeongbuk	25	12.2 (5.6)	2.6	26.2
Gyeongnam	20	12.2 (1.7)	8.9	16.0
Jeju	6	15.2 (3.6)	10.2	20.8
Sejong	1	10.0 (0.0)	10.0	10.0
Total	253	12.1 (3.8)	2.6	26.2

SD, standard deviation; Min, minimum; Max, maximum.

Institute Inc., Cary, NC, USA) and HLM version 7.0 (Scientific Software International Inc., Lincolnwood, IL, USA) was used to conduct the multilevel analyses.

## RESULTS

### Baseline Characteristics of the Study Population

Table 2 shows the distribution of rates of unmet healthcare needs for 253 municipal districts in Korea. The mean rate of unmet healthcare needs in these districts was 12.1%, ranging from a minimum of 2.6% to a maximum of 26.2%, highlighting the disparity across districts in Korea.

Table 3 shows the rates of unmet healthcare needs based on individual-level and contextual-level variables. Unmet need was higher among individuals over 75 years of age, women, individuals with less than elementary education, divorced and widowed individuals, and individuals with lower household income. Individuals residing in the most deprived community were found to experience a greater level of unmet need compared with their counterparts residing in the least deprived community.

### Multilevel Analysis Results

Table 4 displays the results of a multilevel logistic regression analysis performed to examine the association between individuals' unmet healthcare needs and contextual characteristics. In the final model, urbanity among contextual variables was significantly related to unmet healthcare need; urban residents showed a reduced probability of experiencing unmet healthcare need compared to metropolitan residents (odds ratio [OR], 0.89; 95% confidence interval [CI], 0.81 to 0.97). In the case of individual-level variables, all subordinate variables were found to have a significant association with unmet need. In particular, bad or very bad self-reported health was associated with odds of unmet healthcare needs that were 2.98 times greater than very good or good self-reported health (95% CI, 2.80 to 3.17).

Table 5 displays the results of a multilevel analysis performed to test the hypothesis that the association between regional contextual variables and unmet need will vary with the causes of unmet needs. First, in cases where unmet healthcare needs occurred due to "financial burden," regional deprivation and the degree of urbanity were associated with unmet healthcare need in model 1. However, in model 3, in which all individual-level variables were included, the association between regional deprivation and unmet need became non-significant, while urbanity variables remained statistically significant. In this model, the individual-level variable associated with the highest OR was the income level. Compared with the highest income level, the lowest income level showed 5.63 times higher odds of experiencing unmet healthcare need (95% CI, 4.76 to 6.66).

On the other hand, in the case of unmet need due to "service not available when needed," the significant regional variable was the number of physicians per 1000 residents. This showed an inverse relationship with unmet healthcare in only the third quartile (OR, 1.18; 95% CI, 1.04 to 1.32). The individual-level variable associated with the highest OR was bad or very bad self-reported health (OR, 2.53; 95% CI, 2.26 to 2.81), followed in order by two or more complex diseases, and female. Employment status also showed a strong relationship with unmet healthcare: the odds of the unemployed experiencing unmet healthcare was 84% lower than that for wage workers (95% CI, 0.12 to 0.19).

**Table 3.** Distribution of individual and contextual characteristics in relation to unmet healthcare needs among Korean adults (n=228 902)

	Unmet healthcare needs			p-value <sup>1</sup>
	Yes	No	Total	
Individual level				
Unmet healthcare need	27661 (12.1)	201 241 (87.9)	228 902 (100.0)	
Predisposing				
Age (yr)				<0.001
19-39	7734 (12.5)	54 259 (87.5)	61 993 (27.1)	
40-64	12 530 (11.5)	96 285 (88.5)	108 815 (47.5)	
65-74	4132 (11.6)	31 349 (88.4)	35 481 (15.5)	
75+	3265 (14.4)	19 348 (85.6)	22 613 (9.9)	
Sex				<0.001
Male	9987 (9.7)	92 903 (90.3)	102 890 (44.9)	
Female	17 674 (14.0)	108 338 (86.0)	126 012 (55.1)	
Educational attainment				<0.001
Elementary school or less	9547 (14.9)	54 320 (85.1)	63 867 (28.0)	
Middle school	2831 (11.1)	22 753 (88.9)	25 584 (11.2)	
High school	8342 (10.8)	68 938 (89.2)	77 280 (33.8)	
University or higher	6894 (11.2)	54 849 (88.8)	61 743 (27.0)	
Enabling				
Cohabitation with spouse				<0.001
Currently cohabiting	17 896 (11.3)	140 712 (88.7)	158 608 (69.3)	
Separated	664 (13.7)	4193 (86.3)	4857 (2.1)	
Widowed	4175 (16.3)	21 513 (83.7)	25 688 (11.2)	
Divorced	1206 (18.3)	5383 (81.7)	6589 (2.9)	
Never married	3702 (11.2)	29 296 (88.8)	32 998 (14.4)	
Private health insurance				<0.001
Yes	17 531 (11.4)	136 850 (88.6)	154 381 (67.8)	
No	10 015 (13.6)	63 413 (86.4)	73 428 (32.2)	
Employment status				<0.001
Wage worker	10 833 (12.9)	73 148 (87.1)	83 981 (36.8)	
Employer/self-employed	5452 (11.1)	43 594 (88.9)	49 046 (21.5)	
Unpaid family worker	1730 (13.9)	10 740 (86.1)	12 470 (5.5)	
Student	580 (7.7)	6948 (92.3)	7528 (3.3)	
House worker	5414 (11.9)	40 061 (88.1)	45 475 (19.9)	
Unemployed	3583 (12.0)	26 332 (88.0)	29 915 (13.1)	
Income level				<0.001
1st quintile (lowest)	8457 (15.2)	47 048 (84.8)	55 505 (24.2)	
2nd quintile	4941 (12.4)	35 063 (87.6)	40 004 (17.5)	
3rd quintile	5548 (11.3)	43 746 (88.7)	49 294 (21.5)	
4th quintile	4296 (10.6)	36 131 (89.4)	40 427 (17.7)	
5th quintile (highest)	4419 (10.1)	39 253 (89.9)	43 672 (19.1)	
Need				
No. of medical comorbidities				<0.001
0	7901 (9.0)	80 300 (91.0)	88 201 (38.5)	
1	7283 (12.2)	52 577 (87.8)	59 860 (26.2)	
2+	12 477 (15.4)	68 361 (84.6)	80 838 (35.3)	

(Continued to the next page)

**Table 3.** Continued from the previous page

	Unmet healthcare needs			p-value <sup>1</sup>
	Yes	No	Total	
Depression history				<0.001
Yes	1311 (23.3)	4308 (76.7)	5619 (2.5)	
No	26 339 (11.8)	196 879 (88.2)	223 218 (97.5)	
Self-reported health				<0.001
Very good/good	6733 (7.6)	82 189 (92.4)	88 922 (38.9)	
Moderate	11 372 (12.4)	80 072 (87.6)	91 444 (40.0)	
Bad/very bad	9548 (19.7)	38 966 (80.3)	48 514 (21.2)	
Health behaviors				<0.001
Smoking status				
Current smoker	5589 (11.7)	42 221 (88.3)	47 810 (20.9)	
Former smoker	3605 (9.8)	33 223 (90.2)	36 828 (16.1)	
Non-smoker	18 464 (12.8)	125 772 (87.2)	144 236 (63.0)	
Contextual level				
Predisposing				
Regional deprivation				<0.001
1st quintile (least deprived)	5586 (11.9)	41 247 (88.1)	46 833 (20.5)	
2nd quintile	5653 (12.1)	41 008 (87.9)	46 661 (20.4)	
3rd quintile	5411 (11.5)	41 603 (88.5)	47 014 (20.5)	
4th quintile	5229 (11.9)	38 702 (88.1)	43 931 (19.2)	
5th quintile (most deprived)	5782 (13.0)	38 681 (87.0)	44 463 (19.4)	
Urbanity				<0.001
Metropolitan	7701 (12.2)	55 675 (87.8)	63 376 (27.7)	
Urban	10 652 (11.7)	80 210 (88.3)	90 862 (39.7)	
Rural	9308 (12.5)	65 356 (87.5)	74 664 (32.6)	
Enabling				
No. of physicians per 1000 residents				<0.001
1st quartile (lowest)	7064 (12.1)	51 554 (87.9)	58 618 (25.6)	
2nd quartile	6650 (11.9)	49 259 (88.1)	55 909 (24.4)	
3rd quartile	7096 (13.0)	47 684 (87.0)	54 780 (23.9)	
4th quartile (highest)	6851 (11.5)	52 744 (88.5)	59 595 (26.0)	
No. of hospital beds per 1000 residents				<0.001
1st quartile (lowest)	7430 (12.5)	51 911 (87.5)	59 341 (25.9)	
2nd quartile	6975 (12.4)	49 106 (87.6)	56 081 (24.5)	
3rd quartile	6410 (11.4)	49 971 (88.6)	56 381 (24.6)	
4th quartile (highest)	6846 (12.0)	50 253 (88.0)	57 099 (24.9)	

Values are presented as number (%).

Missing data: educational attainment, 428; cohabitation with spouse, 162; private health insurance, 1,093; employment status, 487; number of medical comorbidities, 3; depression history, 65; self-reported health, 28.

<sup>1</sup>p-values are calculated by chi-square test of the difference in unmet healthcare needs across different socio-demographic groups.

## DISCUSSION

The present study began with the hypothesis that contextual characteristics will have a significant association with unmet healthcare need even after controlling for individuals' compositional characteristics. In other words, we expected a region's

socioeconomic status would have a contextual effect beyond individual-level socioeconomic status on unmet healthcare need. KCHS data obtained from 228 902 individuals from 253 districts in Korea were analyzed through a multilevel analysis. Importantly, this study differed from existing, standard logistic regression analyses because the model took into account re-

**Table 4.** Multilevel analysis results of unmet healthcare needs among Korean adults (n=228 902)

	Unmet healthcare needs		
	Model 1	Model 2	Model 3
Individual level			
Predisposing			
Age			0.97 (0.96, 0.97)***
Sex			
Male			1.00 (reference)
Female			1.47 (1.39, 1.55)***
Educational attainment			
University or higher			1.00 (reference)
High school			1.01 (0.96, 1.06)
Middle school			1.04 (0.95, 1.12)
Elementary school or less			1.34 (1.24, 1.43)***
Enabling			
Cohabitation with spouse			
Currently cohabiting			1.00 (reference)
Separated			1.16 (1.01, 1.32)*
Widowed			1.28 (1.19, 1.37)***
Divorced			1.29 (1.17, 1.41)***
Never married			0.75 (0.69, 0.80)***
Private health insurance			
Yes			1.00 (reference)
No			1.12 (1.06, 1.17)***
Employment status			
Wage worker			1.00 (reference)
Employer/self-employed			0.92 (0.86, 0.97)**
Unpaid family worker			0.78 (0.71, 0.85)***
Student			0.51 (0.45, 0.58)***
House worker			0.61 (0.58, 0.64)***
Unemployed			0.63 (0.58, 0.67)***
Income level			
5th quintile (highest)			1.00 (reference)
4th quintile			1.08 (1.00, 1.15)*
3rd quintile			1.16 (1.08, 1.24)***
2nd quintile			1.32 (1.22, 1.41)***
1st quintile (lowest)			1.49 (1.38, 1.61)***
Need			
No. of medical comorbidities			
0			1.00 (reference)
1			1.39 (1.32, 1.45)***
2+			1.60 (1.51, 1.69)***
Depression history			
No			1.00 (reference)
Yes			1.55 (1.39, 1.71)***

(Continued to the next page)

Table 4. Continued from the previous page

	Unmet healthcare needs		
	Model 1	Model 2	Model 3
Self-reported health			
Very good/good			1.00 (reference)
Moderate			1.65 (1.57, 1.72)***
Bad/very bad			2.98 (2.80, 3.17)***
Health behaviors			
Smoking status			
Non-smoker			1.00 (reference)
Former smoker			1.04 (0.97, 1.11)
Current smoker			1.22 (1.15, 1.29)***
Contextual level			
Predisposing			
Regional deprivation			
1st quintile (least deprived)	1.00 (reference)	1.00 (reference)	1.00 (reference)
2nd quintile	0.92 (0.76, 1.10)	0.94 (0.76, 1.15)	0.97 (0.80, 1.16)
3rd quintile	0.95 (0.77, 1.15)	0.96 (0.76, 1.20)	1.01 (0.81, 1.25)
4th quintile	0.92 (0.75, 1.13)	0.92 (0.73, 1.16)	1.00 (0.80, 1.24)
5th quintile (most deprived)	0.97 (0.79, 1.17)	0.99 (0.78, 1.24)	1.08 (0.86, 1.35)
Urbanity			
Metropolitan	1.00 (reference)	1.00 (reference)	1.00 (reference)
Urban	0.91(0.83, 0.99)*	0.92 (0.83, 1.00)	0.89 (0.81, 0.97)*
Rural	0.94 (0.79, 1.10)	0.93 (0.76, 1.13)	0.91 (0.75, 1.09)
Enabling			
No. of physicians per 1,000 residents			
4th quartile (highest)	1.00 (reference)	1.00 (reference)	
3rd quartile	1.15 (1.04, 1.26)**	1.10 (1.00, 1.20)*	
2nd quartile	1.05 (0.92, 1.19)	1.01 (0.89, 1.13)	
1st quartile (lowest)		1.05 (0.93, 1.18)	1.03 (0.91, 1.15)
No. of hospital beds per 1000 residents			
4th quartile (highest)	1.00 (reference)	1.00 (reference)	
3rd quartile	1.02 (0.89, 1.16)	1.01 (0.89, 1.15)	
2nd quartile	1.07 (0.94, 1.21)	1.06 (0.93, 1.20)	
1st quartile (lowest)	1.12 (0.98, 1.28)	1.12 (0.99, 1.27)	

Values are presented as odds ratio (95% confidence interval).

Model 1: individual-level variables and contextual-level enabling variables adjusted.

Model 2: contextual-level variables were simultaneously entered in the model.

Model 3: all variables were simultaneously entered in the model.

\* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ .

gional variation in the dependent variables (level 2), that is, unmet healthcare needs. Another reason for multilevel analysis was that 228 802 individuals belonged to 253 districts, which may not be independent of each other. For example, in the “unmet healthcare needs” model, whether someone experiences such needs may be determined by their district affiliation. This can lead to significant differences between districts,

resulting in errors; this would be contrary to the assumptions in traditional methodology, and multilevel analysis would be required. To determine the fitness of multilevel analysis, an evaluation was performed, based on the null hypothesis that variance in the intercept model of unmet healthcare needs is 0 ( $H_0: \tau_{00} = 0$ ). Since the null hypothesis was rejected, it was confirmed that the multilevel model was fit.

**Table 5.** Difference of multilevel analysis results of unmet healthcare needs due to financial burden and unavailability of service when needed among Korean adults

	Unmet healthcare needs due to financial burden			Unmet healthcare needs due to unavailability of service when needed		
	Model 1	Model 2	Model 3	Model 1	Model 2	Model 3
Individual level						
Predisposing						
Age			1.00 (reference)			1.00 (reference)
			0.98 (0.97, 0.98)***			0.96 (0.95, 0.96)***
Sex						
Male			1.00 (reference)			1.00 (reference)
Female			1.42 (1.26, 1.60)***			1.50 (1.39, 1.61)***
Educational attainment						
University or higher			1.00 (reference)			1.00 (reference)
High school			1.71 (1.46, 1.99)***			0.99 (0.92, 1.06)
Middle school			2.33 (1.98, 2.72)***			1.01 (0.88, 1.15)
Elementary school or less			2.61 (2.24, 3.03)***			1.16 (1.02, 1.31)*
Enabling						
Cohabitation with spouse						
Currently cohabiting			1.00 (reference)			1.00 (reference)
Separated			1.63 (1.33, 1.98)***			1.08 (0.89, 1.30)
Widowed			1.28 (1.15, 1.42)***			0.97 (0.83, 1.11)
Divorced			1.73 (1.48, 2.01)***			1.14 (0.99, 1.31)
Never married			0.90 (0.77, 1.05)			0.71 (0.64, 0.77)***
Private health insurance						
Yes			1.00 (reference)			1.00 (reference)
No			1.67 (1.54, 1.81)***			0.84 (0.77, 0.91)***
Employment status						
Wage worker			1.00 (reference)			1.00 (reference)
Employer/self-employed			0.80 (0.71, 0.90)***			0.92 (0.86, 0.99)*
Unpaid family worker			0.71 (0.58, 0.85)***			0.72 (0.63, 0.82)***
Student			0.38 (0.27, 0.53)***			0.35 (0.29, 0.41)***
House worker			0.83 (0.74, 0.92)***			0.20 (0.18, 0.22)***
Unemployed			0.88 (0.79, 0.97) *			0.16 (0.12, 0.19)***
Income level						
5th quintile (highest)			1.00 (reference)			1.00 (reference)
4th quintile			1.74 (1.43, 2.10)***			1.08 (0.99, 1.18)
3rd quintile			2.55 (2.15, 3.02)***			1.11 (1.01, 1.21)*
2nd quintile			4.24 (3.55, 5.06)***			1.17 (1.04, 1.30)**
1st quintile (lowest)			5.63 (4.76, 6.66)***			0.92 (0.81, 1.03)
Need						
No. of medical comorbidities						
0			1.00 (reference)			1.00 (reference)
1			1.26 (1.13, 1.39)***			1.48 (1.38, 1.57)***
2+			1.44 (1.28, 1.61)***			1.80 (1.67, 1.92)***
Depression history						
No			1.00 (reference)			1.00 (reference)
Yes			1.91 (1.65, 2.20)***			1.49 (1.24, 1.79)***

(Continued to the next page)

Table 5. Continued from the previous page

	Unmet healthcare needs due to financial burden			Unmet healthcare needs due to unavailability of service when needed		
	Model 1	Model 2	Model 3	Model 1	Model 2	Model 3
Self-reported health						
Very good/good			1.00 (reference)			1.00 (reference)
Moderate			1.83 (1.65, 2.01)***			1.73 (1.62, 1.84)***
Bad/very bad			4.54 (4.03, 5.10)***			2.53 (2.26, 2.81)***
Health behaviors						
Smoking status						
Non-smoker			1.00 (reference)			1.00 (reference)
Former smoker			0.97 (0.83, 1.12)			1.04 (0.94, 1.14)
Current smoker			1.33 (1.17, 1.49)***			1.26 (1.16, 1.36)***
Contextual level						
Predisposing						
Regional deprivation						
1st quintile (least deprived)	1.00 (reference)	1.00 (reference)	1.00 (reference)	1.00 (reference)	1.00 (reference)	1.00 (reference)
2nd quintile	0.82 (0.65, 1.02)	0.77 (0.59, 0.98)*	0.92 (0.72, 1.18)	1.03 (0.75, 1.39)	1.07 (0.76, 1.49)	0.98 (0.72, 1.32)
3rd quintile	0.87 (0.68, 1.10)	0.78 (0.59, 1.01)	1.13 (0.87, 1.47)	1.17 (0.84, 1.59)	1.28 (0.89, 1.81)	1.08 (0.77, 1.50)
4th quintile	0.68 (0.52, 0.86)**	0.60 (0.46, 0.79)***	1.01 (0.77, 1.32)	1.19 (0.86, 1.63)	1.28 (0.89, 1.82)	1.05 (0.75, 1.46)
5th quintile (most deprived)	0.70 (0.55, 0.89)**	0.61 (0.45, 0.80)***	1.14 (0.86, 1.50)	1.24 (0.90, 1.69)	1.42 (0.96, 2.08)	1.18 (0.82, 1.68)
Urbanity						
Metropolitan	1.00 (reference)	1.00 (reference)	1.00 (reference)	1.00 (reference)	1.00 (reference)	1.00 (reference)
Urban	0.85 (0.73, 0.97)*	0.81 (0.70, 0.93)**	0.76 (0.65, 0.87)***	0.9 (0.80, 1.00)	0.92 (0.81, 1.03)	0.90 (0.80, 1.00)
Rural	0.70 (0.58, 0.84)***	0.60 (0.47, 0.75)***	0.53 (0.42, 0.66)***	0.85 (0.66, 1.08)	0.91 (0.68, 1.22)	0.91 (0.69, 1.19)
Enabling						
No. of physicians per 1000 residents						
4th quartile (highest)		1.00 (reference)	1.00 (reference)		1.00 (reference)	1.00 (reference)
3rd quartile		1.06 (0.90, 1.24)	0.93 (0.79, 1.09)		1.22 (1.07, 1.39)**	1.18 (1.04, 1.32)**
2nd quartile		1.18 (0.98, 1.42)	1.07 (0.87, 1.30)		1.04 (0.87, 1.23)	1.01 (0.85, 1.19)
1st quartile (lowest)		1.25 (1.02, 1.50)*	1.11 (0.93, 1.31)		0.99 (0.83, 1.17)	0.97 (0.82, 1.15)
No. of hospital beds per 1000 residents						
4th quartile (highest)		1.00 (reference)	1.00 (reference)		1.00 (reference)	1.00 (reference)
3rd quartile		1.06 (0.90, 1.24)	1.08 (0.92, 1.26)		0.97 (0.81, 1.15)	0.95 (0.81, 1.12)
2nd quartile		1.18 (0.99, 1.39)	1.26 (1.06, 1.48)**		0.99 (0.83, 1.18)	0.97 (0.81, 1.14)
1st quartile (lowest)		1.09 (0.91, 1.30)	1.19 (0.99, 1.42)		0.94 (0.78, 1.13)	0.93 (0.78, 1.11)

Values are presented as odds ratio (95% confidence interval).

Model 1: individual-level variables and contextual-level enabling variables adjusted.

Model 2: contextual-level variables were simultaneously entered in the model.

Model 3: all variables were simultaneously entered in the model.

\* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ .

While considerable research has been conducted on unmet healthcare needs in Korea using recent panel data, these studies have focused on individual determinants of unmet needs for healthcare [34–36]. The study conducted by Heo et al. [1] is the sole study examining contextual determinants of unmet

needs in Korea. However, this study utilized only province-level data (from 16 provinces) of healthcare supply factors (number of hospital beds and physicians) as regional variables. The present study is important because 253 districts were analyzed, and contextual predisposing condition variables (com-

posite deprivation index, urbanity) were used in the analysis, in addition to healthcare supply factors. In addition, unlike a number of previous studies on unmet healthcare needs that did not distinguish the cause of unmet healthcare needs [6-10,12,13,27], or only included 'unmet healthcare needs for financial reasons' [34], to the best of our knowledge, this is the only study to perform a multilevel analysis, classifying unmet healthcare needs by main cause.

The primary finding of this study is that, compared to regions with lower deprivation, living in regions with higher material and social deprivation is not significantly associated with an increased probability of unmet healthcare needs. In the case of the model of unmet need due to "financial burden," the statistical significance of the regional deprivation index was observed in models 1 and 2, for which individual level variables were not included. However, such an association became nonsignificant in the final model. The following reasons may explain this finding. First, the association between the deprivation index and unmet need observed in models 1 and 2 may be attributable to individuals' compositional characteristics. This pattern is consistent with some previous work, in which most of the association of deprivation and health was explained by individual rather than area factors [17,22,37]. Second, the use of the composite measure may have offset the influence of subordinate indices [19]. In contrast to the present study, which used a composite deprivation index combining separate indicators into one index, a US study measuring separate indices such as poverty rate [12,13] confirmed that region-level variables and unmet need were correlated even after controlling for individual-level variables. Comparing the present study's results against such findings, it can be inferred that regional deprivation was nonsignificant in the present study because of the use of the composite measure. However, as opposed to focusing on identifying the separate regional variables that influence unmet healthcare need, the present study focused on investigating the effects of the overall nature of the regional socioeconomic context on individuals' unmet needs. Follow-up studies further investigating the potential association between unmet healthcare need and regional deprivation via various measures of regional characteristics would be beneficial in that results can be compared to gain more specific and useful insight into the topic.

Whereas the association between the regional deprivation index and unmet need was nonsignificant, urbanity variables were found to be significantly correlated with unmet need

due to "financial burden." Specifically, living in metropolitan areas increased the likelihood of experiencing unmet healthcare needs over living in rural or urban areas. Such a finding is contrasted against the finding of a US study that reported that living in a metropolitan statistical area was not correlated with unmet healthcare needs [7] or that of another prior study that found living in smaller urban areas (population less than one million) to be more highly correlated with unmet healthcare needs than living in more populous urban areas [13]. Heo et al. [1] suggested several possible causes behind the high unmet need in major Korean cities despite the concentration of healthcare resources. Areas with a surplus of private hospital beds tend to competitively acquire more beds or highly expensive equipment and encourage services not covered by insurance, which consequently increases the cost of care. The high unmet healthcare need in these areas is thought to be attributed to the low insurance coverage and high out-of-pocket payment burden imposed on Korean healthcare consumers. According to these results, it can be inferred that individuals with low socioeconomic status residing in metropolitan areas are at an elevated risk of unmet healthcare need than their counterparts residing in rural areas [18,37].

The numbers of hospital beds and physicians, which were included in the analysis model as contextual enabling conditions either impeding or facilitating healthcare utilization, were not found to be correlated with unmet healthcare need, a finding that is supported by prior studies [3,13,14]. Korea has the highest number of doctor consultations per capita among all Organization for Economic Cooperation and Development (OECD) nations and doctor accessibility is almost twice the OECD average as of 2011. The number of hospital beds per capita is also the highest of all OECD nations, with over nine beds per 1000 people as of 2011 [38]. Therefore, it is possible that the number of physicians or hospital beds is not a significant factor influencing individuals' healthcare accessibility in Korea.

Although the associations between contextual-level variables and unmet healthcare needs were nonsignificant in this study, all of the individual-level variables were significantly associated with unmet needs. In particular, the strongest predictor of unmet need was self-reported health, a finding supported by prior studies [1,5,27]. However, when the reasons for unmet health needs were classified as "financial burden" and "service not available when needed," there were considerable differences in the significance and the strength of correlations.

For unmet healthcare needs caused by “financial burden,” the average monthly household equivalent income and subjective health status had the highest and second highest OR, respectively. This result is consistent with those of a previous study that analyzed factors related to unmet healthcare needs caused by financial reasons, using the Korea Welfare Panel data (between 2005 and 2006), and found that poorer households and poorer objective health status were associated with a higher risk of experiencing unmet healthcare needs [34]. Conversely, where the reason for unmet healthcare was cited as “service not available when needed,” the average monthly household equivalent income was not significant in some bands, though employment status was one of the strongest predictors. More specifically, employed individuals, particularly waged workers, were more likely to experience unmet healthcare needs compared to those who were unemployed. Considering previous reviews regarding the occurrence of unmet need in Korea [39], “service not available when needed” may be attributed to Korea’s labor system, which is characterized by the longest working hours of all OECD nations [40]. In fact, according to a survey conducted by the Korea Health Panel, which incorporated a more clearly defined set of terms in investigating unmet healthcare needs, “no time to visit a clinic or hospital” topped the responses at 39.1%, followed by “financial burden of treatment” at 22.3%, similar to the present study’s findings. As of 2011, the average annual hours worked per Korean worker is 470 hours longer than that of the OECD average. The figure translates into 2.74 more months than the OECD average [40]. Such long average annual working hours are likely to impede workers from getting regular health check-ups or treatment [34,35]. In summary of these findings, ultimately, a different policy approach is needed for each at-risk population group to address the root cause of unmet healthcare need. First of all, for the socioeconomically disadvantaged population, policies that lessen the financial burden associated with healthcare services should be implemented, which include improved health insurance coverage. On the other hand, for workers with financial stability, various interventions and relevant labor policies could be considered to enable them to use the medical services they need. This needs further discussion in future studies.

This study has some limitations. First, not all contextual-level variables that influence unmet healthcare needs could be incorporated in the analysis model. An important criticism of the literature on contextual factors and health is that, despite nu-

merous studies, scant attention has been paid to conceptualizing the particular pathways underlying the associations with ad hoc selection of contextual factors studied [22]. Heeding this criticism, we selected the variables based on Andersen’s new healthcare utilization model, which considers both the individual- and contextual-level characteristics. However, due to issues of available data, a wider range of variables including social capital and physical environment could not be incorporated. This study used an RDI, which included both material and social deprivations, as the variable representing contextual-level characteristics in measuring the associations with unmet healthcare needs. However, deprivation indices based on a population census, such as RDI, have a limitation of not incorporating variables that can reflect the characteristics of rural regions [15]. Therefore, this study added the degree of urbanity variable to the model. However, if future studies include additional contextual characteristics that reflect social support of the community, a more meaningful explanation of the associations with unmet healthcare needs would be possible.

Additionally, when two independent variables and one dependent variable are included in a single model, if the association between one of the independent variables (level 1) and the dependent variable is altered by the other independent variable (level 2), an interaction can be created. However, this study did not examine the interactions between individual and regional level variables.

Third, due to the limitations of KCHS, responses from individuals on the details of which healthcare services were unmet could not be obtained. Since the significance and meaning of unmet needs can change based on whether the unmet service was an examination, a treatment of physical or mental problems, or injury care [21], future surveys that include questions on the type of care required will allow a more systematic analysis.

In Korea, the more urbanized the area in which an individual lived, and the lower that individual’s income was, the higher the possibility of experiencing unmet need due to “financial burden.” Conversely, for individuals participating in economic activities, the possibility of occurrence of unmet healthcare due to “service not available when needed” was higher than for individuals not participating in economic activities. A region’s socioeconomic disadvantage itself, which was represented by the deprivation index, did not seem to have an effect on unmet need above and beyond what would be expected given the residents’ socioeconomic status. Our finding

suggests that different policy interventions should be considered for each at-risk population group to address the root cause of unmet healthcare need.

## ACKNOWLEDGEMENTS

We would like to acknowledge the Korea Centers for Disease Control and Prevention for making available the Community Health Survey data, 2012. This work was supported by a research fund from Chungnam National University.

## CONFLICT OF INTEREST

The authors have no conflicts of interest associated with the material presented in this paper.

## ORCID

Seung Eun Lee <http://orcid.org/0000-0001-5120-6429>

Chul-Woung Kim <http://orcid.org/0000-0002-1958-6000>

## REFERENCES

1. Heo J, Oh J, Kim J, Lee M, Lee JS, Kwon S, et al. Poverty in the midst of plenty: unmet needs and distribution of health care resources in South Korea. *PLoS One* 2012;7(11):e51004.
2. Ministry of Health and Welfare. Health data OECD 2012 [cited 2016 Sep 9]. Available from: <file:///C:/Users/user/Downloads/OECD+Health+Data+2012.pdf> (Korean).
3. Organization for Economic Cooperation and Development. Health at a glance: Europe 2014; 2014 [cited 2016 Sep 9]. Available from: <http://www.oecd.org/health/health-at-a-glance-europe-23056088.htm>.
4. Allin S, Grignon M, Le Grand J. Subjective unmet need and utilization of health care services in Canada: what are the equity implications? *Soc Sci Med* 2010;70(3):465-472.
5. Diamant AL, Hays RD, Morales LS, Ford W, Calmes D, Asch S, et al. Delays and unmet need for health care among adult primary care patients in a restructured urban public health system. *Am J Public Health* 2004;94(5):783-789.
6. Zhang P, Tao G, Anderson LA. Differences in access to health care services among adults in rural America by rural classification categories and age. *Aust J Rural Health* 2003;11(2):64-72.
7. Kirby JB, Kaneda T. Neighborhood socioeconomic disadvantage and access to health care. *J Health Soc Behav* 2005;46(1):15-31.
8. Koolman X. Unmet need for health care in Europe. In: Comparative EU statistics on income and living conditions: issues and challenges; Proceedings of the EU-SILC Conference (Helsinki, Nov 6-8, 2006). Luxembourg: Office for Official Publications of the European Communities; 2007, p. 181-191.
9. Law M, Wilson K, Eyles J, Elliott S, Jerrett M, Moffat T, et al. Meeting health need, accessing health care: the role of neighbourhood. *Health Place* 2005;11(4):367-377.
10. Ahn YH, Kim NH, Kim CB, Ham OK. Factors affecting unmet healthcare needs of older people in Korea. *Int Nurs Rev* 2013;60(4):510-519.
11. Anderson RM, Davidson PL. Improving access to care in America. In: Rice TM, Kominski GF, Affi AA, Rosenstock L, Andersen R, editors. Changing the US health care system: key issues in health services policy and management. 3rd ed. San Francisco: Jossey-Bass; 2007, p. 3-31.
12. Litaker D, Love TE. Health care resource allocation and individuals' health care needs: examining the degree of fit. *Health Policy* 2005;73(2):183-193.
13. Peterson LE, Litaker DG. County-level poverty is equally associated with unmet health care needs in rural and urban settings. *J Rural Health* 2010;26(4):373-382.
14. Pampalon R, Hamel D, Gamache P. Health inequalities in urban and rural Canada: comparing inequalities in survival according to an individual and area-based deprivation index. *Health Place* 2010;16(2):416-420.
15. Jordan H, Roderick P, Martin D. The index of multiple deprivation 2000 and accessibility effects on health. *J Epidemiol Community Health* 2004;58(3):250-257.
16. Gunnell DJ, Peters TJ, Kammerling RM, Brooks J. Relation between parasuicide, suicide, psychiatric admissions, and socioeconomic deprivation. *BMJ* 1995;311(6999):226-230.
17. Reijneveld SA, Schene AH. Higher prevalence of mental disorders in socioeconomically deprived urban areas in the Netherlands: community or personal disadvantage? *J Epidemiol Community Health* 1998;52(1):2-7.
18. Stafford M, Marmot M. Neighbourhood deprivation and health: does it affect us all equally? *Int J Epidemiol* 2003;32(3):357-366.
19. Eibner C, Sturm R. US-based indices of area-level deprivation: results from HealthCare for Communities. *Soc Sci Med* 2006;62(2):348-359.
20. Lang IA, Gibbs SJ, Steel N, Melzer D. Neighbourhood deprivation

- tion and dental service use: a cross-sectional analysis of older people in England. *J Public Health (Oxf)* 2008;30(4):472-478.
21. Curtis S, Copeland A, Fagg J, Congdon P, Almog M, Fitzpatrick J. The ecological relationship between deprivation, social isolation and rates of hospital admission for acute psychiatric care: a comparison of London and New York City. *Health Place* 2006;12(1):19-37.
  22. Hillemeier MM, Lynch J, Harper S, Casper M. Measuring contextual characteristics for community health. *Health Serv Res* 2003;38(6 Pt 2):1645-1717.
  23. Andersen RM. Revisiting the behavioral model and access to medical care: does it matter? *J Health Soc Behav* 1995;36(1):1-10.
  24. Korea Centers for Disease Control and Prevention. Community health survey [cited 2016 Sep 9]. Available from: <https://chs.cdc.go.kr/chs/index.do> (Korean).
  25. Kang YW, Ko YS, Kim YJ, Sung KM, Kim HJ, Choi HY, et al. Korea community health survey data profiles. *Osong Public Health Res Perspect* 2015;6(3):211-217.
  26. Kim CS, Han SY, Lee SE, Kang JH, Kim CW. Dental care utilization for examination and regional deprivation. *J Prev Med Public Health* 2015;48(4):195-202.
  27. Baggett TP, O'Connell JJ, Singer DE, Rigotti NA. The unmet health care needs of homeless adults: a national study. *Am J Public Health* 2010;100(7):1326-1333.
  28. Marshall EG. Do young adults have unmet healthcare needs? *J Adolesc Health* 2011;49(5):490-497.
  29. Kim D, Lee S, Ki M, Kim M, Kim S, Kim Y, et al. Developing health inequalities indicators and monitoring the status of health inequalities in Korea. Seoul: Korea Institute for Health and Social Affairs; 2013, p. 166-179 (Korean).
  30. Yoon TH, Kim SY, Yun MS, Moon KJ. Relationship between local authority deprivation and activity limitation in adults aged 50 or over. *Korean J Health Educ Promot* 2015;32(2):27-37 (Korean).
  31. Kang BS, Lee CW. A study on development strategies for jurisdictional change of local metropolitan governments. *J Soc Sci* 2005;16:1-23 (Korean).
  32. Raudenbush SW, Bryk AS. Hierarchical linear models: applications and data analysis methods. 2nd ed. Thousand Oaks: Sage Publications; 2002, p. 68-98.
  33. Heeringa SG, Liu J. Complex sample design effects and inference for mental health survey data. *Int J Methods Psychiatr Res* 1998;7(1):56-65.
  34. Shin YJ, Shon JI. The prevalence and association factors of unmet medical need-using the 1st and 2nd Korea welfare panel data. *Health Soc Welf Rev* 2009;29(1):111-142 (Korean).
  35. Lee SY, Kim CW, Kang JH, Seo NK. Unmet healthcare needs depending on employment status. *Health Policy* 2015;119(7):899-906.
  36. Ko H. Unmet healthcare needs and health status: panel evidence from Korea. *Health Policy* 2016;120(6):646-653.
  37. Robert SA. Socioeconomic position and health: the independent contribution of community socioeconomic context. *Annu Rev Sociol* 1999;25:489-516.
  38. Organization for Economic Cooperation and Development. Health at a glance 2013: OECD indicators [cited 2016 Sep 9]. Available from: <http://apps.who.int/medicinedocs/documents/s21725en/s21725en.pdf>.
  39. Choi SH. Unmet health care needs among adults in Korea. *Public Health Wkly Rep* 2014;7:197-200 (Korean).
  40. Bae KS. What sustains the working time regime of long hours in South Korea? *Econ Soc* 2012;95:128-162 (Korean).