



Nutritional Quality and Physicochemical Characteristics of Defatted Bovine Liver Treated by Supercritical Carbon Dioxide and Organic Solvent

Sung-Won Kang^{1†}, Hye-Min Kim^{1†}, M. Shafiur Rahman¹, Ah-Na Kim¹, Han-Sul Yang¹, and Sung-Gil Choi*

Division of Food Science and Technology (Institute of Agriculture and Life Sciences), Gyeongsang National University, Jinju 52828, Korea

¹Division of Applied Life Science (BK21 Plus), Gyeongsang National University, Jinju 52828, Korea

Abstract

Defatted bovine liver (DBL) is a potential source of protein and minerals. Supercritical carbon dioxide (SC-CO₂) and a traditional organic solvent method were used to remove lipid from bovine liver, and the quality characteristics of a control bovine liver (CBL), bovine liver defatted by SC-CO₂ (DBLSC-CO₂) at different pressures, and bovine liver defatted by organic solvent (DBL-OS) were compared. The DBLSC-CO₂ samples had significantly higher ($p < 0.05$) protein, amino acid, carbohydrate, and fiber contents than CBL and DBL-OS. There was a higher yield of lipid from CBL when using SC-CO₂ than the organic solvent method. SDS-PAGE analysis demonstrated that the CBL and DBLSC-CO₂ had protein bands of a similar intensity and area, whereas DBL-OS appeared extremely poor bands or no bands due to the degradation of proteins, particularly in the 50 to 75 kDa and 20 to 25 kDa molecular weight ranges. In addition, DBLSC-CO₂ was shown to have superior functional properties in terms of total soluble content, water and oil absorption, and foaming and emulsification properties. Therefore, SC-CO₂ treatment offers a nutritionally and environmentally friendly approach for the removal of lipid from high protein food sources. In addition, SC-CO₂ may be a better substitute of traditional organic solvent extraction for producing more stable and high quality foods with high-protein, fat-free, and low calorie contents.

Keywords bovine liver, quality characteristics, supercritical carbon dioxide, organic solvent, defatted

Received November 15, 2016
Revised December 12, 2016
Accepted December 15, 2016

†These authors contributed equally to this work.

Corresponding author

Sung-Gil Choi

Division of Food Science and Technology (Institute of Agriculture and Life Sciences), Gyeongsang National University, Jinju 660-701, Korea
Tel: +82-55-772-1906
Fax: +82-55-772-1909
E-mail: sgchoi@gnu.ac.kr

Introduction

Bovine liver is a rich source of essential nutrients including proteins, minerals, and vitamins; it is approximately 1-2% of the live weight of the bovine (Li *et al.*, 2014). It is a potential source of nutrients in relation to the safety and adequacy of human nutrition. Desiccated and defatted bovine liver may be a good source of high protein and low calorie food with vitamin A, B₁₂, folate, iron, zinc, copper, and purine. Such foods can be used to prepare a healthy diet for children, the obese and anemic, nyctalopic, low immunity, and diabetic persons (Devatkal *et al.*, 2004). Generally, bovine liver is underutilized in relation to its huge production across the world. This widely available bovine liver can be utilized for processing food formulations to support economically viable meat production systems (Estévez *et al.*, 2005; Li *et al.*, 2014).

© This is an open access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

A traditional organic solvent method is widely used for the extraction of lipid from plant and animal foods (Pariser *et al.*, 1978). This method has several limitations such as loss of the nutrient content and functional properties of foods, expensive and time consuming, toxic, and potential to adversely affect human health and the environment (Hultin, 1994; Staby and Mollerup, 1993).

Supercritical fluid extraction (SFE) method is a promising alternative to conventional organic solvent methods. This method has superior extraction properties including high compressibility, liquid-like density, low viscosity, and high diffusivity (Lim *et al.*, 2002). SFE has many advantages over traditional organic solvent methods such as a lower critical temperature and surface tension, better selectivity, and being nontoxic, nonflammable, inexpensive, and easily removable from the extracts. Removal of lipid from foods using SC-CO₂ at low operating temperature minimizes the thermal degradation of proteins, antioxidants, and other nutritionally valuable components (Imison and Unthank, 2000; Kiran and Zhuang, 1997).

The purpose of this study was to investigate the major quality characteristics, such as proximate composition, amino acid profile, sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) pattern, and functional properties of CBL, DBLSC-CO₂, and DBL-OS.

Materials and Methods

Materials

Bovine liver was purchased from a local market in Jinju, Korea, and was stored at -80°C until used. Diethyl ether (purity 99.0%) and chloroform (purity 99.5%), were obtained from Daejung Chemicals & Metals Co., Ltd., Korea. Methanol and all other chemicals used in the study

were analytical or HPLC grade.

Preparation of defatted bovine liver

Frozen bovine liver was freeze-dried for 72-96 h. It was ground and DIN-sieved to obtain fine powder samples. Samples were stored at 4°C until used for SC-CO₂ and organic solvent extraction.

Defatting bovine liver by SC-CO₂

The SC-CO₂ extraction was performed using a supercritical fluid apparatus supplied by Ilshin Autoclave Co. (Korea). A schematic of the SC-CO₂ extraction process is shown in Fig. 1. Liquid CO₂ contained in the siphoned cylinder was cooled to 4°C by the chiller. Bovine liver powder (300 g) was loaded into the extraction vessel, the top of which was tightly sealed. CO₂ was pumped through the vessel at a flow rate of 50 g/min at a constant temperature of 45°C using pressures of 200, 300, and 450 bar to obtain different DSFSC-CO₂ samples. After completion (3 h), the vessel was gradually depressurized. Extracted oil and defatted powder were collected in a poly pack and maintained at 4°C for analysis. Extraction yield (%) was calculated as the weight of extracted oil obtained from the samples.

Defatting bovine liver by organic solvent

Lipid extraction from 50 g bovine liver samples was performed in a soxhlet apparatus for 14-16 h using ether or a mixture of chloroform/methanol (CM; 2:1 v/v) as organic solvents. After extraction was complete, the sediment was desolventized using air drying under a fume hood until no organic solvent odor was discernible. The defatted samples were stored at 4°C until further analysis and the extraction yield (%) was calculated as the weight




Fig. 1. Schematic of supercritical carbon dioxide (SC-CO₂) treatment system. 1. CO₂ cylinder; 2. electronic balance; 3. chiller; 4. CO₂ pump; 5. controller; 6. cosolvent reservoir; 7. circulation pump; 8. heating bath; 9. circulation pump; 10. extractor; 11. separator 1; 12. separator 2; V-1, valve 1, V-2, valve 2; BPR, back pressure regulator; dotted lines, water line; solid lines, CO₂ line.

of extracted oil obtained from the samples.

Proximate composition

The proximate compositions of CBL, DBLSC-CO₂, and DBL-OS were estimated for their moisture, ash, lipid, carbohydrate, and protein (N×6.25) contents following the method of AOAC (2000).

SDS-PAGE analysis

To establish the SDS-PAGE pattern, a 10 mg bovine liver sample was placed into a tube containing 1 mL lysis buffer (RIPA buffer, Sigma Aldrich Co., USA), and then sonicated five times (2 s each time) in ice. After incubation, the sonicated samples were transferred to room temperature for 30 min and then centrifuged at 12,000 *g* for 10 min at 4°C. The supernatant was used for sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) analysis, following the method of Laemmli (1970), using 5% (w/v) stacking gel and 10% (w/v) separating gel. The electrophoresis was performed using a Mini-PROTEAN Tetra Cell (Bio-rad Laboratories, USA) at a constant voltage of 70 V for 100 min, followed by 10 mA/plate. After electrophoresis, the PAGE gel was stained with 1% Coomassie staining solution (1.0 g Coomassie Brilliant Blue R-250, 300 mL methanol, 600 mL H₂O, and 100 mL acetic acid) for 15 min and destained using the same solution without dye. Protein quantification was performed by the analysis of bands using a scanning densitometer (UMAX PowerLook 1100, Taiwan).

Amino acid analysis

Amino acid analysis was conducted according to the method of Jeong and Shim (2004) with minor modifications. Bovine liver samples of 50 mg were placed in heat-proof screw cap tubes and dissolved in 5 mL of 6 N HCl. The tubes were then flowed with N₂ gas for 1 min and sealed. The samples were incubated in a drying oven at 110°C for 24 h. After reaction was complete, the supernatant was filtered using a glass filter. To remove HCl, the supernatant was concentrated using a rotary evaporator and the sediment was dissolved in 3 mL of sodium citrate buffer (pH 2.2). Finally, the solution was passed through a 0.45-µm PTFE filter (Sigma-Aldrich, Korea) and then analyzed using an auto amino acid analyzer (Biochrom 30, Sweden).

Water absorption capacity

Water absorption capacity (WAC) was determined fol-

lowing the method of Jyothirmayi *et al.* (2006) with minor modifications. Bovine liver samples of 1.0 g were placed into 50 mL polypropylene conical centrifuge tubes and dissolved in 10 mL water, and then vortexed thoroughly and heated at 60°C for 15 min in a water bath (BS-21, Jeio Tech, Korea). The slurry was centrifuged at 3,000 *g* for 15 min. After centrifugation the supernatant was removed, and the sediment was drained for 30 min. The per unit weight gain was reported as the WAC (g/g).

Oil absorption capacity

Oil absorption capacity (OAC) was measured following the method Jyothirmayi *et al.* (2006) with some modifications. Samples of 1.0 g were placed into 50 mL polypropylene conical centrifuge tubes and thoroughly mixed with 10 mL of refined soybean oil. This mixture was heated at 60°C, with shaking, in a water bath (BS-21; Jeio Tech, Korea) for 1 h. The solution was centrifuged at 3,000 *g* for 15 min, then the supernatant was removed, and the sediment was drained for 30 min at room temperature. OAC (g/g) was calculated as the per unit weight gain of the sample.

Total soluble content

Solubility was determined using a modification of the method of Rodriguez *et al.* (2005). Bovine liver sample solutions (1%, w/v) were prepared using a magnetic stirrer at room temperature for 3 h. The solution was then heated to 75°C in a water bath (BS-21; Jeio Tech, Korea) with little shaking at 185 *g* for 1 h. After heating was complete, the solution was allowed to stand for 30 min at ambient temperature and then centrifuged at 4,000 *g* for 25 min. A 10 mL aliquot of the supernatant was placed in a pre-weighed Petri dish and oven-dried at 105°C for 3 h. The total soluble content (%) was calculated as follows:

$$\text{Total soluble content (\%)} = \left[\frac{\text{weight of solids in 10 mL} \times 10}{\text{weight of sample}} \right] \times 100$$

Emulsifying properties

Emulsifying activity (EA) and emulsion stability (ES) were determined following the method of Pedroche *et al.* (2004) with minor modifications. Bovine liver samples of 0.5 g were dispersed in 12.5 mL of deionized water at room temperature and 12.5 mL of refined soybean oil was added. This was then homogenized (D-500 homogenizer; Wiggen Hauser, Germany) at 10,000 *g* for 1 min. The emulsions were then centrifuged at 1,300 *g* for 5

min. The volume was recorded before and after centrifugation and EA was calculated as follows:

$$\text{EA (\%)} = [(\text{height of emulsified layer}) / (\text{height of contents of the tube})] \times 100$$

ES was measured after re-centrifugation at 1,300 g for 5 min following heating in a water bath (BS-21; Jeio Tech, Korea) at 80°C for 30 min; it was calculated as follows:

$$\text{ES (\%)} = [(\text{height of remaining emulsion layer}) / (\text{height of original emulsified layer})] \times 100$$

Foaming properties

Foaming capacity (FC) and foaming stability (FS) were determined based on the method of Ogunwolu *et al.* (2009) with minor modifications. The solutions of bovine liver sample (1%, w/v) were prepared in a 50 mL polypropylene conical tube and then homogenized (D-500 Wiggen Hauser, Germany) at 20,000 g for 2 min. The foaming solutions were immediately transferred to a 100 mL graduated cylinder and the foam volume was recorded before and after homogenization. FC was expressed as the increase in volume (%) after homogenization. To obtain FS, the graduated cylinder was allowed to stand at room temperature, and the foam volume was recorded after 20, 40, 60, 80, and 100 min. FC and FS were calculated as follows:

$$\text{FC (\%)} = [(\text{volume after homogenization} - \text{volume before homogenization}) / \text{volume before homogenization}] \times 100$$

$$\text{FS (\%)} = [(\text{volume of foam after standing}) / (\text{initial foam volume})] \times 100$$

Statistical analysis

All data are presented as mean values \pm standard deviations (SD). The data were analyzed using SAS[®] program,

(versions 9.1; SAS Institute Inc., USA) and analysis of variance was performed following the ANOVA procedures. Duncan's multiple range test was used to determine the difference of means, with $p < 0.05$ as the cut off for statistical significance.

Results and Discussion

Proximate composition

The proximate composition is a fundamental analysis used to evaluate the essential nutrient content of food. The chemical composition of CBL, DBLSC-CO₂, and DBL-OS is shown in Table 1. CBL was higher in fat content than DBLSC-CO₂ and DBL-OS, whereas it was found that all DBLSC-CO₂ samples had significantly higher ($p < 0.05$) protein, carbohydrate, and fiber contents than CBL and DBL-OS. It was also observed that the protein, carbohydrate, and fiber contents among the DBLSC-CO₂ samples prepared using different pressures were not significantly different ($p > 0.05$). Sparks *et al.* (2006) reported similar results that the SC-CO₂-defatted rice bran had higher protein and ash contents than rice bran defatted using hot organic solvent. Khorshid *et al.* (2007) postulated that maximum recovery of soy protein was achieved using a pressurized CO₂ extraction process. Stahl *et al.* (1980) confirmed that defatting using SC-CO₂ had no deleterious effects on the nutritional value of oil seeds. Pariser *et al.* (1978) demonstrated that protein denaturation is associated with conventional liquid solvents because of the removal of lipids.

Extraction yield of lipid

Extraction yield (%) of lipid from CBL was calculated after extraction using the SC-CO₂ and organic solvent methods. The amount of extracted lipid when using the SC-CO₂ method was found to be significantly higher ($p < 0.05$) than organic solvent method, and the yield increased in line with increasing operating pressure between 200 and 450 bar (Fig. 2). Sparks *et al.* (2006) reported similar res-

Table 1. Proximate composition of CBL, DBLSC-CO₂, and DBL-OS samples

Components (%)	CBL	DBLSC-CO ₂			DBL-OS	
		200 bar	300 bar	450 bar	Ether	CM
Moisture	4.78 \pm 0.34 ^{b1}	5.96 \pm 1.23 ^{ab}	6.78 \pm 1.46 ^{ab}	6.49 \pm 0.96 ^{ab}	7.81 \pm 0.23 ^a	7.43 \pm 2.46 ^a
Protein	49.26 \pm 1.93 ^c	59.99 \pm 1.18 ^a	60.04 \pm 3.56 ^a	60.99 \pm 2.40 ^a	55.19 \pm 1.22 ^b	57.06 \pm 1.66 ^{ab}
Fat	32.09 \pm 0.60 ^a	5.04 \pm 1.38 ^c	4.49 \pm 0.38 ^c	3.08 \pm 0.36 ^c	9.50 \pm 0.30 ^b	9.52 \pm 2.47 ^b
Ash	11.61 \pm 0.64 ^{bc}	10.38 \pm 2.87 ^c	11.01 \pm 0.34 ^{bc}	11.57 \pm 2.25 ^{bc}	14.86 \pm 0.21 ^b	12.97 \pm 0.27 ^{ab}
Carbohydrate and fibers	2.26	18.63	17.68	17.87	12.64	13.02

¹ All values are mean \pm SD (n=3). Different letters indicate that means are significantly different ($p < 0.05$) by Duncan's test.




Fig. 2. Yield of lipid using SC-CO₂ and organic solvents extraction methods (%).

ults, i.e., a higher yield of rice bran oil was observed when using SC-CO₂ than when using propane as the organic solvent. Herrero *et al.* (2006) demonstrated that SC-CO₂ has better transport properties and offers faster extraction yields than liquids. Rubio-Rodriguez *et al.* (2012) postulated that SC-CO₂ is effective in obtaining fish oil, including high amounts of ω -3 fatty acids. Thus, in the present study, lipid yields when using SC-CO₂ were closely related to other reported results.

The SDS-PAGE pattern

SDS-PAGE is the most commonly used analytical method for characterizing proteins in terms of molecular weight (Mw), peptide structure, purity, and distribution pattern. Under reducing conditions using 2-mercaptoethanol, the protein patterns of CBL, DBLSC-CO₂, and DBL-OS were observed (Fig. 3). The staining intensity of the protein bands of the DBLSC-CO₂ samples was shown to be similar to that of CBL (Fig. 3A; lane 2, and lanes 3, 4, and 5), whereas the DBL-OS samples appeared very poor or no bands intensity for proteins, particularly in the 50 to 75 kDa and 20 to 25 kDa Mw ranges (Fig. 3B; lanes 3 and 4). The protein content of an electrophoretic band depends on the area and intensity of the band. These parameters indicate that DBLSC-CO₂ had no protein denaturation and had a similar protein content (including quality) to CBL. Similarly, Asaduzzaman and Chun (2015) demonstrated that the gel banding patterns of the protein of SC-CO₂-treated mackerel residues were more prominent than those in n-hexane-treated samples. Uddin *et al.* (2009) and




Fig. 3. The sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) patterns of different bovine liver samples. (A) Lane-1; Molecular weight standard (M), Lane-2; Control bovine liver (CBL), Lane-3; Defatted bovine liver by SC-CO₂ (DBLSC-CO₂) at 200 bar; Lane-4; DBLSC-CO₂ at 300 bar; Lane-5; DBLSC-CO₂ at 450 bar; (B) Lane-1; Molecular weight standard (M), Lane-2; CBL; Lane-3; Defatted bovine liver by organic solvent (DBL-OS) using ether; Lane-4: DBL-OS using chloroform-methanol.

Abdelkarder *et al.* (2012) reported that no protein denaturation was found in SC-CO₂-treated squid viscera and krill samples. Park *et al.* (2008) noted the similar intensity of the electrophoretic patterns of SC-CO₂-treated and control mackerel viscera samples.

Amino acid composition

Amino acids are essential nutrients associated with the growth of infants and strengthen the immunity of the human body. The amino acid compositions of CBL, DBLSC-CO₂, and DBL-OS are listed in Table 2. Seventeen amino acids were identified in freeze-dried bovine liver samples, including large amounts of essential and nonessential amino acids. In all samples, leucine had the highest content, followed by phenylalanine, lysine, arginine, and alanine. The contents of most amino acids in DBLSC-CO₂ samples were significantly higher ($p < 0.05$) than those in DBL-OS samples. Because of the minimization of amino acid denaturation in DBLSC-CO₂, it appears to be a better source of amino acids than DBL-OS. Stahl *et al.* (1980) reported that the SC-CO₂ method had no effect on the proteins of oilseed meal. Park *et al.* (2008) demonstrated that the activity of digestive enzyme proteins and the amino acid content of mackerel viscera were increased to a greater extent after the removal of oil when using SC-CO₂ than when using organic solvent. Christianson *et al.*

Table 2. Amino acid composition of CBL, DBLSC-CO₂, and DBL-OS samples (Unit: mg/100 g)

Amino acids	Control	DBLSC-CO ₂			DBL-OS	
		200 bar	300 bar	450 bar	Ether	CM ¹
Total essential amino acids	9718.68	8033.95	8132.15	7959.40	5624.79	4035.68
Threonine	507.03 ^{a2} ±2.63	367.78 ^b ±0.21	349.34 ^c ±0.62	330.42 ^d ±2.27	260.29 ^e ±1.22	187.02 ^f ±0.26
Valine	1199.25 ^a ±3.16	910.80 ^d ±2.84	941.63 ^c ±3.16	993.85 ^b ±0.21	663.49 ^e ±2.45	452.52 ^f ±2.15
Methionine	491.64 ^a ±1.00	448.43 ^b ±2.94	435.29 ^c ±3.26	399.58 ^d ±0.02	301.97 ^e ±2.51	215.31 ^f ±2.21
Isoleucine	742.43 ^a ±0.36	562.88 ^d ±0.39	585.93 ^c ±1.95	625.82 ^b ±1.49	461.82 ^e ±1.29	290.70 ^f ±1.69
Leucine	2059.96 ^a ±2.51	1571.41 ^d ±0.28	1675.56 ^c ±0.39	1729.67 ^b ±0.19	1003.82 ^e ±0.21	862.16 ^f ±0.35
Phenylalanine	1165.55 ^a ±3.37	912.10 ^d ±2.04	984.65 ^c ±2.10	1001.13 ^b ±4.00	629.12 ^e ±1.95	503.07 ^f ±0.54
Histidine	495.06 ^a ±2.62	432.45 ^b ±0.05	412.63 ^c ±1.39	375.38 ^e ±2.81	406.56 ^d ±1.22	195.96 ^f ±2.09
Lysine	1649.63 ^a ±2.15	1505.79 ^b ±1.69	1443.20 ^c ±0.33	1360.90 ^e ±3.20	1122.59 ^d ±2.43	717.92 ^f ±2.29
Arginine	1408.13 ^a ±3.16	1322.31 ^b ±3.30	1303.92 ^c ±0.32	1142.65 ^d ±2.45	775.13 ^e ±2.64	611.01 ^f ±1.43
Total nonessential amino acids	6263.24	5269.69	5192.50	5073.43	4034.89	3184.98
Aspartic acid	132.24 ^a ±2.61	63.79 ^b ±0.11	62.38 ^b ±0.46	52.76 ^d ±1.32	60.03 ^c ±1.03	59.81 ^c ±2.11
Serine	71.66 ^a ±1.37	40.15 ^b ±3.10	33.40 ^c ±2.63	29.18 ^e ±1.44	24.68 ^d ±2.11	19.80 ^e ±0.24
Glutamic acid	13.73 ^a ±2.12	8.55 ^b ±1.09	7.69 ^b ±2.10	6.93 ^b ±2.87	7.02 ^b ±1.28	7.49 ^b ±3.41
Proline	1371.24 ^a ±1.11	1136.67 ^c ±1.00	1138.77 ^c ±4.10	1210.28 ^b ±2.92	609.66 ^e ±2.11	609.66 ^e ±2.11
Glycine	833.94 ^a ±2.01	675.61 ^c ±2.21	620.10 ^e ±2.08	605.42 ^f ±1.05	730.40 ^b ±1.47	673.05 ^d ±3.33
Alanine	2707.15 ^a ±2.17	2335.73 ^b ±2.01	2328.59 ^c ±4.12	2225.55 ^d ±2.74	1420.21 ^e ±2.19	1106.66 ^f ±3.25
Cysteine	299.02 ^a ±2.96	276.97 ^d ±2.06	278.58 ^c ±0.49	291.87 ^b ±1.93	230.52 ^e ±1.97	141.80 ^f ±2.40
Tyrosine	834.26 ^a ±1.41	732.22 ^b ±3.83	722.99 ^c ±2.01	651.44 ^d ±3.26	640.58 ^e ±2.74	566.71 ^f ±2.57
Total amino acids	13532.60	13573.00	13573.00	12555.52	9659.68	7220.66

¹Chloroform/methanol (2:1, v/v).

²All values are mean±SD (n=3). Different letters indicate that means are significantly different ($p<0.05$) by Duncan's test.

(1984) postulated that peroxidase activity in corn germ flour defatted using SC-CO₂ was decreased to one-tenth than that of hexane-extracted flour.

Functional properties of bovine liver

The functional properties of bovine liver were observed using samples of CBL, DBLSC-CO₂ (at a pressure of 300 bar), and DBL-OS (ether).

Water and oil absorption capacity

Fluid retention is an index of the ability of proteins to absorb and retain water. Protein and carbohydrates are the major chemical components that support the WAC and OAC of food. In this study, DBLSC-CO₂ (3.25 g/g) was found to give superior WAC over CBL (3.01 g/g) and DBL-OS (3.07 g/g), as shown in Fig. 4. Friedman (1996) postulated that the degree of protein denaturation was an important factor in terms of WAC. Siwaporn *et al.* (2008) demonstrated that a lipid content of flour limit the WAC. Chandi and Sogi (2007) noted that a WAC of 1.49 to 4.72 (g/g) was necessary for viscous foods such as soup. The OAC is another important functional property, influencing mouthfeel and flavor retention (Kinsella, 1976). Abdalbasit *et al.* (2010) reported that high oil absorption is es-

sential, particularly for sausages, cake batters, mayonnaises, and salad dressings. It was found that DBLSC-CO₂ (2.04 g/g) had a significantly higher ($p<0.05$) OAC than CBL (1.91 g/g) and DBL-OS (1.82 g/g), as shown in Fig. 4. Ogunwolu *et al.* (2009) demonstrated that the protein content of cashew nut and soy protein products influenced the OAC. Siwaporn *et al.* (2008) reported that defatted macadamia flour exhibited a greater OAC than control




Fig. 4. Water absorption capacity (WAC) and oil absorption capacity (OAC) of different bovine liver (g/g) samples.

samples. Therefore, the higher WAC and OAC suggest that protein denaturation was minimized in DBLSC-CO₂, thus producing a better quality defatted bovine liver than DBL-OS.

Total soluble content

Solubility is the most superior functional property, affecting the functionality of most other food applications. Kinsella *et al.* (1976) reported that protein solubility widely influences the functionalities of the emulsification, foaming, and gelation properties. Konak *et al.* (2014) demonstrated that in alkaline conditions, the solubility of CO₂-treated oat proteins was significantly greater than in acidic conditions. The current study was conducted under neutral conditions at 80°C for 30 min, and it was found that DBLSC-CO₂ (30.59%) had significantly higher ($p < 0.05$) soluble contents than CBL (26.44%) and DBL-OS (26.21%), as shown in Fig. 5. Similarly, Konak *et al.* (2014) demonstrated that SC-CO₂-defatted oat flour had a greater soluble protein content than flours produced using other processing methods. Asaduzzaman and Chun (2015) reported higher protein contents in the water soluble extracts of SC-CO₂ defatted mackerel residues than in samples extracted using hexane as the organic solvent. Adebowale *et al.* (2005) postulated that the better solubility of defatted flour proteins from the *Mucuna* species was an indicator that they had promising food applications.

Emulsifying properties

Emulsifying property is an important functionality that depends on chemical composition, e.g., protein and poly-




Fig. 5. Total soluble contents of different bovine liver samples (%).

saccharide contents. The emulsifying activity of DBLSC-CO₂ (65.62%) was significantly higher ($p < 0.05$) than DBL-OS (60.02%), but was not significantly different ($p > 0.05$) from that of CBL (65.38%). In addition, it was observed that CBL (96.11%) and DBLSC-CO₂ (95.64%) expressed similar ES values, which were significantly higher ($p < 0.05$) than that of DBL-OS (91.50%), as shown in Fig. 6. Siwaporn *et al.* (2008) reported that there was no significant difference ($p > 0.05$) between the emulsifying properties of totally defatted macadamia flour and those of partially defatted flour, and that the presence of fat had no effect on emulsifying properties. Sun *et al.* (2008) postulated that SC-CO₂-defatted canola meal (65%) had significantly higher EC than hexane-extracted canola pressed meal (59%), and it was suggested that SC-CO₂ extraction has potential as an emulsifier in food applications. Konak *et al.* (2014) reported that SC-CO₂-extracted oat flour showed better emulsification properties than oat flour extracted using other methods. Choi *et al.* (2010) reported that the emulsion stability of meat batter indicates the water and fat retention capacity of meat protein, that determine the quality characteristics of meat products.

Foaming properties

Foaming property is one of the major functionality in the food application of protein products. More soluble protein content in an aqueous phase enhances the formation of foam. In this study, the FC and FS of DBLSC-CO₂ (55.94% and 69.42%, respectively) were significantly higher ($p < 0.05$) than those of CBL (29.43% and 26.82%, respectively) and DBL-OS (44.86% and 46.46%, respec-




Fig. 6. Emulsifying activity (EA) and emulsion stability (ES) of CBL, DBLSC-CO₂ and DBL-OS (%).




Fig. 7. (A) Foaming capacity (FC) after homogenization and foaming stability (FS) after 1 h of homogenization (%), (B) FS reduction ratio against time when total volume was considered 100% for each samples.

tively) after homogenization, as shown in Fig. 7A. For CBL and DBL-OS samples, the reduction ratio of FS was found to be more significantly decreased over time than that of DBLSC-CO₂ samples, as shown in Fig. 7B. Sun *et al.* (2008) demonstrated that the foam volume of SC-CO₂-extracted canola meal was higher than that of samples extracted using hexane as the organic solvent. Konak *et al.* (2014) postulated that extracts from oat flour had a lower FC than samples extracted using CO₂. Therefore, our study results confirm those of previously reported studies.

Conclusion

The present study has shown that DBLSC-CO₂ is a potential source of essential nutrients and promising functionalities, making it an useful ingredient in food applications. However, it can be concluded that SC-CO₂ offers a nutritionally and environmentally friendly approach to removing oil from high-protein food sources, and it is a better substitute for conventional organic solvent extraction.

Acknowledgements

This research was supported by the Korea Institute of Planning and Evaluation for Technology in Food and Agriculture, Forestry and Fisheries, Ministry of Agriculture, Food and Rural Affairs (Project No. 315017-05-2-SB150), Korea.

References

1. Abdalbasit, A. M., Bertrand, M., and Maznah, I. (2010) Com-

parison of supercritical fluid and hexane extraction methods in extracting kenaf (*Hibiscus cannabinus*) seed oil lipids. *J. Am. Oil Chem. Soc.* **88**, 931-935.

- Abdelkarder, A. N., Kim, S. B., Lee, Y. B., and Chun, B. S. (2012) Digestive enzymes characterization of krill (*Euphausia superba*) residues deoiled by supercritical carbon dioxide and organic solvents. *J. Ind. Eng. Chem.* **18**, 1314-1319.
- Adebowale, Y. A., Deyemi, I. A. A., and Oshodi, A. A. (2005) Functional and physico-chemical properties of flours of six *Mucuna* species. *Afr. J. Biochem. Res.* **4**, 1461-1468.
- AOAC. (2000) Official methods of analysis of AOAC Intl. 17th ed, Association of Official Analytical Chemists, Washington, DC.
- Asaduzzaman, A. K. M. and Chun, B. S. (2015) Characterization of digestive enzymes from de-oiled mackerel (*Scomber japonicus*) muscle obtained by supercritical carbon dioxide and n-hexane extraction as a comparative study. *J. Food Sci. Technol.* **52**, 3494-3503.
- Chandi, G. K. and Sogi, D. S. (2007) Functional properties of rice bran protein concentrates. *J. Food Eng.* **79**, 592-597.
- Choi, Y. S., Choi, J. H., Han, D. J., Kim, H. Y., Lee, M. A., Jeong, J. Y., Chung, H. J., and Kim, C. J. (2010) Effects of replacing pork back fat with vegetable oils and rice bran fiber on the quality of reduced-fat frankfurters. *Meat Sci.* **84**, 557-563.
- Christianson, D. D., Friedrich, J. P., List, G. R., Warner, K., Bagley, E. B., Stringfellow, A. C., and Inglett, G. E. (1984) Supercritical fluid extraction of dry milled corn germ with carbon dioxide. *J. Food Sci.* **49**, 229-232.
- Devatkal, S., Mendiratta, S. K., Kondaiyah, N., Sharma, M. C., and Anjaneyulu A. S. R. (2004) Physicochemical, functional and microbiological quality of buffalo liver. *Meat Sci.* **68**, 79-86.
- Estévez, M., Ventanas, J., Cava, R., and Puolanne, E. (2005) Characterisation of a traditional Finnish liver sausage and

- different types of Spanish liver pâtés: A comparative study. *Meat Sci.* **71**, 657-669.
11. Friedman, M. (1996) Nutritional value of food proteins from different food sources: A review. *J. Agric. Food Chem.* **44**, 6-29.
 12. Herrero, M., Cifuentes, A., and Ibanez, E. (2006) Sub- and supercritical fluid extraction of functional ingredients from different natural sources: Plants, food-by-products, algae and microalgae: A review. *Food Chem.* **98**, 136-148.
 13. Hultin, H. O. (1994) Oxidation of lipids in seafoods. In: *Seafoods; chemistry, processing technology and quality*. Shahidi, F. and Botta, J. R. (eds) Springer, US, pp. 49-74.
 14. Imison, B. and Daryl, U. (2000) Adding value to essential oils and other natural ingredients. rural industries research and Development Corporation. Available from: <http://www.rirdc.gov.au/reports/EOI/00-40.pdf>. Accessed Sep. 2016.
 15. Jeong, C. H. and Shim, K. H. (2004) Quality characteristics of sponge cakes with addition of *Pleurotus eryngii* mushroom powders. *J. Korean Soc. Food Sci. Nutr.* **33**, 716-722.
 16. Jyothirmayi, T., Prabhakara Rao, P. G., and Walde, S. G. (2006) Nitrogen extractability and functional properties of defatted *Erythrina variegata* flour. *Food Chem.* **96**, 242-247.
 17. Khorshid, N., Hossain, M. M., and Farid, M. M. (2007) Precipitation of food protein using high pressure carbon dioxide. *J. Food Eng.* **79**, 1214-1220.
 18. Kinsella, J. E. (1976) Functional properties of proteins in foods: A survey. *Crit. Rev. Food Sci. Nutr.* **7**, 219-232.
 19. Kiran, E. and Zhuang, W. (1997) Miscibility and phase separation of polymers in near-and supercritical fluids. In *Supercritical fluids: Extraction and pollution prevention*. American Chemical Society, Washington, DC, pp. 2-36.
 20. Konak, U. I., Ercili-Cura, D., Sibakov, J., Sontag-Strohm, T., Certel, M., and Loponen, J. (2014) CO₂-defatted oats: Solubility, emulsification and foaming properties. *J. Cereal Sci.* **60**, 37-41.
 21. Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680-685.
 22. Li, R. R., Yu, Q. L., Han, L., and Cao, H. (2014) Nutritional characteristics and active components in liver from *Wagyu* × *Qinchuan* cattle. *Korean J. Food Sci.* **34**, 214-220.
 23. Lim, G. B., Lee, S. Y., Lee, E. K., Haam, J. S., and Kim, W. S. (2002) Separation of astaxanthin from red yeast *Phaffia rhodozyma* by supercritical carbon dioxide extraction. *Biochem. Eng. J.* **11**, 181-187.
 24. Ogunwolu, S. O., Henshaw, F. O., Mock, H. P., Santros, A., and Awonorin, S. O. (2009) Functional properties of protein concentrates and isolates produced from cashew (*Anacardium occidentale* L.) nut. *Food Chem.* **115**, 852-858.
 25. Pariser, E. R., Wallerstein, M. B., Corkery, C. J., and Brown, N. L. (1978) Fish protein concentrate, panacea for protein malnutrition? In: *International nutrition policy series*. Mass. MIT Press, Cambridge, MA, **3**, 296.
 26. Park, J. Y., Back, S. S., and Chun, B. S. (2008) Protein properties of mackerel viscera extracted by supercritical carbon dioxide. *J. Env. Biol.* **29**, 443-448.
 27. Pedroche, J., Yust, M., Lqari, H., Giron-Calle, J., Alaiz, M., Vioque, J., and Millan, F. (2004) *Brassica carinata* protein isolates: Chemical composition, protein characterization and improvement of functional properties by protein hydrolysis. *Food Chem.* **88**, 337-346.
 28. Rodriguez-Ambriz, S., Martinez-Ayala, A., Millan, F., and Davila-Ortiz, G. (2005) Composition and functional properties of *Lupinus campestris* protein isolates. *Plant Food Hum. Nutr.* **60**, 99-107.
 29. Rubio-Rodriguez, N., Sara, M., de Diego, D., Beltran, S., Jaime I., Sanz, M. T., and Jordi, R. (2012) Supercritical fluid extraction of fish oil from fish by-products: A comparison with other extraction methods. *J. Food Eng.* **109**, 238-248.
 30. Siwaporn, J., Hongsuwankul, J., and Tananuwong, K. (2008) Chemical compositions, functional properties, and microstructure of defatted macadamia flours. *J. Food Chem.* **110**, 23-30.
 31. Sparks, D., Hernandez, R., Zappi, M., Blackwell, D., and Flaming, T. (2006) Extraction of rice bran oil using supercritical carbon dioxide and propane. *JAOCS* **83**, 10.
 32. Staby, A. and Mollerup, J. (1993) Separation of constituents of fish oil using supercritical fluids: A review of experimental solubility, extraction, and chromatographic data. *Fluid Phase Equilib.* **91**, 349-386.
 33. Stahl, E., Erwin Schutz, E., and Mangold, H. K. (1980) Extraction of seed oils with liquid and supercritical carbon dioxide. *J. Agric. Food Chem.* **28**, 1153-1157.
 34. Sun, M., Xu, L., Saldana, M. D. A., and Temelli, F. (2008) Comparison of canola meals obtained with conventional methods and supercritical CO₂ with and without ethanol. *J. Am. Oil Chem. Soc.* **85**, 667-675.
 35. Uddin, M. S., Ahn, H. M., Kishimura, H., and Chun, B. S. (2009) Comparative study of digestive enzymes of squid (*Todarodes pacificus*) viscera after supercritical carbon dioxide and organic solvent extraction. *J. Biotechnol. Bioproc. Eng.* **14**, 338-344.