

Research Article



CrossMark

Open Access

생산단계 잔류허용기준 설정을 위한 딸기 중 bistrifluron과 chlorantraniliprole의 잔류 특성 연구

이재원¹, 김지윤¹, 김희곤¹, 허경진^{1,2}, 권찬혁³, 허장현^{1,2*}

1

2

3

Residual Characteristics of Bistrifluron and Chlorantraniliprole in Strawberry (*Fragaria ananassa* Duch.) for Establishing Pre-Harvest Residue Limit

Jae Won Lee¹, Ji Yoon Kim¹, Hee gon Kim¹, Kyung Jin Hur^{1,2}, Chan Hyeok Kwon³ and Jang Hyun Hur^{1,2*}
(¹School of Natural Resources and Environmental Science, College of Agriculture and Life Sciences, Kangwon National University, Chuncheon 24341, Korea, ²Environment Friendly Agricultural Products Safety Center, Chuncheon 24341, Korea, ³Food Standard Division, Food Safety Policy Bureau, Ministry of Food and Drug Safety, Chungju 28159, Korea)

Received: 1 March 2017 / Revised: 17 March 2017 / Accepted: 25 March 2017

Copyright © 2017 The Korean Society of Environmental Agriculture

This is an Open-Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Jae Won Lee

<http://orcid.org/0000-0001-7303-9617>

Jang-Hyun Hur

<http://orcid.org/0000-0003-0526-9028>

Abstract

BACKGROUND: Pesticide residue analysis is essentially required for safety evaluation of agricultural products. Bistrifluron and chlorantraniliprole have been currently considered as potentials to deeply evaluate their residues in agricultural products because they are frequently found in strawberry. This work was performed to investigate the residual patterns of bistrifluron and chlorantraniliprole in strawberry after harvest.

METHODS AND RESULTS: Strawberry was treated with bistrifluron and chlorantraniliprole 0, 1, 2, 3, 5, 7 and 10 days before harvest under greenhouse conditions. The strawberry samples were subjected to solvent and solid phase extractions followed by LC-MS/MS analysis. There coverly percentages of bistrifluron and chlorantraniliprole for tified in the control samples ranged from approximately 82 to 103% with the method limit of 0.005 mg/kg. The

concentrations of bistrifluron and chlorantraniliprole in strawberry samples decreased significantly in 10 days after treatment, giving the safety levels of 0.04 to 0.06 mg/kg at 10 days after application, as considered maximum residue limit. The half-lives of bistrifluron and chlorantraniliprole based on first order kinetics were determined to 6.3 days and 6.4 days, respectively.

CONCLUSION: Bistrifluron and chlorantraniliprole are suggested to use in strawberry 10 days before harvest to reach residual safety levels.

Key words: Biological half-life, Bistrifluron, Chlorantraniliprole, Pre-harvest residue limit (PHRL), Strawberry

서
론

가



가

*Corresponding author: Jang Hyun Hur

Phone: +82-33-257-6441; Fax: +82-33-241-6640;

E-mail: jhhur@kangwon.ac.kr

Table 1. Physicochemical properties of bistrifluron and chlorantraniliprole

	Bistrifluron	Chlorantraniliprole
Structure		
IUPAC name	1-[2-chloro-3,5-bis(trifluoromethyl)phenyl]-3-(2,6-difluorobenzoyl)urea	3-Bromo-N-[4-chloro-2-methyl-6-(methylcarbamoyl)phenyl]-1-(3-chloropyridin-2-yl)-1H-pyrazole-5-carboxamide
Mol. wt.	446.7	483.15
V.P. (mPa)	2.7 10 ⁻³ mPa (25°C)	6.3 10 ⁻⁹ mPa (20°C) 2.1 10 ⁻⁸ mPa (25°C)
logP	5.74 (pH 7.8, 22°C)	2.76
Solubility in water	<0.03 mg/L (25°C)	0.9-1.0 mg/L (20°C)

(Hwang *et al.*, 2012).MRL
2013).(Kim *et al.*,PHRL
(Lee *et al.*, 2013).

vitamin

C

(Cho *et al.*, 2004;Choi *et al.*, 2013).

가

(Nam *et al.*, 2012).

가

(Claeys *et al.*,

2008).

53

가 (MFDS, 2016).

가

가 bistrifluron chlorantraniliprole

bistrifluron chlorantraniliprole

재료 및 방법**시험농약**

bistrifluron (99.0%)

chlorantraniliprole (99.5%) Dr. Ehrenstorfer GmbH
(Germany)

bistrifluron 5% (: , ())

chlorantraniliprole 5% (: , ())

Table 1

시약, 재료 및 기구

bistrifluron chlorantraniliprole

acetonitrile dichloromethane Junsei
Chemical (Japan) GR , methanol

water Merck (Germany) GR

. Formic acid Sigma-Aldrich (USA)

sodium sulfate
anhydrous, sodium chloride Dae Jung (Korea)NH₂ SPE
cartridge (1 g) Phenomenex (USA)homogenizer (Nissei, Japan),
vacuum rotary evaporator

(Eyela, Japan)

분석기기

bistrifluron chlorantraniliprole

LC-MS/MS , HPLC
(Nanospace SI-2, Shiseido, Japan) (TSQ

Quantum ultra, Tehrmo scientific, USA)	, acetonitrile	1,000 mg/kg
Column bistrifluron Capcell Core-C ₁₈ (150×2.1 mm, 2.7 μm, Shiseido, Japan)	stock solution	
UK-C ₁₈ (100×2.0 mm, 3.0 μm, Imtakt, Japan)	0.05, 0.1, 0.2, 0.5, 1.0, 2.0 mg/kg working solution	
chlorantraniliprole	1 mL , 가 ,	(1 mL) blank
시험포장 및 농약살포	standard . Bistrifluron chlorantraniliprole	
(:) ,	1.0 μL LC-MS/MS , chromatogram	
(1)	peak matrix matched	
(2)		
2015 3 16 12	딸기 중 잔류농약 분석	
1 40 m ² , 2 38.4 m ²	bistrifluron chlorantraniliprole	
3 1 m	가 20 g acetonitrile 100	
	mL 가 homogenizer 3 , 10,000 rpm	
	acetonitrile	
(KS-PK2000, , Korea)	50 mL, 100 mL 가 dichloromethane	
. Bistrifluron chlorantraniliprole	50 mL, 50 mL 2 ,	
Table 2 .	dichloromethane 5 mL .	
시료채취	SPE-NH ₂ cartridge (1 g) , dichloromethane	
10 1 , 2	5 mL pre-washing SPE-NH ₂ cartridge (1 g)	
0 0, 1, 2, 3, 5, 7 10	5 mL loading , (dichloromethane:methanol	
2 kg	=95:5, v/v) 5 mL .	
	acetonitrile 2 mL	
	(0.2 μm membrane filter) , LC-MS/MS	
-20°C 4	MS/MS	
	Table 3 .	
Bistrifluron 및 chlorantraniliprole 표준검량선 작성	딸기 중 회수율 분석	
Bistrifluron (99.0%) 101.0 mg 100 mL	bistrifluron chlorantraniliprole	
chlorantraniliprole (99.5%) 100.5 mg 100 mL		

Table 2. Guidelines for the safe use of bistrifluron and chlorantraniliprole

Pesticide	Formulation	A.I. ^{a)} (%)	MRL ^{b)} (mg/kg)	Safe use guideline		
				PHI ^{c)} (day)	MNA ^{d)} (time)	Dilution
Bistrifluron	SC ^{e)}	5	0.5	2	2	2,000
Chlorantraniliprole	WG ^{f)}	5	1.0	2	2	2,000

^{a)}A.I.: Active ingredient, ^{b)}MRL: Maximum residue limit, established by MFDS, ^{c)}PHI: Pre-harvest interval, ^{d)}MNA: Maximum number of application, ^{e)}SC: Suspension concentrate, ^{f)}WG: Water dispersible granule

Table 3. Selected Reaction Monitoring (SRM) condition of bistrifluron and chlorantraniliprole

Compound	Precursor ion	product ion	CE	Q1 PW	Q3 PW	SRM mode
Bistrifluron	444.724	259.944	43	0.5	0.7	ESI-
		261.857	22	0.5	0.7	
		388.987	17	0.5	0.7	
Chlorantraniliprole	483.836	176.740	45	0.5	0.7	ESI+
		285.603	26	0.5	0.7	
		452.531	18	0.5	0.7	

Table 4. Linear equation of calibration curve for the quantification of the pesticide residues in strawberries

Pesticide	Linear equation	r ²
Bistrifluron	y=1678656.572x+182992.033	0.9958
Chlorantraniliprole	y=8585760.9956x-30434.9973	0.9937

Table 5. Recovery rate and MLOQ for bistrifluron and chlorantraniliprole in strawberries

Pesticide	Fortification level (mg/kg)	Recovery (%)				MLOQ (mg/kg)
		1	2	3	AVG±C.V.	
Bistrifluron	0.005	85.4	88.8	82.8	85.7±1.7	0.005
	0.05	98.5	103.9	95.9	99.5±4.3	
Chlorantraniliprole	0.005	88.0	100.2	89.4	92.5±7.2	0.005
	0.05	85.4	84.7	82.7	84.3±1.7	

bistrifluron chlorantraniliprole
 1 mL, 1 mg/kg 1 mL
 10 (0.05 mg/kg)
 30

20 g
 0.1 mg/kg
 (0.005 mg/kg),
 가 ,
 Table 4 .
 (r²)가 0.99

시험농약의 생물학적 반감기 및 생산단계 잔류허용기준
 (Pre-Harvest Residue Limit, PHRL) 산출
 (PHRL)
 first order kinetics
 , k
 (Hwang et al., 2014).

분석정량한계 및 회수율
 bistrifluron chlorantraniliprole
 0.005 mg/kg ,
 (0.005 mg/kg), 10 (0.05 mg/kg)
 가 3
 bistrifluron 82.8~103.9%, chlorantraniliprole
 82.7~100.2% , 10% (Coefficient of
 variation, C.V.) (Table 5).
 70~120%
 bistrifluron chlorantraniliprole
 (Lee, 2012).

결과 및 고찰

시험기간 중 환경조건과 딸기의 중량
 (1) (2)
 1 12.9~16.7°C,
 2 14.0~20.4°C ,
 51.0%~72.9%, 72.6%~83.4% . 1 2

bistrifluron chlorantraniliprole
 (Retention time, RT) 6.28 min., 3.94 min. .
딸기 생산단계 중 잔류량 변화에 따른 생물학적 반감기

bistrifluron
 26.21~32.02 g, 24.88~29.68 g ,
 chlorantraniliprole 24.40~31.05 g,
 24.89~30.14 g 1, 2
 가 .

bistrifluron 1 0.16 mg/kg,
 2 0.15 mg/kg , 10
 0.05 mg/kg 66.67%, 68.75%
 Chlorantraniliprole
 0.17 mg/kg , 10 1 0.04
 mg/kg, 2 0.06 mg/kg
 76.47%, 64.70% ,
 (bistrifluron: 0.5 mg/kg, chlorantraniliprole: 1.0
 mg/kg)

Bistrifluron 및 chlorantraniliprole 표준검량선 작성
 ,
 LC-MS/MS
 matrix effect
 matrix matched ,

Table 6. PHRL of bistrifluron and chlorantraniliprole in strawberries

Pesticides	PHRL (mg/kg)											
	Field	10 day	9 day	8 day	7 day	6 day	5 day	4 day	3 day	2 day	1 day	0 day
Bistrifluron	1	1.10	1.02	0.94	0.87	0.80	0.74	0.69	0.63	0.59	0.54	0.50
	2	1.23	1.13	1.03	0.94	0.86	0.78	0.72	0.66	0.60	0.55	0.50
Chlorantraniliprole	1	3.07	2.74	2.45	2.19	1.96	1.75	1.57	1.40	1.25	1.12	1.00
	2	2.29	2.10	1.94	1.78	1.64	1.51	1.39	1.28	1.18	1.09	1.00

bistrifluron 1 $y=0.1521e^{-0.1093x}$ ($r^2=0.9459$),
2 $y=0.1604e^{-0.1101x}$ ($r^2=0.9760$)

MRL

Chlorantraniliprole 1 $y=0.158e^{-0.1484x}$
($r^2=0.9570$), 2 $y=0.1649e^{-0.1076x}$ ($r^2=0.9610$)

요약

Kim (2007) 1 4.7, 2 6.4, bistrifluron (1), chlorantraniliprole (2)

2 0, 0, 1, 2, 3, 5, 7, 10

LC-MS/MS

Kim (2012) 0.005 mg/kg, bistrifluron 82.8~103.9%,
가 가 chlorantraniliprole 82.7~100.2%

5.4

, 10% (Coefficient of variation, C.V.)

bistrifluron chlorantraniliprole

1 0.16 mg/kg, 0.17 mg/kg

Lee (2009)

, 2 0.15 mg/kg, 0.17 mg/kg MRL
(bistrifluron: 0.5 mg/kg, chlorantraniliprole: 1.0 mg/kg)

, 25, 10, 1 0.05

1.27, 가 mg/kg, 0.04 mg/kg, 2 0.05 mg/kg, 0.06 mg/kg, 66.67%, 76.49%

bistrifluron chlorantraniliprole

bistrifluron

6.3, chlorantraniliprole 4.7 (1) 6.4 (

2)

가

bistrifluron chlorantraniliprole 10

(Son *et al.*, 2009).

1.10 mg/kg, 2.29 mg/kg

생산단계 잔류허용기준 설정

PHRL

MRL

bistrifluron chlorantraniliprole

Table 6

bistrifluron

MRL 0.5 mg/kg, chlorantraniliprole 1.0

mg/kg, bistrifluron 10

1 1.10 mg/kg, 2 1.23 mg/kg

Chlorantraniliprole 10

1 3.07 mg/kg, 2 2.29 mg/kg

bistrifluron chlorantraniliprole

10 1.10 mg/kg, 2.29 mg/kg

Acknowledgement

This study is supported by 2015 Research Grant from Kangwon National University, also This work was carried out by support of "Establishment of Pre-Harvest Residue limit of Pesticides for Agricultural Products in 2015", Ministry of Food and Drug Safety (MFDS).

References

Claeys, W. L., De Voghel, S., Schmit, J. F., Vromman, V.,

- & Pussemier, L. (2008). Exposure assessment of the Belgian population to pesticide residues through fruit and vegetable consumption. *Food Additives and Contaminants*, 25(7), 851-863.
- Cho, J. I., Ha, S. D., & Kim, K. S. (2004). Inhibitory effects of temperature, pH, and potassium sorbate against natural microflora in strawberry paste during storage. *Korean Journal of Food Science and Technology*, 36(2), 355-360.
- Choi, H. G., Kang, N. J., Moon, B. Y., Kwon, J. K., Rho, I. R., Park, K. S., & Lee, S. Y. (2013). Changes in fruit quality and antioxidant activity depending on ripening levels, storage temperature, and storage periods in strawberry cultivars. *Korean Journal of Horticultural Science and Technology*, 31(2), 194-202.
- Hwang, K. W., Kim, T. W., Yoo, J. H., Park, B. S., & Moon, J. K. (2012). Dissipation pattern of amisulbrom in cucumber under greenhouse condition for establishing pre-harvest residue limit. *The Korean Journal of Pesticide Science*, 16(4), 288-293.
- Hwang, J. I., & Kim, J. E. (2014). Residual Patterns of Acaricides, Etoxazole and Flufenoxuron in Apples. *The Korean Journal of Pesticide Science*, 18(2), 61-68.
- Kim, H. Y., Hwang, J. I., Lee, E. H., Jeon, Y. H., Kim, J. H., Ahn, J. W., Park, H. J., Chung, C. K., Kim, S. Y., Lee, S. H., & Kim, J. E. (2012). Residue patterns of insecticide flubendiamide by varieties of peaches. *Korean Journal of Environmental Agriculture*, 31(2), 152-156.
- Kim, K. J., Kim, D. S., Heo, S. J., Ham, H. J., & Hur, J. H. (2013). Establishment of pre-harvest residue limit (PHRL) of emamectin benzoate during cultivation of amaranth. *The Korean Journal of Pesticide Science*, 17(2), 77-83.
- Kim, K., Chang, H. R., Yang, K. W., Jeong, B. J., & Kim, Y. H. (2007). Physicochemical properties of bistrifluron, benzoylphenylurea insecticide. *Korean Journal of Environmental Agriculture*, 26(3), 254-258.
- Lee, D. Y., Kim, Y. J., Kim, S. G., & Kang, K. Y. (2013). Residual characteristics of insecticides used for oriental tobacco budworm control of paprika. *Korean Journal of Environmental Agriculture*, 32(1), 84-93.
- Lee, D. Y., Kim, Y. J., Park, M. H., Lee, S. H., Kim, S. G., Kang, N. J., & Kang, K. Y. (2013). Establishment of pre-harvest residue limit (PHRL) of fungicides azoxystrobin and difenoconazole on prunus mume fruits. *The Korean Journal of Pesticide Science*, 17(4), 307-313.
- Lee, E. Y., Noh, H. H., Park, Y. S., Kang, K. W., Lee, K. H., Park, H. K., Yun, S. S., Jin, C. W., Han, S. K., & Kyung, K. S. (2009). Residual characteristics of Neonicotinoid insecticide dinotefuran and thiacloprid in cucumber. *The Korean Journal of Pesticide Science*, 13(2), 98-104.
- Nam, M. H., Jeon, Y. N., Lee, H. C., Lee, H. D., & Kang, H. K. (2012). Comparative analysis between healthy and powdery mildew-infected plants of strawberry cultivar Seolhyang. *Research in Plant Disease*, 18(2), 80-85.
- Son, K., Im, G. J., Hong, S. M., Kim, J. B., Ihm, Y. B., Ko, H. S., & Kim, J. E. (2012). Comparison of pesticide residues in perilla leaf, lettuce and kale by morphological characteristics of plant. *The Korean Journal of Pesticide Science*, 16(4), 336-342.