



Assessment of Microbial and Radioactive Contaminations in Korean Cold Duck Meats and Electron-Beam Application for Quality Improvement

Kyung-A An^{1,2}, Yunhee Jo¹, Muhammad Sajid Arshad^{1,3}, Gui-Ran Kim¹, Cheorun Jo⁴, and Joong-Ho Kwon^{1*}

¹School of Food Science and Biotechnology, Kyungpook National University, Daegu 41566, Korea

²Hazardous Substances Analysis Division, Daegu Regional Food and Drug Administration, Daegu 42612, Korea

³Institute of Home and Food Sciences, Government College University, Faisalabad 36000, Pakistan

⁴Division of Animal and Food Biotechnology, Seoul National University, Seoul 08826, Korea

Abstract

Animal-origin food products pose serious threat to public food safety due to high microbial loads. The microbial and radioactive contaminations in commercial cold duck meat products were evaluated. Ten different lots of commercial samples (C₁-C₁₀) were classified based on type and smoking process. All samples were highly contaminated (< 4-7 Log CFU/g) with total aerobic bacteria (TAB), yeasts and molds (Y&M), and 7 samples (C₁-C₇) were positive for coliforms. Furthermore, three samples were contaminated with *Listeria monocytogenes* (C₄-C₆) and one with *Salmonella typhimurium* (C₆). No radionuclides (¹³¹I, ¹³⁷Cs, and ¹³⁴Cs) were detected in any sample. The results of DEFT (direct epifluorescent filter technique)/APC (aerobic plate count), employed to screen pre-pasteurization treatments of products, indicated that smoked samples were positive showing DEFT/APC ratios higher than 4. Notably, the samples showed a serious threat to microbial safety, thus were irradiated with electron-beam (e-beam). The D₁₀ values for *S. typhimurium* and *L. monocytogenes* were 0.65 and 0.42 kGy, respectively. E-beam application at 3 and 7 kGy resulted in reduction of initial TAB, Y&M, and coliform populations by 3 and 6 log cycles, respectively. Thus, e-beam was proven to be a good decontamination approach to improve the hygiene of cold duck meat.

Keywords cold duck meat, microbial contamination, radioactive contamination, DEFT/APC, electron beam irradiation

Introduction

The consumption of duck meat is increasing worldwide because of reduced intake of red meat, which causes many cardiovascular disorders (Adzitey *et al.*, 2012). Duck meat contains essential elements, such as selenium and iron, and is a good source of proteins (Kim *et al.*, 2016). Therefore, after chicken and turkey, the production of duck meat is increasing worldwide (Matitaputty *et al.*, 2015). In Korea, the consumption of duck meat is continuously increasing and has increased 5-folds from 1997 to 2012 (Korea Duck Association, 2014). Although, duck meat is very famous, the risk of food-borne and meat-borne pathogens is quite high due to higher levels of microbial loads, including pathogens, on a commercial scale (Adzitey *et al.*, 2012). Therefore, use of some technologies to assess the microbial

Received January 3, 2017
Revised March 20, 2017
Accepted March 22, 2017

*Corresponding author

Joong-Ho Kwon
School of Food Science and Biotechnology, Kyungpook National University, Daegu 41566, Korea
Tel: +82-53-950-5775
Fax: +82-53-950-6772
E-mail: jhkwon@knu.ac.kr

© This is an open access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

decontamination in food products is currently necessary.

Radioisotope is one of the contaminants can be found in food products. Radioactive materials can be found anywhere, in different kind of meat and meat products, water, and animal feed. In addition, anthropogenic radioisotopes, such as ^{134}Cs and ^{137}Cs , might be released into the atmosphere (water or air) in normal practice or found on a higher scale after an incident. (Asano *et al.*, 2001). The average level of radionuclide Cesium in meat was 0.0000005 Bq/kg before atomic bomb testing (Eisler, 2003). Apart from radioactive contamination, the microbial contamination can also be assessed by different kind of techniques. The total number of contaminating microorganisms was assessed regardless of viability by direct epifluorescent filter technique (DEFT), while aerobic plate count (APC) determines the number of viable microorganisms. The assessment of previous decontamination treatments is possible by comparing the two counts (i.e., DEFT/APC ratios) (Ahn *et al.*, 2013).

Food irradiation, without using heat and chemicals, is an effective method and has an ability to prevent food spoilage by controlling pathogenic and/or spoilage microorganisms (Kim *et al.*, 2014). Ionizing radiations enhance the shelf life and quality of meat by killing the pathogens by targeting their DNA (Akram and Kwon, 2010). Different kinds of pathogens in meat, such as *Yersinia enterocolitica*, *Escherichia coli* O157:H7, *Listeria monocytogenes*, and *Salmonella* spp., can be easily destroyed by irradiation (Cárcel *et al.*, 2015). The overall microbial quality of the

food can be easily increased by irradiation and the dose for meat is up to 7 kGy as recommended by the United States Department of Agriculture (Lung *et al.*, 2015). Irradiation is a promising technology used for preservation of meat without affecting the nutritional as well as sensory attributes (Grolichova *et al.*, 2004). There are different sources of radiations. Electron beam (e-beam) uses high-energy electrons generated from an electron accelerator machine, which is different from gamma rays from radioisotope sources in terms of safety concerns (Ahn *et al.*, 2013). The e-beam also has an ability of 5-fold reduction in *E. coli* O157:H7 in inoculated carcasses but it can be vary by different kinds of meat and meat products (Maxim *et al.*, 2014).

The main objectives of this study were to determine hygienic status of commercial cold duck meat samples available in different supermarkets from different regions in Daegu city by monitoring the radioactive contamination as well as microbial loads. In addition, e-beam sensitivity (D_{10} value) was determined from the linear regression model for the log of surviving bacterial cells and irradiation dose.

Materials and Methods

Materials

Commercial cold duck meats were purchased from three different supermarkets in different regions in Daegu, Korea. Ten different samples (1 kg/package: 3 replications)

Table 1. Conditions for enrichment, isolation, and confirmation of 9 kinds of foodborne pathogenic microorganisms

Microorganisms	Enrichment		Isolation		Confirmation
	Medium	Condition	Medium	Condition	
<i>B. cereus</i>	-	-	Mannitol egg yolk polymyxin agar	30°C, 24 h	VITEC 2 –compact
<i>C. perfringens</i>	Cooked meat agar	37°C, 24 h, anaerobic	Perfringens agar base	37°C, 24 h	VITEC 2 –compact
<i>C. jejuni</i>	Bolton broth	37°C, 5 h, microaerophilic	Modified Campy blood free agar base	42°C, 48 h, microaerophilic	VITEC 2 –compact
<i>E. coli</i> O157:H7	mTSB*	42°C, 48 h, microaerophilic	TC-SMAC*, BCIG*	37°C, 24 h	VITEC 2 –compact
<i>L. monocytogenes</i>	UVM-modified Listeria selective agar base	37°C, 24 h	Oxford agar	30°C, 24 h	API Listeria
<i>Salmonella</i> spp.	Peptone water, Rappaport-Vassiliadis	37°C, 24 h, 42°C, 24 h	MacConkey	37°C, 24 h	VITEC 2 –compact
<i>S. aureus</i>	TSB*	37°C, 24 h	Baird-Parker agar	37°C, 24 h	VITEC 2 –compact
<i>Y. enterocolitica</i>	PSBB*	10°C, 10 d	MacConkey, CIN*	30°C, 24 h	VITEC 2 –compact
<i>V. parahaemolyticus</i>	Alkaline peptone water	37°C, 24 h	TCBS*	37°C, 24 h	VITEC 2 –compact

*Modified Tryptic Soy Broth (mTSB), Peptone Sorbitol Bile Broth (PSBB), Thiosulfate-citrate-bile salts-sucrose (TCBS), MacConkey Agar with Sorbitol, Cefixime and Tellurite (TC-SMAC), 5-bromo-4-chloro-3-indolyl- β -D-glucuronide (BCIG), Cefsulodin, Irgasan, Novobiocin (CIN).

were purchased having the same manufacturing date within circulation date. Ten samples (C1-C10) were classified according to their product types (whole, boneless whole/BW, and boneless slice/BS) and smoking process (smoked and non-smoked). The samples were stored in a refrigerator (4°C) for 1 wk prior to use and all experiments were completed within each circulation dates.

Microbial and foodborne pathogen analyses

Microbial counts of the total aerobic bacteria (TAB), yeasts & molds (Y&M), and coliforms were evaluated according to the guidelines of the Association of Official Analytical Chemists (AOAC, 2000) and expressed as colony forming units (CFU)/g. Nine kinds of foodborne pathogens were analyzed according to Korea Food Code (MFDS, 2011). The condition for enrichment, isolation, and confirmation of 9 kinds of foodborne pathogens is given in Table 1. Meat sample was added to each enrichment broth and incubated at optimal conditions. Colonies isolated from each selective medium were finally identified using VITEK2 Compact (BioMerieux Ind., France).

Measurement of radioactive contamination

The commercial cold-duck meat samples were analyzed by gamma-ray spectrometry counting system by using standard analysis method suggested by Association of Analytical Communities with method number 996.05 (AOAC, 1998). The minimum detectable activity for radionuclides was quantified at 1 Bq/kg per fresh weight after considering the size and the counting time of the sample.

DEFT/APC analysis

Analysis was performed according to EN 13783 (2001) to screen whether the samples have been pre-pasteurized. Samples of 10 g were added to peptone saline and diluted 10 times. The solution was diluted by logarithmic dilution series (10^1 - 10^7). Each diluted solution was transferred to the filtration manifold tower containing 10 μ m polypropylene filter above 0.6 μ m polycarbonate filter for DEFT. After that, staining and rinsing steps were performed. For APC (aerobic plate count), 1.0 mL of each dilution was transferred to a Petri dish containing plate count agar (Difco, Laboratories, USA). The plates were incubated at 30°C for 72 h and then expressed as \log_{10} CFU/g. The DEFT count (X) per gram of meat was calculated by using the mean number of DEFT units per microscope field (N/n), the dilution factor (DF) of the sample, and the microscope factor (MF) of the sample Eq. (1).

$$X = \text{DEFT count/g} = (N \times \text{MF} \times \text{DF})/n \quad (1)$$

The DEFT count was then converted to a logarithmic value. The difference between the DEFT count and APC count is then obtained by subtracting the APC count (logarithmic value) from the DEFT count (logarithmic value).

Electron beam irradiation

E-beam irradiation was carried out using an electron accelerator (High Energy Linear Accelerator, 10 MeV, EB Tech, Korea) at doses of 0, 1, 3, and 7 kGy, which were applied for determining the reduction effect of the initial microbial populations. The absorbed doses were measured using an alanine-electron paramagnetic resonance dosimetry system, with an EMS 104 EPR analyzer (Bruker Biospin, Germany).

Determining e-beam sensitivity (D_{10} value) of pathogens

Two pathogens, *Salmonella typhimurium* (KCTC 1916) and *Listeria monocytogenes* (KCTC 3569), were grown in a tryptic soy broth (Difco, Laboratories, USA) at 30°C for 48 h. The pathogens were cultured to a cell density of approximately 10^6 - 10^7 CFU/mL levels. One gram of sterile meat samples (30 kGy) were inoculated with cell suspension (200 mL) of the two pathogens, respectively. Then, it was kept in a sterile workstation for 1 min to allow it to be absorbed. The inoculated samples in the stomacher bag were e-beam irradiated from 0 to 3.5 kGy. Ten gram sample was aseptically homogenized in a sterile stomacher bag containing 90 mL sterile saline solution. After serial dilutions, 100 μ L aliquot from an appropriate dilution was plated on to the medium. Medium used for the microbial count was tryptic soy agar (Difco, Laboratories, USA). Plates were incubated at the optimal growing temperature of the bacteria for 48 h and the CFU per gram were counted at 30-300 CFU per plate. D_{10} values (the dose required to inactivate 90% of a population) for each of the organisms tested were determined by the linear fit of the logarithmic survivors versus irradiation dose points (Kim *et al.*, 2007).

Statistical analysis

All experiments were performed in triplicate. Data were analyzed by using SPSS 19.0. Statistical significance was set to $p < 0.05$. The comparison of the means was conducted using the Duncan's Multiple Range test.

Results and Discussion

Microbial contamination of commercial cold duck meats

Ten different cold commercial duck meat samples (C₁-C₁₀) were monitored for total aerobic bacteria (TAB), yeasts & molds (Y&M), and coliform counts, which ranged from 5.18 to 7.23 Log CFU/g, 4.85 to 6.56 Log CFU/g, and 3.72 to 6.04 Log CFU/g, respectively, as shown in Table 2. TAB and Y&M posed a serious microbial threat, and maximum microbial contamination was found in the BS (C₄-C₆) duck meat samples. Seven different samples (C₁-C₇) were highly contaminated with coliforms, whereas the smoked BW (C₈-C₁₀) samples were not contaminated. The contamination level in duck meat has been studied by many researchers. Kim *et al.* (2016) reported that the refrigerated whole raw duck meat showed lower TAB than that showed by sliced duck meat. The results for TAB and coliforms of whole duck were also in consistent with the findings of Chae *et al.* (2006) and Sung *et al.* (2013). Szosland-Faltyn *et al.* (2014) suggested that yeasts and molds in whole duck meat were up to 3.77 and 4.45 Log CFU/g, respectively.

The nine foodborne pathogens were also monitored from different cold duck meat samples (Table 2). The pathogen, *L. monocytogenes*, was detected in all BS duck meat samples (C₄-C₆), while *Salmonella* spp. was only found in BS (C₆). The other pathogens were not detected in all the

10 different cold duck meat samples. Haslia *et al.* (2015) depicted that *L. monocytogenes* and *Salmonella* spp. pathogens were detected in the raw duck meat products, which is consistent with the results of the present study. Some other studies are also consistent with the present study and demonstrated that some samples out of 32 cooked and raw meat samples showed contamination with *Salmonella* spp. (Jalali *et al.*, 2008). Szosland-Faltyn *et al.* (2014) reported that about 25% and 6% of *L. monocytogenes* and *Salmonella* spp. pathogens, respectively, were present in the duck meat sample. Higher contamination of microorganisms, including *L. monocytogenes* and *Salmonella* spp., in commercial cold duck meat may be very lethal for the safety status of meat and in return may seriously threaten the human health.

Radioactive contamination

The contents of radionuclides, such as ¹³¹I, ¹³⁷Cs, and ¹³⁴Cs, from different commercial cold duck meat samples are given in Table 3. It is evident from the results that there were no radionuclides detected from any of the 10 different samples. The acceptable limit for each radionuclide in foods is different. The maximum permitted concentration of ¹³⁴Cs and ¹³⁷Cs is less than 100 Bq/kg in general foods according to the Korean Food Code (MFDS, 2011). According to the Ministry of Agriculture, Forestry, and Fisheries, Japan (MAFF), concentration less than 10 Bq/kg in drinking water, less than 50 Bq/Kg in milk and infant

Table 2. Sample classification and microbial quality of 10 kinds of commercial cold duck meat products

Sample Classification	Sample No.									
	C ₁	C ₂	C ₃	C ₄	C ₅	C ₆	C ₇	C ₈	C ₉	C ₁₀
Type ¹⁾	W	W	W	BS	BS	BS	BW	BW	BW	BW
Smoking Process	None	None	None	None	None	None	Smoked	Smoked	Smoked	Smoked
Hygiene microorganism ²⁾										
Total aerobic bacteria	6.04	5.65	7.00	6.98	7.23	7.08	5.88	5.54	5.18	5.79
Yeasts & molds	4.90	4.87	4.85	6.08	6.15	6.56	5.69	5.67	-	5.89
Coliforms	4.90	3.72	6.04	5.70	5.86	5.92	4.95	-	-	
Foodborne pathogen microorganism ²⁾										
<i>B. cereus</i>	-	-	-	-	-	-	-	-	-	-
<i>C. perfringens</i>	-	-	-	-	-	-	-	-	-	-
<i>C. jejuni</i>	-	-	-	-	-	-	-	-	-	-
<i>E. coli</i> O157:H7	-	-	-	-	-	-	-	-	-	-
<i>L. monocytogenes</i>	-	-	-	+	+	+	-	-	-	-
<i>Salmonella</i> spp.	-	-	-	-	-	+	-	-	-	-
<i>S. aureus</i>	-	-	-	-	-	-	-	-	-	-
<i>Y. enterocolitica</i>	-	-	-	-	-	-	-	-	-	-
<i>V. parahaemolyticus</i>	-	-	-	-	-	-	-	-	-	-

¹⁾W: whole; BS: boneless sliced; BW: boneless whole. ²⁾Log CFU/g. Values are means of triplicate experiments; +: Positive; -: Negative.

Table 3. Presence of radionuclides in commercial cold duck meat products

Sample No.	Radionuclides (Bq/kg fresh weight)	
	^{131}I	$^{134+137}\text{Cs}$
C ₁	- ¹⁾	-
C ₂	-	-
C ₃	-	-
C ₄	-	-
C ₅	-	-
C ₆	-	-
C ₇	-	-
C ₈	-	-
C ₉	-	-
C ₁₀	-	-

¹⁾-: Negative (below the minimum detectable level).

foods, and less than 100 Bq/Kg in meat and fish are acceptable for human consumption (Manabe *et al.*, 2016). The results are somehow consistent with the findings of Brandhoff *et al.* (2016), who reported that different food samples, including dairy and meat, were subjected to the analysis of selected radionuclides in meat but are less than the acceptable limit. The results were further supported by the findings of Miyazaki *et al.* (2013) who depicted that about 300 samples were determined for the presence of radioactive iodine (^{131}I) and cesium (^{137}Cs and ^{134}Cs) and it was found that only few samples showed some contamination, but below the regulatory limits.

DEFT/APC

DEFT shows the number of both viable and non-viable cells from the samples, while APC shows the number of viable cells only from the sample (Akram *et al.*, 2012). DEFT/APC technique confirms the microbial decontamination by some previous processing operations. When there is no decontamination treatment, the DEFT count

Table 4. DEFT/APC results of commercial cold duck meat samples

Sample No.	DEFT	APC	D/A ratio
C ₁	9.48	6.04	3.44
C ₂	9.41	5.65	3.76
C ₃	10.37	7.00	3.37
C ₄	10.50	6.98	3.52
C ₅	10.55	7.23	3.32
C ₆	10.10	7.08	3.02
C ₇	10.46	5.88	4.58
C ₈	11.31	5.54	5.77
C ₉	11.35	5.15	6.20
C ₁₀	11.27	5.79	5.48

can be similar to that of APC. However, mostly the APC count will be lesser than the DEFT count after decontamination treatments because heat or non-heat processes have an ability to inhibit the growth of viable microorganisms (Ahn *et al.*, 2013). If DEFT/APC ratio is higher than 4, it can be an indication of some processing treatment (EN 13783, 2001). The DEFT/APC results of commercial cold duck meat products are shown in Table 4.

The results showed that DEFT count (9.41-11.35) of commercial duck meat samples was higher than the APC (5.15-7.08). The ratios of DEFT/APC for commercial duck meat samples ranged from 3.02 to 6.20. The cold duck meats (C₁-C₆) showed DEFT/APC ratios below 3.76; however, the smoked duck meat samples (C₇-C₁₀) presented DEFT/APC ratio higher than 4. Higher DEFT/APC ratios were demonstrated by smoked duck meat samples constituting significantly higher difference in microbial contamination, which was attributed to pre-pasteurization process, such as smoking (EN 13783, 2001). Osman *et al.* (2013) reported that DEFT/APC is a microbiological screening method for irradiation treatments based on comparison of counts. They discriminated the fresh and deboned chicken meat samples based on microbial population. Previously, Sommers and Boyd (2006) reported that this method clearly distinguishes the samples having different level of APC among different samples, which is consistent with the findings of our study where different duck meat samples have different value of APC and DEFT.

E-beam D₁₀ values for *Salmonella typhimurium* and *Listeria monocytogenes*

The microbial qualities of 10 (C₁-C₁₀) commercial cold duck meat samples were evaluated and the sample BS (C₆) showed presence of *Salmonella typhimurium* and *Listeria monocytogenes* among all the samples as shown in Table 2. We calculated D₁₀ values from graphical analysis and regression equation and determination of D₁₀ values was carried out in accordance with the reported method of Kortei *et al.* (2014). The sample BS (C₆) was designated as model sample and treated with different doses (0.5, 1, 1.5, 2, 2.5, 3, 3.5 kGy) of e-beam and D₁₀ values (kGy) of *S. typhimurium* and *L. monocytogenes* were determined. The results showed that the D₁₀ values for *S. typhimurium* and *L. monocytogenes* were 0.65 kGy and 0.42 kGy, respectively. The regression (R²) for *L. monocytogenes* and *S. typhimurium* was 0.9859 and 0.9615, respectively. The level of contamination decreased with increasing the dose level of e-beam irradiation. *L. monocytogenes* could not




Fig. 1. D₁₀ values (kGy) of *Salmonella typhimurium* and *Listeria monocytogenes* inoculated into commercial cold duck meats.

be detected after increasing the dose to 2 kGy, but *Salmonella* was detected even up to 3.5 kGy as shown in Fig. 1. The results are consistent with the findings of Medina *et al.* (2009) who reported that the loads of *L. monocytogenes* in cold smoked salmon were reduced with increasing e-beam irradiation. In another study, application of 3 kGy e-beam irradiation resulted in the reduction of the *Salmonella* load in beef by 2-4 Log CFU/g, which supports findings of the present study (Li *et al.*, 2015).

Microbial reduction by electron beam irradiation

The microbial quality upon e-beam irradiation of the BS (C₆) cold duck meat is shown in Fig. 2. Four doses (0, 1, 3, 7 kGy) of e-beam were applied to the BS (C₆) cold duck meat sample and the findings indicated that TAB, Y&M, and coliforms were significantly reduced from 9.36 to 3.13 Log CFU/g, 9.19 to 3.00 Log CFU/g, and 3.38 Log CFU/




Fig. 2. Microbial qualities of commercial cold duck meat by e-beam irradiation.

g to the non-detected level, respectively. It is obvious that the initial microbial loads of TAB in the meat sample were effectively reduced in a dose-dependent manner and similar patterns were observed for Y&M and coliforms. Lewis *et al.* (2002) reported that the use of e-beam at 1.8 kGy significantly reduced the total plate count from 4.60 to 1.62 Log CFU/200 mL in poultry meat. This showed that e-beam irradiation significantly reduced the TAB count. The results are in agreement with the findings of Kim *et al.* (2013) who reported that application of e-beam dose to pork jerky led to the reduction in TAB count from 4.54 to 2.81 Log CFU/g. Recently, the results were further supported by the reports of An *et al.* (2017) who depicted that there was reduction in TAB and coliforms were obtained in duck meat.

Conclusions

This study aimed to assess the hygienic quality of commercial cold duck meat samples by determining their microbial and radioactive contaminations. The status of pre-pasteurization treatment was also confirmed by counting the DEFT/APC ratios of each sample. The 10 commercial cold duck meat samples have serious threat of contamination with TAB and YAM (4-7 Log CFU/g). The pathogens, *S. typhimurium* and *L. monocytogenes*, were also detected in 3 kinds of duck meat samples. However, there was no radioactive contamination in all commercial duck meat samples. The smoked samples presented DEFT/APC ratios higher than 4, which was due to pre-pasteurization process. E-beam of 7 kGy reduced 6 log cycles of TAB and Y&M populations in a model sample. Conclusively, the commercial cold duck meat samples have a serious threat of microbial contamination, and e-beam irradiation showed a potential to be used for the improvement of microbiological quality of commercial duck meat products. It is recommended that relevant perspective studies should be conducted in future for commercial duck meat products.

References

- Adzitey, F., Huda, N., and Rusul, G. (2012) Prevalence and antibiotic resistance of *Campylobacter*, *Salmonella*, and *L. monocytogenes* in ducks: A review. *Foodborne Pathog. Dis.* **9**, 498-505.
- Ahn, D. U., Kim, I. S., and Lee, E. J. (2013) Irradiation and additive combinations on the pathogen reduction and quality of poultry meat. *Poult. Sci.* **92**, 534-545.

3. Ahn, J. J., Akram, K., Kwak, J. Y., Jeong, M. S., and Kwon, J. H. (2013) Reliable screening of various foodstuffs with respect to their irradiation status: A comparative study of different analytical techniques. *Rad. Phys. Chem.* **91**, 186-192.
4. Akram, K., Ahn, J. J., and Kwon, J. H. (2012) Analytical methods for the identification of irradiated foods. In: *Ionizing radiation: Applications, sources and biological effects*. Belotserkovsky, E. and Ostaltsov, Z. (eds) NOVA publishers, NY, pp. 1-36.
5. Akram, K. and Kwon, J. H. (2010) Food irradiation for mushrooms: A review. *J. Kor. Soc. App. Biol. Chem.* **53**, 257-265.
6. An, K. A., Arshad, M. S., Jo, Y., Chung, N., and Kwon, J. H. (2017) E-beam irradiation for improving the microbiological quality of smoked duck meat with minimum effects on physicochemical properties during storage. *J. Food Sci.* **82**, 865-872.
7. AOAC. (1998) In P. Cunniff (Ed.), *Official methods of analysis* (16th ed.). USA: Association of Official Analytical Chemists.
8. AOAC. (2000) *Official methods of analysis*. 17th ed, Association of Official Analytical Chemists, Washington, DC.
9. Asano, T., Sato, K., and Onodera, J. I. (2001) United nations scientific committee on the effects of atomic radiation 2000 report. *Hoken Butsuri* **36**, 149-158.
10. Brandhoff, P. N., van Bourgondiën, M. J., Onstenk, C. G. M., van Avezathe, A. V., and Peters, R. J. B. (2016) Operation and performance of a national monitoring network for radioactivity in food. *Food Control* **64**, 87-97.
11. Cárcel, J. A., Benedito, J., Cambero, M. I., Cabeza, M. C., and Ordóñez, J. A. (2015) Modeling and optimization of the E-beam treatment of chicken steaks and hamburgers, considering food safety, shelf-life, and sensory quality. *Food Bioprod. Process.* **96**, 133-144.
12. Chae, H. S., Yoo, Y. M., Ahn, C. N., Jeong, S. G., Ham, J. S., Lee, J. M., and Singh, N. K. (2006) Effect of singeing time on physico-chemical characteristics of duck meat. *Kor. J. Poultr. Sci.* **33**, 273-281.
13. Eisler, R. (2003) The Chernobyl nuclear power plant reactor accident: ecotoxicological update. In: *Handbook of ecotoxicology*. 2nd ed. Hoffman, D. J., Rattner, B. A., Burton Jr, G. A., and Cairns Jr, J. (Ed). Lewis publisher, FL, pp. 703-736.
14. EN 13783 (2001) Foodstuffs - Detection of irradiated food using direct epifluorescent filter technique/aerobic plate count (DEFT/APC) - screening method. European Committee of Standardization (CEN), Brussels, Belgium.
15. Grolichova, M., Dvoak, P., and Musilova, H. (2004) Employing ionizing radiation to enhance food safety. *Acta. Vet. Brno.* **73**, 143-149.
16. Haslia, F., Adzitey, F., Huda, N., and Ali, G. R. R. (2015) Effect of temperature on the growth and survival of pathogens in duck and quail meatballs. *J. Life Sci. Biomed.* **5**, 48-52.
17. Jalali, M., Abedi, D., Pourbakhsh, S. A., and Ghoukasin, K. (2008) Prevalence of *Salmonella* spp. in raw and cooked foods in Isfahan-Iran. *J. Food Saf.* **28**, 442-452.
18. Kim, D., Song, H., Lim, S., Yun, H., and Chung, J. (2007) Effects of gamma irradiation on the radiation-resistant bacteria and polyphenol oxidase activity in fresh kale juice. *Rad. Phys. Chem.* **76**, 1213-1217.
19. Kim, H. J., Kang, M., Yong, H. I., Bae, Y. S., Jung, S., and Jo, C. (2013) Synergistic effects of electron-beam irradiation and leek extract on the quality of pork jerky during ambient storage. *Asian-Austr. J. Anim. Sci.* **26**, 596-602.
20. Kim, H. J., Yong, H. I., Lee, H. J., Jung, S., Kwon, J. H., Heo, K. N. and Jo, C. (2016) Identification of microorganisms in duck meat products available in Korea and the effect of high hydrostatic pressure. *Kor. J. Food Sci. Anim. Res.* **36**, 283-288.
21. Kim, H. W., Choi, J. H., Choi, Y. S., Kim, H. Y., Lee, M. A., Hwang, K. E., and Kim, C. J. (2014) Effects of *Kimchi* and smoking on quality characteristics and shelf life of cooked sausages prepared with irradiated pork. *Meat Sci.* **96**, 548-553.
22. Korea Duck Association. (2014) General Statistics: Per capita duck meat consumption. Available from: <http://www.korea-duck.org/>
23. Kortei, N. K., Odamtten, G. T., Appiah, V., Obodai, M., Adu-Gyamfi, A., Annan, T. A., and Mills, S. W. O. (2014). Microbiological quality assessment of gamma irradiated fresh and dried mushrooms (*Pleurotus ostreatus*) and determination of D10 values of *Bacillus cereus* in storage packs. *Eur. J. Biotechnol. Biosci.* **2**, 28-34.
24. Lewis, S. J., Velasquez, A., and Cuppett, S. L. (2002) Effect of electron beam irradiation on poultry meat safety and quality. *Poult. Sci.* **81**, 896-903.
25. Li, S., Kundu, D., and Holley, R. A. (2015) Use of lactic acid with electron beam irradiation for control of *Escherichia coli* O157: H7, non-O157 VTEC *E. coli*, and *Salmonella serovars* on fresh and frozen beef. *Food Microbiol.* **46**, 34-39.
26. Lung, H. M., Cheng, Y. C., Chang, Y. H., Huang, H. W., Yang, B. B., and Wang, C. Y. (2015) Microbial decontamination of food by electron beam irradiation. *Trends Food Sci. Technol.* **44**, 66-78.
27. Manabe, N., Takahashi, T., Endo, M., Piao, C., Li, J., Kokado, H., and Nakanishi, T. M. (2016) Effects of "clean feeding" management on livestock products contaminated with radioactive Cesium due to the Fukushima Daiichi nuclear power plant accident. In: *Agricultural implications of the Fukushima nuclear accident: The first three years*. Nakanishi, T. M. and Tanoi, K. (ed.) Springer, Japan. pp. 77-90.
28. Matitaputty, P. R., Wijaya, C. H., Bansi, H., Laudadio, V., and Tufarelli, V. (2015) Influence of duck species and cross-breeding on sensory and quality characteristics of Alabio and Cihateup duck meat. *CyTA-J. Food.* **13**, 522-526.
29. Maxim, J. E., Neal, J. A., and Castillo, A. (2014) Development of a novel device for applying uniform doses of electron beam irradiation on carcasses. *Meat Sci.* **96**, 373-378.
30. Medina, M., Cabeza, M. C., Bravo, D., Cambero, I., Montiel, R., Ordóñez, J. A., and Hoz, L. (2009) A comparison between E-beam irradiation and high pressure treatment for cold-smoked salmon sanitation: microbiological aspects. *Food Microbiol.* **26**, 224-227.
31. MFDS. (2011) Korea Food Standard Code. Korea Food &

- Drug Administration. Seoul. Korea.
32. Miyazaki, H., Tsuchiyama, T., and Terada, H. (2013) Examination of radioactive contamination in foods. *J. Food Hygienic Soc. Jap.* **54**, 156-164.
 33. Osman, M. E., Abo El-Nasr, A., Abo El Nour, S. A., Hammad, A., and Ibrahim, H. M. A. (2013) Microbiological method (DEFT/APC) for the detection of irradiated foods in Egypt. *Isot. Radiat. Res.* **45**, 531-544.
 34. Sommers, C. H. and Boyd, G. (2006) Variations in the radiation sensitivity of foodborne pathogens associated with complex ready-to-eat food products. *Radiat. Phys. Chem.* **75**, 773-778.
 35. Sung, S. H., Bae, Y. S., Oh, S. H., Lee, J. C., Kim, H. J., and Jo, C. (2013) Possibility of instrumental differentiation of duck breast meat with different processing and storage conditions. *Kor. J. Food Sci. Anim. Res.* **33**, 96-102.
 36. Szosland-Fałtyn, A., Bartodziejska, B., Królasik, J., and Paziaik-Domańska, B. (2014) Occurrence of *Campylobacter* sp. and *Salmonella* sp. in Polish duck meat in the presence of other spoilage microbial groups. *Acta Sci. Pol., Medicina Veterinaria* **13**, 27-36.