

# Serological and Molecular Detection of *Toxoplasma gondii* and *Babesia microti* in the Blood of Rescued Wild Animals in Gangwon-do (Province), Korea

Sung-Hee Hong<sup>1</sup>, Hee-Jong Kim<sup>2</sup>, Young-Il Jeong<sup>1</sup>, Shin-Hyeong Cho<sup>1</sup>, Won-Ja Lee<sup>1</sup>, Jong-Tak Kim<sup>2,\*</sup>, Sang-Eun Lee<sup>1,\*</sup>

<sup>1</sup>Division of Malaria and Parasitic Diseases, Korea National Institute of Health, Korea Center for Disease Control and Prevention, Cheongju 28159, Korea; <sup>2</sup>College of Veterinary Medicine, Gangwon National University, Chuncheon 24341, Korea

**Abstract:** Infections of *Toxoplasma gondii* and *Babesia microti* are reported in many wild animals worldwide, but information on their incidence and molecular detection in Korean wild fields is limited. In this study, the prevalence of *T. gondii* and *B. microti* infection in blood samples of 5 animal species (37 Chinese water deer, 23 raccoon dogs, 6 roe deer, 1 wild boar, and 3 Eurasian badgers) was examined during 2008-2009 in Gangwon-do (Province), the Republic of Korea (=Korea) by using serological and molecular tests. The overall seropositivity of *T. gondii* was 8.6% (6/70); 10.8% in Chinese water deer, 4.3% in raccoon dogs, and 16.7% in roe deer. PCR revealed only 1 case of *T. gondii* infection in Chinese water deer, and phylogenetic analysis showed that the positive isolate was practically identical to the highly pathogenetic strain type I. In *B. microti* PCR, the positive rate was 5.7% (4/70), including 2 Chinese water deer and 2 Eurasian badgers. Phylogenetic analysis results of 18S rRNA and the  $\beta$ -tubulin gene showed that all positive isolates were US-type *B. microti*. To our knowledge, this is the first report of *B. microti* detected in Chinese water deer and Eurasian badger from Korea. These results indicate a potentially high prevalence of *T. gondii* and *B. microti* in wild animals of Gangwon-do, Korea. Furthermore, Chinese water deer might act as a reservoir for parasite infections of domestic animals.

**Key words:** *Toxoplasma gondii*, *Babesia microti*, wild animal, zoonotic pathogen, blood, Korea

Transmission of zoonotic pathogens between animals and humans is well documented, occurring through direct routes, such as contact or consumption of meat products, or indirect routes via an intermediate vector such as an arthropod [1,2]. Wild animals also play a role in the transmission of zoonosis as intermediators, reservoirs, amplifiers, or final hosts for pathogens. Because it is hard to recognize specific clinical symptoms or representative signs of zoonosis in wild animals, continuous surveys of the prevalence of zoonotic pathogens in wild animals are important from the perspectives of public health, to serve as a warning and to establish prevention strategies for the outbreak of critical zoonotic diseases [3]. In this regard, among various types of zoonotic pathogens, focusing on parasitic infections in the wild animal population is particularly important, because parasites tend to have longer periods of infection establishment with more complex transmission

routes compared to those of bacteria and viruses [4].

*Toxoplasma gondii*, *Babesia microti*, *Dirofilaria immitis*, *Dirofilaria repens*, and *Cryptosporidium* are well known as representative zoonotic parasites [5]. Previous studies of zoonotic parasitoses in wild animals in the Republic of Korea (=Korea) have reported the prevalence of *T. gondii* and *Trichinella* in wild boars, *B. microti*-like parasites in wild raccoon dogs [6,7]. Toxoplasmosis is a zoonotic disease caused by the protozoan parasite *T. gondii* [8], and babesiosis, a well-known disease of domestic animals caused by *Babesia* spp., has attracted increased attention as an emerging zoonotic disease [9]. The prevalence of *T. gondii* infection in wild boars in Korea was found to be relatively high compared to that reported in Japan, Austria, and Germany [10]. *B. microti* infections were detected in raccoon dogs and small wild mammals in Korea [6,11,12]. Although the consumption of undercooked meat may expose humans to a high risk of *T. gondii* infection [13], tick vectors are essential for the transmission of *B. microti*, thereby requiring implementation of efficient vector control strategies [14]. In particular, several wild animals, such as raccoons, foxes, and deer, are important for the maintenance of these ticks [15].


However, surveillance of pathogen infections in wild animals

•Received 16 November 2016, revised 25 February 2017, accepted 18 March 2017.

\*Corresponding authors (kimjt@kangwon.ac.kr; ondalgl@korea.kr)

© 2017, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Fig. 1.** Location surveyed for *Toxoplasma gondii* and *Babesia microti* infections in Gangwon-do (Province), Korea.

has thus far been very limited in Korea. Therefore, the aim of the present survey was to investigate the prevalence of *T. gondii* and *B. microti* among wild animals in Gangwon-do (Province), via evaluation of blood samples provided by the Gangwon branch of the Korea Wildlife Rescue and Management Association in 2008-2009, as a representative mountainous region in Korea. The results of this study will prove to be valuable for evaluating the risk of exposure to these parasites from wild animals to humans and other animals in Korea.

Total 70 blood samples of 5 animal species (37 Chinese water deer, 23 raccoon dogs, 6 roe deer, 1 wild boar, and 3 Eurasian badgers) were collected from the jugular vein of each animal in Gangwon-do (Fig. 1). Sera were separated by centrifugation at 2,000 g for 5 min. Genomic DNA was also extracted from blood samples using a DNeasy tissue kit (Qiagen, Hilden, Germany) according to the manufacturer instructions. Genomic DNA was resolved in 100 µL Tris-EDTA buffer and stored at -20°C until used. *T. gondii* was detected by PCR targeting the *GRA5* and *GRA6* gene according to a previously described method [16,17]. *B. microti* was detected using nested PCR targeting the 18S rRNA and β-tubulin genes as previously described [11]. Amplified products were size-fractionated by electrophoresis on agarose gels containing SafePinky DNA gel staining solution (GenDEPOT, Katy, Texas, USA). The PCR products were then purified using an agarose gel DNA purification kit (Qiagen). TA cloning was performed using the

TOPO TA cloning kit with isolated PCR products for sequencing (Invitrogen, Carlsbad, California, USA). These samples were sequenced using an ABI PRISM 3730xl Analyzer (Applied Biosystems, Foster City, California, USA). In addition, antibodies against *T. gondii* were detected using a commercial toxoplasmosis multi-species ELISA kit (ID Vet, Montpellier, France). All procedures were carried out according to the manufacturer's instructions. The samples were tested twice for the most part, and any samples showing inconsistent results were examined once more. The reference sequences of *GRA6* of *T. gondii* strains and of 18S rRNA and β-tubulin for *B. microti* strains were obtained from GenBank [17,18]. Sequence alignment was performed using CLUSTAL W (Multiple sequence alignment computer programs Histon, Cambridgeshire, UK). Phylogenetic trees were constructed using the neighbor-joining method [19] with maximum composite likelihood distance correction in the molecular evolutionary genetics analysis (MEGA6) program [20]; the robustness of groupings was assessed using 1,000 bootstrap replicates [21].

Six out of 70 serum samples examined in wild animals were positive for *T. gondii* antibodies, with the highest positive rate observed in roe deer (16.7%, 1/6), despite the small sample number, followed by Chinese water deer (10.8%, 4/37), and raccoon dog (4.3%, 1/23) in ELISA (Table 1). Only 1 sample of the positives for *T. gondii* antibodies was positive for *GRA5* and *GRA6* genes in PCR which was from a Chinese water deer

(2.7%, 1/37). The analysis of sequence polymorphism about *GRA6* showed the highest identities with the GT1, VEL, and RH (Type I) strains in the phylogenetic analysis (Fig. 2). This result revealed that the *T. gondii* isolate detected in Chinese water deer belonged to a high-virulence type (type I).


The population structure of *T. gondii* in Europe and North America consists of 3 different clonal lineages known as types I, II, and III [22]. The type I has been detected in wild boars and foxes, which is a significant finding, because this genotype is not frequently reported in animals in Europe [23,24]. This finding is similar to previous reports showing that the *T. gondii* antigen detected from small mammals captured in Gyeonggi-do and Gangwon-do of Korea belonged to the clonal type I group [12]. Moreover, the possible overlap between the wild and domestic cycles of *T. gondii* could have implications for the possible infection of farm animals and humans. Although specific transmission routes could not be identified in the present survey, stray cats may be presumed to be one of the main sources of *T. gondii* transmission. Most of the recent investigations of *T. gondii* have focused on humans or domestic

animals [25-27]. The increasing urbanization of Korea has resulted in more interactions between humans and wild animals, including raccoons, wild boars, and Chinese water deer, which can survive well in close contact with humans. Furthermore, a recent review summarized the worldwide prevalence of *T. gondii* infection in wild pigs and deer [28], and human infections have been shown to be related to the consumption of the meat of wild animals [13,29]. Recently, consumption of raw pork was also identified as a critical dietary risk factor contributing to the seropositivity of *T. gondii* infection [27]. In addition, the seropositivity of *T. gondii* infection among veterinarians in a public professional activities group was significantly related to contact with animals infected with zoonotic pathogens [27]. Hence, the generally high prevalence of *T. gondii* in wild animals suggests that transmission from wild animals via consumption of raw or undercooked meat or direct contact with wild animals may be the main route of human infections.

*B. microti* can infect a variety of animal species, but the reservoir competence of numerous wildlife hosts in Korea is not well known. *B. microti* or *B. microti*-like infection has been observed in other widespread mammalian species of the eastern United States, such as short-tailed shrews [30], eastern cottontail rabbits, eastern chipmunks [31], raccoons [32], and foxes [33]. In Korea, *B. microti* infection has been reported in wild raccoon dogs [7] and small wild mammals [12]. However, the present results represented the first detection of *B. microti* infection in Chinese water deer (5.4%, 2/37) and Eurasian badgers (66.7%, 2/3). Thus, *B. microti* has been recognized as a genetically various species complex that consists of several clusters.

**Table 1.** Positive number of serologic and molecular detection for *T. gondii* and *B. microti* in wild animals

Wild animal species	No.	<i>T. gondii</i>		<i>B. microti</i>
		ELISA (%)	PCR (%)	PCR (%)
Chinese water deer	37	4 (10.8)	1 (2.7)	2 (5.4)
Raccoon dog	23	1 (4.3)	-	-
Roe deer	6	1 (16.7)	-	-
Wild boar	1	-	-	-
Eurasian badger	3	-	-	2 (66.7)
Total	70	6 (8.6)	1 (1.4)	4 (5.7)




**Fig. 2.** Phylogenetic relationships among *Toxoplasma gondii* species and genotypes according to neighbor-joining analysis with maximum composite likelihood distance correction, implemented with MEGA (version 6), using a fragment from the partial *GRA6* gene. Sequences of other *T. gondii* isolates and genotypes were obtained from GenBank (accession numbers are indicated in parentheses). The genotype of 0809-32 in this study was type I.

Analysis of *B. microti* genes showed sequence variations from geographically distant areas, such as Kobe (Japan), Hobetsu (Japan), G1 (USA), Gray (USA), and HK (Germany) [11].

Phylogenetic reconstruction based on 18S rRNA and  $\beta$ -tubulin sequence data of 4 *B. microti*-positive samples showed that all positive samples of 18S rRNA (0809-26, 0809-62, 0809-64, and 0809-69) were most similar to the US-type *B. microti* (Fig. 3A). In addition, the  $\beta$ -tubulin sequence in our positive samples was strongly related to those previously re-

ported in Korea and Russia (Fig. 3B). Similar to the results for 18S rRNA,  $\beta$ -tubulin sequence analysis showed that *B. microti* detected in small mammals in Korea most closely aligned with the US-type strain. Furthermore, US-type *B. microti* is commonly distributed among small wild mammals in China and Russia [11]. Therefore, our report shows that the US-type of *B. microti* in Korea may distribute in wild animals, and that both small and large zoonotic mammals can be reservoirs of *B. microti*, facilitating maintenance of the pathogen in nature. More-



**Fig. 3.** Phylogenetic relationships among *Babesia* species and genotypes according to neighbor-joining analysis with maximum composite likelihood distance correction, implemented with MEGA (version 6), using a fragment of the partial 18S rRNA (A) and  $\beta$ -tubulin (B) sequences of *Babesia microti*. Sequences of other *Babesia* species and genotypes were obtained from GenBank (accession numbers are indicated in parentheses). All positive isolates of *B. microti* are identified as the Gray (US-type).

over, ixodid ticks can transmit the pathogen to zoonotic and domestic hosts, and incidentally to humans [30]; thus, further studies are needed to identify the specific tick species involved in its transmission.

One of the main limitations of the present study is that these data may not be representative of the overall prevalence of *T. gondii* and *B. microti* infections in wild animals of Korea because of insufficient sample numbers and the limited study area. However, in spite of these limitations, this is the first record of *T. gondii* and *B. microti* infections in wild animals, including Chinese water deer, raccoon dog, roe deer, and Eurasian badger, inhabiting a mountainous region of Gangwon-do, as detected by serological and molecular screening. Genotyping of the *T. gondii* isolate revealed a type I genotype, whereas *B. microti* more closely aligned with the US-type. To our knowledge, this is the first report of *B. microti* detected in Chinese water deer and Eurasian badgers from Korea. These results suggested that these species might serve as an important reservoir for the transmission of *T. gondii* and *B. microti*, highlighting the need for closer monitoring of zoonotic infections in wild animals of Korea.

## ACKNOWLEDGMENTS

This research was performed in collaboration with the Korea CDC and Mongolia NCCD and funded by a grant (no. 4847-302-210-13, 2013) from the Korea National Institute of Health, Korea Centers for Disease Control and Prevention.

## CONFLICT OF INTEREST

We have no conflict of interest related to this work.

## REFERENCES

- Chomel BB, Belotto A, Meslin FX. Wildlife, exotic pets, and emerging zoonoses. *Emerg Infect Dis* 2007; 13: 6-11.
- Mackenstedt U, Jenkins D, Romig T. The role of wildlife in the transmission of parasitic zoonoses in peri-urban and urban areas. *Int J Parasitol Parasites Wildl* 2015; 4: 71-79.
- Cutler SJ, Fooks AR, van der Poel WH. Public health threat of new, reemerging, and neglected zoonoses in the industrialized world. *Emerg Infect Dis* 2010; 16: 1-7.
- Fèvre EM, Bronsvoort BM, Hamilton KA, Cleaveland S. Animal movements and the spread of infectious diseases. *Trends Microbiol* 2006; 14: 125-131.
- Youn HJ. Review of Zoonotic Parasites in Medical and Veterinary Fields in the Republic of Korea. *Korean J Parasitol* 2009; 47: 133-141.
- Kang SW, Doan HT, Noh JH, Choe SE, Yoo MS, Kim YH, Reddy KE, Nguyen TT, Van, Quyen D, Nguyen LT, Kweon CH, Jung SC. Seroprevalence of *Toxoplasma gondii* and *Trichinella spiralis* infections in wild boars (*Sus scrofa*) in Korea. *Parasitol Int* 2013; 62: 583-585.
- Han JI, Lee SJ, Jang HJ, and Na KJ. Asymptomatic *Babesia microti*-like parasite infection in wild raccoon dogs (*Nyctereutes procyonoides*) in South Korea. *J Wildl Dis* 2010; 46: 632-635.
- Dubey JP, Jones JL. *Toxoplasma gondii* infection in humans and animals in the United States. *Int J Parasitol* 2008; 38: 1257-1278.
- Kjemtrup AM, Conrad PA. Human babesiosis: an emerging tick-borne disease. *Int J Parasitol* 2000; 30: 1323-1337.
- Jeong W, Yoon H, Kim YK, Moon OK, Kim DS, An DJ. Prevalence of Antibodies to *Toxoplasma gondii* in South Korean Wild Boar (*Sus scrofa coreanus*). *J Wildl Dis* 2014; 50: 902-905.
- Zamoto A, Tsuji M, Wei Q, Cho SH, Shin EH, Kim TS, Leonova GN, Hagiwara K, Asakawa M, Kariwa H, Takashima I, Ishihara C. Epizootiologic survey for *Babesia microti* among small wild mammals in Northeastern Eurasia and a geographic diversity in the  $\beta$ -tubulin gene sequences. *J Vet Med Sci* 2004; 66: 785-792.
- Hong SH, Lee SE, Jeong YI, Kim HC, Chong ST, Klein TA, Song JW, Gu SH, Cho SH, Lee WJ. Prevalence and molecular characterizations of *Toxoplasma gondii* and *Babesia microti* from small mammals captured in Gyeonggi and Gangwon Provinces, Republic of Korea. *Vet Parasitol* 2014; 205: 512-517.
- Choi WY, Nam HW, Kwak NH, Huh W, Kim YR, Kang MW, Cho SY, Dubey JP. Foodborne outbreaks of human toxoplasmosis. *J Infect Dis* 1997; 175: 1280-1282.
- Morzaria S, Katende J, Kairo A, Nene V, Musoke A. New methods for the diagnosis of *Babesia bigemina* infection. *Mem Inst Oswaldo Cruz* 1992; 87 (Suppl): 201-205.
- Kawabuchi T, Tsuji M, Sado A, Matoba Y, Asakawa M, Ishihara C. *Babesia microti*-like parasites detected in feral raccoons (*Procyon lotor*) captured in Hokkaido, Japan. *J Vet Med Sci* 2005; 67: 825-827.
- Chen J, Li ZY, Zhou DH, Liu GH, Zhu XQ. Genetic diversity among *Toxoplasma gondii* strains from different hosts and geographical regions revealed by sequence analysis of *GRA5* gene. *Parasit Vectors* 2012; 5: 279.
- Su C, Zhang X, Dubey JP. Genotyping of *Toxoplasma gondii* by multilocus PCR-RFLP markers: a high resolution and simple method for identification of parasites. *Int J Parasitol* 2006; 36: 841-848.
- Tsuji M, Zamoto A, Kawabuchi T, Kataoka T, Nakajima R, Asakawa M, Ishihara C. *Babesia microti*-like parasites detected in Eurasian red squirrels (*Sciurus vulgaris orientis*) in Hokkaido, Japan. *J Vet Med Sci* 2006; 68: 643-646.
- Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 1987; 4: 406-425.

20. Kumar S, Nei M, Dudley J, Tamura K. MEGA: A biologist-centric software for evolutionary analysis of DNA and protein sequences. *Brief Bioinform* 2008; 9: 299-306.
21. Felsenstein J. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 1985; 39: 783-791.
22. Sibley LD, Boothroyd JC. Virulent strains of *Toxoplasma gondii* comprise a single clonal lineage. *Nature* 1992; 359: 82-85.
23. Berger-Schoch AE, Herrmann DC, Schares G, Müller N, Bernet D, Gottstein B, Frey CF. Prevalence and genotypes of *Toxoplasma gondii* in feline faeces (oocysts) and meat from sheep, cattle and pigs in Switzerland. *Vet Parasitol* 2011; 177: 290-297.
24. De Craeye S, Speybroeck N, Ajzenberg D, Dardé ML, Collinet E, Tavernier P, Van Gucht S, Dorny P, Dierick K. *Toxoplasma gondii* and *Neospora caninum* in wildlife: common parasites in Belgian foxes and Cervidae? *Vet Parasitol* 2011; 178: 64-69.
25. Hong SH, Jeong YI, Kim JY, Cho SH, Lee WJ, Lee SE. Prevalence of *Toxoplasma gondii* infection in household cats in Korea and risk factors. *Korean J Parasitol* 2013; 51: 357-361.
26. Jung BY, Gebeyehu EB, Lee SH, Seo MG, Byun JW, Oem JK, Kim HY, Kwak D. Detection and determination of *Toxoplasma gondii* seroprevalence in native Korean goats (*Capra hircus coreanae*). *Vector Borne Zoonotic Dis* 2014; 14: 374-377.
27. Lee SE, Hong SH, Jeong YI, Lee JH, Yoo SJ, Lim HS, Lee WJ, Cho SH. Cross-sectional analysis of the seropositivity and risk factors of *Toxoplasma gondii* infection among veterinarians, in relation to their public professional activities. *Vet Parasitol* 2014; 203: 29-34.
28. Dubey JP. Review of "Toxoplasmosis of Animals and Humans (Second Edition)" by J.P. Dubey. *Parasit Vectors* 2010; 3: 112.
29. Ross RD, Stec LA, Werner JC, Blumenkranz MS, Glazer L, Williams GA. Presumed acquired ocular toxoplasmosis in deer hunters. *Retina* 2001; 21: 226-229.
30. Telford SR 3rd, Mather TN, Adler GH, Spielman A. Short-tailed shrews as reservoirs of the agents of lyme disease and human babesiosis. *J Parasitol* 1990; 76: 681-683.
31. Spielman A, Etkind P, Piesman J, Ruebush TK 2nd, Juranek DD, Jacobs MS. Reservoir hosts of human babesiosis on Nantucket Island. *Am J Trop Med Hyg* 1981; 30: 560-565.
32. Birkenheuer AJ, Marr HS, Hladik N, Acton AE. Molecular evidence of prevalent dual piroplasma infections in North American raccoons (*Procyon lotor*). *Parasitology* 2008; 135: 33-37.
33. Birkenheuer AJ, Horney B, Bailey M, Scott M, Sherbert B, Catto V, Marr HS, Camacho AT, Ballman AE. *Babesia microti*-like infections are prevalent in North American foxes. *Vet Parasitol* 2010; 172: 179-182.