http://e-nrp.org

# Instant noodle consumption is associated with cardiometabolic risk factors among college students in Seoul

In Sil Huh<sup>1\*</sup>, Hyesook Kim<sup>2\*</sup>, Hee Kyung Jo<sup>3</sup>, Chun Soo Lim<sup>3</sup>, Jong Seung Kim<sup>3</sup>, Soo Jin Kim<sup>4</sup>, Oran Kwon<sup>2</sup>, Bumjo Oh<sup>5§</sup> and Namsoo Chang<sup>2§</sup>

<sup>1</sup>Seoul National University Medical Research Center, Seoul 03080, Korea

<sup>2</sup>Department of Nutritional Science and Food Management, Ewha Womans University, 52, Ewhayeodae-gil, Seodaemun-gu, Seoul 03760, Korea

<sup>3</sup>Seoul National University Health Service Center, Seoul 08826, Korea

<sup>4</sup>Health Promotion Center, Chung-Ang University Hospital, Seoul 06973, Korea

<sup>5</sup>Department of Family Medicine, SMG - SNU Boramae Medical Center, 20, Boramae-ro 5-gil, Dongjak-gu, Seoul 07061, Korea

**BACKGROUND/OBJECTIVES:** Increased consumption of instant noodles has recently been reported to be positively associated with obesity and cardiometabolic syndrome in South Korea, which has the highest per capita instant noodle consumption worldwide. This study aimed to investigate the association between instant noodle consumption and cardiometabolic risk factors among college students in Seoul.

**SUBJECTS/METHODS:** The study subjects consisted of 3,397 college students (1,782 male; 1,615 female) aged 18-29 years who participated in a health checkup. Information on instant noodle consumption was obtained from the participants' answers to a question about their average frequency of instant noodle intake over the 1 year period prior to the survey.

**RESULTS:** Statistical analysis using a general linear model that adjusted for age, body mass index, gender, family income, health-related behaviors, and other dietary factors important for cardiometabolic risk, showed a positive association between the frequency of instant noodle consumption and plasma triglyceride levels, diastolic blood pressure, and fasting blood glucose levels in all subjects. Compared to the group with the lowest frequency of instant noodle intake ( $\leq$  1/month), the odds ratio for hypertriglyceridemia in the group with an intake of  $\geq$  3/week was 2.639 [95% confidence interval (Cl), 1.393-5.000] for all subjects, while it was 2.149 (95% Cl, 1.045-4.419) and 5.992 (95% Cl, 1.859-21.824) for male and female students, respectively. In female students, diastolic blood pressure was also higher among more frequent consumers of instant noodles.

**CONCLUSIONS:** Our results suggest that frequent consumption of instant noodles may be associated with increased cardiometabolic risk factors among apparently healthy college students aged 18-29 years.

Nutrition Research and Practice 2017;11(3):232-239; https://doi.org/10.4162/nrp.2017.11.3.232; pISSN 1976-1457 eISSN 2005-6168

Keywords: Overweight, blood pressure, hypertriglyceridemia, metabolic factors

### INTRODUCTION

Overconsumption of instant noodles has recently received special attention, owing to its association with obesity and cardiometabolic syndrome among adults in South Korea [1]. The high-calorie content and the high concentration of refined carbohydrates, fats, and sodium [2] in instant noodles, contribute to an increased risk of metabolic disease [1]. Consumption of instant noodles as a staple food source is rising in many Asian countries [3] but South Korea is ranked the world's number one per capita in instant noodle consumption (72.8 servings of instant noodles per annum) [4]. A recent survey [5] reported that college students consume instant noodles more frequently than adults in other age groups because of the convenience of preparing instant noodles on school premises.

The fact that college students with unhealthy dietary habits frequently eat instant noodles and may be at increased risk of negative health outcomes should receive special attention because of a recent survey that reported an increasing trend of early premature cardiovascular diseases (CVD) death among adults as young as 20-49 years of age in Korea [6]. There is also an increasing trend in the proportion of individuals in the same group of young adults with increased cardiometabolic risk factors, such as overweight or obesity, hypertension, and elevated blood levels of glucose and lipids [7].

Many cardiometabolic risk factors are known to be modified by lifestyle behaviors including diet. Among many diet-related behaviors, avoiding instant foods and/or fast food and increasing

<sup>§</sup> Corresponding Authors: Namsoo Chang, Tel. 82-2-3277-3468, Fax. 82-2-3277-2862, Email. nschang@ewha.ac.kr

Received: February 28, 2017, Revised: April 28, 2017, Accepted: May 1, 2017

\* These authors contributed to this article equally.

This work was supported by the Second Stage of Brain Korea 21 Plus.

Bumjo Oh, Tel. 82-2-870-2682, Fax. 82-2-831-0714, Email. bumjo.oh@gmail.com

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (http://creativecommons.org/licenses/by-nc/3.0/) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

the intake of basic and unprocessed foods, is considered a simple but pivotal strategy. Cultivation of healthy lifestyle behaviors including good dietary practices among young adults, such as college students, is crucial for health promotion and disease prevention on both an individual and national level. Among young adults, modification of undesirable dietary habits and a reduction in CVD risk factors, is becoming more important for the prevention of CVD.

Recent studies that report a positive relationship between instant noodles and obesity and cardiometabolic syndrome among Korean adults [1,8], have focused on the general population, including all adult age groups. Despite an increasing trend in the proportion of young Korean adults in their 20s and 30s, with elevated metabolic markers, no study, to our knowledge, has previously investigated this association, by specifically targeting young adults, such as college students [9], which are known to be the most frequent consumers of instant noodles. Therefore, the current study was conducted to investigate the association between instant noodle consumption and cardiometabolic risk factors among young college students in Korea.

# SUBJECTS AND METHODS

#### Study subjects

A student health checkup was conducted at the health center of a university located in Seoul, Korea. A total of 3,481 students, aged 18-29 years, participated in the health checkup program from 24/04/2013-16/05/2013. Of the 3,481 subjects, 81 students who did not respond to a food questionnaire were excluded from the analysis. Three other students were excluded because they were receiving treatment for chronic disorders such as hypertension, dyslipidemia, or diabetes. In total, 3,397 subjects (1,782 male; 1,615 female) were included in the analysis. The study protocol was approved by the 2012 Seoul National Bioethics Committee and the 2013 Institutional Review Board of Seoul National University College of Medicine/Seoul National University Hospital (No. C-1304-062-481). All individuals provided informed consent to participate in the study.

General characteristics and instant noodle consumption survey All of the students completed an online survey that ascertained demographic and socioeconomic factors, and health-related behaviors including age, height, weight, family income, alcohol consumption, smoking behavior, and physical activity. Family monthly income was classified into < \$2000 USD, \$2000-4000 USD, and > \$4000 USD. Alcohol consumption was categorized as non-drinker, moderate drinker [weekly alcohol consumption < 14 drinks (males) or < 7 drinks (females), where 1 drink constitutes 14 g pure alcohol], and heavy drinker [weekly alcohol consumption > 14 drinks (males) or > 7 drinks (females)]. Smoking behavior was classified as non-smokers, ex-smokers, or current smokers, depending on current smoking status. Physical activity was classified as highly active (1500 MET-min/week, for any strenuous physical activity practiced > 3 d/week or 3000 MET-min/week, for any activity conducted daily), moderately active (intense physical activity for > 20 min/week or moderate activity or walking 5 d/week for > 30 min/d), or no physical activity/inactive (below a moderate level of physical activity).

Information on instant noodle consumption was obtained from the participants' answers to a question about their average frequency of instant noodle intake over the 1 year period prior to the survey. The questionnaire listed several potential candidate foods important for cardiometabolic risk in addition to instant noodles (fruits [1,10], vegetables [1,10], milk and dairy products [11], high-fat fish [1,12], high-fat and processed meats [1,13], sweets and confectionery [1,14], and carbonated beverages [1,15]), which was divided into nine frequency categories: barely eat, once/month, 2-3 times/month, 1-2 times/week, 3-4 times/ week, 5-6 times/week, once/day, twice/day, and 3 times/day. Those food groups were fruits, vegetables (except Kimchi), milk and dairy products (yogurt/cheese, etc.), high-fat fish (mackerel/ sardines/tuna/herring/salmon, etc.), high-fat and processed meats (fresh bacon/beef ribs/charcoal grilled meat/ tripe/ham/ sausage/bacon), instant noodle/cup noodle, sweets and confectionery (cakes/cookie/pies/candies/chocolates), and carbonated beverages (coke/sprite/soft drinks/fruit-flavored drinks except 100% fruit juice). In this study, the consumption of these foods as well as instant noodle was divided into 4 categories ( $\leq 1$ time/month, 2-3 times/month, 1-2 times/week, and  $\geq$  3 times/ week) according to frequency distribution.

#### Anthropometric and clinical examination

Anthropometric measurements were taken by a trained nurse. Standing height was measured to the nearest 0.1 cm with a stadiometer (BSM 330, Biospace Co. Seoul, South Korea). Body weight was measured to the nearest 0.1 kg using an Inbody 520 (Biospace Co., Ltd., Seoul, Korea). Body mass index (BMI) was calculated as weight (kg) divided by the square of the height (m<sup>2</sup>). Waist circumference was measured at the midpoint between the top of the iliac crest and the lowest rib, without any garments on. Blood pressure was taken when participants had been at rest for > 5 min, using an automatic blood pressure measurement system (CK-301, Spirit Medical Co., Taiwan). The second measured value was recorded for above-normal range blood pressure. When two preceding measurements were within the normal range, another measurement was conducted manually. Blood samples were drawn after a minimum 12-h overnight fast, collected in EDTA-containing tubes, and centrifuged at 3,000 rpm, at 4°C, for 20 min (Hanil Science Industrial Co., Ltd., Seoul, Korea). Fasting plasma levels of glucose, triglycerides, total cholesterol, and high density lipoprotein (HDL) cholesterol were assessed with an autoanalyzer (Cobas 6000, Roche Diagnostics International Ltd., Rotkreuz, Switzerland). Low density lipoprotein (LDL) cholesterol levels were calculated using the following equation, proposed by Friedwald [16] and Lauer [17]:

LDL-cholesterol = total cholesterol - HDL-cholesterol - (triglycerides/5)

#### Definition of metabolic syndrome

Metabolic syndrome was diagnosed based on the Adult Treatment Panel III of the National Cholesterol Education Program's (NCEP ATP III) standard diagnostic criteria [18]. Central obesity was defined as an absolute waist circumference > 90 cm (males) or > 85 cm (females), based on the guidelines of the Korean Society for the Study of Obesity [19]. The criteria for metabolic syndrome were as follows: (1) abdominal obesity: waist circumference > 90 cm (males) or > 85 cm (females), (2) high blood pressure: systolic blood pressure  $\geq$  130 mmHg or diastolic blood pressure  $\geq$  85 mmHg, (3) impaired fasting glucose: fasting blood sugar  $\geq$  100 mg/dL, (4) high triglycerides: triglycerides  $\geq$  150 mg/dL, and (5) low HDL- cholesterol: < 40 mg/dL (males) or < 50 mg/dL (females). Subjects who met three or more of these criteria, were diagnosed with metabolic syndrome.

## Statistical analysis

Data were expressed as the mean and standard deviations (SD) or as percentages (categorical), where appropriate. Differences in sociodemographic and behavioral characteristics among the instant noodle intake groups were analyzed using the Chi-square  $(\chi^2)$  test. General characteristics, such as age and BMI, were compared by one-way analysis of variance (ANOVA). The  $\chi^2$  test was applied to determine differences in the distribution of the number of metabolic syndrome components, according to instant noodle consumption. Potential confounders in this study included age, BMI, gender, family income (< \$2000 USD/month, \$2000-4000 USD/month, > \$4000 USD/month), alcohol consumption (non-drinker, moderate drinker, heavy drinker), smoking behavior (non-smoker, ex-smoker, current smoker), physical activity (low, moderate, high), and consumption frequency of fruits, vegetables, milk and dairy products, high-fat fish, high-fat and processed meats, sweets and confectionery and carbonated beverages ( $\leq 1$ time/month, 2-3 times/month, 1-2 times/week,  $\geq$  3 times/week). After adjusting for these confounders, a general linear model (GLM) was constructed to examine the differences in the metabolic parameters among the instant noodle consumption groups. Multiple logistic regression analysis was also performed to estimate the odds ratios (ORs) and 95% confidence intervals (Cls) for the cardiometabolic risk factors depending on instant noodle consumption. While performing the GLM or multiple logistic regression analyses, subjects with missing data for alcohol consumption (N = 187), smoking behavior (N = 52), and physical activity (N = 103) were excluded. All statistical analyses were performed using SAS 9.4 (SAS Inc., Cary, NC, USA). The significance level was set at P < 0.05.

# RESULTS

Characteristics of study participants according to the frequency of instant noodle consumption

The mean age of all subjects was  $22.4 \pm 2.9$  years, and did not differ among the instant noodle intake groups. Males were more likely to frequently consume instant noodles ( $\geq$  3 times/ week) than females. Subjects who frequently consumed instant noodles, had a lower family income level than infrequent consumers. Frequent consumers of instant noodles, drank more alcohol, were more likely to be current smokers, and were more physically active (Table 1). Frequent consumers of instant noodles also frequently consumed high-fat fish, high-fat and processed meats, sweets and confectionery, and carbonated beverages, and had less frequent consumption of fruits and vegetables (Table 2).

Metabolic parameters according to the frequency of instant noodle consumption

For all subjects, plasma triglyceride levels (P = 0.0016), diastolic

Table 1. Characteristics of study participants according to the frequency of instant noodle consumption<sup>1)</sup>

		Tota	al (N = 3,397)				Male	e (N = 1,782)		Female (N = 1,615)					
	$\leq$ 1 time/month (N = 937)	2-3 times/month (N = 1,012)	1-2 times/week (N = 1,050)	$\geq$ 3 times/week (N = 398)	P-value <sup>2)</sup>	$\leq$ 1 time/month (N = 365)	2-3 times/month (N = 478)	1-2 times/week (N = 657)	$\geq$ 3 times/week (N = 282)	P-value <sup>2)</sup>	$\leq$ 1 time/month (N = 572)	2-3 times/month (N = 534)	1-2 times/week (N = 393)	$\geq$ 3 times/week (N = 116)	P-value <sup>2)</sup>
Age (yrs)	$22.3\pm2.8$	22.3 ± 2.9	$22.4\pm2.9$	$22.6\pm3.0$	0.3640	22.6 ± 2.9	22.7 ± 3.1	22.6 ± 3.1	$22.6\pm3.0$	0.9814	22.2 ± 2.7	21.9 ± 2.7	22.1 ± 2.7	22.4 ± 3.0	0.2674
BMI (kg/m²)	$21.2\pm2.6$	$21.3\pm2.6$	$21.8\pm2.8$	21.9 ± 3.1	< 0.0001	$22.5 \pm 2.7$	$22.5 \pm 2.7$	$22.8\pm2.7$	22.7 ± 3.1	0.3232	20.4 ± 2.1	$20.2\pm2.1$	$20.3\pm2.1$	$20.2\pm2.2$	0.5145
Sex					< 0.0001										
Male	365 (39.0)	478 (47.2)	657 (62.6)	282 (70.9)											
Female	572 (61.0)	534 (52.8)	393 (37.4)	116 (29.1)											
Family income					0.0061					0.0371					0.5646
< 2000 US\$/mo	102 (10.9)	100 (9.9)	79 (7.5)	56 (14.1)		45 (12.3)	50 (10.5)	54 (8.2)	44 (15.6)		57 (10.0)	50 (9.4)	25 (6.4)	12 (10.3)	
2000-4000 US\$/mo	281 (30.0)	311 (30.7)	349 (33.2)	130 (32.7)		124 (34.0)	158 (33.1)	229 (34.9)	98 (34.8)		157 (27.5)	153 (28.7)	120 (30.5)	32 (27.6)	
>4000 US\$/mo	554 (59.1)	601 (59.4)	622 (59.2)	212 (53.3)		196 (53.7)	270 (56.5)	374 (56.9)	140 (49.7)		358 (62.6)	331 (62.0)	248 (63.1)	72 (62.1)	
Alcohol consumptio	n				< 0.0001					< 0.0001					< 0.0001
Non-drinker	122 (13.7)	75 (7.9)	61 (6.1)	27 (7.2)		42 (12.2)	30 (6.7)	35 (5.5)	16 (6.0)		80 (14.7)	45 (9.0)	26 (7.1)	11 (10.2)	
Moderate drinker	695 (78.0)	752 (79.4)	782 (78.3)	266 (71.3)		276 (80.0)	366 (81.5)	507 (79.8)	191 (72.1)		419 (76.7)	386 (77.5)	275 (75.6)	75 (69.4)	
Heavy drinker	74 (8.3)	120 (12.7)	156 (15.6)	80 (21.5)		27 (7.8)	53 (11.8)	93 (14.7)	58 (21.9)		47 (8.6)	67 (13.5)	63 (17.3)	22 (20.4)	
Smoking behavior					< 0.0001					0.0018					0.1060
Non-smoker	867 (93.8)	901 (90.5)	900 (87.4)	324 (82.0)		313 (87.9)	391 (83.4)	541 (83.4)	218 (77.6)		554 (97.5)	510 (96.8)	359 (94.2)	106 (93.0)	
Ex-smoker	28 (3.0)	49 (4.9)	61 (5.9)	23 (5.8)		21 (5.9)	40 (8.5)	50 (7.7)	19 (6.8)		7 (1.2)	9 (1.7)	11 (2.9)	4 (3.5)	
Current smoker	29 (3.1)	46 (4.6)	69 (6.7)	48 (12.2)		22 (6.2)	38 (8.1)	58 (8.9)	44 (15.7)		7 (1.2)	8 (1.5)	11 (2.9)	4 (3.5)	
Physical activity					0.0127					0.2366					0.0008
Low	87 (9.6)	67 (6.8)	98 (9.7)	39 (10.0)		46 (13.1)	41 (8.9)	83 (13.0)	34 (12.4)		41 (7.3)	26 (5.0)	15 (4.0)	5 (4.4)	
Moderate	592 (65.0)	623 (63.4)	605 (59.8)	225 (57.8)		224 (63.6)	304 (65.9)	394 (61.9)	162 (58.9)		368 (65.8)	319 (61.2)	211 (56.3)	63 (55.3)	
High	232 (25.5)	292 (29.7)	309 (30.5)	125 (32.1)		82 (23.3)	116 (25.2)	160 (25.1)	79 (28.7)		150 (26.8)	176 (33.8)	149 (39.7)	46 (40.4)	

BMI: body mass index

<sup>10</sup> Data are presented as mean ± SD or N(%). Alcohol consumption, smoking behavior, and physical activity data were analyzed in 3,210, 3,345, and 3,294 subjects, respectively.

 $^{2)}\ensuremath{\textit{P}}\xspace$  value by one-way analysis of variance or  $\chi^2$  test.

235

 $\textbf{Table 2.} Other food consumption of study participants according to the frequency of instant noodle consumption^{1)}$ 

		Tota	l (N = 3,397)			Mal	e (N = 1,782)		Female (N = 1,615)						
	≤ 1	2-3	1-2	$\geq$ 3	2	≤ 1	2-3	1-2	$\geq$ 3	2	≤ 1	2-3	1-2	$\geq$ 3	2)
	time/month	times/month $(N - 1.012)$	times/week	times/week	P-value <sup>2)</sup>	time/month	times/month	times/week	times/week	P-value <sup>2)</sup>	time/month	times/month	times/week	times/week	P-value <sup>2)</sup>
Fundation	(N = 937)	(N = 1,012)	(N = 1,050)	(IN = 398)	. 0.0001	(IN = 305)	(IN = 478)	(10 = 057)	(N = 282)	+ 0.0001	(N = 572)	(IN = 534)	(N = 393)	(N = 116)	0.0257
Fruits	00 (0 E)			(10.0)	< 0.0001					< 0.0001	aa (= a)			e (= e)	0.0257
$\leq$ 1 time/month	89 (9.5)	58 (5.7)	76 (7.2)	55 (13.8)		56 (15.3)	34 (7.1)	57 (8.7)	49 (17.4)		33 (5.8)	24 (4.5)	19 (4.8)	6 (5.2)	
2-3 times/month	110 (11.7)	149 (14.7)	185 (17.6)	74 (18.6)		45 (12.3)	81 (16.9)	129 (19.6)	48 (17.0)		65 (11.4)	68 (12.7)	56 (14.2)	26 (22.4)	
1-2 times/week	213 (22.7)	286 (28.3)	316 (30.1)	109 (27.4)		91 (24.9)	152 (31.8)	209 (31.8)	79 (28.0)		122 (21.3)	134 (25.1)	107 (27.2)	30 (25.9)	
$\geq$ 3 times/week	525 (56.0)	519 (51.3)	473 (45.1)	160 (40.2)		173 (47.4)	211 (44.1)	262 (39.9)	106 (37.6)		352 (61.5)	308 (57.7)	211 (53.7)	54 (46.5)	
Vegetables (excep	t Kimchi)				0.0335					0.1276					0.0528
$\leq$ 1 time/month	32 (3.4)	26 (2.6)	28 (2.7)	24 (6.0)		19 (5.2)	12 (2.5)	20 (3.0)	17 (6.0)		13 (2.3)	14 (2.6)	8 (2.0)	7 (6.0)	
2-3 times/month	42 (4.5)	54 (5.3)	53 (5.0)	22 (5.5)		20 (5.5)	32 (6.7)	28 (4.3)	15 (5.3)		22 (3.9)	22 (4.1)	25 (6.4)	7 (6.0)	
1-2 times/week	161 (17.2)	190 (18.8)	210 (20.0)	83 (20.9)		76 (20.8)	88 (18.4)	133 (20.2)	60 (21.3)		85 (14.9)	102 (19.1)	77 (19.6)	23 (19.8)	
$\geq$ 3 times/week	702 (74.9)	742 (73.3)	759 (72.3)	269 (67.6)		250 (68.5)	346 (72.4)	476 (72.5)	190 (67.4)		452 (79.0)	396 (74.2)	283 (72.0)	79 (68.1)	
Milk and dairy pr	oducts				0.2142					0.0840					0.4652
$\leq$ 1 time/month	57 (6.1)	44 (4.4)	46 (4.4)	21 (5.3)		35 (9.6)	28 (5.9)	32 (4.9)	20 (7.1)		22 (3.8)	16 (3.0)	14 (3.6)	1 (0.9)	
2-3 times/month	77 (8.2)	78 (7.7)	75 (7.1)	38 (9.5)		30 (8.2)	41 (8.6)	54 (8.2)	31 (11.0)		47 (8.2)	37 (6.9)	21 (5.3)	7 (6.0)	
1-2 times/week	206 (22.0)	271 (26.8)	269 (25.6)	93 (23.4)		73 (20.1)	125 (26.1)	171 (26.0)	66 (23.4)		133 (23.2)	146 (27.3)	98 (24.9)	27 (23.3)	
$\geq$ 3 times/week	596 (63.7)	619 (61.2)	660 (62.9)	246 (61.8)		226 (62.1)	284 (59.4)	400 (60.9)	165 (58.5)		370 (64.7)	335 (62.7)	260 (66.2)	81 (69.8)	
High-fat fish					0.0366					0.0381					0.0720
$\leq 1$ time/month	218 (23.3)	198 (19.6)	196 (18.7)	95 (23.9)		96 (26.3)	87 (18.2)	121 (18.4)	65 (23.1)		122 (21.3)	111 (20.8)	75 (19.1)	30 (25.9)	
2-3 times/month	320 (34.1)	345 (34.1)	331 (31.5)	116 (29.1)		116 (31.8)	164 (34.3)	213 (32.4)	88 (31.2)		204 (35.7)	181 (33.9)	118 (30.0)	28 (24.1)	
1-2 times/week	286 (30.5)	347 (34.3)	390 (37.1)	133 (33.4)		107 (29.3)	165 (34.5)	254 (38.7)	96 (34.0)		179 (31.3)	182 (34.1)	136 (34.6)	37 (31.9)	
> 3 times/week	113 (12.1)	122 (12.1)	133 (12.7)	54 (13.6)		46 (12.6)	62 (13.0)	69 (10.5)	33 (11.7)		67 (11.7)	60 (11.2)	64 (16.3)	21 (18.1)	
High-fat and proc	essed meats	(,		- (,	< 0.0001		(,	,		< 0.0001	,	,	- ( ,	(,	< 0.0001
< 1 time/month	143 (153)	60 (59)	33 (31)	15 (3.8)		50 (137)	20 (4.2)	19 (29)	8 (28)		93 (163)	40 (75)	14 (3.6)	7 (60)	
2-3 times/month	264 (28.2)	281 (27.8)	187 (17.8)	71 (17.8)		94 (25.7)	135 (28.2)	113 (172)	47 (167)		170 (297)	146 (27.3)	74 (18.8)	24 (20.7)	
1.2 times/month	207 (20.2)	201 (27.0)	474 (45 1)	125 (22.0)		110 (22.6)	100 (20.2)	207 (46.7)	100 (25.5)		102 (22.0)	100 (27.3)	167 (10.0)	25 (20.7)	
	202 (24.2)	202 (20.3)	256 (22.0)	177 (44.5)		102 (27.0)	122 (27.0)	210 (22 2)	100 (33.3)		105 (32.0)	150 (37.1)	120 (25.1)	50 (42.1)	
≥ 5 times/week	220 (24.3)	265 (26.0)	330 (33.9)	177 (44.5)	< 0.0001	102 (27.9)	155 (27.6)	219 (55.2)	127 (45.0)	< 0.0001	120 (22.0)	150 (26.1)	130 (33.1)	50 (45.1)	< 0.0001
sweets and come		26 (26)	21 (2.0)	21 (5.2)	< 0.0001	67 (10 4)	25 (5 2)	24 (27)	17 (6 0)	< 0.0001	21 (E A)	11 (2.1)	7 (1.0)	4 (2 5)	< 0.0001
$\leq$ 1 time/month	98 (10.5)	30 (3.0)	31 (2.9)	21 (5.3)		67 (18.4)	25 (5.2)	24 (3.7)	17 (0.0)		31 (5.4)	11 (Z.1)	7 (1.8)	4 (3.5)	
2-3 times/month	128 (13.7)	151 (14.9)	71 (0.8)	22 (5.5)		09 (18.9) 70 (21.4)	99 (20.7)	58 (8.8)	21 (7.4)		59 (10.3)	52 (9.7)	13 (3.3)	T (0.9)	
I-2 times/week	220 (23.5)	291 (28.7)	322 (30.7)	61 (15.3)		/9 (21.6)	141 (29.5)	226 (34.4)	49 (17.4)		141 (24.6)	150 (28.1)	96 (24.4)	12 (10.3)	
$\geq$ 3 times/week	491 (52.4)	534 (52.8)	626 (59.6)	294 (73.9)		150 (41.1)	213 (44.6)	349 (53.1)	195 (69.1)		341 (59.6)	321 (60.1)	2// (/0.5)	99 (85.3)	
Carbonated bever	ages				< 0.0001					< 0.0001					< 0.0001
$\leq$ 1 time/month	354 (37.8)	130 (12.8)	79 (7.5)	25 (6.3)		93 (25.5)	40 (8.4)	34 (5.2)	16 (5.7)		261 (45.6)	90 (16.9)	45 (11.5)	9 (7.8)	
2-3 times/month	189 (20.2)	266 (26.3)	118 (11.2)	36 (9.0)		79 (21.6)	110 (23.0)	55 (8.4)	23 (8.2)		110 (19.2)	156 (29.2)	63 (16.0)	13 (11.2)	
1-2 times/week	242 (25.8)	346 (34.2)	438 (41.7)	83 (20.8)		110 (30.1)	174 (36.4)	276 (42.0)	56 (19.9)		132 (23.1)	172 (32.2)	162 (41.2)	27 (23.3)	
$\geq$ 3 times/week	152 (16.2)	270 (26.7)	415 (39.5)	254 (63.8)		83 (22.7)	154 (32.2)	292 (44.4)	187 (66.3)		69 (12.1)	116 (21.7)	123 (31.3)	67 (57.8)	

Food groups were fruits, vegetables (except Kimchi), milk and dairy products (yogurt/cheese, etc.), high-fat fish (mackerel/sardines/tuna/herring/salmon, etc.), high-fat and processed meats (fresh bacon/beef ribs/charcoal grilled meat/tripe/ham/sausage/bacon), instant noodle/cup noodle, sweets and confectionery (cakes/cookie/pies/candies/chocolates), and carbonated beverages (coke/sprite/soft drinks/fruit-flavored drinks except 100% fruit juice).

 $^{1)}$  Data are presented as N(%).  $^{2)}$  /P-value by  $\chi^2$  test.

Table 3. Metabolic parameters according to the frequency of instant noodle consumption  $^{1\!\mathrm{l}}$ 

		Tota	l (N = 3,397)				Mal	e (N = 1,782)			Female (N = 1,615)					
	$\leq$ 1 time/month (N = 937)	2-3 times/month (N = 1,012)	1-2 times/week (N = 1,050)	$\geq 3 \\ times/week \\ (N = 398)$	P-value <sup>2)</sup>	$\leq 1$ time/month (N = 365)	2-3 times/month (N = 478)	1-2 times/week (N = 657)	$\geq 3 \\ times/week \\ (N = 282)$	P-value <sup>2)</sup>	$\leq 1 \\ time/month \\ (N = 572)$	2-3 times/month (N = 534)	1-2 times/week (N = 393)	$\geq 3 \\ times/week \\ (N = 116)$	P-value <sup>2)</sup>	
WC (cm)	$71.5\pm8.0$	$72.2\pm8.6$	$74.2\pm8.6$	$75.2\pm9.0$	0.8453	$77.6\pm7.6$	$77.7\pm7.5$	$\textbf{78.2} \pm \textbf{7.8}$	$\textbf{78.4} \pm \textbf{8.1}$	0.7553	$67.7\pm5.4$	$67.2\pm6.3$	67.6 ± 5.1	$67.4\pm5.7$	0.7445	
SBP (mmHg)	113.8 ± 14.1	115.5 ± 13.8	119.2 ± 14.0	120.6 ± 13.5	0.2243	124.4 ± 11.6	$124.5\pm11.6$	125.4 ± 12.0	125.6 ± 11.6	0.4457	107.1 ± 11.0	107.5 ± 10.3	108.7 ± 10.4	108.6 ± 9.3	0.5137	
DBP (mmHg)	$64.2\pm9.1$	$65.3\pm9.2$	$67.0\pm9.0$	$68.1\pm8.7$	0.0216	$68.3\pm8.8$	$69.3\pm9.0$	$69.1\pm8.7$	$70.0\pm8.5$	0.1235	$61.7\pm8.3$	$61.7\pm7.7$	$63.4\pm8.4$	$63.5\pm7.3$	0.0167	
FBS (mg/dL)	83.7 ± 7.6	$84.5\pm7.4$	$84.9\pm7.5$	$86.0\pm7.7$	0.0238	$85.7\pm7.4$	$85.8\pm7.6$	$85.7\pm7.6$	$87.0 \pm 7.9$	0.0687	$82.5\pm7.4$	$83.4\pm7.0$	$83.6\pm7.3$	$83.4\pm6.4$	0.5753	
TG (mg/dL)	$75.4\pm31.6$	$80.3\pm33.0$	81.8 ± 36.1	90.6 ± 53.4	0.0016	83.6 ± 39.6	$88.8\pm39.0$	87.8 ± 39.9	95.3 ± 58.1	0.0482	70.1 ± 23.9	$72.7\pm24.0$	71.7 ± 25.8	$79.1\pm37.4$	0.0198	
HDL-cholesterol (mg/dL)	70.6 ± 15.3	68.3 ± 14.3	66.8 ± 14.7	65.6 ± 14.7	0.1862	62.6 ± 12.9	62.3 ± 12.8	62.5 ± 13.1	62.1 ± 13.0	0.8780	75.7 ± 14.5	73.7 ± 13.5	74.1 ± 14.4	74.0 ± 15.3	0.1452	
LDL-cholesterol (mg/dL)	100.4 ± 27.9	$99.9\pm25.8$	99.9 ± 26.1	99.4 ± 25.9	0.7900	97.6 ± 26.3	100.5 ± 26.3	101.2 ± 27.1	99.8 ± 26.4	0.2866	102.3 ± 28.7	99.4 ± 25.3	97.7 ± 24.1	98.5 ± 24.9	0.3141	
No. of MS components	$0.27\pm0.57$	0.30 ± 0.61	0.41 ± 0.68	0.48 ± 0.76	0.3723	0.57 ± 0.73	$0.54\pm0.75$	0.59 ± 0.77	0.62 ± 0.82	0.6297	0.08 ± 0.31	$0.10\pm0.33$	0.11 ± 0.33	0.15 ± 0.44	0.3720	

WC, waist circumference; SBP, systolic blood pressure; DBP, diastolic blood pressure; FBS, fasting blood glucose; TG, triglycerides; MS, metabolic syndrome.

 $^{1)}\,\text{Data}$  are presented as mean  $\pm\,\text{SD}$ 

<sup>2)</sup> Adjusted Avalue from the general linear model after controlling for age, BMI, gender, family income (< 2000 US \$/mo, 2000-4000 US \$/mo, > 4000 US \$/mo, alcohol consumption (non-drinker, moderate drinker, heavy drinker), smoking behavior (non-smoker, ex-smoker, current smoker), physical activity (low, moderate, high), and consumption frequency of fruits, vegetables, milk and dairy products, high-fat fish, high-fat and processed meats, sweets and confectionery, carbonated beverages (< 1 time/month, 2-3 times/month, 1-2 times/week).</p>

#### Instant noodle and cardiometabolic risk factors



Fig. 1. Number of metabolic syndrome components according to the frequency of instant noodle consumption in all subjects. Reference values for MS components according to the Adult Treatment Panel III of the National Cholesterol Education Program (abdominal obesity: males  $\geq$  90 cm WC, females  $\geq$  85 cm; elevated blood pressure: SBP  $\geq$  130 mmHg or DBP  $\geq$  85 mmHg, hyperglycemia: FBS  $\geq$  100 mg/dL, hypertriglycemia: TG  $\geq$  150 mg/dL, low HDL cholesterol: males < 40 mg/dL, females < 50 mg/dL), MS, metabolic syndrome; WC, waist circumference; SBP, systolic blood pressure; DBP, diastolic blood pressure; FBS, fasting blood glucose; TG, triglycerides, *P*-value by  $\chi^2$  test,

blood pressure (P = 0.0216), and fasting blood glucose levels (P = 0.0238) were higher in the frequent instant noodle consumer group ( $\geq 3$  times/week) than the infrequent consumer group ( $\leq 1$  time/month) after adjusting for the covariates of age, BMI, gender, family income, alcohol consumption, smoking behavior, physical activity, and other dietary factors important for cardiometabolic risk. In female students, triglyceride levels (P = 0.0198) and diastolic blood pressure (P = 0.0167) were higher among the frequent consumers of instant noodles. In male students, triglyceride levels (P = 0.0482) were also higher among the frequent consumers (Table 3).

## Association between the frequency of instant noodle consumption and cardiometabolic risk factors

College students who consumed instant noodles frequently were more likely to have multiple cardiometabolic risk factors (Fig. 1) (P < 0.0001). The proportion of subjects with three or more risk factors was 0.8% among those who consumed instant noodles  $\leq 1$  time/month and 2.5% among those who consumed instant noodles  $\geq 3$  times/week.

As shown in Table 4, multiple logistic regression analysis with covariates revealed that frequent consumption of instant noodles

Table 4. Multivariate odds ratios (ORs) and 95% confidence intervals (CIs) for cardiometabolic risk factors according to the frequency of instant noodle consumption<sup>11</sup>

		Tota	al (N = 3,397)				Mal	e (N = 1,782)		Female (N = 1,615)					
	$\leq 1$ time/month	2-3 times/month	1-2 times/week	$\geq$ 3 times/week	P-value for	$\leq 1$ time/month	2-3 times/month	1-2 times/week	$\geq$ 3 times/week	P-value for	$\leq 1$ time/month	2-3 times/month	1-2 times/week	≥ 3 times/week	P-value for
	(N = 937)	(N = 1,012)	(N = 1,050)	(N = 398)	trend	(N = 365)	(N = 478)	(N = 657)	(N = 282)	trend	(N = 572)	(N = 534)	(N = 393)	(N = 116)	trend
Abdominal obes	ity														
No. of case (%)	28 (3.0)	36 (3.6)	58 (5.5)	29 (7.3)		25 (6.9)	33 (6.9)	56 (8.5)	27 (10.0)		3 (0.5)	3 (1.0)	2 (0.5)	2 (1.7)	
Multivariate OR (95% CI)	1	1.316 (0.753-2.300)	1.582 (0.922-2.715)	1.662 (0.887-3.117)	0.3404	1	1.383 (0.763-2.507)	1.679 (0.948-2.972)	1.779 (0.922-3.435)	0.2808	1	0.953 (0.170-5.361)	0.655 (0.084-5.125)	0.882 (0.071-10.964)	0.9798 )
Elevated blood	pressure														
No. of case (%)	156 (16.7)	172 (17.1)	260 (24.8)	109 (27.4)		139 (38.2)	158 (33.2)	241 (36.8)	105 (37.2)		17 (3.0)	14 (2.6)	19 (4.9)	4 (3.5)	
Multivariate OR (95% CI)	1	0.852 (0.639-1.137)	1.036 (0.782-1.372)	0.940 (0.664-1.331)	0.4732	1	0.849 (0.623-1.157)	0.990 (0.734-1.335)	0.930 (0.647-1.338)	0.6519	1	0.799 (0.338-1.899)	1.560 (0.674-3.610)	0.995 (0.254-3.904)	0.4381
Hyperglycemia															
No. of case (%)	23 (2.5)	27 (2.7)	26 (2.5)	14 (3.5)		17 (4.7)	19 (4.0)	18 (2.8)	13 (4.6)		6 (1.1)	8 (1.5)	8 (2.0)	1 (0.9)	
Multivariate OR (95% CI)	1	1.092 (0.582-2.050)	1.019 (0.527-1.971)	1.368 (0.631-2.963)	0.8434	1	1.017 (0.473-2.187)	0.834 (0.375-1.856)	1.460 (0.612-3.484)	0.5790	1	1.254 (0.377-4.173)	2.050 (0.599-7.011)	0.639 (0.064-6.366)	0.5571
Hypertriglycemia	1														
No. of case (%)	21 (2.3)	38 (3.8)	62 (5.9)	32 (8.1)		16 (4.4)	30 (6.3)	56 (8.6)	26 (9.3)		5 (1.0)	8 (1.5)	6 (1.5)	6 (5.2)	
Multivariate OR (95% CI)	1	1.685 (0.944-3.011)	2.163 (1.229-3.805)	2.639 (1.393-5.000)	0.0192	1	1.651 (0.844-3.228)	2.089 (1.103-3.955)	2.149 (1.045-4.419)	0.1245	1	2.033 (0.607-6.810)	2.179 (0.566-8.384)	5.992 (1.859-21.824)	0.0346 )
Low HDL-choles	terol														
No. of case (%)	26 (2.8)	35 (3.5)	24 (2.3)	8 (2.0)		10 (2.8)	16 (3.4)	1 (2.1)	4 (1.4)		16 (2.8)	19 (3.6)	10 (2.6)	4 (3.5)	
Multivariate OR (95% CI)	1	1.487 (0.834-2.651)	1.010 (0.527-1.935)	0.786 (0.321-1.923)	0.2948	1	1.473 (0.594-3.654)	0.807 (0.308-2.113)	0.430 (0.114-1.623)	0.1742	1	1.452 (0.676-3.121)	1.152 (0.466-2.851)	1.542 (0.450-5.287)	0.7690
High LDL-choles	terol														
No. of case (%)	116 (12.4)	113 (11.2)	126 (12.0)	48 (12.1)		44 (12.1)	57 (12.0)	96 (14.7)	39 (13.8)		72 (12.6)	56 (10.5)	30 (7.7)	9 (7.8)	
Multivariate OR (95% CI)	1	0.912 (0.674-1.234)	0.920 (0.675-1.254)	0.842 (0.556-1.276)	0.8632	1	1.135 (0.719-1.791)	1.296 (0.840-1.999)	1.230 (0.729-2.075)	0.6874	1	0.824 (0.544-1.246)	0.580 (0.347-0.968)	0.406 (0.162-1.017)	0.0881
Overweight (BM	I ≥ 23)														
No. of case (%)	190 (20.3)	246 (24.3)	315 (30.0)	130 (32.7)		128 (35.1)	197 (41.2)	278 (42.3)	121 (42.9)		62 (10.8)	49 (9.2)	37 (9.4)	9 (7.8)	
Multivariate OR (95% CI)	1	1.095 (0.854-1.405)	1.096 (0.853-1.407)	1.058 (0.770-1.454)	0.8837	1	1.235 (0.910-1.676)	1.217 (0.904-1.638)	1.232 (0.860-1.765)	0.5233	1	0.928 (0.599-1.437)	0.922 (0.564-1.508)	0.613 (0.259-1.452)	0.7443
Metabolic syndr	ome														
No. of case (%)	7 (0.7)	12 (1.2)	16 (1.5)	10 (2.5)		6 (1.6)	12 (2.5)	16 (2.4)	10 (3.6)		1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	
Multivariate OR (95% Cl)	1	1.565 (0.557-4.401)	1.438 (0.508-4.064)	2.218 (0.713-6.899)	0.5667	1	1.797 (0.602-5.358)	1.665 (0.553-5.009)	2.600 (0.791-8.553)	0.4591	1	-	-	-	-

<sup>1)</sup> Determined by multivariate logistic regression analysis after controlling for age, BMI, gender, family income (< 2000 US \$/mo, 2000-4000 US \$/mo, > 4000 US \$/mo), alcohol consumption (non-drinker, moderate drinker, heavy drinker), smoking behavior (non-smoker, ex-smoker, current smoker), physical activity (low, moderate, high), and consumption frequency of fruits, vegetables, milk and dairy products, high-fat fish, high-fat and processed meats, sweets and confectionery, carbonated beverages (< 1 time/month, 2-3 times/meth, 2-3 times/week, ≥ 3 times/week).</p>
Reference values according to the Adult Treatment Panel III of the National Cholesterol Education Program (abdominal obesity: males ≥ 90 cm, females ≥ 85 cm; elevated blood pressure: SBP ≥ 130 mmHg or DBP ≥ 85 mmHg, hyperglycemia: FBS ≥ 100 mg/dL, hyperfriglycemia: TG ≥ 150 mg/dL, low HDL cholesterol: males < 40 mg/dL, females < 50 mg/dL, high LDL cholesterol: ≥ 130 mg/dL).</p>

was associated with a higher prevalence of hypertriglyceridemia. Compared to the group with the lowest frequency of instant noodle consumption ( $\leq 1$  time/month), the OR for hypertriglyceridemia was significantly higher in the groups with a frequency consumption of 1-2 times/week (OR: 2.163, 95% CI: 1.229-3.805) and  $\geq$  3 times/week (OR: 2.693, 95% CI: 1.393-5.000) (*P* for trend = 0.0192) in all subjects. This association was also observed for males alone (OR: 2.149, 95% CI: 1.045-4.419) and females alone (OR: 5.992, 95% CI: 1.859-21.824), with a frequency of instant noodle consumption  $\geq$  3 times/week compared to  $\leq$  1 time/month. The OR for a high LDL-cholesterol was significantly lower in the group with a consumption frequency of 1-2 times/week (OR: 0.580, 95% CI: 0.347-0.968) compared to the group with the lowest frequency of instant noodle consumption ( $\leq$  1 time/month) in female students only.

## DISCUSSION

In this study, we found a positive association between the frequency of instant noodle consumption and plasma triglyceride levels, diastolic blood pressure, and fasting blood glucose levels in Korean college students. Study subjects with a higher frequency of instant noodle consumption were more likely to have multiple cardiometabolic risk factors. The OR for hypertriglyceridemia was significantly higher in the group that consumed instant noodles  $\geq$  3 times/week compared to the group with the lowest consumption frequency ( $\leq$  1 time/month). This association existed in male and female students respectively, as well as in the combined subjects.

The positive association between instant noodle consumption and triglyceride levels that we observed among college students is supported by the results of another recent study conducted on Korean adults aged 19-64 years. Yeon & Bae [8] showed that the group with high instant noodle consumption ( $\geq$  1 serving/ week) had significantly higher triglyceride levels compared with the low instant noodle consumption group (< 1 serving/week). We also found that college students who reported consuming instant noodles  $\geq$  3 times/week were at a higher risk for hypertriglyceridemia. We are unable to provide a mechanistic interpretation of these findings but it is probably due to the large amounts of carbohydrates and fat contained in instant noodles. Excess carbohydrate intake is known to increase triglyceride levels [20].

In our study, diastolic but not systolic blood pressure was increased, according to the frequency of instant noodle consumption and the effect was more apparent in female students. Diastolic blood pressure is considered an index of salt sensitivity [21]. Instant noodles made in Korea contain a significant amount of sodium in one serving (600-2,770 mg) [22]. According to the Korea National Health and Nutrition Examination Survey [23], noodles are one of the main sources of sodium intake in Korea, and instant noodles comprised the largest proportion of noodles eaten by South Korean adults in their 20s. If diastolic blood pressure is an index of salt sensitivity, our data indicate a gender difference in salt sensitively, such that women may be more salt-sensitive than men. This needs to be verified though further studies.

In the present study, a positive association was observed

between instant noodle consumption and fasting blood glucose levels. This observation is partially consistent with a previous report [8], which found that instant noodle consumption was associated with an increased prevalence of hyperglycemia in Korean women only. Instant noodles are generally high in refined carbohydrates but low in fiber [2]. This correlates with the high glycemic index (GI) or the high glycemic load of instant noodles. Some studies [24,25] have reported that the instant noodles produced in Korea have a GI of 71-87 and instant noodles are categorized as a high-GI food.

Although there was no difference in the proportion of subjects with metabolic syndrome between groups, students with a higher frequency of instant noodle consumption were more likely to have multiple cardiometabolic risk factors. Significant differences were observed in the proportion of subjects with two risk factors (17.7 vs. 27.6%) and three or more risk factors (0.7 vs. 2.5%), and between those who consumed instant noodles  $\leq 1$  time/month vs.  $\geq 3$  times/week. Our findings suggest that instant noodle consumption might be associated with the presence of multiple risk factors, even if metabolic syndrome has not yet manifested because the study subjects consisted of relatively young individuals.

In this study, we found unexpectedly that the OR for a high LDL-cholesterol was significantly lower in female students who consumed instant noodle 1-2 times/week than  $\leq$  1 time/month. At this point, we do not know whether this is a chance observation or artifact, therefore, it needs to be explored further in other studies with different study subjects. Blood levels of LDL-cholesterol, a well-established cardiometabolic risk factor, however, has been reported to be influenced by both genes and the environment, with the genetic factor being stronger than environmental variables, such as diet [26]. This is one of the reasons why LDL-cholesterol level is excluded from the diagnostic criteria of metabolic syndrome.

Researchers at Harvard University reported a 68% higher risk of metabolic syndrome among women who consume instant noodles  $\geq$  twice/week [1], but not in men. We found that frequent consumption of instant noodles was associated with increased diastolic blood pressure in women. These gender discrepancies might result from differences in food group compositions in the dietary intake patterns between males and females [27,28] and the biological differences between males and females, such as sex hormones and metabolism [27,29]. Previous authors suggested an interaction between bisphenol A and the estrogen receptor, as a possible contributor to the gender difference. Bisphenol A is known to be a selective modulator of estrogen receptors that can accelerate adipogenesis [30]. Considering that women may be more sensitive to instant noodle-associated metabolic changes is intriguing. Whether this is due to bisphenol A found in packaging or to other substances in instant noodles, warrants further investigation. Meanwhile, it is advisable to guide young women to minimize their instant noodle consumption to decrease future CVD metabolic risk.

Unlike fresh ramen noodles, most instant noodles are deep-fried to dry them, so they are high in calories, refined carbohydrates, saturated fat, and sodium [2]. The most consumed [9] type of instant noodles in Korea, for example, contains 505 kcal and provides 24, 31, 53, and 90% of the

recommended daily values for carbohydrates, fat, saturated fat and sodium, respectively [2]. Several studies have suggested that the high energy density, glycemic load (due to the refined carbohydrates), saturated fat content, and sodium content of instant noodles, may contribute to increased cardiometabolic risk factors. South Korea leads the world in per capita consumption of instant noodles, at 72.8 servings/year, which is almost six times higher than the world average of 13.5 servings in 2015 [31], with per capita consumption showing an increasing trend [32]. The convenience and low price of instant noodles have intensified their popularity in recent decades [32]. The preference for instant noodles is particularly strong among college students, with time or financial constraints. It has been reported that among adults of all age groups, younger adults aged 20-49 years, consume greater amounts of instant noodles (22.2 g/d) than those aged 50-64 years (7.6 g/d) [33].

The limitations of this study are as follows: first, it was impossible to define causal associations with this data set because of the cross-sectional study design. Further prospective studies are warranted to investigate the causal associations between instant noodle consumption and cardiometabolic risk factors. Second, there is a possibility that those who often consumed instant noodles as a snack, also consumed more calories overall, which may have led to a higher risk of metabolic syndrome. However, a previous study [34] observed that most adults consume instant noodles at lunch or dinner time as a substitute for a meal instead of a snack. For more accurate results, controlling participants' total daily caloric intake and determining the percentage of total caloric intake fulfilled by instant noodles is necessary. Despite these limitations, this study is significant, as it presents homogeneous data collected over a short duration, which includes data on a large number of young adults. Although various studies have been conducted on unhealthy eating habits developed over a long duration in subjects or patients in their late 40s [35] regarding an increased risk of cardiometabolic syndrome, insufficient studies have been conducted to show whether the same results will be seen when investigating instant noodle intake in younger individuals.

In conclusion, we found that frequent consumption of instant noodles may be associated with increased cardiometabolic risk factors among apparently healthy college students aged 18-29 years. Considering the significant, harmful association of multiple risk factors with increased CVD risk and later development of CVD, this issue requires immediate attention. It is necessary to encourage reduced consumption of instant noodles but there is also a need to educate individuals and for the food industry to improve the nutritional quality of instant noodles or develop healthy instant noodles.

# CONFLICT OF INTEREST

The authors declare no potential conflicts of interests.

## REFERENCES

 Shin HJ, Cho E, Lee HJ, Fung TT, Rimm E, Rosner B, Manson JE, Wheelan K, Hu FB. Instant noodle intake and dietary patterns are associated with distinct cardiometabolic risk factors in Korea. J Nutr 2014;144:1247-55.

- Kim MJ, Shin SN, Kim SK. Proximate composition and calorie of Korean instant noodles. Korean J Food Sci Technol 2000;32:1043-50.
- Srinath Reddy K, Katan MB. Diet, nutrition and the prevention of hypertension and cardiovascular diseases. Public Health Nutr 2004;7:167-86.
- Korea Bizwire. South Korea ranked no.1 in instant noodle consumption [Internet]. Seoul: Kobiz Media Co., Ltd.; 2014 [cited 2014 September 13]. Available from: http://koreabizwire.com/southkorea-ranked-no-1-in-instant-noodle-consumption/19034.
- Nelson MC, Kocos R, Lytle LA, Perry CL. Understanding the perceived determinants of weight-related behaviors in late adolescence: a qualitative analysis among college youth. J Nutr Educ Behav 2009;41:287-92.
- Kim HK, Ahn Y. Mortality trends of cardiovascular disease in Korea; big challenges in ischemic heart disease. Korean Circ J 2015;45: 192-3.
- Ministry of Health and Welfare, Korea Centers for Disease Control and Prevention. Korea Health Statistics 2013: Korea National Health and Nutrition Examination Survey (KNHANES VI-1). Cheongju: Korea Centers for Disease Control and Prevention; 2014.
- Yeon JY, Bae YJ. Association of instant noodle intake with metabolic factors in Korea: based on 2013-2014 Korean National Health and Nutrition Examination Survey. J Nutr Health 2016;49:247-57.
- Gallup Korea. The public opinion census about instant noodle [Internet]. Seoul: Gallup Korea; 2013 [cited 2015 November 11]. Available from: http://www.gallup.co.kr/gallupdb/reportDownload. asp?seqNo=494.
- Shin JY, Kim JY, Kang HT, Han KH, Shim JY. Effect of fruits and vegetables on metabolic syndrome: a systematic review and meta-analysis of randomized controlled trials. Int J Food Sci Nutr 2015;66:416-25.
- Chen GC, Szeto IM, Chen LH, Han SF, Li YJ, van Hekezen R, Qin LQ. Dairy products consumption and metabolic syndrome in adults: systematic review and meta-analysis of observational studies. Sci Rep 2015;5:14606.
- Tørris C, Molin M, Cvancarova Småstuen M. Associations between fish consumption and metabolic syndrome. A large cross-sectional study from the Norwegian Tromsø Study: Tromsø 4. Diabetol Metab Syndr 2016;8:18.
- Babio N, Sorlí M, Bulló M, Basora J, Ibarrola-Jurado N, Fernández-Ballart J, Martínez-González MA, Serra-Majem L, González-Pérez R, Salas-Salvadó J; Nureta-PREDIMED Investigators. Association between red meat consumption and metabolic syndrome in a Mediterranean population at high cardiovascular risk: cross-sectional and 1-year follow-up assessment. Nutr Metab Cardiovasc Dis 2012;22:200-7.
- 14. Stanhope KL. Sugar consumption, metabolic disease and obesity: the state of the controversy. Crit Rev Clin Lab Sci 2016;53:52-67.
- Crichton G, Alkerwi A, Elias M. Diet soft drink consumption is associated with the metabolic syndrome: a two sample comparison. Nutrients 2015;7:3569-86.
- Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. Clin Chem 1972;18: 499-502.
- Lauer RM, Lee J, Clarke WR. Factors affecting the relationship between childhood and adult cholesterol levels: the Muscatine Study. Pediatrics 1988;82:309-18.

- Grundy SM, Brewer HB Jr, Cleeman Jl, Smith SC Jr, Lenfant C; National Heart, Lung, and Blood Institute; American Heart Association. Definition of metabolic syndrome: report of the National Heart, Lung, and Blood Institute/American Heart Association conference on scientific issues related to definition. Arterioscler Thromb Vasc Biol 2004;24:e13-8.
- Lee S, Park HS, Kim SM, Kwon HS, Kim DY, Kim DJ, Cho GJ, Han JH, Kim SR, Park CY, Oh SJ, Lee CB, Kim KS, Oh SW, Kim YS, Choi WH, Yoo HJ. Cut-off points of waist circumference for defining abdominal obesity in the Korean population. Korean J Obes 2006;15:1-9.
- 20. Kamada C, Yoshimura H, Okumura R, Takahashi K, Iimuro S, Ohashi Y, Araki A, Umegaki H, Sakurai T, Yoshimura Y, Ito H; Japanese Elderly Diabetes Intervention Trial Study Group. Optimal energy distribution of carbohydrate intake for Japanese elderly patients with type 2 diabetes: the Japanese Elderly Intervention Trial. Geriatr Gerontol Int 2012;12 Suppl 1:41-9.
- 21. Costa Ede A, Rose G, Klein CH, Achutti AC. Diastolic pressure as an index of salt sensitivity. J Hum Hypertens 1994;8:703-9.
- Kim YY, Bae YJ, Choi MK. Sodium content and daily intake of instant noodle in Korean adolescents and adults. Trace Elem Electrolytes 2015;32:197-203.
- 23. Lee HS, Duffey KJ, Popkin BM. Sodium and potassium intake patterns and trends in South Korea. J Hum Hypertens 2013;27: 298-303.
- Yun EK, Heo YR, Lim HS. Effects of Sasa Borealis leaf extract on the glucose tolerance of major foods for carbohydrate. Korean J Nutr 2010;43:215-23.
- Song S, Choi H, Lee S, Park JM, Kim BR, Paik HY, Song Y. Establishing a table of glycemic index values for common Korean foods and an evaluation of the dietary glycemic index among the Korean adult population. Korean J Nutr 2012;45:80-93.
- 26. Jee SH, Yun JE, Nam CM, Suh I. Heritability and segregation analysis of the level of LDL-cholesterol. Korean Circ J 2005;35:233-9.

- 27. Wirfält E, Hedblad B, Gullberg B, Mattisson I, Andrén C, Rosander U, Janzon L, Berglund G. Food patterns and components of the metabolic syndrome in men and women: a cross-sectional study within the Malmö diet and cancer cohort. Am J Epidemiol 2001;154: 1150-9.
- Woo HD, Shin A, Kim J. Dietary patterns of Korean adults and the prevalence of metabolic syndrome: a cross-sectional study. PLoS One 2014;9:e111593.
- Ding EL, Song Y, Malik VS, Liu S. Sex differences of endogenous sex hormones and risk of type 2 diabetes: a systematic review and meta-analysis. JAMA 2006;295:1288-99.
- Elobeid MA, Allison DB. Putative environmental-endocrine disruptors and obesity: a review. Curr Opin Endocrinol Diabetes Obes 2008;15: 403-8.
- World Instant Noodles Association (JP). Instant noodles at a glance [Internet]. Tokyo: World Instant Noodles Association; 2015 [cited 2016 January 13]. Available from: http://instantnoodles.org/en/ noodles/report.html.
- World Instant Noodles Association (JP). National trends in instant noodles demands [Internet]. Tokyo: World Instant Noodles Association; 2009 [cited 2009 October 25]. Available from: http://instantnoodles. org/en/noodles/report.html.
- Park J, Lee JS, Jang YA, Chung HR, Kim J. A comparison of food and nutrient intake between instant noodle consumers and non-instant noodle consumers in Korean adults. Nutr Res Pract 2011;5:443-9.
- 34. Yu KJ, Jung HS, Yoon HH. A comprehensive study on the intake patterns and expenditures on ramyun among adults in metropolitan areas of Korea. Korean J Culinary Res 2013;19:204-14.
- Lakka HM, Laaksonen DE, Lakka TA, Niskanen LK, Kumpusalo E, Tuomilehto J, Salonen JT. The metabolic syndrome and total and cardiovascular disease mortality in middle-aged men. JAMA 2002; 288:2709-16.