

RESEARCH NOTE

***Pseudoperonospora urticae* Occurring on *Urtica angustifolia* in Korea**Young-Joon Choi^{1*}, Hyang Burm Lee², Hyeon-Dong Shin³¹Department of Biology, College of Natural Sciences, Kunsan National University, Gunsan 54150, Korea²Division of Food Technology, Biotechnology and Agrochemistry, Chonnam National University, Gwangju 61186, Korea³Division of Environmental Science and Ecological Engineering, Korea University, Seoul 02841, Korea

*Corresponding author: yjchoi@kunsan.ac.kr

Abstract

The genus *Pseudoperonospora* (Peronosporales, Oomycota) comprises six accepted species, including *Ps. cubensis*, which causes downy mildew on many economically important cucurbitaceous crops, and *Ps. humuli*, which occurs on hops. During a survey of downy mildew flora in Korea, a previously unreported species of *Pseudoperonospora* was found on *Urtica angustifolia*. Based on molecular phylogenetic and morphological analyses, the causal agent was identified as *Pseudoperonospora urticae*. This is the first report of *Pseudoperonospora urticae* occurring on *Urtica angustifolia* in Korea.

Keywords: *cox2* mtDNA, Internal transcribed spacer rDNA, Oomycetes, *Pseudoperonospora cubensis*, *Pseudoperonospora humuli*

 OPEN ACCESS

Kor. J. Mycol. 2017 June, 45(2): 160-166
<https://doi.org/10.4489/KJM.20170020>

pISSN : 0253-651X
 eISSN : 2383-5249

Received: 13 May, 2017

Accepted: 22 May, 2017

© The Korean Society of Mycology



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Downy mildews (Peronosporaceae; Oomycota) are an obligate biotrophic group that infects a wide range of mono- and dicotyledonous plants, including many economically relevant crops [1]. *Pseudoperonospora cubensis* is a notorious species infesting many cucurbitaceous crops, such as cucumber, gourd, pumpkin, and watermelon [2, 3], and *Ps. humuli* is one of the most important threats to the cultivation of hops (Cannabaceae) [4, 5]. Given their association with high economic losses, many recent studies have focused on the biology, host specificity, population structure, detection, and control of *Pseudoperonospora* species [3, 6-11], as well as their taxonomy and phylogeny [12, 13].

To date, four species of *Pseudoperonospora* have been reported in Korea [14, 15], *Ps. cannabina*, *Ps. celtidis*, *Ps. cubensis*, and *Ps. humuli*. In September 2009, symptoms typical of downy mildew were found on the leaves of *Urtica angustifolia* Fisch. ex Hornem. (Urticaceae) growing near Jangjeon valley in Pyeongchang, Korea (N37°29'36"; E128°32'33"). *Urtica angustifolia* is distributed in the wastelands, grasslands, valleys, wet places, and ridges of East Asian countries, including China, Korea, and Russia [16], and it is used as a traditional medicinal plant due to its high hypoglycemic activity [17]. The downy mildew

infection resulted in slight discoloration of the leaf tissues, with yellow or pale green spots on the upper leaf surfaces that developed dark grey fungal growth on the lower surfaces. The lesions were poly-angular, and were delimited by the leaf veins (Fig. 1A, 1B). As the disease progressed, the spots turned blackish and often merged to cover larger areas. A representative sample was deposited in the National Institute of Biological Resources (KZITFG000000017) and the Korea University Herbarium (KUS-F24488).




Fig. 1. *Pseudoperonospora urticae* occurring on *Urtica angustifolia*. A, symptoms on the upper surface of *Urtica angustifolia* leaves; B, focus on the vein-limited spots; C, D, sporangiophores; E, F, ultimate branchlets; G, H, sporangia (scale bars: C, D = 100 μ m, E~H = 20 μ m).

A detailed microscopic examination was performed using an Olympus BX53 microscope (Olympus, Tokyo, Japan), and DIC micrographs were captured with a DigiRetina 16M camera (Tucsen, Fuzhou, China). The following morphological characteristics were observed at 100–200 × for sporangiophores and at 400 × for sporangia and ultimate branchlets. The sporangiophores emerging through the stomata were tree-like, hyaline, straight, slender, and 120–400 μm in height (Fig. 1C, 1D). Trunks were straight, 4–6 μm wide below the first branch, of nearly uniform width, with no callose plug, and slightly swollen at the bases. Branching was monopodial, ramified 3–5 orders, but sometimes appeared subdichotomous. Ultimate branchlets were mostly in pairs but rarely single, straight to substraight, 7–12 μm long, and 1.5–2.5 μm wide at the base, with obtuse or subtruncate tips (Fig. 1E, 1F). Sporangia were pale brown to violet, ellipsoidal, 22–32 (~42) μm long, and 14–20 μm wide, with a round or gradually narrowing tip and base and a somewhat protruding pedicel. The length to width ratio of the sporangia was 1.4–1.8 (n = 69), with the greatest width mostly at the median, but rarely supra-median. In the dehiscence apparatus, the inner layer of the wall was discontinuous, with a pore of 3–5 μm diameter and a papilla of 1.5–2.3 μm thick (Fig. 1G, 1H). Resting organs were not seen. The morphological observations revealed that this fungus unequivocally belongs to the genus *Pseudoperonospora*, and were well consistent with the known characteristics of *Ps. urticae* (Lib.) E.S. Salmon and Ware [18–20], except for the slight differences in sporangial size (Table 1). Waterhouse and Brothers [19] noted that the sporangia of *Ps. urticae* (maximum 40 μm and average 30 μm) were larger than those of other *Pseudoperonospora* species; however, others reported smaller sporangia: 19–32 × 13–21 (ave. 25.5 × 16.6) μm by Ito [21], 18–25 (~30) × 12–16 (~20) μm by Kochman and Majewski [22]; 19–33.5 × 12.5–23 μm by Vanev et al. [23]; 16–28 × 14–21 (mostly 23–25 × 18–20) μm by Ul'yanishchev et al. [24]; 22–32 × 14–22 μm by Mazelaitis and

Table 1. Morphological comparison of the Korean specimen and previously reported *Pseudoperonospora urticae*

	The present study	Salmon and Ware [18]	Constantinescu [20]
Sporangiophores			
shape	simple, straight	simple, straight	almost straight
branching	branched acute angle	branched acute angle	-
Ultimate branchlet's tip	obtuse or subtruncate	-	round to subacute
Sporangia			
shape	ellipsoidal	ovate	-
tip	round or gradually narrowing	apiculate	-
operculum	clearly present	-	present
base	papillate	papillate	papillate
length	22–32 (~42) μm long,	22–40 (average 27) μm	25–30 μm
width	14–20 μm wide	14–22 (average 18) μm	17–19 μm
Host plant	<i>U. angustifolia</i>	<i>U. dioica</i> , <i>U. urens</i> *	<i>U. dioica</i> , <i>U. kioviensis</i>

*Constantinescu [20] stated that *U. urens* is not a host plant of *Ps. urticae*.

Staneviciene [25]; and 25–30 × 17–19 μm by Constantinescu [20]. In the Korean sample, such large sporangia were quite rare, but unambiguously present among mature sporangia with darker color, sympathizing with the opinion of Waterhouse and Brothers [19], along with a study of Yu [26] (maximum 40 μm).

Genomic DNA was extracted from the infected plant tissue of the herbarium specimen using the DNeasy Plant Mini Kit (Qiagen, Germantown, MD, USA). PCR amplification was performed with the primers ITS1-O & LR-0 for internal transcribed spacer (ITS) rDNA [27] and *cox2*-F & *cox2*-RC4 for *cox2* mtDNA [28]. Amplicons were sequenced by a DNA sequencing service (Macrogen, Seoul, Korea) with the primers used for amplification. The resulting sequences were deposited in GenBank (accession numbers: KY986684 for ITS rDNA and *cox2* mtDNA for KY986683). The sequences were edited using the DNASTAR Lasergene software package (DNASTAR, Madison, WI, USA), version 5.05. Alignments of each locus were generated using MAFFT 7 [29] with the Q-INS-1 algorithm. Minimum evolution (ME) and maximum likelihood (ML) methods were used to construct two different trees. ME analysis was done using MEGA 7.0 [30], with the default settings of the program, except for replacement with the Tamura-Nei model. For ML analysis, 1,000 rounds of random addition of sequences as well as 500 fast bootstrap replicates were performed with RAxML 7.0.3 [31] using the GTRCAT model. In the ITS rDNA and *cox2* mtDNA regions, the barcoding loci of oomycetes [28], the Korean isolate from *Urtica angustifolia* exhibited a high similarity of 99.5% (4 out of 750 characters are different) with *Ps. urticae* sensu stricto from *Urtica dioica* (AY198307, HM636048, HM636049) for the ITS sequence, but 98% (12 out of 550) with three sequences (DG3657644, HM635952, HM635953) for the *cox2* sequences. The phylogenetic trees for a combined alignment of ITS rDNA and *cox2* mtDNA were inferred using the ME and ML methods. As the two trees were congruent, only a ME tree is shown in Fig. 2. The Korean sample was a close sister-lineage to *Ps. urticae* s. s., with a maximum support in both ME and ML trees. The phylogenetic divergence between the Korean sample and *Ps. urticae* s. s. may be due to either the different host species (*Urtica angustifolia* vs. *Urtica dioica*) or the distant geographic origins (Korea vs Europe), as in this study, no morphological differences were found between them (data not shown). Further study with additional collections is needed to investigate the precise relationship of the two lineages.

Based on the morphological and phylogenetic analyses, the pathogen was identified as *Pseudoperonospora urticae*. This fungus has now been reported on six species of *Urtica*; *U. angustifolia*, *U. dioica*, *U. fissa*, *U. gracilis* (often regarded as a subspecies of *U. dioica*), *U. kioviensis*, and *U. urens* [32]. After a detailed review of downy mildews parasitic to *Urtica* spp., Constantinescu [20] suggested that both *U. gracilis* and *U. urens* are not the host plant of *Ps. urticae*, but instead they are infected by *Peronospora debaryi*, another downy mildew species occurring on *Urtica* spp. The present study confirmed that *U. angustifolia* is a rare host plant of *Ps. urticae*, and infection by *Pseudoperonospora* has




Fig. 2. Minimum evolution tree of *Pseudoperonospora* species using a combined alignment of internal transcribed spacer (ITS) rDNA and *cox2* mtDNA sequences. Bootstrapping values (minimum evolution BP/ maximum likelihood BP) higher than 60% are shown above the branches (1,000 replicates). The scale bar equals the number of nucleotide substitutions per site.

been reported only once in Far Eastern Russia [33]. To our knowledge, this is the first report of *Ps. urticae* occurring on *U. angustifolia* in Korea.

Acknowledgements

This study was supported by a grant from the National Institute of Biological Resources (NIBR), funded by of the Ministry of Environment (MOE) and the National Research Foundation of Korea (NRF), funded by the Ministry of Science, ICT & Future Planning (2016R1C1B2008013), Republic of Korea.

REFERENCES

1. Thines M, Choi YJ. Evolution, diversity, and taxonomy of the Peronosporaceae, with focus on the genus *Peronospora*. *Phytopathology* 2016;106:6-18.
2. Cohen Y. Downy mildew of cucurbits. In: Spencer DM, editor. *The downy mildews*. London: Academic Press; 1981. p. 341-54.
3. Lebeda A, Cohen Y. Cucurbit downy mildew (*Pseudoperonospora cubensis*)-biology, ecology, epidemiology, host-pathogen interaction and control. *Eur J Plant Pathol* 2011;

- 129:157-92.
4. Francis SM. *Pseudoperonospora humuli*. CMI Descr Pathog Fungi Bact 1983;769:1-2.
 5. Royle DJ, Kremheller HT. Downy mildew of the hop. In: Spencer DM, editor. The downy mildew. London: Academic Press; 1981. p. 395-419.
 6. Summers CF, Adair NL, Gent DH, McGrath MT, Smart CD. *Pseudoperonospora cubensis* and *P. humuli* detection using species-specific probes and high definition melt curve analysis. Can J Plant Pathol 2015;37:315-30.
 7. Kitner M, Lebeda A, Sharma R, Runge F, Dvořák P, Tahir A, Choi YJ, Sedláková B, Thines M. Coincidence of virulence shifts and population genetic changes of *Pseudoperonospora cubensis* in the Czech Republic. Plant Pathol 2015;64:1461-70.
 8. Polat I, Baysal O, Mercati F, Kitner M, Cohen Y, Lebeda A, Carimi F. Characterization of *Pseudoperonospora cubensis* isolates from Europe and Asia using ISSR and SRAP molecular markers. Eur J Plant Pathol 2014;139:641-53.
 9. Quesada-Ocampo LM, Granke LL, Olsen J, Gutting HC, Runge F, Thines M, Lebeda A, Hausbeck MK. The genetic structure of *Pseudoperonospora cubensis* populations. Plant Dis 2012;96:1459-70.
 10. Savory EA, Granke LL, Quesada-Ocampo LM, Varbanova M, Hausbeck MK, Day B. The cucurbit downy mildew pathogen *Pseudoperonospora cubensis*. Mol Plant Pathol 2011;12:217-26.
 11. Mitchell MN, Ocamo CM, Grünwald NJ, Mancino LE, Gent DH. Genetic and pathogenic relatedness of *Pseudoperonospora cubensis* and *P. humuli*. Phytopathology 2011;101:805-18.
 12. Runge F, Choi YJ, Thines M. Phylogenetic investigations in the genus *Pseudoperonospora* reveal overlooked species and cryptic diversity in the *P. cubensis* species cluster. Eur J Plant Pathol 2011;129:135-46.
 13. Choi YJ, Hong SB, Shin HD. A re-consideration of *Pseudoperonospora cubensis* and *P. humuli* based on molecular and morphological data. Mycol Res 2005;109:841-8.
 14. Shin HD, Choi YJ. A first check-list of Peronosporaceae from Korea. Mycotaxon 2003; 86:249-67.
 15. Shin HD, Choi YJ. Peronosporaceae of Korea. Suwon: National Institute of Agricultural Science and Technology; 2006.
 16. Zhang H, Yan X, Jiang Y, Han Y, Zhou Y. The extraction, identification and quantification of hypoglycemic active ingredients from stinging nettle (*Urtica angustifolia*). Afr J Biotechnol 2011;10:9428-37.
 17. Stepanova TA, Stusenko OV. Medicinal plants of the Russian Far East. Aust J Med Herb 2008;20:142-5.
 18. Salmon ES, Ware WM. The downy mildew of the hop and its epidemic occurrence in 1924. Ann Appl Biol 1925;12:121-51.
 19. Waterhouse GM, Brothers MP. The taxonomy of *Pseudoperonospora*. Mycol Pap 1981;148:1-28.
 20. Constantinescu O. Notes on *Pseudoperonospora*. Mycotaxon 1985;24:301-11.
 21. Ito S. Mycological flora of Japan. Tokyo: Yokendo; 1936.
 22. Kochman J, Majewski T. Głonowce (Phycomycetes); Wroslikowe (Peronosporales). Warsaw: Panik Scientific Publishing House; 1970.

23. Vanev SG, Dimitrova EG, Ilieva EI. Razred Peronosporales. Sofia: In the aedibus Academiae Scientiarum Bulgaricae; 1993.
24. Ulyanishchev VI, Osipyan LL, Kanchaveli LA, Akhundov TM. Peronosporovye Griby. In: Osipyan LL, editor. Erevan: Erevan University; 1985.
25. Mazelaitis J, Stanevicienė S. Gleivunai (Myxomycota), Peronosporiečiai (Peronosporales). Vilnius: Science and Encyclopedia Publishing House; 1995.
26. Yu Y. Flora fungorum sinicorum Vol. 6: Peronosporales. Beijing: Science Press; 1998.
27. Choi YJ, Klosterman SJ, Kummer V, Voglmayr H, Shin HD, Thines M. Multi-locus tree and species tree approaches toward resolving a complex clade of downy mildews (Straminipila, Oomycota), including pathogens of beet and spinach. *Mol Phylogenet Evol* 2015;86:24-34.
28. Choi YJ, Beakes G, Glockling S, Kruse J, Nam B, Nigrelli L, Ploch S, Shin HD, Shivas RG, Telle S, et al. Towards a universal barcode of oomycetes - a comparison of the *cox1* and *cox2* loci. *Mol Ecol Resour* 2015;15:1275-88.
29. Katoh K, Standley DM. MAFFT Multiple sequence alignment software version 7: Improvements in performance and usability. *Mol Biol Evol* 2013;30:772-80.
30. Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol* 2016;33:1870-4.
31. Stamatakis A. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 2006;22:2688-90.
32. Farr DF, Rossman AY. Fungal databases, U.S. National Fungus Collections, ARS, USDA [Internet]. Beltsville: Systematic Mycology and Microbiology Laboratory; 2016 [cited 2016 Jan 24]. Available from: <http://nt.ars-grin.gov/fungaldatabases/>.
33. Gannibal PB, Gasich EL, Berestetskiy AO, Gagkaeva TY, Khlopunova LB, Bilder IV, Levitin MM, Kolombet LV. Materials to the study of micromycetes of weeds and wild herbaceous plants in the south of Russian Far East (Primorie and Khabarovsk territories). *Nov Sist Nizs Rast* 2010;44:105-17.