

Mobilization of Heavy Metals induced by Button Mushroom Compost in Sunflower

Chang-Hoon Han and Min-Ho Yoon*

Department of Bio-Environmental Chemistry, College of Agriculture and Life Sciences, Chungnam National University, Daejeon, 34134, Korea.

ABSTRACT: This study focused on evaluating the phytoextraction of heavy metals (Co, Pb, and Zn) induced by bioaugmentation of button mushroom compost (BMC) in *Helianthus annuus* (sunflower). When the potential ability of BMC to solubilize heavy metals was assessed in a batch experiment, the inoculation with BMC could increase more the concentrations of water-soluble Co, Pb, and Cd by 35, 25, and 45% respectively, compared to those of non-inoculated soils. BMC-assisted growth promotion and metal uptake in *H. annuus* was also evaluated in a pot experiment. In comparison with non-inoculated seedlings, the inoculation led to an increase in the growth of *H. annuus* by 27, 25, and 28% in Co-, Pb-, and Zn-contaminated soils, respectively. Moreover, enhanced accumulation of Co, Pb, and Zn in the shoot and root systems was observed in inoculated plants, where metal the translocation from root to the above-ground tissues was also found to be enhanced by the BMC. Evidently, these results suggest that the BMC could be effectively employed in enhancing the phytoextraction of Co, Pb, and Zn from contaminated soils.

KEYWORDS: Mobilization, heavy metals, button mushroom compost, bioaugmentation, sunflower

Introduction

Metal contamination of soils has become one of the most significant environmental problems today. Excessive metal uptake by crop plants from the contaminated agricultural lands can result in decreased crop yield due to the inhibition of plant metabolic processes (Singh and Aggarwal, 2006). Apart from the metals with unknown biological functions (Cd, Cr, Pb, Co, Ag, Se, and Hg), essential elements (Fe, Mn, Zn, Cu, Mg, Mo, and Ni) also keep accumulating in agricultural soils by means of wastewater irrigation, animal manures and sewage sludge application, use of

fertilizer and agrochemicals (Thomas *et al.*, 2012).

The remediation of metal contaminated soils receives increasing attention (Cao *et al.*, 2007) due to the fact that metals are not easily degraded. Depending on the resource availability and nature of metal contaminated soil, different methods such as bioremediation and physical/chemical remediation have been employed in restoring the contaminated lands (Arunakumara *et al.*, 2013, Luo, 2009). However, the physical and chemical methods such as physical separation, acid leaching or electrochemical processes, are considered to be ineffective because of high cost, low efficiency, and destruction of soil structure and fertility (Jing *et al.*, 2007). In contrast, phytoremediation, a method which uses plants to extract, sequester and detoxify pollutants has received considerable attention (Arunakumara, 2011). However, the wider application of the technology has been restricted due to the limitations such as low soil thickness that can be treated, low translocation rate of metals from roots to shoots, and the slowness of the treatment (Lebeau *et al.*, 2008).

The amount of heavy metals uptake in plants varies with the mobility and the concentration of metals in soil (Chen *et al.*, 2010) and the interface between soil microbes and plant roots (rhizosphere) is displayed to

J. Mushrooms 2017 June, 15(2):61-68
<http://dx.doi.org/10.14480/JM.2017.15.2.61>
Print ISSN 1738-0294, Online ISSN 2288-8853
© The Korean Society of Mushroom Science

*Corresponding author
E-mail : mhyoon@cnu.ac.kr
Tel : +82-42-821-6733, Fax : +82-42-823-9241

Received June 13, 2017
Revised June 23, 2017
Accepted June 30, 2017

This is an Open-Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

have a great influence on the uptake of nutrients as well as on the decrease of metal toxicity (McNear, 2013). Since soil microbes could alter the metal status of the soil (Fazal and Bano, 2010), exploitation of such microbes to reduce the metal toxicity in plants is worth investigating (Rajkumar and Freitas, 2008). It was known that some metal resistant bacterial strains were proved exceptional at enhancing the growth of the host plant through different mechanisms such as the production of plant growth promoting substances, nitrogen fixation and phosphate solubilization (Hemambika *et al.*, 2013). As reported by Rajkumar *et al.* (2008), heavy metal tolerance of the microbes may be attributed to one or several mechanisms including exclusion, active removal, biosorption, and precipitation or bioaccumulation of metals both in external and intracellular spaces. Therefore, microbes having remarkable metal tolerance and plant growth-promoting abilities could play a significant role in remediation of metal-contaminated soils, because bioaugmentation with such microbes could promote phytoextraction of metals (Prapagdee *et al.*, 2013). Previous report was the results on mobilization of heavy metals induced by button mushroom compost collected in Boryeong area, South Korea (Lee *et al.*, 2015).

In the present study, button mushroom compost collected in Buyeo-gun area was employed in assessing the potential of mobilization of Co, Pb and Zn in soils and it was estimated that the effect of inoculation with BMC on plant growth and uptake of Co, Pb and Zn by *Helianthus annuus* (sunflower).

Materials and Methods

Preparation of Button mushroom compost

The waste mushroom bed from *Agaricus bisporus* (button mushroom) collected in Chungcheongnam-Do, Buyeo-gun area, South Korea was used for preparation of button mushroom compost (BMC). The waste mushroom bed was composted with pig and fowl manure as nitrogen sources and sawdust as carbon source, respectively. The BMC was prepared by mixing to the ratio of waste mushroom bed(4) : sawdust(8) : pig and fowl manure(1) and fermenting the mixture for 40 days at 30°C.

Isolation of microorganisms from BMC

Heavy metal contaminated soils were collected from

the sediment tailing in abandoned mines of Boryeong area, South Korea. Aliquots of serially diluted soil samples were spread on Tryptic soy broth (TSB, 30 g L⁻¹) agar media which was adjusted to pH 6.5 ± 0.1. The colonies on agar plate were purely isolated by repeated sub culturing at 30°C for 5 days. The preculture of each colonies was inoculated in 50mL TSB containing 0.2g - 0.4 g L⁻¹ of CoCl₂·6H₂O, 2PbCO₃·Pb(OH)₂, and Cd(NO₃)₂ at 30°C for 7 days on a horizontal shaker at 150 rpm and the cells were identified and used for further experiments.

Strain identification

The partial sequencing of 16S rRNA for the bacterial strain was done with the help of DNA sequencing service, SOLGENT, Daejeon, South Korea using universal primers, 27F (5'-AGAGTTTGATCCTGGCTCAG-3') and 1492R (5'-GGTTACCTTGTTACGACTT-3'). The online program BLAST was used in identifying the related sequences with known taxonomic information available at the databank of NCBI (<http://www.ncbi.nlm.nih.gov/BLAST>). A Phylogenetic tree was constructed using CLUSTAL X program (Thompson *et al.*, 1997), which involved sequence alignment by neighbor joining method (Saitou and Nei, 1987) and maximum parsimony using the MEGA4 program (Kumar *et al.*, 2001). Grouping of sequences was based on confidence values obtained by bootstrap analysis of 1,000 replicates. Gaps were edited in the BioEdit program and evolutionary distances were calculated using Kimura two parameter model. Reference sequences were retrieved from GenBank under the accession numbers indicated in the trees.

Mobility of the metals in soil

The impact of BMC inoculation on the mobility of metals in soil was investigated under laboratory conditions with 50 mL scaled polypropylene centrifuge tubes. Artificially contaminated soil (5 g) in the centrifuge tubes was inoculated with 10 ml TSB supplemented with 5 g BMC. After taking the weight of the tubes, they were wrapped with brown paper and placed on an orbital shaker at 200 rpm at 25°C. At the end of the period of 10 d, the weight of the tubes was recorded and 10 ml of sterile water were added to each tube to extract the soil water soluble heavy metals. The extracts were centrifuged at 10000 ×g for 10 min and filtered through a 0.45 µm nylon syringe filter (Watman,

Table 1. Change of microbial population in the compost prepared with spent mushroom from *Agaricus bisporus*. (cfu/g)

Treatments		Aerobic bacteria	Gram negative	Bacillus species	Fluorescence bacteria	Actionomycetes	Fungi
PiM+SD	Before	4.3×10^6	3.4×10^7	4.5×10^6	2.9×10^4	6.1×10^4	2.1×10^5
	After	6.8×10^6	2.2×10^7	6.3×10^6	9.7×10^4	9.4×10^4	7.3×10^4
ChM+SD	Before	3.3×10^7	1.3×10^7	2.9×10^6	5.2×10^4	4.9×10^4	6.7×10^5
	After	7.5×10^7	3.5×10^7	8.3×10^6	1.9×10^5	3.1×10^5	2.6×10^5

The compost was prepared by mixing to the ratio of waste mushroom bed(4) : sawdust(8) : pig or fowl manure(1).

PiM: Pig manure, Ch: Fowl manure, SD: Sawdust, M: spent mushroom from *Agaricus bisporus*

England) and acidified with HNO_3 to minimize an interference by organic matters. The metal contents (Co, Pb, and Zn) in the filtrate were determined using an ICP (Perkinelmer, Analyst 800, USA). Artificially contaminated soil without inoculation with the strain served as the control after centrifugation.

Effect of bioaugmentation on growth and metal uptake by *H. annuus*

A pot experiment was conducted under green house conditions at the College of Agriculture, Chungnam National University in April 2016. Several locations of soils collected from abandoned mines of Boryeong-gun as contaminated soil and a button mushroom compost in Buyeo-gun area, were respectively mixed with the ratio of 1:1, air dried and sieved (2 mm). Sterilized forest soil (by steaming at 100°C for three consecutive days) was amended with aqueous solutions of different heavy metals (Co, Pb and Zn) to achieve the final concentrations of 200 mg/kg soil. They were then kept for 2 weeks in a greenhouse for metal stabilization and used in filling the plastic pots (25 cm diameter, 35 cm height). Seeds of *Helianthus annuus* were surface sterilized by immersing in alcohol (70%) for 40 s, NaClO (1.0%) for 15 min followed by rinsing several times with sterile distilled water. Seeds sown in germination trays containing sterilized non-contaminated soil were provided with 14/10 light/dark regime and kept at 25°C for germination. Three weeks old seedlings were carefully uprooted from the germination bed and were transplanted into the plastic pots (five plants/pot) containing 300 g of metal contaminated or non-contaminated soil and allowed to grow at 25°C and 14/10 light/dark regime. The average pH of soil at the time of planting was recorded as 6.8. Three weeks later, the plants were carefully uprooted and cleaned the root surface thoroughly with distilled water. As growth parameters, fresh and dry biomass was measured and

accumulation of metals in plant biomass was quantified as described by Freitas *et al.* (2004). Each treatment had three replicates.

Results

Microbial diversity of BMC

The number of microorganisms in BMC were higher in the fowl manure than those in the pig manure as nitrogen source. BMC comprised diverse microorganisms ranging of aerobic bacteria 6.2×10^7 cfu g^{-1} , Gram negative bacteria 3.5×10^7 cfu g^{-1} , genus *Bacillus* 8.3×10^6 cfu g^{-1} , genus *Pseudomonas* 1.9×10^5 cfu g^{-1} , actinomycetes 3.1×10^5 cfu g^{-1} , and fungi 2.6×10^4 cfu g^{-1} (Table 1). Besides actinomycetes and fungi, the microbial population in the process of composting was increased to the level of $1 \times 10^1 \sim 1 \times 10^2$ g^{-1} after fermentation, compared with those of before fermentation.

Isolation and identification of microorganisms

The major group of bacterial strains from BMC were isolated and the strains with the potential ability to alleviate heavy metals were screened based on estimating the metal concentration alleviated by each bacterium in medium supplemented with 3 different heavy metals (Co, Pb, Zn). The selected strains were identified by systematic analysis using phylogenetic tree according to 16S rRNA sequence analysis. The main bacteria were identified with the close promixsity to each type strains as Genus *Bacillus*, *Pseudomonas*, *Pantoea*, *Burkholderia* and *Enterobacter* (Table 2). In addition, to estimate the metal resistance potential of the bacterial strains isolated from BMC, the amount of metals remaining in Tryptic soy broth medium was estimated 5 days after the inoculation with each strain. As was represented in Table 2, they showed the resistance of the range from 10% to 27% for heavy metals Co, Pb, and Zn, respectively. This result

Table 2. Identification of metals resistance bacteria isolated from button mushroom compost

Genus of isolated strains	Identity with type strain (%)	*Ratios of metals resistance (%)			Most closet species (Accession number)
		Co	Pb	Zn	
<i>Bacillus</i>	99.7	16	10	28	<i>Bacillus subtilis</i> (AJ276351.1)
<i>Pseudomonas</i>	99.3	15	08	20	<i>Pseudomonas koreensis</i> (AF468452.1)
<i>Enterobacter</i>	98.9	17	09	23	<i>Enterobacter ludwigii</i> (AJ853891.1)
<i>Pantoea</i>	99.36	14	10	21	<i>Pantoea agglomerans</i> (AJ233423.1)
<i>Burkholderia</i>	99.71	14	11	23	<i>Burkholderia stabilis</i> (AF148554.1)

*The amount of heavy metals alleviated by the strains in TSB medium supplemented with metals (Co, Pb and Zn) at the concentration of 200 mg L⁻¹. Values are the means of three replicates.

demonstrated that BMC with diverse microorganisms could be useful as microbial inoculant for bioremediation of metal contaminated soils.

Mobility of the metals in soil

To investigate the impact of BMC inoculation on the mobility of metals in soil, metal mobilization potential of BMC with the strains was assessed in a batch experiment with artificially contaminated soil. When alleviation effect of heavy metals was estimated by measuring the amount of metals remaining in the soil, BMC was shown to be capable of reducing the amount of metal in the order of Zn, Co and Pb. As was given in Fig. 1, the inoculation of BMC could increase the contents of water soluble metals from the soil extract, representing that heavy metals could be solubilized by BMC inoculation. The mobilization of Co, Pb and Zn was respectively 35, 25 and 45% higher than those of the control soil.

Effect of bacterial strain on growth and metal uptake by *H. annuus*

As shown in Table 1, inoculation with BMC into sunflower (*Helianthus annuus*) pots resulted in increased fresh and dry biomass of *H. annuus* plants compared to non-inoculated plants. In case of the non-inoculated plants exposed by heavy metal stress, the growth of plant was inhibited in a significant level with $p < 0.05$. For instance, Pb toxicity caused 30 and 24% reductions in fresh and dry weight of the plant, respectively. Inoculation however led to increase in plant fresh and dry weight in the presence of heavy metals. The fresh




Fig. 1. Effect on mobilization of Co, Pb and Zn in soil by inoculation with button mushroom compost. Soil without inoculation of BMC served as the control. Values are the means of three replicates. Error bars represent standard deviation.

and dry weight of the plants grown in Co contaminated soils were respectively 26 and 31% higher than those of non-inoculated plants. Similarly, in Pb contaminated soil, the percent increments were recorded as 29 and 23% respectively, and in Zn contaminated soil, the corresponding figures were 29 and 27%.

The amounts of Co, Pb and Zn accumulated in the roots and shoots of *H. annuus* grown under inoculated and non-inoculated conditions are given in Table 2. Inoculation with BMC resulted in increased accumulation of metals both in the shoots and roots. The accumulations of Co, Pb and Zn in shoots were respectively 22, 16 and 25% higher than those of non-

Table 3. Effect of inoculation with button mushroom compost on shoot and root weight of *Helianthus annuus*

Metal	Treatment	Fresh weight (g/plant)		Dry weight (g/plant)	
		Shoot	Root	Shoot	Root
Metal free soil	control	1.65 (\pm 0.035)	0.109 (\pm 0.011)	0.083 (\pm 0.006)	0.041 (\pm 0.004)
	with strain	1.89 (\pm 0.041)	0.128 (\pm 0.01)	0.109 (\pm 0.005)	0.055 (\pm 0.005)
Co	control	1.35 (\pm 0.033)	0.069 (\pm 0.005)	0.076 (\pm 0.004)	0.029 (\pm 0.004)*
	with strain	1.68 (\pm 0.045)	0.089 (\pm 0.006)	0.094 (\pm 0.004)	0.040 (\pm 0.004)*
Pb	control	1.23 (\pm 0.031)	0.058 (\pm 0.005)	0.052 (\pm 0.002)*	0.029 (\pm 0.004)*
	with strain	1.51 (\pm 0.033)	0.089 (\pm 0.004)	0.077 (\pm 0.004)*	0.041 (\pm 0.004)*
Zn	control	1.39 (\pm 0.038)	0.074 (\pm 0.003)	0.084 (\pm 0.004)*	0.030 (\pm 0.002)
	with strain	1.73 (\pm 0.041)	0.096 (\pm 0.002)	0.098 (\pm 0.003)*	0.045 (\pm 0.003)

Values are means ($n=3$) \pm standard deviation. Within each column, the means indexed by * are not significantly different at $p > 0.05$ between inoculated and non-inoculated plants according to Duncan's multiple range test.

Table 4. Effect of inoculation with button mushroom compost on accumulation and translocation of Co, Pb and Zn in *Helianthus annuus*

Metal	Treatment	Metal content (mg/kg dry weight)		Bioconcentration Factor (BCF) ^a	Translocation Factor (TF) ^b
		Shoot	Root		
Co	control	18.23 (\pm 3.75)	71.39 (\pm 4.37)	0.356	0.255
	with strain	23.27 (\pm 4.09)	89.48 (\pm 4.77)	0.447	0.260
Pb	control	11.25 (\pm 1.58)*	48.35 (\pm 5.69)	0.241	0.233
	with strain	16.71 (\pm 1.69)*	65.36 (\pm 6.12)	0.326	0.256
Zn	control	20.58 (\pm 6.02)	97.73 (\pm 6.14)*	0.488	0.210
	with strain	31.57 (\pm 7.82)	109.23 (\pm 7.12)*	0.546	0.289

inoculated plants. The corresponding accumulations for Co, Pb and Zn in roots were 36, 42 and 15% higher than those of non-inoculated plants. Regardless of inoculation or non-inoculation, the accumulation of metals in root system was found to be considerably higher than that of in shoots, which has been further confirmed by the low translocation factor (TF) for all the metals. However, TF of Co was somewhat higher than that of the other two metals. Reversely low bioconcentration factor (BCF) was also recorded from Co and Pb, compared to Zn. However, the results showed a good agreement and demonstrated that inoculation of the bacterial strain led to increase both TF and BCF of the three metals distinctly.

Discussion

Generally microorganisms isolated from heavy metals contaminated soils possess the ability to withstand against multiple pollutants as they have adapted to such environments (Pal *et al.*, 2005; Abou-Shanab *et al.*, 2007). Remedation of heavy metal contaminated soil

have been employed with both microorganism and phytoextraction. The effectiveness of the strain for phytoextraction was assessed with *Helianthus annuus*, (sunflower) was known to have an ability to accumulate biomass rapidly and take up substantial amounts of metals (Turgut *et al.*, 2004; Prapagdee *et al.*, 2013). As reported by Ouzounidou *et al.* (2005) and El-Tayeb *et al.* (2006), accumulation of plant biomass could be affected by excessive concentrations of heavy metals, which exert adverse impacts on growth and function of root system resulting in poor uptake of water and nutrients. As reported by Jiang *et al.* (2008), inoculation with *Burkholderia* sp. J62 led to increase shoot and root dry weights of corn and tomato plants. Inoculation with *Pseudomonas fluorescens* PsIA12 resulted in enhanced growth of *Zea mays* and its uptake of N, P and K (Egamberdiyeva *et al.*, 2002). Most recently, Prapagdee *et al.* (2013) reported that growth of *H. annuus* could be enhanced by the inoculation of *Micrococcus* sp. MU1 and *Klebsiella* sp. BAM1 under Cd contaminated conditions. Belimov *et al.* (2001) also observed bacterial-assisted growth enhancement in *Brassica napus*

grown in a soil contaminated with Cd. The plant growth-promoting potential of the present strain could be attributed at least partly to the phosphate solubilization ability of the strain under metal stress conditions.

Regardless of inoculation or non-inoculation, the accumulation of metals in root systems was found to be considerably higher than that of in shoots. This could primarily be attributed to the poor translocation of heavy metals from roots to shoots (Rajkumar *et al.*, 2006). However, as shown in Table 2, translocation factor of the each metal was increased with the inoculation of the strain, which was of enormous practical significance. Furthermore, metal accumulations in both shoots and roots were found to be higher in inoculated plants than those of non-inoculated plants. Similar observations were made by Rajkumar *et al.* (2008) for Zn accumulation in *H. annuus* inoculated with *Bacillus weihenstephanensis*. However, according to Wani *et al.* (2007), inoculation of *Bradyrhizobium* sp. on surface sterilized seeds of *Vigna radiate* reduced the concentration of Ni in roots, shoots and grains by 15, 19 and 22%, respectively, compared with non-inoculated plants.

The increased accumulation of metals in the presence of bacterial strain might be due to the increased uptake of metals under acidic soil conditions created by the phosphate solubilization (Rajkumar *et al.*, 2008). Inoculation of Cd-resistant bacterial strains to *Brassica napus* to a metal contaminated soil significantly increased the plant uptake of Cd when compared with the non-inoculated controls, as a result of pH reduction (Sheng and Xia, 2006). The present findings of metal mobilization are in agreement with Wu *et al.* (2006) and Prapagdee *et al.* (2012) who also reported bacteria-assisted increase in heavy metal mobilization. Generally, the low amount of metals extracted by plants from a soil is attributed mainly to the low availability of metals. As reported by several authors, the available metal content in a soil is less than 1% of the total metal content (Whiting *et al.*, 2001; Braud *et al.*, 2006). Metal availability is influenced by the nature of the metal and soil characteristics such as pH, CEC and organic matter (Kayser *et al.*, 2001; Lebeau *et al.*, 2008). Bioaugmentation could enhance metal bioavailability by increasing the concentration of the available fractions. As revealed by the present results, the release of heavy metals from the non-soluble phases to soluble phases could be facilitated

by the bacterial strain. Therefore, increased accumulation of metals, in particular Co in both the shoots and roots of *H. annuus* could be attributed to the higher water soluble metal contents in soil inoculated with bacterial strain. As reported by the results of previous studies, *H. annuus* is capable of accumulating high amounts of Pb, Cd, Cu, Zn and Co, in both the shoots and the roots (Boonyapookana *et al.*, 2005; Marchiol *et al.*, 2007). According to Braud *et al.* (2006), inoculation of *Pseudomonas aeruginosa* and *Pseudomonas fluorescens* has resulted in 113% increment of Pb content in the exchangeable fraction of the soil. However, the Pb concentration bound to free Mn oxides, organic matter and in the residual fraction remained stable. Abou-Shanab *et al.* (2006) observed an increase of extractable Ni with *Microbacterium arabinogalactanolyticum* by a factor up to 15. As reported by Baum *et al.* (2006), the concentrations in NH_4NO_3 -extractable Cd, Cu, Pb and Zn in a soil bioaugmented with ectomycorrhizal fungus *Paxillus involutus*, were 1.22-, 1.11-, 1.33- and 1.33-fold higher than those of non-bioaugmented soil, depending on the soil composition.

Conclusion

The button mushroom compost, which was prepared with waste button mushroom, fowl manure and sawdust, compromised diverse microorganisms and was found to be capable of solubilizing heavy metals (Co, Pb and Zn). Metal mobilization potential of the BMC showed that inoculation of BMC could increase the concentrations of water soluble Co, Pb and Zn in soil than those of non-inoculated soils. Inoculation with the BMC also resulted in increased shoot and root biomass and enhanced accumulation of Co, Pb and Zn in *Helianthus annuus* plants. Furthermore BMC was found to be capable of promoting metal translocation from the roots to the shoots of *H. annuus*. Therefore, BMC could be identified as an effective promoter of phytoextraction of Co, Pb and Zn from metal-contaminated soils.

Acknowledgment

This study was supported by a grant from the research project of National Institute of Horticultural & Herbal Science, Rural Development Administration, Republic of Korea.

References

- Abou-Shanab RAI, Angle JS, Chaney RL. 2006. Bacterial inoculants affecting nickel uptake by *Alyssum murale* from low, moderate and high Ni soils, *Soil Biol Biochem.* 38: 2882-2889.
- Abou-Shanab RAI, van Berkum P, Angle JS. 2007. Heavy metal resistance and genotypic analysis of metal resistance genes in gram-positive and gram-negative bacteria present in Ni-rich serpentine soil and in the rhizosphere of *Alyssum murale*, *Chemosphere* 68: 360-367.
- Arunakumara KKIU. 2011. Use of Crop Plants for Removal of Toxic Metals, pp. 439-457. In Khan MS, Zaidi A, Goel R, and Mussarrat J. (eds.), Bio-management of Metal Contaminated Soils- 2011. Springer, Springer Science + Business Media B.V.
- Arunakumara KKIU, Walpola BC, Yoon MH. 2013. Agricultural methods for toxicity alleviation in metal contaminated soils, *Korean J Soil Sci Fert.* 46: 73-80.
- Baum C, Hryniewicz K, Leinweber P, Meissner R. 2006. Heavy-metal mobilization and uptake by mycorrhizal and nonmycorrhizal willows (*Salix dasyclados*). *J Plant Nutr Soil Sci.* 169:516-522.
- Belimov AA, Hontzeas N, Safronova VI, Demchinskaya SV, Piluzza G, Bullitta S, Glick BR. 2005. Cadmium-tolerant plant growth-promoting bacteria associated with the roots of Indian mustard (*Brassica juncea* L. Czern.), *Soil Biol Biochem.* 37: 241-250.
- Belimov AA, Kunakova AM, Safronova VI, Stepanok VV, Yudkin LY, Alekseev YV, Kozhemyakov AP. 2004. Employment of rhizobacteria for the inoculation of barley plants cultivated in soil contaminated with lead and cadmium, *Microbiology* 73: 99-106.
- Belimov AA, Safronova VI, Sergeeva TA, Egorova TN, Matveyeva VA, Tsyganov VE, Borisov AY, Tikhonovich IA. 2001. Characterisation of plant growth-promoting rhizobacteria isolated from polluted soils and containing 1-aminocyclopropane-1-carboxylate deaminase, *Can J Microbiol.* 47: 642-652.
- Boonyapookana B, Parkpian P, Techapinyawat S, DeLaun RD, Jugsujinda A. 2005. Phytoaccumulation of lead by sunflower (*Helianthus annuus*), tobacco (*Nicotiana tabacum*), and vetiver (*Vetiveria zizanioides*), *J Environ Sci Health, Part A*.40: 117-137.
- Borgmann U. 2000. Methods for assessing the toxicological significance of metals in aquatic ecosystems: bioaccumulation-toxicity relationships, water concentrations and sediment spiking approaches, *Aquat Ecosyst Health.* 3: 277-289.
- Cao A, Carucci A, Lai T, La Colla P, Tamburini E. 2007. Effect of biodegradable chelating agents on heavy metals phytoextraction with *Mirabilis jalapa* and on its associated bacteria. *Euro J Soil Biol.* 43, 200-206.
- Cervantes C, Chavez J, Cardova NA, de la Mora P, Velasco JA. 1986. Resistance to metal by *Pseudomonas aeruginosa* clinical isolates, *Microbios.* 48: 159-163.
- Chen YE, Yuan S, Su YQ, Wang L. 2010. Comparison of heavy metal accumulation capacity of some indigenous mosses in Southwest China cities: a case study in Chengdu city, *Plant Soil Environ.* 56: 60-66.
- Chen YX, Wang YP, Lin Q, Luo YM. 2005. Effect of copper-tolerant rhizosphere bacteria on mobility of copper in soil and copper accumulation by *Elsholtzia splendens*, *Environ Int.* 31: 861-866.
- Egamberdiyeva D, Juraeva D, Gafurova L, Hořflich G. 2002. Promotion of plant growth of maize by plant growth promoting bacteria in different temperature and soils. In van Santen, E. (ed.), Making Conservation Tillage Conventional: Building a Future on 25 Years of Research. Proceedings of 25th Annual Southern Conservation Tillage Conference for Sustainable Agriculture. Auburn, AL 24-26 June 2002. Special Report No. 1. Alabama Agricultural Experiment Station and Auburn University, AL 36849, USA.
- El-Tayeb MA, El-Enany AE, Ahmed NL. 2006. Salicylic acid-induced adaptive response to copper stress in sunflower (*Helianthus annuus* L.), *Plant Growth Regul.* 50: 191-199.
- Fazal H, Bano A. 2010. The effect of diazotrophs (*rhizobium* and *azatobactor*) on growth and biomass of maize in lead (Pb) polluted soil, and accumulation of the lead in different parts of plant, *Pak J Bot.* 42: 4363-4370.
- Freitas H, Prasad MNV, Pratas J. 2004. Analysis of serpentinophytes from north-east of Portugal for trace metal accumulation-relevance to the management of mine environment, *Chemosphere* 54: 1625-1642.
- Gadd GM. 2004. Microbial influence on metal mobility and application for bioremediation, *Geoderma.* 122: 109-119.
- Hemambika B, Balasubramanian V, Kannan VR, James RA. 2013. Screening of chromium-resistant bacteria for plant growth-promoting activities, *Soil Sediment Contam.* 22: 717-736.
- Jiang CY, Sheng XF, Qian M, Wang QY. 2008. Isolation and characterization of a heavy metal-resistant *Burkholderia* sp. from heavy metal-contaminated paddy field soil and its potential in promoting plant growth and heavy metal accumulation in metal-polluted soil, *Chemosphere* 72: 157-164.
- Jing YD, He ZL, Yang XE. 2007. Role of soil rhizobacteria in phytoremediation of heavy metal contaminated soils, *J Zhejiang Univ SC B*.8: 192-207.
- Kayser G, Korckritz T, Markert B. 2001. Bioleaching for the decontamination of heavy metals, *Wasser Boden.* 53: 54-58.
- Klopper JW. 2003. A review of mechanisms for plant growth promotion by PGPR, pp. 81-92. In Sixth International PGPR Workshop, Calicut, India.
- Kumar S, Tamura K, Jakobsen IB, Nei M. 2001. MEGA2: molecular evolutionary genetics analysis software, *Bioinformatics* 17: 1244-1245.
- Kyeong KC, Kim YG, Lee CJ, Lee HH, Yoon MH. 2014. Effect of button mushroom compost on mobilization of heavy metals by sunflower. *J Mushrooms.* 12(3): 163-175.
- Lebeau T, Braud A, Je'ze'quel K. 2008. Performance of bioaugmentation-assisted phytoextraction applied to metal contaminated soils: A review, *Environ Pollut.* 153: 497-

- 522.
- Lee JJ, Lee HH, Kim SC, Yoo JA, Lee CJ, Yoon MH. 2015. Mobilization of heavy metals induced by button mushroom compost in sunflower. *Korean J Soil Sci Fert* 48: 469-476.
- Luo L, Ma Y, Zhang S, Wei D, Zhu YG. 2009. An inventory of trace element inputs to agricultural soils in China. *J Environ Manage.* 90: 2524-2530.
- Marchiol L, Fellet G, Perosa D, Zerbi G. 2007. Removal of trace metals by *Sorghum bicolor* and *Helianthus annuus* in a site polluted by industrial wastes: a field experience. *Plant Physiol Biochem.* 45: 379-387.
- McNear Jr DH. 2013. The rhizosphere-roots, soil and everything in between. *Nature Education Knowledge* 4: 1.
- Nautiyal CS. 1999. An efficient microbiological growth medium for screening of phosphate solubilizing microorganisms. *FEMS Microbiol Lett.* 170: 265-270.
- Ouzounidou G, Ilias I. 2005. Hormone-induced protection of sunflower photosynthetic apparatus against copper toxicity. *Bio Plantarum.* 49: 223-228.
- Pal A, Dutta S, Mukherjee PK, Paul AK. 2005. Occurrence of heavy metal resistance in microflora from serpentine soil of Andaman. *J Basic Microbiol.* 45: 207-218.
- Prapagdee B, Chumphonwong N, Khonsue N. 2012. Influence of cadmium resistant bacteria on promoting plant root elongation and increasing cadmium mobilization in contaminated soil. *Frese Eviron Bull.* 21: 1186-1191.
- Prapagdee B, Chanprasert M, Mongkolsuk S. 2013. Bioaugmentation with cadmium-resistant plant growth-promoting rhizobacteria to assist cadmium phytoextraction by *Helianthus annuus*. *Chemosphere* 92:659-666.
- Rajkumar M, Ying M, Freitas H. 2008. Characterization of metal-resistant plant-growth promoting *Bacillus weihenstephanensis* isolated from serpentine soil in Portugal. *J basic microbial.* 48:500-508.
- Saitou N, Nei M. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol.* 4:406-425.
- Sheng XF, Xia JJ. 2006. Improvement of rape (*Brassica napus*) plant growth and cadmium uptake by cadmium-resistant bacteria. *Chemosphere* 64:1036 - 1042
- Singh S, Aggarwal PK. 2006. Effect of heavy metals on biomass and yield of different crop species. *Indian J Agric Sci.* 76:688-691.
- Thomas EY, Omuetti JAI, Ogundayomi O. 2012. The effect of phosphate fertilizer on heavy metal in soils and *Amaranthus caudatu*. *Agric Biol J N Am.* 3:145-149.
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG. 1997. The Clustal X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* 24:4876-4882.
- Turgut C, Katie Pepe M, Cutright TJ. 2004. The effect of EDTA and citric acid on phytoremediation of Cd, Cr, and Ni from soil using *Helianthus annuus*. *Environ Pollut.* 131:147-154.
- Ryan PR, Dessaux Y, Thomashow LS, Weller DM. 2009. Rhizosphere engineering and management for sustainable agriculture. *Plant Soil.* 321: 363-383.
- Wani PA, Khan MS, Almas Z. 2007. Synergistic effects of the inoculation with nitrogen-fixing and phosphate-solubilizing rhizobacteria on the performance of field-grown. *J Plant Nutr Soil Sci.* 170, 283-287.
- Whiting SN, MP de Souza, Terry N. 2001. Rhizosphere Bacteria Mobilize Zn for Hyperaccumulation by *Thlaspi caerulescens*. *Environ Sci Technol.* 35:3144-3150.
- Zaidi S, Usmani S, Singh BR. 2006. Significance of *Bacillus subtilis* strain SJ-101 as a bioinoculant for concurrent plant growth promotion and nickel accumulation in *Brassica juncea*. *Chemosphere* 64:991-997.