

ON THE GAUSS MAP OF HELICOIDAL SURFACES

DONG-SOO KIM, WONYONG KIM, AND YOUNG HO KIM

ABSTRACT. We study the Gauss map G of helicoidal surfaces in the 3-dimensional Euclidean space \mathbb{E}^3 with respect to the so called Cheng-Yau operator \square acting on the functions defined on the surfaces. As a result, we completely classify the helicoidal surfaces with Gauss map G satisfying $\square G = AG$ for some 3×3 matrix A .

1. Introduction

We consider a surface M of the Euclidean 3-space \mathbb{E}^3 . The map $G : M \rightarrow S^2$ which sends each point of M to the unit normal vector to M at the point is called the Gauss map of the surface M , where S^2 is the unit sphere in \mathbb{E}^3 centered at the origin. The theory of Gauss map of a surface in a Euclidean space and a pseudo-Euclidean space is always one of interesting topics and it has been investigated from the various viewpoints by many differential geometers ([8, 9, 10, 11, 14, 16, 17, 18, 21, 23]).

It is well known that a surface M in the Euclidean 3-space \mathbb{E}^3 has constant mean curvature if and only if $\Delta G = \|dG\|^2 G$, where Δ is the Laplace operator on M corresponding to the induced metric on M from \mathbb{E}^3 ([25]). Surfaces whose Gauss map is an eigenfunction of Laplacian, that is, $\Delta G = \lambda G$ for some constant $\lambda \in \mathbb{R}$, are the planes, circular cylinders and spheres ([6]).

Generalizing the equation $\Delta G = \lambda G$, F. Dillen et al. ([12]) and C. Baikoussis et al. ([2]), respectively, studied surfaces of revolution and ruled surfaces in the Euclidean 3-space \mathbb{E}^3 such that its Gauss map G satisfies the condition

$$(1.1) \quad \Delta G = AG, \quad A \in R^{3 \times 3}.$$

As a result, they proved ([2, 12]):

Received September 13, 2016; Accepted December 22, 2016.

2010 *Mathematics Subject Classification.* 53A05, 53B25.

Key words and phrases. Gauss map, helicoidal surfaces, Laplace operator, Cheng-Yau operator.

The first author was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2015020387).

The third author was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIP) (2016R1A2B1006974).

Proposition 1.1. 1) Among the surfaces of revolution in \mathbb{E}^3 , the only ones whose Gauss map satisfies (1.1) are the planes, the spheres and the circular cylinders.

2) Among the ruled surfaces in \mathbb{E}^3 , the only ones whose Gauss map satisfies (1.1) are the planes and the circular cylinders.

A natural generalization of rotation surfaces are the helicoidal surfaces ([13]). For helicoidal surfaces C. Baikoussis et al. proved ([4]):

Proposition 1.2. Let M be a helicoidal surface in \mathbb{E}^3 . Then the Gauss map G of M satisfies (1.1) if and only if it is an open part of either a plane, a sphere or a circular cylinder.

A natural extension of the Laplace operator Δ is the so-called Cheng-Yau operator \square (or, L_1) introduced by Cheng and Yau ([7]) for the study of hypersurfaces with constant scalar curvature. For an isometric immersion $X : M \rightarrow \mathbb{R}^3$ of a surface M , Alías et al. established the following classification theorem ([1]).

Proposition 1.3. The only surfaces in \mathbb{E}^3 , which satisfy the condition $\square X = AX + b$ for some constant 3×3 matrix A and some constant vector b are either flat or an open part of a sphere.

In fact, they classified hypersurfaces in the Euclidean space \mathbb{E}^n satisfying the condition $\square X = AX + b$ for some constant $n \times n$ matrix A and some constant vector b ([1]), which extends the classification theorem for hypersurfaces in \mathbb{E}^n satisfying $\Delta X = AX + b$ given by Chen and Petrovic ([5]) and Hasanis and Vlachos ([15]).

Hence, following the condition (1.1), it is natural to ask as follows.

Question 1.4. Among helicoidal surfaces in the Euclidean 3-space \mathbb{E}^3 , which one satisfies the following condition?

$$(1.2) \quad \square G = AG, \quad A \in R^{3 \times 3}.$$

In this paper, we give a complete answer to the above question.

For surfaces of revolution whose Gauss map satisfies (1.2), we refer to [19].

The notion of generalized slant cylindrical surfaces (GSCS's) is also a natural extended one of surfaces of revolution ([20]). Surfaces of revolution, cylindrical surfaces and tubes along a plane curve are special cases of GSCS's. In [22], the first author and B. Song proved that among the GSCS's in \mathbb{E}^3 , the only ones whose Gauss map satisfies (1.1) are the planes, the spheres and the circular cylinders.

Throughout this paper, we assume that all objects are smooth and connected, unless otherwise mentioned.

2. Preliminaries

A natural generalization of rotation surfaces are the helicoidal surfaces that can be defined as follows. Let \mathbb{R}^3 have coordinates (x, y, z) . Consider the one-parameter subgroup $g_t : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ of the group of rigid motions of \mathbb{R}^3 given by

$g_t : \mathbb{R}^3 \rightarrow \mathbb{R}^3$, $g_t(x, y, z) = (x \cos t - y \sin t, x \sin t + y \cos t, z + ht)$. The motion g_t is called a helicoidal motion with axis Oz and pitch h . A helicoidal surface with axis Oz and pitch h is a surface that is invariant by g_t , for all t . When $h = 0$, they reduce to rotation surfaces.

For a surface M in \mathbb{E}^3 with Gauss map G , we denote by S the shape operator of M with respect to the Gauss map G . For each $k = 0, 1$, we put $P_0 = I, P_1 = \text{tr}(S)I - S$, where I is the identity operator acting on the tangent bundle of M . Let us define an operator $L_k : C^\infty(M) \rightarrow C^\infty(M)$ by $L_k(f) = -\text{tr}(P_k \circ \nabla^2 f)$, where $\nabla^2 f : \chi(M) \rightarrow \chi(M)$ denotes the self-adjoint linear operator metrically equivalent to the hessian of f . Then, up to signature, L_k is the linearized operator of the first variation of the $(k + 1)$ -th mean curvature arising from normal variations of the surface. Note that the operator L_0 is nothing but the Laplace operator acting on M , i.e., $L_0 = \Delta$ and $L_1 = \square$ is called the Cheng-Yau operator introduced in [7].

Now, we state a useful lemma as follows ([1]).

Lemma 2.1. *Let M be an oriented surface in \mathbb{E}^3 with Gaussian curvature K and mean curvature H . Then, the Gauss map G of M satisfies*

$$(2.1) \quad \square G = \nabla K + 2HKG,$$

where ∇K denotes the gradient of K .

Next, we need the well-known classification theorem for isoparametric surfaces in \mathbb{E}^3 , which are surfaces with constant principal curvatures.

Proposition 2.2 ([24]). *Let M be an isoparametric surface in \mathbb{E}^3 . Then M is an open part of either a plane, a sphere or a circular cylinder.*

Finally in this section, using Lemma 2.1 we give some examples of surfaces with Gauss map satisfying (1.2).

Example 2.3. (1) Flat surfaces. In this case, we have $\square G = 0$, and hence flat surfaces satisfy $\square G = AG$ for some 3×3 matrix A . Note that the matrix A must be singular.

(2) Spheres: $(x - a)^2 + (y - b)^2 + (z - c)^2 = r^2$. In this case, we have $G = \frac{1}{r}(x - a, y - b, z - c)$ so the sphere satisfies $\square G = AG$ with $A = -\frac{2}{r^3}I$, where I denotes the identity matrix.

(3) Circular cylinders: $x^2 + y^2 = r^2$. In this case, we have $G = \frac{1}{r}(x, y, 0)$. Hence, the surface M satisfies $\square G = AG$ for some nonzero matrix A of the following form:

$$A = \begin{pmatrix} 0 & 0 & * \\ 0 & 0 & * \\ 0 & 0 & * \end{pmatrix}.$$

3. Gauss map of helicoidal surfaces

We consider a regular plane curve $\alpha(s) = (x(s), 0, z(s))$ with $x(s) > 0$ in the xz plane which is defined on an interval I . A surface M in the Euclidean space

\mathbb{E}^3 defined by

$$(3.1) \quad X(s, t) = (x(s) \cos t, x(s) \sin t, z(s) + ht),$$

where h is a constant, is said to be the helicoidal surface with axis Oz , pitch h and profile curve α . If $h = 0$, then a helicoidal surface is just a surface of revolution. If $h \neq 0$, we call M a genuine helicoidal surface. For the helicoidal rigid motion $g_t : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by $g_t(x, y, z) = (x \cos t - y \sin t, x \sin t + y \cos t, z + ht)$, $t \in \mathbb{R}$, the helicoidal surface (3.1) is invariant under g_t for all t .

We assume that the profile curve $\alpha(s) = (x(s), 0, z(s))$ is of unit speed with $x(s) > 0$. The adapted frame field $\{e_1, e_2, G\}$ on the helicoidal surface M are given by

$$(3.2) \quad \begin{aligned} e_1 &= X_s = (x' \cos t, x' \sin t, z'), \\ e_2 &= \frac{1}{Q} \{-hz'X_s + X_t\}, \\ G &= e_1 \times e_2 = \frac{1}{Q}(-xz' \cos t + hx' \sin t, -hx' \cos t - xz' \sin t, xx'), \end{aligned}$$

where X_s and X_t denote the derivative of X with respect to s and t , respectively and $'$ means the derivative with respect to s . We put

$$(3.3) \quad Q = Q(s) = \{x^2 + h^2(x')^2\}^{1/2}.$$

The metric (g_{ij}) and the second fundamental form (h_{ij}) on the helicoidal surface M are, respectively, given by

$$(3.4) \quad g_{11} = 1, \quad g_{12} = g_{21} = hz', \quad g_{22} = x^2 + h^2$$

and

$$(3.5) \quad h_{11} = \frac{x\kappa}{Q}, \quad h_{12} = h_{21} = -\frac{(x')^2 h}{Q}, \quad h_{22} = \frac{x^2 z'}{Q},$$

where $\kappa = \kappa(s)$ denotes the curvature function of the curve α .

It follows from (3.4) and (3.5) that the Gaussian curvature K and the mean curvature H of M are given by

$$(3.6) \quad \begin{aligned} K &= \frac{\det(h_{ij})}{\det(g_{ij})} = \frac{x^3 z' \kappa - h^2 (x')^4}{Q^4}, \\ 2H &= \frac{g_{11} h_{22} + g_{22} h_{11} - 2g_{12} h_{12}}{\det(g_{ij})} \\ &= \frac{\kappa x(x^2 + h^2) + x^2 z' + 2h^2 (x')^2 z'}{Q^3}. \end{aligned}$$

Note that both of the curvature functions K and H depend only on the parameter s . Using (3.2), ∇K can be computed as follows:

$$(3.7) \quad \nabla K = e_1(K)e_1 + e_2(K)e_2 = PX_s + RX_t,$$

where we put

$$(3.8) \quad P = P(s) = \frac{x^2 + h^2}{Q^2} K'(s), \quad R = R(s) = -\frac{hz'}{Q^2} K'(s).$$

Now, we suppose that the Gauss map G on the helicoidal surface M satisfies $\square G = AG$ for some matrix $A = (a_{ij})$. Then, it follows from Lemma 2.1 and (3.7) that

$$(3.9) \quad PX_s + RX_t + 2HKG = AG.$$

Putting $X_s, X_t = (-x \sin t, x \cos t, h)$ and G in (3.2) into the equation (3.9), we get the following:

$$(3.10) \quad \begin{aligned} & QPx' \cos t - QRx \sin t + 2HK(hx' \sin t - xz' \cos t) \\ & = a_{11}(hx' \sin t - xz' \cos t) + a_{12}(-hx' \cos t - xz' \sin t) + a_{13}xx', \end{aligned}$$

$$(3.11) \quad \begin{aligned} & QPx' \sin t + QRx \cos t + 2HK(-hx' \cos t - xz' \sin t) \\ & = a_{21}(hx' \sin t - xz' \cos t) + a_{22}(-hx' \cos t - xz' \sin t) + a_{23}xx' \end{aligned}$$

and

$$(3.12) \quad \begin{aligned} & QPz' + QRh + 2HKxx' \\ & = a_{31}(hx' \sin t - xz' \cos t) + a_{32}(-hx' \cos t - xz' \sin t) + a_{33}xx'. \end{aligned}$$

Since $\cos t, \sin t$ and 1 are linearly independent and their coefficients in (3.10) depend only on the parameter s , we get from (3.10)

$$(3.13) \quad QPx' - 2HKxz' = -a_{11}xz' - a_{12}hx',$$

$$(3.14) \quad -QRx + 2HKhx' = a_{11}hx' - a_{12}xz'$$

and

$$(3.15) \quad a_{13}xx' = 0.$$

Similarly, we obtain from (3.11)

$$(3.16) \quad QRx - 2HKhx' = -a_{21}xz' - a_{22}hx',$$

$$(3.17) \quad QPx' - 2HKxz' = a_{21}hx' - a_{22}xz'$$

and

$$(3.18) \quad a_{23}xx' = 0.$$

From (3.12), we also get the following:

$$(3.19) \quad a_{31}hx' - a_{32}xz' = 0,$$

$$(3.20) \quad -a_{31}xz' - a_{32}hx' = 0$$

and

$$(3.21) \quad QPz' + QRh + 2HKxx' = a_{33}xx'.$$

It follows from (3.13) and (3.17) that

$$(3.22) \quad (a_{12} + a_{21})hx' = (a_{22} - a_{11})xz'.$$

From (3.14) and (3.16) we also get

$$(3.23) \quad (a_{12} + a_{21})xz' = -(a_{22} - a_{11})hx'.$$

Combining (3.22) and (3.23), we have

$$(3.24) \quad \{(a_{12} + a_{21})^2 + (a_{22} - a_{11})^2\}hxx'z' = 0.$$

Let us put $J = \{s \in I \mid x'(s)z'(s) \neq 0\}$. We divide by two cases as follows.

Case 1. J is empty. In this case, $x'(s)z'(s)$ identically vanishes on the domain I of s . If $x(s)$ is a constant r , then $X(s, t)$ is nothing but a parametrization of the circular cylinder M given by $x^2 + y^2 = r^2$, which is flat.

If $z(s)$ is a constant, then we may assume $x(s) = s + c$, and hence M is a helicoid. In this case, from (3.3), (3.6) and (3.8) we have

$$(3.25) \quad Q = \{(s + c)^2 + h^2\}^{1/2}, K = -\frac{h^2}{Q^4}, H = 0, P = K'(s), R = 0.$$

Hence, it follows from (3.17) that

$$(3.26) \quad QP = a_{21}h,$$

which contradicts to the equations in (3.25). Therefore, this case cannot occur.

Case 2. Suppose that the subinterval J is nonempty. Then, from (3.15), (3.18) and (3.24) we see that the matrix A is of the following form:

$$A = \begin{pmatrix} \lambda & \mu & 0 \\ -\mu & \lambda & 0 \\ 0 & 0 & \nu \end{pmatrix}.$$

Hence, (3.14) and (3.21), respectively, reduce to

$$(3.27) \quad -QRx + 2HKhx' = \lambda hx' - \mu xz'$$

and

$$(3.28) \quad QPz' + QRh + 2HKxx' = \nu xx'.$$

On the other hands, (3.8) shows that

$$(3.29) \quad QPz' + QRh = \frac{x^2 z'}{Q} K'(s) = -\frac{x^2}{h} QR.$$

Hence, (3.28) becomes

$$(3.30) \quad -QRx + 2HKhx' = \nu hx'.$$

Together with (3.27), this equation implies

$$(3.31) \quad (\lambda - \nu)hx' = \mu xz'.$$

We divide by two subcases as follows.

Subcase 2-1. Suppose that $\lambda = \nu$. In this case, $\mu = 0$. Hence we have $A = \lambda I$, where I is the identity matrix. Thus, Lemma 2.1 yields that K is constant and

$2HK = \lambda$. If the Gaussian curvature K is nonzero, then the mean curvature H is also constant. Hence, Proposition 2.2 implies that M is an open part of a sphere, which leads to a contradiction. Thus, the Gaussian curvature K vanishes identically, that is, M is flat.

Subcase 2-2. Suppose that $\lambda \neq \nu$. In this case, (3.31) shows that $\mu \neq 0$ because J is not empty. It follows from (3.31) that

$$(3.32) \quad z = a \ln x + b,$$

where $a = (\lambda - \nu)h/\mu$ and b is a constant. Without loss of generality, we may assume that $b = 0$.

Next, we claim the following.

Claim. Let M denote a helicoidal surface given by

$$X(s, t) = (s \cos t, s \sin t, a \ln s + ht).$$

Then, the Gauss map G of the surface M does not satisfy $\square G = AG$ for any matrix A .

Proof. Note that the generating curve is not of unit speed. For simplicity, we assume that $a = 1$. Then, just as in the above argument we proceed as follows. The adapted frame field $\{e_1, e_2, G\}$ on the helicoidal surface M are given by

$$(3.33) \quad \begin{aligned} e_1 &= \frac{s}{\sqrt{s^2 + 1}}X_s, \\ e_2 &= \frac{1}{Q} \left\{ \frac{-h}{\sqrt{s^2 + 1}}X_s + \frac{\sqrt{s^2 + 1}}{s}X_t \right\}, \\ G &= e_1 \times e_2 = \frac{1}{Q}(-\cos t + h \sin t, -h \cos t - \sin t, s), \end{aligned}$$

where we put

$$(3.34) \quad Q = Q(s) = \{s^2 + h^2 + 1\}^{1/2}.$$

The metric (g_{ij}) and the second fundamental form (h_{ij}) on the helicoidal surface M are, respectively, given by

$$(3.35) \quad g_{11} = 1 + \frac{1}{s^2}, \quad g_{12} = g_{21} = \frac{h}{s}, \quad g_{22} = s^2 + h^2$$

and

$$(3.36) \quad h_{11} = \frac{-1}{sQ}, \quad h_{12} = h_{21} = \frac{-h}{Q}, \quad h_{22} = \frac{s}{Q}.$$

It follows from (3.35) and (3.36) that the Gaussian curvature K and the mean curvature H of M are, respectively, given by

$$(3.37) \quad K = \frac{-(h^2 + 1)}{Q^4}, \quad 2H = \frac{(h^2 + 1)}{sQ^3}.$$

The gradient ∇K is also given by

$$(3.38) \quad \begin{aligned} \nabla K &= e_1(K)e_1 + e_2(K)e_2 \\ &= PX_s + RX_t, \end{aligned}$$

where we put

$$(3.39) \quad P = \frac{s^2 + h^2}{Q^2}K'(s), \quad R = \frac{-h}{sQ^2}K'(s).$$

Suppose that the Gauss map G of the surface M satisfies $\square G = AG$ for a matrix A . Then the matrix A must be of the form in Case 2 and we get

$$(3.40) \quad QP - 2HK = -\lambda - \mu h$$

and

$$(3.41) \quad -QRs + 2HKh = \lambda h - \mu.$$

Hence, combining (3.40) and (3.41) we obtain

$$(3.42) \quad Q(Ph - Rs) = -\mu(h^2 + 1).$$

Since $Ph - Rs = hK'(s)$, (3.42) shows that

$$(3.43) \quad K'(s) = \frac{-\mu(h^2 + 1)}{Qh},$$

which contradicts (3.37). This completes the proof of Claim. \square

Finally, combining Cases 1 and 2 we get the following classification theorem.

Theorem 3.1. *Let M be a genuine helicoidal surface in the Euclidean 3-space \mathbb{E}^3 . Then the Gauss map G of M satisfies $\square G = AG$ for some 3×3 matrix A if and only if it is flat.*

For flat helicoidal surfaces, we refer to [3].

For a genuine helicoidal surface M with Gauss map G satisfying $\square G = AG$ for some nonzero 3×3 matrix A , as a corollary we get the following.

Corollary 3.2. *Let M be a genuine helicoidal surface in the Euclidean 3-space \mathbb{E}^3 . Then the Gauss map G of M satisfies $\square G = AG$ for some nonzero 3×3 matrix A if and only if it is an open part of a circular cylinder.*

Proof. Suppose that $\square G = AG$ for some nonzero 3×3 matrix A . Then, it follows from Theorem 3.1 that the surface M is flat, hence we have $AG = \square G = 0$. If we denote by V the kernel space of the matrix A , the image of the Gauss map G lies in the space V . Since A is nonzero, there exists a unit vector $a = (a_1, a_2, a_3)$ which is orthogonal to V . It follows from (3.2) that

$$(3.44) \quad a_1(-xz' \cos t + hx' \sin t) + a_2(-hx' \cos t - xz' \sin t) + a_3xx' = 0,$$

which shows that

$$(3.45) \quad a_1xz' + a_2hx' = a_1hx' - a_2xz' = a_3xx' = 0.$$

The equations in (3.45) reduce to

$$(3.46) \quad (a_1^2 + a_2^2)xz' = a_3xx' = 0.$$

If $J = \{s \in I \mid x'(s)z'(s) \neq 0\}$ is nonempty, then (3.46) implies that $a = 0$. This contradiction shows that J is empty. Hence, it follows from Case 1 in the proof of Theorem 3.1 that the surface M is an open part of a circular cylinder.

The converse follows from Example 2.3. \square

With the help of the classification theorem for surfaces of revolution ([19]), we obtain:

Theorem 3.3. *Let M be a helicoidal surface in the Euclidean 3-space \mathbb{E}^3 . Then we have the following.*

1) *The Gauss map G of M satisfies $\square G = AG$ for some 3×3 matrix A if and only if it is either a flat surface or an open part of a sphere.*

2) *The Gauss map G of M satisfies $\square G = AG$ for some nonzero 3×3 matrix A if and only if it is isoparametric, that is, an open part of either a plane, a sphere or a circular cylinder.*

References

- [1] L. J. Alias and N. Gürbüz, *An extension of Takahashi theorem for the linearized operators of the higher order mean curvatures*, *Geom. Dedicata* **121** (2006), 113–127.
- [2] C. Baikoussis and D. E. Blair, *On the Gauss map of ruled surfaces*, *Glasgow Math. J.* **34** (1992), no. 3, 355–359.
- [3] C. Baikoussis and T. Koufogiorgos, *Helicoidal surfaces with prescribed mean or Gaussian curvature*, *J. Geom.* **63** (1998), no. 1-2, 25–29.
- [4] C. Baikoussis and L. Verstraelen, *On the Gauss map of helicoidal surfaces*, *Rend. Sem. Mat. Messina Ser. II* **2(16)** (1993), 31–42.
- [5] B.-Y. Chen and M. Petrovic, *On spectral decomposition of immersions of finite type*, *Bull. Austral. Math. Soc.* **44** (1991), no. 1, 117–129.
- [6] B.-Y. Chen and P. Piccinni, *Submanifolds with finite type Gauss map*, *Bull. Austral. Math. Soc.* **35** (1987), no. 2, 161–186.
- [7] S. Y. Cheng and S. T. Yau, *Hypersurfaces with constant scalar curvature*, *Math. Ann.* **225** (1977), no. 3, 195–204.
- [8] M. Choi, D.-S. Kim, and Y. H. Kim, *Helicoidal surfaces with pointwise 1-type Gauss map*, *J. Korean Math. Soc.* **46** (2009), no. 1, 215–223.
- [9] M. Choi, D.-S. Kim, Y. H. Kim, and D. W. Yoon, *Circular cone and its Gauss map*, *Colloq. Math.* **129** (2012), no. 2, 203–210.
- [10] S. M. Choi, *On the Gauss map of surfaces of revolution in a 3-dimensional Minkowski space*, *Tsukuba J. Math.* **19** (1995), no. 2, 351–367.
- [11] ———, *On the Gauss map of ruled surfaces in a 3-dimensional Minkowski space*, *Tsukuba J. Math.* **19** (1995), no. 2, 285–304.
- [12] F. Dillen, J. Pas, and L. Verstraelen, *On the Gauss map of surfaces of revolution*, *Bull. Inst. Math. Acad. Sinica* **18** (1990), no. 3, 239–246.
- [13] M. P. do Carmo and M. Dajczer, *Helicoidal surfaces with constant mean curvature*, *Tohoku Math. J. (2)* **34** (1982), no. 3, 425–435.
- [14] U. Dursun, *Flat surfaces in the Euclidean space \mathbb{E}^3 with pointwise 1-type Gauss map*, *Bull. Malays. Math. Sci. Soc. (2)* **33** (2010), no. 3, 469–478.
- [15] T. Hasanis and T. Vlachos, *Hypersurfaces of E_{n+1} satisfying $\Delta x = Ax + B$* , *J. Austral. Math. Soc. Ser. A* **53** (1992), no. 3, 377–384.

- [16] U.-H. Ki, D.-S. Kim, Y. H. Kim, and Y.-M. Roh, *Surfaces of revolution with pointwise 1-type Gauss map in Minkowski 3-space*, Taiwanese J. Math. **13** (2009), no. 1, 317–338.
- [17] D.-S. Kim, *On the Gauss map of quadric hypersurfaces*, J. Korean Math. Soc. **31** (1994), no. 3, 429–437.
- [18] ———, *On the Gauss map of hypersurfaces in the space form*, J. Korean Math. Soc. **32** (1995), no. 3, 509–518.
- [19] D.-S. Kim, J. R. Kim, and Y. H. Kim, *Cheng-Yau operator and Gauss map of surfaces of revolution*, Bull. Malays. Math. Sci. Soc. **39** (2016), no. 4, 1319–1327.
- [20] D.-S. Kim and Y. H. Kim, *Surfaces with planar lines of curvature*, Honam Math. J. **32** (2010), no. 4, 777–790.
- [21] D.-S. Kim, Y. H. Kim and D. W. Yoon, *Extended B-scrolls and their Gauss maps*, Indian J. Pure Appl. Math. **33** (2002), no. 7, 1031–1040.
- [22] D.-S. Kim and B. Song, *On the Gauss map of generalized slant cylindrical surfaces*, J. Korea Soc. Math. Educ. Ser. B: Pure Appl. Math. **20** (2013), no. 3, 149–158.
- [23] Y. H. Kim and N. C. Turgay, *Surfaces in \mathbb{E}^3 with L_1 -pointwise 1-type Gauss map*, Bull. Korean Math. Soc. **50** (2013), no. 3, 935–949.
- [24] T. Levi-Civita, *Famiglie di superficie isoparametriche nell'ordinario spazio euclideo*, Atti. Accad. naz. Lincei. Rend. Cl. Sci. Fis. Mat. Natur. **26** (1937), 355–362.
- [25] E. A. Ruh and J. Vilms, *The tension field of the Gauss map*, Trans. Amer. Math. Soc. **149** (1970), 569–573.

DONG-SOO KIM
 DEPARTMENT OF MATHEMATICS
 CHONNAM NATIONAL UNIVERSITY
 KWANGJU 500-757, KOREA
E-mail address: dosokim@chonnam.ac.kr

WONYONG KIM
 DEPARTMENT OF MATHEMATICS
 CHONNAM NATIONAL UNIVERSITY
 KWANGJU 500-757, KOREA
E-mail address: yong4625@naver.com

YOUNG HO KIM
 DEPARTMENT OF MATHEMATICS
 KYUNGPOOK NATIONAL UNIVERSITY
 TAEGU 702-701, KOREA
E-mail address: yhkim@knu.ac.kr