

ON WEAKLY 2-ABSORBING PRIMARY SUBMODULES OF MODULES OVER COMMUTATIVE RINGS

AHMAD YOUSEFIAN DARANI, FATEMEH SOHEILNIA, UNSAL TEKIR,
AND GULSEN ULUCAK

ABSTRACT. Assume that M is an R -module where R is a commutative ring. A proper submodule N of M is called a weakly 2-absorbing primary submodule of M if $0 \neq abm \in N$ for any $a, b \in R$ and $m \in M$, then $ab \in (N : M)$ or $am \in M\text{-rad}(N)$ or $bm \in M\text{-rad}(N)$. In this paper, we extended the concept of weakly 2-absorbing primary ideals of commutative rings to weakly 2-absorbing primary submodules of modules. Among many results, we show that if N is a weakly 2-absorbing primary submodule of M and it satisfies certain condition $0 \neq I_1I_2K \subseteq N$ for some ideals I_1, I_2 of R and submodule K of M , then $I_1I_2 \subseteq (N : M)$ or $I_1K \subseteq M\text{-rad}(N)$ or $I_2K \subseteq M\text{-rad}(N)$.

1. Introduction

Throughout this paper, we suppose that all rings are commutative with $1 \neq 0$. Let M be an R -module. A submodule N of M is called a proper submodule if $N \neq M$. Let N be a proper submodule of M . $(N : M)$ is the set of all $r \in R$ such that $rM \subseteq N$ for any submodule N of M . Then the radical of N , denoted by $M\text{-rad}(N)$, is defined as the intersection of all prime submodules of M containing N . If M has no prime submodule containing N , then $M\text{-rad}(N) = M$.

Following the concept of 2-absorbing ideals of commutative rings as in [2] and [10] and the concept of weakly prime ideals of commutative rings as in [8], Badawi and Darani introduced the concept of weakly 2-absorbing ideals in commutative rings as in [3]. Afterwards, in [5], Badawi, Tekir and Yetkin introduced the concept of 2-absorbing primary ideals as a generalization of primary ideals. Also, the concept of weakly 2-absorbing primary ideals, which is a generalization of weakly primary as in [1], was studied extensively by Badawi, Tekir and Yetkin, see [6]. A proper ideal I of R is called 2-absorbing (weakly 2-absorbing) primary ideal if $abc \in I$ ($0 \neq abc \in I$) for any $a, b, c \in R$,

Received August 16, 2016; Revised December 6, 2016; Accepted March 7, 2017.

2010 *Mathematics Subject Classification.* 13C99, 13C13.

Key words and phrases. 2-absorbing submodule, 2-absorbing primary submodule, weakly 2-absorbing primary submodule.

then $ab \in I$ or $ac \in \sqrt{I}$ or $bc \in \sqrt{I}$. Recently, Mostafanasab and Darani generalized the concept of ϕ -2-absorbing primary ideals of commutative rings as in [4] to ϕ - n -absorbing primary ideals in commutative rings, see [13].

The concept of 2-absorbing submodule and weakly 2-absorbing submodules, generalizations of prime submodules and weakly prime submodules, respectively, were studied inclusively by Moradi, Azizi and other authours, see for example, [7]-[12]. A proper submodule N of M is called 2-absorbing (weakly 2-absorbing) submodule if $abm \in N$ ($0 \neq abm \in N$) for some $a, b \in R$ and $m \in M$, then $ab \in (N : M)$ or $am \in N$ or $bm \in N$. For general informations of ideals and submodules, we refer the reader to [9] and [15].

In this study, we investigate weakly 2-absorbing primary submodules. Recall that a proper submodule N of M is said to be 2-absorbing (weakly 2-absorbing) primary submodule of M if $abm \in N$ ($0 \neq abm \in N$) for any $a, b \in R$ and $m \in M$, then $ab \in (N : M)$ or $am \in M\text{-rad}(N)$ or $bm \in M\text{-rad}(N)$.

Our main aim is to answer the question: If N is a weakly 2-absorbing primary submodule of an R -module M and $0 \neq I_1I_2K \subseteq N$ for some ideals I_1, I_2 of R and some submodule K of M , does it follow that $I_1I_2 \subseteq (N : M)$ or $I_1K \subseteq M\text{-rad}(N)$ or $I_2K \subseteq M\text{-rad}(N)$? (see, Theorem 2 and Theorem 3). Among some results of this paper, it is shown that (Theorem 7) if $M\text{-rad}(0)$ is a prime submodule, then N is a weakly 2-absorbing primary submodule if and only if N is a 2-absorbing primary submodule. In Theorem 8 (Theorem 9), it is seen that if N is a weakly 2-absorbing primary submodule, then $(N : M)$ is a weakly 2-absorbing primary ideal (hence if N is a weakly 2-absorbing primary submodule, then $(N : M)$ is a weakly 2-absorbing primary ideal). In Theorem 11, it is obtained that if $N = N_1 \times N_2$ is a weakly 2-absorbing primary submodule, then $N = 0$ or N is 2-absorbing primary.

2. On weakly 2-absorbing primary submodules

Definition 1. A proper submodule N of an R -module M is called a weakly 2-absorbing primary submodule of M if $0 \neq abm \in N$ for any $a, b \in R$ and $m \in M$, then $ab \in (N : M)$ or $am \in M\text{-rad}(N)$ or $bm \in M\text{-rad}(N)$.

Proposition 1. *Let N be a weakly 2-absorbing primary submodule of an R -module M . Assume that K is a submodule of M with $N \subsetneq K$. Then N is a weakly 2-absorbing primary submodule of K .*

Proof. Let $a, b \in R$ and $k \in K$ with $0 \neq abk \in N$. Then $ab \in (N : M)$ or $ak \in M\text{-rad}(N)$ or $bk \in M\text{-rad}(N)$ as N is a weakly 2-absorbing primary. Thus $ab \in (N : K)$ or $ak \in K\text{-rad}(N)$ or $bk \in K\text{-rad}(N)$ since $(N : M) \subseteq (N : K)$. \square

The following result is an analogue of [6, Theorem 2.18].

Proposition 2. *Let N, K be submodules of an R -module M with $K \subseteq N$. If N is a weakly 2-absorbing primary submodule of M , then N/K is a weakly*

2-absorbing primary submodule of M/K . The converse is true when K is a weakly 2-absorbing primary submodule.

Proof. Assume that N is a weakly 2-absorbing primary submodule of M . Let $a, b \in R$ and $m + K \in M/K$ where $0_{M/K} \neq ab(m + K) \in N/K$. Since $ab(m + K) \neq 0_{M/K}$, we get $abm \in N$ and $abm \notin K$. If $abm = 0$, we obtain $abm + K = 0_{M/K}$. So $abm \neq 0$. Thus $ab \in (N : M)$ or $am \in M\text{-rad}(N)$ or $bm \in M\text{-rad}(N)$ as N is weakly 2-absorbing primary. Consequently, we get $ab \in (N/K : M/K)$ or $am + K = a(m + K) \in M\text{-rad}(N)/K = M/K\text{-rad}(N/K)$ or $bm + K = b(m + K) \in M\text{-rad}(N)/K = M/K\text{-rad}(N/K)$. Conversely, let K be a weakly 2-absorbing primary submodule. Assume that N/K is a weakly 2-absorbing primary submodule of M/K . Let $a, b \in R$ and $m \in M$ where $0 \neq abm \in N$. Then we have $abm + K \in N/K$. If $abm + K = 0_{M/K}$, then $abm \in K$. Thus $ab \in (K : M)$ or $am \in M\text{-rad}(K)$ or $bm \in M\text{-rad}(K)$, since K is weakly 2-absorbing primary. Therefore, $ab \in (N : M)$ or $am \in M\text{-rad}(N)$ or $bm \in M\text{-rad}(N)$, since $K \subseteq N$. Let $abm + K = ab(m + K) \neq 0_{M/K}$. Then $ab \in (N/K : M/K)$ or $a(m + K) \in M/K\text{-rad}(N/K)$ or $b(m + K) \in M/K\text{-rad}(N/K)$. Thus $ab \in (N : M)$ or $am \in M\text{-rad}(N)$ or $bm \in M\text{-rad}(N)$. \square

The following result is an analogue of [6, Theorem 2.20].

Proposition 3. *Let M be an R -module and S be a multiplicatively closed subset of R . If N is a weakly 2-absorbing primary submodule of M where $(N : M) \cap S = \emptyset$, then $S^{-1}N$ is a weakly 2-absorbing primary submodule of $S^{-1}M$.*

Proof. Let $0 \neq \frac{r}{s} \frac{t}{k} \frac{m}{l} = \frac{rtm}{skl} \in S^{-1}N$ where $r, t \in R$, $s, k, l \in S$ and $m \in M$. Then there is an element u of S such that $0 \neq urtm \in N$. Hence we get $urt \in (N : M)$ or $urm \in M\text{-rad}(N)$ or $tm \in M\text{-rad}(N)$ since N is weakly 2-absorbing primary. Then $\frac{r}{s} \frac{t}{k} = \frac{urt}{usk} \in S^{-1}(N : M) \subseteq (S^{-1}N : S^{-1}M)$ or $\frac{r}{s} \frac{m}{l} = \frac{urm}{ust} \in S^{-1}(M\text{-rad}(N)) = S^{-1}M\text{-rad}(S^{-1}N)$ or $\frac{t}{k} \frac{m}{l} = \frac{tm}{kl} \in S^{-1}(M\text{-rad}(N)) = S^{-1}M\text{-rad}(S^{-1}N)$. \square

Definition 2. Let N be a weakly 2-absorbing primary submodule of M . (a, b, m) is called a triple-zero of N if $abm = 0$, $ab \notin (N : M)$, $am \notin M\text{-rad}(N)$ and $bm \notin M\text{-rad}(N)$.

Note that if N is a weakly 2-absorbing primary submodule of M and there is no triple-zero of N , then N is a 2-absorbing primary submodule of M .

Proposition 4. *Let N be a weakly 2-absorbing primary submodule of M and K be a proper submodule of M with $K \subseteq N$. Then for any $a, b \in R$ and $m \in M$, (a, b, m) is a triple-zero of N if and only if $(a, b, m + K)$ is a triple-zero of N/K .*

Proof. Let (a, b, m) be a triple-zero of N for some $a, b \in R$ and $m \in M$. Then $abm = 0$, $ab \notin (N : M)$, $am \notin M\text{-rad}(N)$ and $bm \notin M\text{-rad}(N)$. By Proposition 2, we get that N/K is a weakly 2-absorbing primary submodule of M/K . Thus $ab(m + K) = K$, $ab \notin (N/K : M/K)$, $a(m + K) \notin M\text{-rad}(N)/K$

and $b(m + K) \notin M\text{-rad}(N)/K$. Hence $(a, b, m + K)$ is a triple-zero of N/K . Conversely, assume that $(a, b, m + K)$ is a triple-zero of N/K . Suppose that $abm \neq 0$. Then $abm \in N$ since $ab(m + K) = K$. Thus $ab \in (N : M)$ or $am \in M\text{-rad}(N)$ or $bm \in M\text{-rad}(N)$ as N is weakly 2-absorbing primary, a contradiction. So it must be $abm = 0$. Consequently, (a, b, m) is a triple-zero of N . \square

The following result is an analogue of [6, Theorem 2.9].

Theorem 1. *Let N be weakly 2-absorbing primary submodule of M and (a, b, m) be a triple-zero of N for some $a, b \in R$ and $m \in M$. Then the followings hold.*

- (1) $abN = am(N : M) = bm(N : M) = 0$.
- (2) $a(N : M)N = b(N : M)N = m(N : M)^2 = 0$.

Proof. Suppose that (a, b, m) is a triple-zero of N for some $a, b \in R$ and $m \in M$.

(1) Assume that $abN \neq 0$. Then there is an element $n \in N$ where $abn \neq 0$. Thus $ab(m + n) = abm + abn = abn \neq 0$. So $a(m + n) \in M\text{-rad}(N)$ or $b(m + n) \in M\text{-rad}(N)$ since $ab \notin (N : M)$ and N is weakly 2-absorbing primary. Hence $am \in M\text{-rad}(N)$ or $bm \in M\text{-rad}(N)$, which is a contradiction. Now, we suppose that $am(N : M) \neq 0$. Thus there exists an element $r \in (N : M)$ where $arm \neq 0$. Hence $a(r + b)m = arm + abm = arm \neq 0$ and since $am \notin M\text{-rad}(N)$, $bm \in M\text{-rad}(N)$ or $ab \in (N : M)$, a contradiction. Similarly, it can be easily seen that $bm(N : M) = 0$.

(2) Assume that $a(N : M)N \neq 0$. Then there are $r \in (N : M)$, $n \in N$ such that $arn \neq 0$. By (1), we get $a(b + r)(m + n) = arn \neq 0$ and so $a(b + r) \in (N : M)$ or $a(m + n) \in M\text{-rad}(N)$ or $(b + r)(m + n) \in M\text{-rad}(N)$. Therefore, we obtain $ab \in (N : M)$ or $am \in M\text{-rad}(N)$ or $bm \in M\text{-rad}(N)$, a contradiction. In a similar way, one can easily seen that $b(N : M)N = 0$. Let $m(N : M)^2 \neq 0$. Thus there exist $r, s \in (N : M)$ where $mrs \neq 0$. By (1), we get $(a + r)(b + s)m = rsm \neq 0$. Thus we have $(a + r)m \in M\text{-rad}(N)$ or $(b + s)m \in M\text{-rad}(N)$ or $(a + r)(b + s) \in (N : M)$ and so we get $am \in M\text{-rad}(N)$ or $bm \in M\text{-rad}(N)$ or $ab \in (N : M)$, a contradiction. Consequently, $m(N : M)^2 = 0$. \square

The following two results are an analogue of [6, Theorem 2.10].

Lemma 1. *Assume that N is a weakly 2-absorbing primary submodule of an R -module M that is not 2-absorbing primary. Then $(N : M)^2N = 0$. In particular, $(N : M)^3 \subseteq \text{Ann}(M)$.*

Proof. Suppose that N is a weakly 2-absorbing primary submodule of an R -module M that is not 2-absorbing primary. Then there is a triple-zero (a, b, m) of N for some $a, b \in R$ and $m \in M$. Assume that $(N : M)^2N \neq 0$. Thus there exist $r, s \in (N : M)$ and $n \in N$ with $rsn \neq 0$. By Theorem 1, we get $(a + r)(b + s)(n + m) = rsn \neq 0$. Then we have $(a + r)(b + s) \in (N : M)$ or $(a + r)(n + m) \in M\text{-rad}(N)$ or $(b + s)(n + m) \in M\text{-rad}(N)$ and so $ab \in (N : M)$ or $am \in M\text{-rad}(N)$ or $bm \in M\text{-rad}(N)$, which is a contradiction. Thus we

get $(N : M)^2N = 0$. We get $(N : M)^3 \subseteq ((N : M)^2N : M) = (0 : M) = \text{Ann}(M)$. □

Proposition 5. *Let M be a multiplication R -module and N be a weakly 2-absorbing primary submodule of M that is not 2-absorbing primary. Then $N^3 = 0$.*

Proof. We have that $(N : M)M = N$ since M is multiplication module. Then $N^3 = (N : M)^3M = (N : M)^2N = 0$. Consequently, $N^3 = 0$. □

Definition 3. Let N be a weakly 2-absorbing primary submodule of an R -module M and let $0 \neq I_1I_2K \subseteq N$ for some ideals I_1, I_2 of R and some submodule K of M . N is called free triple-zero in regard to I_1, I_2, K if (a, b, m) is not a triple-zero of N for every $a \in I_1, b \in I_2$ and $m \in K$.

The following result and its proof are analogues of [6, Theorem 2.29] and its proof.

Lemma 2. *Let N be a weakly 2-absorbing primary submodule of M . Assume that $abK \subseteq N$ for some $a, b \in R$ and some submodule K of M where (a, b, m) is not a triple-zero of N for every $m \in K$. If $ab \notin (N : M)$, then $aK \subseteq M\text{-rad}(N)$ or $bK \subseteq M\text{-rad}(N)$.*

Proof. Assume that $aK \not\subseteq M\text{-rad}(N)$ and $bK \not\subseteq M\text{-rad}(N)$. Then there are $x, y \in K$ such that $ax \notin M\text{-rad}(N)$ and $by \notin M\text{-rad}(N)$. We get $bx \in M\text{-rad}(N)$ since N is a weakly 2-absorbing primary submodule, (a, b, x) is not a triple-zero of N , $ab \notin (N : M)$ and $ax \notin M\text{-rad}(N)$. In a similar way, $ay \in M\text{-rad}(N)$. Now, we obtain $a(x+y) \in M\text{-rad}(N)$ or $b(x+y) \in M\text{-rad}(N)$ since $(a, b, x+y)$ is not a triple-zero of N , $ab(x+y) \in N$ and $ab \notin (N : M)$. Assume that $a(x+y) = ax + ay \in M\text{-rad}(N)$. As $ay \in M\text{-rad}(N)$, we get $ax \in M\text{-rad}(N)$, which is a contradiction. Assume that $b(x+y) = bx + by \in M\text{-rad}(N)$. As $bx \in M\text{-rad}(N)$, we get $by \in M\text{-rad}(N)$, a contradiction again. Hence we obtain that $aK \subseteq M\text{-rad}(N)$ or $bK \subseteq M\text{-rad}(N)$. □

Let N be a weakly 2-absorbing primary submodule of an R -module M and $I_1I_2K \subseteq N$ for some ideals I_1, I_2 of R and some submodule K of M where N is free triple-zero in regard to I_1, I_2, K . Note that if $a \in I_1, b \in I_2$ and $m \in K$, then $ab \in (N : M)$ or $am \in M\text{-rad}(N)$ or $bm \in M\text{-rad}(N)$.

The following result and its proof are analogues of [6, Theorem 2.30] and its proof.

Theorem 2. *Assume that N is a weakly 2-absorbing primary submodule of an R -module M and $0 \neq I_1I_2K \subseteq N$ for some ideals I_1, I_2 of R and some submodule K of M where N is free triple-zero in regard to I_1, I_2, K . Then $I_1I_2 \subseteq (N : M)$ or $I_1K \subseteq M\text{-rad}(N)$ or $I_2K \subseteq M\text{-rad}(N)$.*

Proof. Let N be a weakly 2-absorbing primary submodule of M and $0 \neq I_1I_2K \subseteq N$ for some ideals I_1, I_2 of R and some submodule K of M where

N is free triple-zero in regard to I_1, I_2, K . Suppose that $I_1I_2 \not\subseteq (N : M)$. Now, we show that $I_1K \subseteq M\text{-rad}(N)$ or $I_2K \subseteq M\text{-rad}(N)$. Assume that $I_1K \not\subseteq M\text{-rad}(N)$ and $I_2K \not\subseteq M\text{-rad}(N)$. Then $xK \not\subseteq M\text{-rad}(N)$ and $yK \not\subseteq M\text{-rad}(N)$ where $x \in I_1$ and $y \in I_2$. By Lemma 2, we get $xy \in (N : M)$ since $xyK \subseteq N$, $xK \not\subseteq M\text{-rad}(N)$ and $yK \not\subseteq M\text{-rad}(N)$. By our assumption, there are $a \in I_1$ and $b \in I_2$ such that $ab \notin (N : M)$. By Lemma 2, we get $aK \subseteq M\text{-rad}(N)$ or $bK \subseteq M\text{-rad}(N)$ as $abK \subseteq N$ and $ab \notin (N : M)$. We investigate three cases. **First case:** Assume that $aK \subseteq M\text{-rad}(N)$ and $bK \not\subseteq M\text{-rad}(N)$. Since $xbK \subseteq N$, $xK \not\subseteq M\text{-rad}(N)$ and $bK \not\subseteq M\text{-rad}(N)$, then we get $xb \in (N : M)$, by Lemma 2. We have $(x+a)K \not\subseteq M\text{-rad}(N)$ as $(x+a)bK \subseteq N$, $xK \not\subseteq M\text{-rad}(N)$ and $aK \subseteq M\text{-rad}(N)$. Since $bK \not\subseteq M\text{-rad}(N)$ and $(x+a)K \not\subseteq M\text{-rad}(N)$, then we obtain $b(x+a) \in (N : M)$, by Lemma 2. Thus since $b(x+a) = xb + ab \in (N : M)$ and $xb \in (N : M)$, then $ab \in (N : M)$, which is a contradiction. **Second case:** Assume that $aK \not\subseteq M\text{-rad}(N)$ and $bK \subseteq M\text{-rad}(N)$. It is easily shown similarly to the first case. **Third case:** Suppose that $aK \subseteq M\text{-rad}(N)$ and $bK \subseteq M\text{-rad}(N)$. Then $(y+b)K \not\subseteq M\text{-rad}(N)$ as $yK \not\subseteq M\text{-rad}(N)$ and $bK \subseteq M\text{-rad}(N)$. By Lemma 2, $x(y+b) \in (N : M)$ since $x(y+b)K \subseteq N$, $xK \not\subseteq M\text{-rad}(N)$ and $(y+b)K \not\subseteq M\text{-rad}(N)$. Then $xb \in (N : M)$ since $x(y+b) \in (N : M)$ and $xy \in (N : M)$. As $aK \subseteq M\text{-rad}(N)$ and $xK \not\subseteq M\text{-rad}(N)$, then $(x+a)K \not\subseteq M\text{-rad}(N)$. As $(x+a)yK \subseteq N$, $yK \not\subseteq M\text{-rad}(N)$ and $(x+a)K \not\subseteq M\text{-rad}(N)$, then $(x+a)y = xy + ay \in (N : M)$ by Lemma 2. As $xy \in (N : M)$ and $ay + xy \in (N : M)$, then $ay \in (N : M)$. By Lemma 2, we get $(x+a)(y+b) = xy + xb + ay + ab \in (N : M)$ since $(x+a)(y+b)K \subseteq N$, $(x+a)K \not\subseteq M\text{-rad}(N)$ and $(y+b)K \not\subseteq M\text{-rad}(N)$. As $xb, ay, xy \in (N : M)$, then $xy + xb + ay \in (N : M)$. Thus, $ab \in (N : M)$ since $xy + xb + ay + ab \in (N : M)$ and $xy + xb + ay \in (N : M)$, a contradiction. Hence $I_1K \subseteq M\text{-rad}(N)$ or $I_2K \subseteq M\text{-rad}(N)$. \square

Lemma 3. *Let N be a weakly 2-absorbing primary submodule of M . If $abK \subseteq N$ and $0 \neq 2abK$ for some submodule K of M and for some $a, b \in R$, then $ab \in (N : M)$ or $aK \subseteq M\text{-rad}(N)$ or $bK \subseteq M\text{-rad}(N)$.*

Proof. Assume that $ab \notin (N : M)$. Now, we show that $K \subseteq (M\text{-rad}(N) : a) \cup (M\text{-rad}(N) : b)$. Let $x \in K$. If $0 \neq abx$, then $ax \in M\text{-rad}(N)$ or $bx \in M\text{-rad}(N)$ since $ab \notin (N : M)$. Thus $x \in (M\text{-rad}(N) : a) \cup (M\text{-rad}(N) : b)$. Suppose that $abx = 0$. As $0 \neq 2abK$, there is $y \in K$ such that $2aby \neq 0$. Then we get $0 \neq aby \in N$. So $ay \in M\text{-rad}(N)$ or $by \in M\text{-rad}(N)$ since N is weakly 2-absorbing primary. Let $z = x + y$. Then $0 \neq abz \in N$. Since $ab \notin (N : M)$, then $az \in M\text{-rad}(N)$ or $bz \in M\text{-rad}(N)$. Now, we consider three cases. **First case:** Assume that $ay \in M\text{-rad}(N)$ and $by \notin M\text{-rad}(N)$. Let $ax \notin M\text{-rad}(N)$. Then $az \notin M\text{-rad}(N)$. Thus $bz \in M\text{-rad}(N)$. So $a(z+x) \notin M\text{-rad}(N)$ and $b(z+x) \notin M\text{-rad}(N)$. Hence $0 = ab(z+x) = 2abz$ since N is a weakly 2-absorbing primary submodule and $ab \notin (N : M)$. This is a contradiction. Consequently, $ax \in M\text{-rad}(N)$. **Second case:** Assume that $ay \notin M\text{-rad}(N)$ and $by \in M\text{-rad}(N)$. It is proved by a similar way to first case. **Third case:**

Assume that $ay \in M\text{-rad}(N)$ and $by \in M\text{-rad}(N)$. Since $az \in M\text{-rad}(N)$ or $bz \in M\text{-rad}(N)$, $ax \in M\text{-rad}(N)$ or $bx \in M\text{-rad}(N)$. \square

Lemma 4. *Let N be a weakly 2-absorbing primary submodule of an R -module M . If $aJK \subseteq N$ and $0 \neq 4aJK$ for some submodule K of M , for any ideal J of R and for some $a \in R$, then $aJ \subseteq (N : M)$ or $aK \subseteq M\text{-rad}(N)$ or $JK \subseteq M\text{-rad}(N)$.*

Proof. Suppose that $aJ \not\subseteq (N : M)$. Then there is $j \in J$ such that $aj \notin (N : M)$. Our claim is that there is $b \in J$ such that $0 \neq 4abK$ and $ab \notin (N : M)$. As $0 \neq 4aJK$, then there is $j' \in J$ such that $0 \neq 4aj'K$. Assume that $aj' \notin (N : M)$ and $0 \neq 4aj'K$. Then we have the result for $b = j'$ or $b = j$. Now, let $aj' \in (N : M)$ and $4aj'K = 0$. Then $0 \neq 4a(j + j')K \subseteq N$ and $a(j + j') \notin (N : M)$ since $aj \notin (N : M)$ and $aj' \in (N : M)$. Hence we get $b = j + j' \in J$ such that $0 \neq 4abK \subseteq N$ and $ab \notin (N : M)$. Thus $0 \neq 2abK$ and so $K \subseteq (M\text{-rad}(N) : a) \cup (M\text{-rad}(N) : b)$ by Lemma 3. If $K \subseteq (M\text{-rad}(N) : a)$, the proof is completed. Suppose that $K \not\subseteq (M\text{-rad}(N) : a)$. Then $K \subseteq (M\text{-rad}(N) : b)$, that is, $bK \subseteq M\text{-rad}(N)$. Let $c \in J$. Assume that $0 \neq 2acK$. By Lemma 3, $ac \in (N : M)$ or $cK \subseteq M\text{-rad}(N)$ since $K \not\subseteq (M\text{-rad}(N) : a)$. Then $c \in ((N : M) : a) \cup (M\text{-rad}(N) : K)$. Now, let $2acK = 0$. Then $0 \neq 2a(b+c)K \subseteq N$. By Lemma 3, we have $a(b+c) \in (N : M)$ or $(b+c)K \subseteq M\text{-rad}(N)$ since $K \not\subseteq (M\text{-rad}(N) : a)$. Thus we get $b+c \in ((N : M) : a) \cup (M\text{-rad}(N) : K)$. If $b+c \in (M\text{-rad}(N) : K)$, then $c \in (M\text{-rad}(N) : K)$. Hence $JK \subseteq M\text{-rad}(N)$. Let $b+c \in ((N : M) : a) \setminus (M\text{-rad}(N) : K)$. Note that $2a(b+c+b)K = 4abK \neq 0$ and $2a(b+c+b)K \subseteq N$. Since $ab \notin (N : M)$ and $a(b+c) \in (N : M)$, $a(b+c+b) \notin (N : M)$. Thus by Lemma 3, $K \subseteq (M\text{-rad}(N) : a) \cup (M\text{-rad}(N) : (b+c+b))$. As $b+c \notin (M\text{-rad}(N) : K)$ and $b \in (M\text{-rad}(N) : K)$, then we get $(b+c+b) \notin (M\text{-rad}(N) : K)$ and so $K \subseteq (M\text{-rad}(N) : a)$, a contradiction. Thus $b+c \in (M\text{-rad}(N) : K)$. Since $b \in (M\text{-rad}(N) : K)$, $c \in (M\text{-rad}(N) : K)$. Hence $J \subseteq ((N : M) : a) \cup (M\text{-rad}(N) : K)$. Consequently $J \subseteq (M\text{-rad}(N) : K)$ since $aJ \not\subseteq (N : M)$. \square

Theorem 3. *Let I_1, I_2 be ideals of R and N, K be submodules of an R -module M . Assume that N is a weakly 2-absorbing primary submodule. If $0 \neq I_1I_2K$ and $0 \neq 8(I_1I_2 + (I_1 + I_2)(N : K))(K + N)$, then $I_1I_2 \subseteq (N : M)$ or $I_1K \subseteq M\text{-rad}(N)$ or $I_2K \subseteq M\text{-rad}(N)$.*

Proof. It is clear from Lemma 4 and [12, Theorem 2.3]. \square

Theorem 4. *Let M be a finitely generated multiplication R -module whose every proper submodule is weakly 2-absorbing primary. Then there exist at most three maximal ideals of R containing $\text{Ann}(M)$.*

Proof. Assume that R has four maximal ideals $\mathfrak{S}_1, \mathfrak{S}_2, \mathfrak{S}_3$ and \mathfrak{S}_4 containing $\text{Ann}(M)$. Let $K = \mathfrak{S}_1 \cap \mathfrak{S}_2 \cap \mathfrak{S}_3$ and $N = KM$. Note that $\mathfrak{S}_iM \neq M$ for every index i . Indeed, if $\mathfrak{S}_iM = M$ for some index i , then there is an element $r \in \mathfrak{S}_i$ such that $r - 1 \in \text{Ann}(M) \subseteq \mathfrak{S}_i$, contradiction. It is easily seen that

$\mathfrak{S}_i = (\mathfrak{S}_i M : M)$ as $\mathfrak{S}_i \subseteq (\mathfrak{S}_i M : M)$. We get $K \subseteq (N : M) \subseteq \bigcap_{i=1}^3 (\mathfrak{S}_i M : M) = K$. Thus $\sqrt{(N : M)} = \sqrt{K} = \sqrt{\mathfrak{S}_1} \cap \sqrt{\mathfrak{S}_2} \cap \sqrt{\mathfrak{S}_3}$. Hence we get that $(N : M)$ is not a 2-absorbing primary ideal by [13, Corollary 2.7]. Then N is not a 2-absorbing primary submodule by [14, Theorem 2.6]. By Lemma 1, $(N : M)^3 \subseteq \text{Ann}(M) \subseteq \mathfrak{S}_4$. Then $\mathfrak{S}_i = \mathfrak{S}_4$ for $i \in \{1, 2, 3\}$, a contradiction. Therefore, there are at most three maximal ideals of R containing $\text{Ann}(M)$. \square

Corollary 1. *Let M be a finitely generated multiplication R -module whose every proper submodule is weakly 2-absorbing primary. Then $(J(R))^3 M = 0$.*

Proof. We have that $K^3 \subseteq \text{Ann}(M)$ by the proof of Theorem 4. Thus we obtain $(J(R))^3 M = 0$. \square

Let M be a multiplication R -module and K, L be R -submodules of M . Then there exist ideals I and J of R such that $K = IM$ and $L = JM$. Hence $KL = IJM = IL$ and $KL = IJM = KJ$. In particular, we obtain that $KM = IM = K$, $LM = JM = L$ and $Km = KRm$ for every $m \in M$. Therefore, $Km = IRm = Im$.

Lemma 5. *Let M be a multiplication R -module. If $(N : M)$ is a weakly primary ideal of R , then N is a weakly primary submodule of M .*

Proof. Assume that $0 \neq am \in N$ for some $a \in R$ and $m \in M$ with $a \notin \sqrt{(N : M)}$. Since M is multiplication, there exists an ideal I of R such that $m = Rm = IM$, then $0 \neq RaIM \subseteq N$. Since $(N : M)$ is weakly primary and $a \notin \sqrt{(N : M)}$, we have $((N : M) : Ra) = (N : M)$ or $((N : M) : Ra) = (0 : Ra)$, by [1, Proposition 2.1]. We claim that $Ra \subseteq (N : M)$ or $IM \subseteq N$. Suppose that $Ra \not\subseteq (N : M)$. Then since $0 \neq RaIM$, we get that $I \subseteq (N : RaM) = ((N : M) : Ra) = (N : M)$. Hence $IM \subseteq N$ and so $m \in N$, as needed. \square

Corollary 2. *Let M be a finitely generated multiplication R -module. Suppose that every proper ideal of R is weakly primary such that not primary ideal. Then $(J(R))^3 M = 0$.*

Proof. Assume that $(N : M)$ is a proper weakly primary ideal of R . Thus $(N : M)M = N$ is a weakly 2-absorbing primary submodule. By Corollary 1, we get $(J(R))^3 M = 0$. \square

The following result is an analogue of [6, Theorem 2.21].

Lemma 6. *Let $R = R_1 \times R_2$ and $M = M_1 \times M_2$ where $0 \neq M_1$ is a multiplication R_1 -module and $0 \neq M_2$ is a multiplication R_2 -module. Then the followings hold:*

- (1) *The following statements are equivalent:*
 - i. $N_1 \times M_2$ is a weakly 2-absorbing primary submodule of $M_1 \times M_2$;
 - ii. $N_1 \times M_2$ is a 2-absorbing primary submodule of $M_1 \times M_2$;

- iii. N_1 is a 2-absorbing primary submodule of M_1 .
- (2) The following statements are equivalent:
- i. $M_1 \times N_2$ is a weakly 2-absorbing primary submodule of $M_1 \times M_2$;
 - ii. $M_1 \times N_2$ is a 2-absorbing primary submodule of $M_1 \times M_2$;
 - iii. N_2 is a 2-absorbing primary submodule of M_2 .

Proof. (1) (i) \Rightarrow (ii): Assume that $N_1 \times M_2$ is not a 2-absorbing primary submodule of $M_1 \times M_2$. By Lemma 1, $(0, 0) = ((N_1 \times M_2) : (M_1 \times M_2))^2(N_1 \times M_2) = ((N_1 : M_1) \times (M_1 : M_2))^2(N_1 \times M_2) = (N_1 : M_1)^2 N_1 \times M_2$. Then $M_2 = 0$, a contradiction.

(ii) \Rightarrow (i): Is obvious.

(ii) \Rightarrow (iii): It is clear.

(2) The other part of the lemma can be seen in a similar way to the proof of the first part. \square

Lemma 7. Let $R = R_1 \times R_2$ and $M = M_1 \times M_2$ where $0 \neq M_1$ is a multiplication R_1 -module and $0 \neq M_2$ is a multiplication R_2 -module. If $N_1 \times N_2$ is a weakly 2-absorbing primary submodule of M , $N_1 \neq M_1$, $N_2 \neq 0$ and $N_2 \neq M_2$, then N_1 is a weakly 2-absorbing primary submodule of M_1 .

Proof. Assume that $N_1 \neq M_1$, $N_2 \neq 0$ and $N_2 \neq M_2$. Let $a_1, b_1 \in R_1$ and $x_1 \in M_1$ where $a_1 b_1 x_1 \in N_1$ and let $0 \neq x_2 \in N_2$. Then $(0, 0) \neq (a_1, 1)(b_1, 1)(x_1, x_2) \in N_1 \times N_2$. Since $N_1 \times N_2$ is a weakly 2-absorbing primary submodule, then $(a_1, 1)(b_1, 1) \in ((N_1 \times N_2) : M)$ or $(a_1, 1)(x_1, x_2) \in M\text{-rad}(N_1 \times N_2) = M_1\text{-rad}(N_1) \times M_2\text{-rad}(N_2)$ or $(b_1, 1)(x_1, x_2) \in M\text{-rad}(N_1 \times N_2) = M_1\text{-rad}(N_1) \times M_2\text{-rad}(N_2)$. As $1 \notin (N_2 : M_2)$, then we get $(a_1, 1)(x_1, x_2) \in M_1\text{-rad}(N_1) \times M_2\text{-rad}(N_2)$ or $(b_1, 1)(x_1, x_2) \in M_1\text{-rad}(N_1) \times M_2\text{-rad}(N_2)$. Thus $a_1 x_1 \in M_1\text{-rad}(N_1)$ or $b_1 x_1 \in M_1\text{-rad}(N_1)$. The proof is completed. \square

By [15], it is said that a commutative ring R is a u -ring if an ideal of R contained in a finite union of ideals must be contained in one of those ideals; and a um -ring is a ring R with the property that an R -module which is equal to a finite union of submodules must be equal to one of them.

Theorem 5. Let N be a weakly 2-absorbing primary submodule of M . Then the following statements hold:

- (1) If $ab \notin (N : M)$ for some $a, b \in R$, then $(N : ab) \subseteq (M\text{-rad}(N) : a) \cup (M\text{-rad}(N) : b) \cup (0 : ab)$.
- (2) Let R be a um -ring. If $ab \notin (N : M)$ for some $a, b \in R$, then $(N : ab) \subseteq (M\text{-rad}(N) : a)$ or $(N : ab) \subseteq (M\text{-rad}(N) : b)$ or $(N : ab) = (0 : ab)$.

Proof. Assume that N is a weakly 2-absorbing primary submodule of M .

(1) Let $m \in (N : ab)$. Assume that $abm \neq 0$. Then $abm \in N$ and thus $am \in M\text{-rad}(N)$ or $bm \in M\text{-rad}(N)$ since $ab \notin (N : M)$. Hence $m \in (M\text{-rad}(N) : a)$ or $m \in (M\text{-rad}(N) : b)$. Therefore, $m \in (M\text{-rad}(N) : a) \cup (M\text{-rad}(N) : b)$. Assume that $abm = 0$. Then $m \in (0 : ab)$. Consequently, $m \in (M\text{-rad}(N) : a) \cup (M\text{-rad}(N) : b) \cup (0 : ab)$.

(2) Assume that R is a um -ring. It is easily obtained by part (1). \square

Theorem 6. *Let N be a weakly 2-absorbing primary submodule of M . Then the following statements hold:*

- (1) *If $abm \notin N$ for some $a, b \in R$ and $m \in M$, then $(N : abm) \subseteq (M\text{-rad}(N) : am) \cup (M\text{-rad}(N) : bm) \cup (0 : abm)$.*
- (2) *Let R be a u -ring. If $abm \notin N$ for some $a, b \in R$ and $m \in M$, then $(N : abm) \subseteq (M\text{-rad}(N) : am)$ or $(N : abm) \subseteq (M\text{-rad}(N) : bm)$ or $(N : abm) = (0 : abm)$.*

Proof. Assume that N is a weakly 2-absorbing primary submodule of M .

(1) Let $r \in (N : abm)$. Assume that $rabm \neq 0$. Then $rabm \in N$ and thus $ram \in M\text{-rad}(N)$ or $rbm \in M\text{-rad}(N)$ since $ab \notin (N : M)$. Hence $r \in (M\text{-rad}(N) : am)$ or $r \in (M\text{-rad}(N) : bm)$. Therefore, $r \in (M\text{-rad}(N) : am) \cup (M\text{-rad}(N) : bm)$. Assume that $rabm = 0$. Then $r \in (0 : abm)$. Consequently, $r \in (M\text{-rad}(N) : am) \cup (M\text{-rad}(N) : bm) \cup (0 : abm)$.

(2) Assume that R is a u -ring. The result is easily seen by part (1). \square

Proposition 6. *Let M be a faithful multiplication R -module and N be a submodule of M . If N is a weakly 2-absorbing primary submodule but it is not 2-absorbing primary, then $N \subseteq M\text{-rad}(0)$. In addition $M\text{-rad}(N) = M\text{-rad}(0)$.*

Proof. By Lemma 1, $(N : M)^3 \subseteq \text{Ann}(M)$. Since M is a faithful module, $(N : M)^3 = 0$. Now suppose that $a \in (N : M)$. Then $a^3 = 0$ and so $a \in \sqrt{0}$. Hence $(N : M) \subseteq \sqrt{0}$ and thus $N = (N : M)M \subseteq M\text{-rad}(0)$. In addition, by Lemma 1, $(N : M)^2N = 0$. Then $(N : M)^3 = (N : M)^2(N : M) \subseteq ((N : M)^2N : M) = \text{Ann}(M)$ and so $(N : M) \subseteq \sqrt{\text{Ann}(M)}$. Thus $\sqrt{(N : M)} = \sqrt{\text{Ann}(M)}$. Hence $M\text{-rad}(N) = \sqrt{(N : M)M} = \sqrt{\text{Ann}(M)M} = M\text{-rad}(0)$. \square

Theorem 7. *Let M be an R -module and N be a submodule of M . Suppose that $M\text{-rad}(0)$ is a prime submodule. Then N is a weakly 2-absorbing primary submodule if and only if N is a 2-absorbing primary submodule.*

Proof. Assume that N is weakly 2-absorbing primary. Suppose that $abm \in N$ for some $a, b \in R$ and $m \in M$. If $0 \neq abm \in N$, then either $ab \in (N : M)$ or $am \in M\text{-rad}(N)$ or $bm \in M\text{-rad}(N)$. If $abm = 0$, we may suppose that $ab \notin (N : M)$. Since that $M\text{-rad}(0)$ is a prime submodule, we can conclude that $a \in (M\text{-rad}(0) :_R M)$ or $bm \in M\text{-rad}(0)$. Since $M\text{-rad}(0) \subseteq M\text{-rad}(N)$, we obtain that $am \in M\text{-rad}(N)$ or $bm \in M\text{-rad}(N)$. Therefore N is 2-absorbing primary. The converse is obviously. \square

Theorem 8. *Let M be a multiplication R -module and N be a submodule of M . Suppose that M is a P -primary module. If N is a weakly 2-absorbing primary submodule, then $(N : M)$ is a weakly 2-absorbing primary ideal.*

Proof. Assume that $0 \neq abc \in (N : M)$ for some $a, b, c \in R$ with $ab \notin (N : M)$. Then $abcm \in N$ for each $m \in M$. If $abcm = 0$, then $abc \in (0 :_R M) = P$.

Since M is a P -primary module, we conclude that $c^n \in P \subseteq (N : M)$ for some positive integer n . Thus $ac \in \sqrt{(N : M)}$, as needed. Suppose that $0 \neq abcm \in N$. Since N is weakly 2-absorbing primary, we can conclude that $acm \in M\text{-rad}(N)$ (so $ac \in ((M\text{-rad}(N) : M))$) or $bcm \in M\text{-rad}(N)$ (so $bc \in ((M\text{-rad}(N) : M))$). Hence $ac \in \sqrt{(N : M)}$ or $bc \in \sqrt{(N : M)}$. Therefore $(N : M)$ is a weakly 2-absorbing primary ideal. \square

Theorem 9. *Let M be a multiplication R -module and N be a submodule of M . Suppose that M is a P -primary module. If N is a weakly 2-absorbing primary submodule, then $\sqrt{(N : M)}$ is a weakly 2-absorbing ideal.*

Proof. Assume that $0 \neq abc \in \sqrt{(N : M)}$ with $ab \notin \sqrt{(N : M)}$ for some $a, b, c \in R$. Then $(abc)^n m \in N$ for some positive integer n and every $m \in M$. If $(abc)^n m = 0$ since M is a P -primary module, then $abc \in (0 :_R m) = (0 :_R M) = P \subseteq (N : M)$. Hence $ac \in \sqrt{(N : M)}$, as needed. Suppose that $0 \neq (abc)^n m = a^n b^n (c^n m) \in N$. Since N is a weakly 2-absorbing primary submodule and $ab \notin \sqrt{(N : M)}$, we conclude that $a^n (c^n m) \in M\text{-rad}(N)$ (so $a^n c^n \in (M\text{-rad}(N) : M)$) or $b^n (c^n m) \in M\text{-rad}(N)$ (so $b^n c^n \in (M\text{-rad}(N) : M)$). Since $(M\text{-rad}(N) : M) = \sqrt{(N : M)}$, we can conclude that $ac \in \sqrt{(N : M)}$ or $bc \in \sqrt{(N : M)}$. Therefore $\sqrt{(N : M)}$ is a weakly 2-absorbing ideal. \square

Theorem 10. *Let M be an R -module and N be a submodule of M . If N is a weakly 2-absorbing submodule, then $(N : m)$ is a weakly 2-absorbing ideal for every $m \in M \setminus N$ with $\text{Ann}(m) = 0$.*

Proof. Assume that $0 \neq abc \in (N : m)$ for some $a, b, c \in R$. Then $abcm \in N$. If $abcm = 0$, then $abc \in (0 :_R m) = 0$ and we are done. Suppose that $0 \neq abcm \in N$. Since N is a weakly 2-absorbing submodule, we conclude that $ab \in (N :_R M)$ (so $ab \in (N : m)$) or $bcm \in N$ (so $bc \in (N : m)$) or $acm \in N$ (so $ac \in (N : m)$). Therefore $(N : m)$ is a weakly 2-absorbing ideal for every $m \in M \setminus N$. \square

We invite the reader to compare the following two results with [6, Theorem 2.23].

Theorem 11. *Let $R = R_1 \times R_2$ and $M = M_1 \times M_2$ be a finitely generated multiplication R -module where M_1 and M_2 are multiplication R_1 -module and R_2 -module, respectively. If $N = N_1 \times N_2$ is a weakly 2-absorbing primary submodule, then $N = 0$ or N is 2-absorbing primary.*

Proof. Assume that $0 \neq N = N_1 \times N_2$ and show that N is a 2-absorbing primary submodule. Without loss of generality we may suppose that $N_2 \neq 0$. Then there exists a nonzero element $y \in N_2$. Suppose that $a \in (N_1 : M_1)$ and $x \in M_1$. Then $0 \neq (a, 1)(1, 1)(x, y) \in N = N_1 \times N_2$. Since N is a weakly 2-absorbing primary submodule, we conclude that $(a, 1)(1, 1) \in (N_1 \times N_2 : M)$ or $(a, 1)(x, y) \in M_1\text{-rad}(N_1) \times M_2\text{-rad}(N_2)$ or $(1, 1)(x, y) \in M_1\text{-rad}(N_1) \times$

$M_2\text{-rad}(N_2)$. Then $1 \in (N_2 : M_2)$ or $ax \in M_1\text{-rad}(N_1)$ or $y \in M_2\text{-rad}(N_2)$. Suppose that $1 \in (N_2 : M_2)$. If $N_1 = 0$, then $0 \times M_2 \subseteq N$ and so we may assume that $N = N_1 \times M_2$. Let $ax \in M_1\text{-rad}(N_1)$. Here we will have two cases; if $x \in N_1$, then since $N_2 \neq 0$, we can assume that $0 \times M_2 \subseteq N$ and so $N_1 \times M_2 = N$. If $x \notin N_1$ ($x \in M_1 \setminus N_1$), then since $N_1 \neq M_1$ and $N_2 \neq 0$, we may suppose that $N = N_1 \times M_2$. Then we show that if $N = N_1 \times M_2$, then N_1 is 2-absorbing primary and if $N = M_1 \times N_2$, then N_2 is 2-absorbing primary. For beginning we assume that $N = N_1 \times M_2$ and show that N_1 is 2-absorbing primary. Let $rsm \in N_1$ for some $r, s \in R_1$ and $m \in M_1$. Then $(0, 0) \neq (r, 1)(s, 1)(m, y) \in N_1 \times M_2$. Thus either $(r, 1)(s, 1) \in (N_1 \times N_2 :_R M)$ or $(r, 1)(m, y) \in M_1\text{-rad}(N_1) \times M_2\text{-rad}(M_2)$ or $(s, 1)(m, y) \in M_1\text{-rad}(N_1) \times M_2\text{-rad}(M_2)$ and so either $rs \in N_1$ or $rm \in M_1\text{-rad}(N_1)$ or $sm \in M_1\text{-rad}(N_1)$. Therefore $N = N_1 \times N_2$ is a 2-absorbing primary submodule, by Lemma 6. By similar way we can show that N_2 is a 2-absorbing primary submodule and so $N = M_1 \times N_2$ is 2-absorbing primary, by Lemma 6. Now we show that N_1 and N_2 are primary submodules. Assume that $N_2 \neq M_2$. Let $t \in R_1$ and $n \in M_1$ such that $tn \in N_1$ with $0 \neq n' \in N_2$. Then $(0, 0) \neq (t, 1)(1, 1)(n, n') \in N_1 \times N_2$. Since $N_1 \times N_2$ is a weakly 2-absorbing primary submodule, we conclude that $(t, 1)(1, 1) \in (N_1 \times N_2 :_R M)$ or $(t, 1)(n, n') \in M_1\text{-rad}(N_1) \times M_2\text{-rad}(N_2)$ or $(1, 1)(n, n') \in M_1\text{-rad}(N_1) \times M_2\text{-rad}(N_2)$. But $1 \notin (N_2 : M_2)$. Then $tn \in M_1\text{-rad}(N_1)$ and so N_1 is primary. Similarly we can show that N_2 is a primary submodule. Therefore N is a 2-absorbing primary submodule, by [14, Theorem 2.28]. \square

Theorem 12. *Let M_1 and M_2 be multiplication modules. Suppose that $N_1 \neq M_1$ and $M_2 \neq 0$. The submodule $N_1 \times 0$ is a weakly 2-absorbing primary submodule if one of the following statements hold:*

- (1) N_1 is a weakly primary submodule of M_1 and 0 is a primary submodule of M_2 and $M_1\text{-rad}(N_1) \neq 0$;
- (2) N_1 is a weakly primary submodule of M_1 and 0 is a primary submodule of M_2 and $M_1\text{-rad}(N_1) = 0$;
- (3) $N_1 = 0$.

Proof. Assume that $(0, 0) \neq (a, b)(c, d)(x, y) \in N_1 \times 0$ with $(a, b)(c, d) \notin (N_1 \times 0 : M)$. Then $0 \neq acx \in N_1$ and $bdy = 0$. Since N_1 is weakly primary, we have $a \in \sqrt{(N_1 : M_1)}$ or $cx \in N_1$. Since 0 is a primary submodule, we get that $b \in \sqrt{(0 : M_2)}$ or $dy = 0$. Then $ax \in M_1\text{-rad}(N_1)$ or $cx \in M_1\text{-rad}(N_1)$ and $by \in M_2\text{-rad}(0)$ or $dy \in M_2\text{-rad}(0)$. Hence $(a, b)(x, y) \in M_1\text{-rad}(N_1) \times M_2\text{-rad}(0)$ or $(c, d)(x, y) \in M_1\text{-rad}(N_1) \times M_2\text{-rad}(0)$, as needed. Now suppose that (2) holds. Then since N_1 is weakly primary and $0 \neq acx \in N_1$, we get that $a \in \sqrt{(N_1 : M_1)}$ or $cx \in N_1$. If $a \in \sqrt{(N_1 : M_1)}$, then $ax \in M_1\text{-rad}(N_1) = 0$. Thus as $cx \in M_1\text{-rad}(N_1) = 0$, $acx = 0$ which is a contradiction. Then neither $(a, b)(x, y) \in M_1\text{-rad}(N_1) \times M_2\text{-rad}(0)$ nor

$(c, d)(x, y) \in M_1\text{-rad}(N_1) \times M_2\text{-rad}(0)$. Therefore $N_1 \times 0$ is weakly 2-absorbing primary. The latest statement is obviously. \square

Example 1. Suppose that $M = \mathbb{Z} \times \mathbb{Z}$ is an $R = \mathbb{Z} \times \mathbb{Z}$ -module and $N = p\mathbb{Z} \times \{0\}$ is a submodule of M where $p\mathbb{Z}$ is a weakly primary submodule and $\{0\}$ is weakly primary. Then $\mathbb{Z}\text{-rad}(p\mathbb{Z}) = p\mathbb{Z}$ and $(N : M) = 0$. Assume that $(0, 0) \neq (p, 1)(1, 0)(1, 1) \in \mathbb{Z} \times \{0\}$. Then neither $(p, 1)(1, 0) \in (N : M)$ nor $(p, 1)(1, 1) \in M\text{-rad}(N)$ nor $(1, 0)(1, 1) \in M\text{-rad}(N)$. Hence N is not weakly 2-absorbing primary. Notice that M is not a multiplication module.

Proposition 7. Let M be a multiplication R -module and N_1, \dots, N_n be weakly 2-absorbing primary submodules with $M\text{-rad}(N_i) = P$ where P is a prime submodule. Then $N = \bigcap_{i=1}^n N_i$ is weakly 2-absorbing primary.

Proof. Assume that $0 \neq abm \in N$ for some $a, b \in R$ and $m \in M$ with $ab \notin (N : M)$. By [5, Proposition 2.14], $M\text{-rad}(N) = \bigcap_{i=1}^n M\text{-rad}(N_i)$. Then $ab \notin (N_i : M)$ for some $1 \leq i \leq n$. Since N_i is a weakly 2-absorbing primary submodule, $am \in M\text{-rad}(N_i) = P$ or $bm \in M\text{-rad}(N_i) = P$. Hence $am \in M\text{-rad}(N)$ or $bm \in M\text{-rad}(N)$, as needed. \square

Definition 4. Let N be a weakly 2-absorbing primary submodule of an R -module M and $M\text{-rad}(N) = P$, by Proposition 7. We say that N is a P -weakly 2-absorbing primary submodule.

Let R be a ring with identity and M be an R -module. Then $R(M) = R(+M)$ with multiplication $(a, m)(b, n) = (ab, an + bm)$ and with addition $(a, m) + (b, n) = (a + b, m + n)$ is a commutative ring with identity and $0(+M)$ is a nilpotent ideal of index 2. The ring $R(+M)$ is said to be the *idealization* of M or *trivial extension* of R by M . We view R as a subring of $R(+M)$ via $r \rightarrow (r, 0)$. An ideal H is said to be *homogeneous* if $H = I(+N)$ for some ideal I of R and some submodule N of M ; whence $IM \subseteq N$, [9] and [11, Sec, 25]. Let R_1 and R_2 be commutative rings, M_1 and M_2 be R -modules. Then $(R_1 \times R_2)(+)(M_1 \times M_2) \approx (R_1(+M_1)) \times (R_2(+M_2))$, by [9, Theorem 4.4]. Now we use it to characterize one of the result of weakly 2-absorbing primary ideals by the idealization method.

Proposition 8. Let R_1 and R_2 be commutative rings, M_1 be R_1 -module and M_2 be R_2 -module, respectively. Suppose that $H = I(+M_1)$ is an ideal of $R_1(M_1) = R_1(+M_1)$ and $J = h(+M_2)$ is an ideal of $R_2(M_2) = R_2(+M_2)$. Then the following statement are equivalent:

- (1) $H \times R_2(M_2)$ is a 2-absorbing (resp., weakly 2-absorbing) primary ideal of $R_1(M_1) \times R_2(M_2)$;
 - i. $R_1(M_1) \times J$ is a 2-absorbing (resp., weakly 2-absorbing) primary ideal of $R_1(M_1) \times R_2(M_2)$;
- (2) H is a 2-absorbing (resp., weakly 2-absorbing) primary ideal of $R_1(M_1)$;
 - ii. J is a 2-absorbing (resp., weakly 2-absorbing) primary ideal of $R_2(M_2)$;

Proof. (1) \Rightarrow (2) Assume that $(a, x)(b, y)(c, z) \in H = I(+M_1)$ for some $a, b, c \in R$ and $x, y, z \in M_1$. Thus $abc \in I$. Since $H \times R_2(M_2) = (I(+M_1) \times (R_2(+M_2)) = (I_1 \times R_2)(+)(M_1 \times M_2)$ is a 2-absorbing primary ideal, $H = I(+M_1)$ is 2-absorbing primary and hence I is a 2-absorbing primary ideal of R_1 , by [6, Theorem 2.21]. Then $ab \in I$ or $bc \in \sqrt{I}$ or $ac \in \sqrt{I}$. Therefore $(a, x)(b, y) \in I(+M_1)$ or $(b, y)(c, z) \in \sqrt{I(+M_1)} = \sqrt{I}(+M_1)$ or $(a, x)(c, z) \in \sqrt{I(+M_1)} = \sqrt{I}(+M_1)$, as needed.

(2) \Rightarrow (1) Let H be 2-absorbing primary. Assume that $(u_1, v_1)(u_2, v_2)(u_3, v_3) \in H \times R_2(M_2)$ such that $u_1, u_2, u_3 \in H$ and $v_1, v_2, v_3 \in R_2(M_2)$. Then $u_1 u_2 u_3 \in H = I(+M_1)$ where $u_1 = (a, x)$, $u_2 = (b, y)$ and $u_3 = (c, z)$. Since H is a 2-absorbing primary ideal, I is 2-absorbing primary, by [6, Theorem 2.21]. Then $ab \in I$ or $bc \in \sqrt{I}$ or $ac \in \sqrt{I}$. Hence $(a, x)(b, y) = u_1 u_2 \in I(+M_1)$ or $(b, y)(c, z) = u_2 u_3 \in \sqrt{I(+M_1)} = \sqrt{I}(+M_1)$ or $(a, x)(c, z) = u_1 u_3 \in \sqrt{I(+M_1)} = \sqrt{I}(+M_1)$. Therefore $(u_1, v_1)(u_2, v_2) \in H \times R_2(M_2)$ or $(u_2, v_2)(u_3, v_3) \in \sqrt{H \times R_2(M_2)} = (\sqrt{I}(+M_1) \times (R_2(+M_2))$ or $(u_1, v_1)(u_3, v_3) \in \sqrt{H \times R_2(M_2)} = (\sqrt{I}(+M_1) \times (R_2(+M_2))$. Then $H \times R_2(M_2)$ is a 2-absorbing primary ideal of $R_1(M_1) \times R_2(M_2)$.

The proof of (i) if and only if (ii) is clear by similar arguments as previously shown, and hence we omit the proof. \square

References

- [1] S. E. Atani and F. Farzalipour, *On weakly primary ideal*, Georgian Math. J. **12** (2005), no. 3, 423–429.
- [2] A. Badawi, *On 2-absorbing ideals of commutative rings*, Bull. Aust. Math. Soc. **75** (2007), no. 3, 417–429.
- [3] A. Badawi and A. Y. Darani, *On weakly 2-absorbing ideals of commutative rings*, Houston J. Math. to appear.
- [4] A. Badawi, U. Tekir, E. Ugurlu, G. Ulucak, and E. Yetkin, *Generalizations of 2-absorbing primary ideals of commutative rings*, Turk. J. Math. **40** (2016), no. 3, 703–717.
- [5] A. Badawi, U. Tekir, and E. Yetkin, *On 2-absorbing primary ideals in commutative rings*, Bull. Korean Math. Soc. **51** (2014), no. 4, 1163–1173.
- [6] ———, *On weakly 2-absorbing primary ideals of commutative rings*, J. Korean Math. Soc. **52** (2015), no. 1, 97–111.
- [7] A. Yousefian Darani and F. Soheilnia, *2-absorbing and weakly 2-absorbing submodules*, Thai J. Math. **9** (2011), no. 3, 577–584.
- [8] D. D. Anderson and E. Smith, *Weakly prime ideals*, Houston J. Math. **29** (2003), no. 4, 831–840.
- [9] D. D. Anderson and M. Winders, *Idealization of a module*, J. Commut. Algebra **1** (2009), no. 1, 3–56.
- [10] D. F. Anderson and A. Badawi, *On n-absorbing ideals of commutative rings*, Comm. Algebra **39** (2011), no. 5, 1646–1672.
- [11] J. A. Huckaba, *Commutative Rings with Zero-Divisors*, Monographs and Textbooks in Pure and Applied Mathematics, 117, Marcel Dekker, Inc., Now York, 1988.
- [12] S. Moradi and A. Azizi, *Weakly 2-absorbing submodules of modules*, Turk. J. Math. **40** (2016), no. 2, 350–364.
- [13] H. Mostafanasab and A. Y. Darani, *On ϕ -n-absorbing primary ideals in commutative rings*, arXiv:1503.00108v1(2015).

- [14] H. Mostafanasab, E. Yetkin, U. Tekir, and A. Y. Darani, *On 2-absorbing primary submodules of modules over commutative rings*, An. Şt. Univ. Ovidius Constanţa **24** (2016), no. 1, 335–351.
- [15] P. Quattararo and H. S. Butts, *Finite unions of ideals and modules*, Proc. Amer. Math. Soc. **52** (1975), 91–96.

AHMAD YOUSEFIAN DARANI
DEPARTMENT OF MATHEMATICS
FACULTY OF SCIENCE
UNIVERSITY OF MOHAGHEGH ARDABILI
P.O. BOX 179, ARDEBIL, IRAN
E-mail address: yousefian@uma.ac.ir, youseffian@gmail.com

FATEMEH SOHEILNIA
DEPARTMENT OF MATHEMATICS
FACULTY OF SCIENCE
UNIVERSITY OF MOHAGHEGH ARDABILI
P.O. BOX 179, ARDEBIL, IRAN
E-mail address: soheilnia@gmail.com

UNSAI TEKIR
DEPARTMENT OF MATHEMATICS
MARMARA UNIVERSITY
ZIVERBEY, 34722, GOZTEPE, ISTANBUL, TURKEY
E-mail address: utekir@marmara.edu.tr

GULSEN ULUCAK
DEPARTMENT OF MATHEMATICS
GEBZE TECHNICAL UNIVERSITY
P.K 41400, GEBZE-KOCAELI, TURKEY
E-mail address: gulsenulucak@gtu.edu.tr