

Genetic Diversity and Phylogenetic Analysis of the Iranian *Leishmania* Parasites Based on HSP70 Gene PCR-RFLP and Sequence Analysis

Sara Nemati¹, Asghar Fazaeli^{1,*}, Homa Hajjaran², Ali Khamesipour³, Mohsen Falahati Anbaran⁴,
Arezoo Bozorgomid², Fatah Zarei⁵

¹Department of Medical Parasitology and Mycology, School of Medicine, Zanjan University of Medical Sciences, Zanjan 4515613191, Iran;

²Department of Medical Parasitology and Mycology, School of Public Health, Tehran University of Medical Sciences, Tehran 14155-6446, Iran;

³Center for Research and Training in Skin Diseases and Leprosy, Tehran University of Medical Sciences, Tehran 1416613675, Iran; ⁴School of Biology, University of Tehran, Tehran 14155-6455, Iran; ⁵Department of Animal Sciences, Faculty of Life Sciences and Biotechnology, Shahid Beheshti University, Tehran 1983963113, Iran

Abstract: Despite the broad distribution of leishmaniasis among Iranians and animals across the country, little is known about the genetic characteristics of the causative agents. Applying both HSP70 PCR-RFLP and sequence analyses, this study aimed to evaluate the genetic diversity and phylogenetic relationships among *Leishmania* spp. isolated from Iranian endemic foci and available reference strains. A total of 36 *Leishmania* isolates from almost all districts across the country were genetically analyzed for the HSP70 gene using both PCR-RFLP and sequence analysis. The original HSP70 gene sequences were aligned along with homologous *Leishmania* sequences retrieved from NCBI, and subjected to the phylogenetic analysis. Basic parameters of genetic diversity were also estimated. The HSP70 PCR-RFLP presented 3 different electrophoretic patterns, with no further intraspecific variation, corresponding to 3 *Leishmania* species available in the country, *L. tropica*, *L. major*, and *L. infantum*. Phylogenetic analyses presented 5 major clades, corresponding to 5 species complexes. Iranian lineages, including *L. major*, *L. tropica*, and *L. infantum*, were distributed among 3 complexes *L. major*, *L. tropica*, and *L. donovani*. However, within the *L. major* and *L. donovani* species complexes, the HSP70 phylogeny was not able to distinguish clearly between the *L. major* and *L. turanica* isolates, and between the *L. infantum*, *L. donovani*, and *L. chagasi* isolates, respectively. Our results indicated that both HSP70 PCR-RFLP and sequence analyses are medically applicable tools for identification of *Leishmania* species in Iranian patients. However, the reduced genetic diversity of the target gene makes it inevitable that its phylogeny only resolves the major groups, namely, the species complexes.

Key words: *Leishmania tropica*, *Leishmania major*, *Leishmania infantum*, leishmaniasis, HSP70, phylogenetic analysis, PCR-RFLP, Iran

INTRODUCTION

Leishmaniasis, with diverse clinical manifestations, is a health problem in many countries, including Iran. There are 2 types of leishmaniasis in the country, cutaneous leishmaniasis (CL) and visceral leishmaniasis (VL). CL consists of 2 epidemiological forms, zoonotic CL caused by *Leishmani major* and anthroponotic CL caused by *Leishmania tropica* [1]. They are prevalent throughout the country, with tens of thousands of cases every year. In recent years, due to new settlements, popu-

lation increase, urbanization, migration, and increase in sandfly populations, they have emerged in some new districts [2]. VL is caused by *Leishmania infantum* and has been reported sporadically throughout Iran; however, it is more prevalent in northwestern and southern parts of the country with approximately 100 to 300 reported cases annually [3,4].

The exact identification of *Leishmania* species is of great importance with concern to diagnostic, disease prognosis, epidemiological surveillance, and clinical studies. Many diagnostic techniques, including microscopy, in vitro culture, biochemical, immunological, and molecular approaches have been applied for detection and identification of *Leishmania* parasites. Clinical features and microscopic examinations are not suited for species identification of parasites [5]. Multi-locus enzyme electrophoresis (MLEE) has been known as the reference method for identification of *Leishmania* spp. [6]. However, this

•Received 19 March 2017, revised 5 May 2017, accepted 23 May 2017.

*Corresponding author (fazaeli@zums.ac.ir; fazaeli2017@gmail.com)

© 2017, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

technique requires mass cultivation of parasites, and it lacks of power to differentiate the parasites below the species level [7,8]. On the other hand, PCR-based methods, including PCR-RFLP and DNA sequence analyses are highly informative [9]. RFLP technique, has allowed direct and rapid identification of *Leishmania* species [10-12]. For phylogenetic reconstruction, sequence analysis is preferred over RFLP or fingerprinting patterns [5]. During the past 2 decades, various DNA sequences have been used as targets to reconstruct the phylogeny within

the genus *Leishmania*, including genes encoding the catalytic polypeptide of DNA polymerase a (polA) [13], the largest subunit of RNA polymerase II (rpoIIIS) [13], 7SL RNA [14], ribosomal internal transcribed spacer (ITS) [15-17], the N-acetylglucosamine-1-phosphate transferase (NAGT) gene [18], the mitochondrial cytochrome *b* gene (*cytb*) [19], and recently, sequences of the heat-shock protein 70 gene (HSP70) subfamily [20].

The HSP70 is conserved across prokaryotes and eukaryotes,

Table 1. Details of the original and GenBank *Leishmania* isolates used in the present study

No. ^a	Sample ID	Species	Locality ^b	Host	Disease	Sample ID	Species	Locality ^c
1	LM4	<i>L. major</i>	Esfahan	Human	CL	JN628986.1	<i>L. infantum</i>	Georgia
2	LM5	<i>L. major</i>	Ardestan	Human	CL	JN676923	<i>L. infantum</i>	Georgia
3	LM6	<i>L. major</i>	Mashhad	Human	CL	JX021443	<i>L. turanica</i>	China
4	LM7	<i>L. major</i>	Sistan Baluchestan	Human	CL	JX021433	<i>L. infantum</i>	China
5	LM8	<i>L. major</i>	Shiraz	Human	CL	JX021432	<i>L. infantum</i>	China
6	LM10	<i>L. major</i>	Agha Aliabas	Human	CL	FN395025	<i>L. tropica</i>	India
1	LM12	<i>L. major</i>	Esfahana	Human	CL	FN395026	<i>L. tropica</i>	Kenya
5	LM13	<i>L. major</i>	Shiraz	Human	CL	FN395022	<i>L. major</i>	Sudan
5	LM14	<i>L. major</i>	Shiraz	Human	CL	FN395023	<i>L. major</i>	Israel
7	LM28	<i>L. major</i>	Dehloran	Human	CL	FN395024	<i>L. major</i>	Kenya
8	LM29	<i>L. major</i>	Ilam	Human	CL	XM001684512	<i>L. major</i>	Israel
8	LM30	<i>L. major</i>	Ilam	Human	CL	FN395031	<i>L. infantum</i>	Malta
9	LM31	<i>L. major</i>	Golestan	Human	CL	FN395032	<i>L. infantum</i>	Portugal
10	LM36	<i>L. major</i>	Bandar_Abbas	Human	CL	FN395033	<i>L. infantum</i>	Malta
11	LT27	<i>L. major</i>	Kermanshah	Human	CL	XM001470287	<i>L. infantum</i>	Spain
3	LT	<i>L. tropica</i>	Mashhad	Human	CL	GU071173	<i>L. braziliensis</i>	Brazil
12	LT2	<i>L. tropica</i>	Bam	Human	CL	KG905366	<i>L. donovani</i>	India
12	LT3	<i>L. tropica</i>	Bam	Human	CL	GU071178	<i>L. guyanensis</i>	Brazil
3	LT16	<i>L. tropica</i>	Mashhad	Human	CL	HF586354	<i>L. amazonensis</i>	Panama
17	LT9	<i>L. tropica</i>	Afghanistan	Human	CL	HF586353	<i>L. amazonensis</i>	Colombia
17	LT19	<i>L. tropica</i>	Afghanistan	Human	CL	JX312712	<i>L. donovani</i>	China
17	LT21	<i>L. tropica</i>	Afghanistan	Human	CL	FN669773	<i>L. donovani</i>	Ethiopia
9	LT22	<i>L. tropica</i>	Golestan	Human	CL	GU071174	<i>L. lainsoni</i>	Brazil
3	LT25	<i>L. tropica</i>	Mashhad	Human	CL	HF586413	<i>L. mexicana</i>	Ecuador
5	LT26	<i>L. tropica</i>	Shiraz	Human	CL	LN907842	<i>L. mexicana</i>	Ecuador
13	L.VT5	<i>L. tropica</i>	Ardebil	Human	CL	FN395056	<i>L. naiffi</i>	Brazil
14	L.VT6	<i>L. tropica</i>	Tehran	Human	CL	FN395055	<i>L. panamensis</i>	Panama
15	LI.1	<i>L. infantum</i>	Meshkin Shahr	Canine	VL	HF586367	<i>L. panamensis</i>	Panama
15	LI.2	<i>L. infantum</i>	Meshkin Shahr	Canine	VL	HF586368	<i>L. peruviana</i>	Peru
15	LI.3	<i>L. infantum</i>	Meshkin Shahr	Canine	VL	FN395044	<i>L. peruviana</i>	Peru
15	LI.7	<i>L. infantum</i>	Meshkin Shahr	Canine	VL	GU071175	<i>L. shawi</i>	Brazil
15	LI.8	<i>L. infantum</i>	Meshkin Shahr	Canine	VL	HF586356	<i>L. turanica</i>	China
16	L4	<i>L. major</i>	Hamedan	Rodent		JX021442	<i>L. turanica</i>	China
16	L5	<i>L. major</i>	Hamedan	Rodent		FN395037	<i>L. chagasi</i>	Brazil
16	L38	<i>L. major</i>	Hamedan	Rodent		FN395036	<i>L. chagasi</i>	Brazil
16	L40	<i>L. major</i>	Hamedan	Rodent		FN395035	<i>L. chagasi</i>	Brazil
	JN628988	<i>L. infantum</i>	Georgia ^c			HF586355	<i>L. gerbilli</i>	China
	JN628987	<i>L. infantum</i>	Georgia ^c			EF108422	<i>T. rangeli</i>	Honduras

^aLocality number on Fig. 1.

^bOriginal *Leishmania* strains from different districts of Iran.

^cStrains studied by other researchers, taken from NCBI.

and the protein as well as its encoding gene have been applied in phylogenetic studies of many parasites [5], including *Leishmania* [20,21]. Despite the broad distribution of leishmaniasis among Iranians, little is known about the genetic characteristics of the causative agents across the country [22-24]. Plus, most of the studies, targeted the HSP70 gene sequences to date, have covered only small geographical areas, with limited number of isolates [22,24]. Applying both HSP70 PCR-RFLP and DNA, the present study was designed to evaluate the genetic diversity and phylogenetic relationships among *Leishmania* spp. isolated from Iranian patients across the country and available reference strains.

MATERIALS AND METHODS

Parasite strains and DNA isolation

Samples positive for leishmaniasis were taken from 24 Iranian (from almost all districts of Iran) and 3 Afghan patients (who lived in Iran) referred to either Leishmaniasis Laboratory at the School of Public Health or Center for Research and Training in Skin Diseases and Leprosy, Tehran University of Medical Sciences. Four rodents and 5 canine isolates were also included in the sample (Table 1; Fig. 1). For DNA isolation, samples were transferred to RPMI-1640 culture medium (Gibco, Frankfurt, Germany) supplemented with 10-15% fetal bovine serum (Gibco), 100 U/ml penicillin, and 100 µg/ml streptomycin (Gibco)

and incubated at 24-26°C. *Leishmania* promastigotes were harvested from RPMI cultures and washed twice by PBS buffer and once by sterile distilled water. The pellets were kept frozen at -20°C until use. Genomic DNA was extracted from isolated parasites using the QIAamp DNA mini kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions.

PCR amplification of HSP70 gene

Amplification was performed with 2 specific primers, HSP70sen (5'-GACGGTGCCTGCCTACTTCAAG-3') and HSP70ant (5'-CCGCCCATGCTCTGTACATCC-3'), previously used by Garcia et al. [21]. They were synthesized by the Bioneer Corporation (Daejeon, South Korea). The PCR reaction mix (50 µl) contained 20 µl premix (Roche, Mannheim, Germany), 4 µl forward and reverse primers (10 pmol), 3 µl DNA (50-100 ng) template, and 20 µl double distilled water. The thermal cycling conditions consisted of initial denaturation at 94°C for 5 min, 30 cycles of 94°C for 5 min, 61°C for 1 min, and 72°C for 1 min 30 sec; and a final extension of 5 min at 72°C. Five µl of each PCR amplicon was run on a 1.2% agarose gel (Invitrogen, Carlsbad, California, USA), stained with ethidium bromide and visualized under UV light.

In silico and RFLP analysis of HSP70

The HSP70 gene PCR products (10 µl each) were digested with the HaeIII (BsuR1) restriction enzyme using the conditions




Fig. 1. Map of Iran, showing our sampling localities. Locality numbers are the same as in Table 1.

recommended by the manufacturer (MBI Fermentas, Vilnius, Lithuania). At first, this enzyme was tested against sequences of reference strains (Fasta format genes downloaded from GenBank) and searched for restriction sites using Restriction Mapper 3 (www.restrictionmapper.org). Theoretically, obtained fragments from digestion with HaeIII would be expected to create different patterns as follows: *L. major* (351, 307, 246, 152, 99, 47, 41, 40, 34, and 2 bp), *L. tropica* (354, 338, 246, 150, 99, 80, 41, 40, 21, 13, and 8 bp), and *L. infantum* (338, 307, 246, 152, 99, 80, 53, 47, 41, 40, and 13 bp). Digestion was performed in a total of 30 ml 1× optimal buffer, using 1U HaeIII restriction enzyme. Reactions were incubated at 37°C and completely analyzed by electrophoresis in a 3% small fragment agarose gel. The gels were subsequently ethidium bromide stained and subjected to electrophoresis along with the Gene Ruler™ 50 bp DNA Ladder (MBI Fermentas) as a reference DNA size marker.

DNA typing and sequence analysis

PCR products of the HSP70 gene related to 36 *Leishmania* isolates were purified and sequenced using an ABI Prism™ 3730 Genetic Analyzer (Applied Biosystems, Foster City, California, USA) by the Macrogen Company (Seoul, South Korea). Sequencing was carried out by the same primer pair. Consensus sequences from forward and reverse reads were trimmed and edited using the BioEdit 5.0.9 [25]. Furthermore, HSP70 sequences of 37 reference strains available in GenBank database were taken from NCBI (<http://www.ncbi.nlm.nih.gov>) and included in the analyses. All the original and retrieved sequences were aligned together, first applying the ClustalW algorithm implemented in the MEGA 6.0 [26], and then manually. Applying the corrected Akaike Information Criterion (AICc) [27] in jModelTest 2.1.10 [28], the best-fit models of nucleotide substitution was estimated. The maximum likelihood (ML) and neighbor joining (NJ) phylogenetic trees were constructed using the MEGA software. The HSP70 gene sequence of *Trypanosoma rangeli* (GenBank accession No. EF108422) was used as an outgroup. Using Arlequin 3.11 [29], basic parameters of genetic diversity were estimated. Using the same software, demographic histories were examined based on 2 neutrality tests, Tajima's D [30] and Fu's Fs [31].

RESULTS

RFLP analysis

PCR amplification of the HSP70 gene resulted in produc-




Fig. 2. Agarose gel (3%) showing 3 HSP70 PCR-RFLP profiles for *Leishmania* reference strains after digestion with HaeIII. Lane 1, *L. tropica*; lane 2, *L. tropica*; lane 3, *L. major*; lane 4, *L. major*; lane 5, *L. infantum*; lane 6, *L. infantum*; lane M, 100-bp size marker.

tion of about 1,420 bp fragment in all 36 *Leishmania* isolates. The amplicons of 20 isolates were subjected to PCR-RFLP analysis. Banding patterns were created with accordance to the expected profiles that were theoretically obtained from restriction map. The results of HSP70 PCR-RFLP presented 3 different electrophoretic patterns (Fig. 2), each identical to that of 1 of the 3 *Leishmania* species endemic in Iran, *L. tropica* (35%), *L. major* (50%), and *L. infantum* (15%). No intraspecific variation within the isolates of each species was observed.

Phylogenetic analyses

The best-fit models of nucleotide substitution, GTR+I+G, was applied to reconstruct phylogenetic relationships among all HSP70 sequences based on ML and NJ methods. The ML analysis (Fig. 3) grouped all the HSP70 sequences into 2 major clades, *L. (Leishmania)* clade comprising all Old World species and *L. mexicana* complex from the New World, and *L. (Viannia)* consisting of only New World species. The NJ tree (not shown) had similar subdivision. Phylogenetic analysis of the Iranian HSP70 gene sequences yielded a phylogenetic grouping in complete conformity with the primary PCR-RFLP identification. Iranian lineages, including *L. major*, *L. tropica*, and *L. infantum* were distributed among 3 complexes *L. major*, *L. tropica*, and *L. donovani*. However, within the *L. major* and *L. donovani* complexes, the HSP70 phylogeny was not able to distinguish clearly between the *L. major* and *L. turanica* and between the *L. infantum*, *L. donovani*, and *L. chagasi* isolates, respectively. Furthermore, no evidence for phylogeographic structure and sub-structure was evident within the phylogenetic tree.




Fig. 3. ML tree summarizing the relationship between HSP70 sequences of *Leishmania* species. Bootstrap resampling values are provided at each fork (given only if $\geq 40\%$). Iranian isolates are represented by circle (*L. major*), triangle (*L. tropica*), and square (*L. infantum*) symbols.

Table 2. Basic parameters of genetic diversity and neutrality tests, for 3 Iranian *Leishmania* species

Species	Diversity parameters							Neutrality tests	
	N ^a	S ^b	H ^c	Parsimony informative sites	Singleton variable sites	h ± SD ^d	π ^e	Tajima's D	Fu's Fs
<i>L. major</i>	20	10	3	1	9	0.195 ± 0.11	0.00093	-2.14 ^f	1.42
<i>L. tropica</i>	11	2	2	2	0	0.436 ± 0.13	0.00075	0.85	2.01
<i>L. infantum</i>	5	0	1	0	0	0.00 ± 0.00	0.0000	N.A. ^g	N.A.

^aNo. of sequences.
^bNo. of polymorphic sites.
^cNo. of haplotypes.
^dHaplotype diversity (h ± SD).
^eNucleotide diversity.
^fP < 0.05.
^gNot available.

Genetic diversity and demographic history

The sequence analysis of 1,167 bp fragment of the HSP70 gene among 36 *Leishmania* detected 12 polymorphic sites (9 singletons and 12 parsimony informative sites) among sequences (Table 2). Genetic diversity estimates of 3 *Leishmania* species are summarized in Table 2. In total, 6 haplotypes were identified, 3 in *L. major*, 2 in *L. tropica*, and 1 in *L. infantum*. Haplotype diversity (h) was ranging from 0.53 ± 0.136 for *L. tropica* to 0 for *L. infantum*. Nucleotide diversity was ranging from 0.436 ± 0.13 for *L. tropica* to 0 for *L. infantum*. Neutrality tests presented different results, including significant and negative Tajima's D and non-significant Fu's Fs values for *L. major*, and non-significant and positive Tajima's D and Fu's Fs values for *L. tropica*. Accordingly, these results provided support for rejecting the hypothesis of population growth in each species.

DISCUSSION

To date, most of the studies, targeting the HSP70 gene sequence in Iran, have covered only small geographical areas, with limited number of samples [22,24]. In the present study, using both PCR-RFLP and sequence analysis methods, we have extended previous observations by analyzing the HSP70 gene sequence in a larger cohort consisting of samples from almost all districts across the country.

Totally, 36 human and animal cases, including 27 CL, 5 VL, and 4 rodents were examined. Application of PCR-RFLP method on HSP70 gene clearly presented 3 different electrophoretic patterns, each belonged to 1 of the 3 *Leishmania* species in Iran. From 27 culture-positive samples prepared from CL cases, 15 (55.5%) were infected with *L. major* and 12 (44.5%) with *L. tropica*. All 5 canine specimens were infected with *L. infantum*, and all 4 rodents were infected with *L. major*. These results are in agreement with the results from the previous study on Iranians by Hajjaran et al. [23]. In their study, from 112 culture-positive samples prepared from CL cases, 75 (67%) were infected with *L. major* and 37 (33%) with *L. tropica*; from 25 rodents, 21 were infected with *L. major*; and from 28 culture-positive samples prepared from VL cases, 26 were infected with *L. infantum*.

Genus *Leishmania* is considerable for a large number of described species. They have been described as distinct species mainly according to biological, clinical, geographical, immunological, epidemiological, and biochemical measurements [5]. Though numerous molecular techniques and markers

have been proposed for resolving the taxonomy of *Leishmania* [13-20], describing a *Leishmania* species or embracing all of the described taxa is still not straightforward. The MLEE method have been considered as the reference technique by many authors [6]. Application of this technique resulted in the present *Leishmania* classification system, comprising 17 described species [8,32-34]. Based on this classification, the genus *Leishmania* consists of the 2 subgenera, *L. (Leishmania)* comprising all Old World species and *L. mexicana* complex from the New World, and *L. (Viannia)* consisting of only New World species [5]. Analysis of several DNA sequences as targets during the past 2 decades [13-20] have been consistent in that each of the 2 subgenera formed a distinct monophyletic clade and that a deep phylogenetic gap within *L. (Leishmania)* separated species of the Old and New World. Furthermore, in all of these studies, *L. (Viannia)* was closest to the root, while *L. (Leishmania)* formed the crown of the trees, an indication of New World origin of *Leishmania* [5,6].

Phylogenetic analysis of the genus *Leishmania* based on the HSP70 gene sequence by Fraga et al. [20] which also supported the above grouping, has included by far the largest number of species. Interestingly, only 8 monophyletic clades were deciphered in their phylogenetic trees and networks, 4 in *L. (Leishmania)* and 4 in *L. (Viannia)*. Accordingly, the authors concluded that the HSP70 phylogeny supports only 8 species, namely, *L. donovani*, *L. major*, *L. tropica*, *L. mexicana* within *L. (Leishmania)* and *L. lainsoni*, *L. guyanensis*, *L. naiffi* and *L. braziliensis* within *L. (Viannia)*, and that the concept of species complexes should be abandoned. Subsequently, inspired by these results, some authors voted for revising the taxonomy of the genus *Leishmania* [6,35].

In our study, application of PCR-RFLP method on HSP70 gene clearly presented 3 different electrophoretic patterns, each belonged to 1 of the 3 *Leishmania* species in Iran. However, PCR-RFLP method was not able to detect any intraspecific variations. The efficiency of this method for identification of *Leishmania* species has already been demonstrated by some authors [21,23,24,36]. On the other hand, analysis of HSP70 DNA sequences of *Leishmania* isolates was more informative. Despite the low levels of genetic diversity among Iranian isolates, sequence analysis was able to identify 6 haplotypes, 3 belonged to *L. major*, 2 belonged to *L. tropica*, and 1 haplotype belonged to *L. infantum*. Within *L. (Leishmania)*, Iranian lineages, including *L. major*, *L. tropica*, and *L. infantum*, were distributed among 3 species complexes *L. major*, *L. tropica*, and *L.*

donovani. However, within both *L. major* and *L. donovani* complexes, the HSP70 phylogeny could not show clear discrimination between previously described species. HSP70 is under stabilizing selective pressure for conservation of function since synonymous substitutions (6.3%) are favored over non-synonymous (1.9%) [20]. The reduced genetic diversity of the HSP70, makes it inevitable that its phylogeny only resolves the species complexes and prohibits valid deductions regarding the species and subspecies levels [35].

Overall, given that only a limited number of *Leishmania* species causes leishmaniasis among Iranians (namely, *L. major*, *L. tropica*, and *L. infantum*), our results indicated that both HSP70 PCR-RFLP and DNA sequence analysis are medically applicable tools for identification of *Leishmania* species in Iranian patients. However, in agreement with previous authors [6,20,35], our results support the idea of revising the taxonomy of the genus *Leishmania*. Otherwise, in the future assessments, species assignment in *Leishmania* should be according to highly variable DNA sequences with considerable congruency [35]. For resolving taxonomic ambiguities within the major groups, including *Leishmania*, highly discriminatory methods such as multi-locus sequence typing (MLST) and genome-wide single nucleotide polymorphisms (SNPs) are probably better suited.

ACKNOWLEDGMENTS

The authors would like to thank the staff of the School of Public Health and Center for Research and Training in Skin Diseases and Leprosy, Tehran University of Medical Sciences. This research was supported by a grant from Zanjan University of Medical Sciences, and was a part of the first author's doctoral dissertation.

CONFLICT OF INTEREST

We have no conflict of interest related to this study.

REFERENCES

1. Yaghoobi-Ershadi MR, Hanafi-Bojd AA, Javadian E, Jafari R, Zahraei-Ramazani AR, Mohebbali M. A new focus of cutaneous leishmaniasis caused by *Leishmania tropica*. Saudi Med J 2002; 23: 291-294.
2. Edrisian GhH, Nadim A, Alborzi AV, Aredehali S. Visceral leishmaniasis: the Iranian experience. Arch Iran Med 1998; 1: 22-26.
3. Mohebbali M, Edrisian GhH, Nadim A, Hajjaran H, Akhoundi B, Hooshmand B, Zarei Z, Arshi Sh, Mirsamadi N, Naeini KM, Mamishi S, Sanati AA, Moshfe AA, Charehadar S, Fakhar M. Application of direct agglutination test (DAT) for the diagnosis and seroepidemiological studies of visceral leishmaniasis in Iran. Iran J Parasitol 2006; 1: 15-25.
4. Nadim A, Navid-Hamidi A, Javadian E, Tahvildari Bidruni GH, Amini H. Present status of kala-azar in Iran. Am J Trop Med Hyg 1978; 27: 25-28.
5. Schönian G, Cupolillo E, Mauricio I. Molecular evolution and phylogeny of *Leishmania*. In Ponte-Sucré A, Diaz E, Padron-Nieves M eds, Drug Resistance in *Leishmania* Parasites. London, UK. Springer. 2013, pp. 15-44.
6. Schönian G, Mauricio I, Cupolillo E. Is it time to revise the nomenclature of *Leishmania*? Trends Parasitol 2010; 26: 466-469.
7. Cupolillo E, Grimaldi G, Momen H, Beverley SM. Intergenic region typing (IRT): a rapid molecular approach to the characterization and evolution of *Leishmania*. Mol Biochem Parasitol 1995; 73: 145-155.
8. Cupolillo E, Grimaldi Jr G, Momen H. A general classification of new world *Leishmania* using numerical zymotaxonomy. Am J Trop Med Hyg 1994; 50: 296-311.
9. Montalvo AM, Fraga J, Maes I, Dujardin JC, Van der Auwera G. Three new sensitive and specific heat-shock protein 70 PCRs for global *Leishmania* species identification. Eur J Clin Microbiol Infect Dis 2012; 31: 1453-1461.
10. Degraeve W, Fernandes O, Campbell D, Bozza M, Lopes U. Use of molecular probes and PCR for detection and typing of *Leishmania*: a mini-review. Mem Inst Oswaldo Cruz 1994; 89: 463-469.
11. Schönian G, Nasereddin A, Dinse N, Schweynoch C, Schallig HDFH, Presber W, Jaffe CL. PCR diagnosis and characterization of *Leishmania* in local and imported clinical samples. Diagn Microbiol Infect Dis 2003; 47: 349-358.
12. Wilson SM. DNA-based methods in the detection of *Leishmania* parasites: field applications and practicalities. Ann Trop Med Parasitol 1995; 89: 95-100.
13. Croan DG, Morrison DA, Ellis JT. Evolution of the genus *Leishmania* revealed by comparison of DNA and RNA polymerase gene sequences. Mol Biochem Parasitol 1997; 89: 149-159.
14. Zelazny AM, Fedorko DP, Li L, Neva FA, Fischer SH. Evaluation of 7SL RNA gene sequences for the identification of *Leishmania* spp. Am J Trop Med Hyg 2005; 72: 415-420.
15. Berzunza-Cruz M, Cabrera N, Crippa-Rossi M, Sosa Cabrera T, Pérez-Montfort R, Becker I. Polymorphism analysis of the internal transcribed spacer and small subunit of ribosomal RNA genes of *Leishmania mexicana*. Parasitol Res 2002; 88: 918-925.
16. Dávila AM, Momen H. Internal-transcribed-spacer (ITS) sequences used to explore phylogenetic relationships within *Leishmania*. Ann Trop Med Parasitol 2000; 94: 651-654.
17. Spanakos G, Piperaki ET, Menounos PG, Tegos N, Fliemetakis A, Vakalis NC. Detection and species identification of Old World *Leishmania* in clinical samples using a PCR-based method. Trans R Soc Trop Med Hyg 2008; 102: 46-53.

18. Waki K, Dutta S, Ray D, Kolli BK, Akman L, Kawazu SI, Lin CP, Chang KP. Transmembrane molecules for phylogenetic analyses of pathogenic protists: *Leishmania*-specific informative sites in hydrophilic loops of trans-endoplasmic reticulum *N*-acetylglucosamine-1-phosphate transferase. *Eukaryot Cell* 2007; 6: 198-210.
19. Asato Y, Oshiro M, Myint CK, Yamamoto Y, Kato H, Marco JD, Mimori T, Gomez EA, Hashiguchi Y, Uezato H. Phylogenetic analysis of the genus *Leishmania* by cytochrome *b* gene sequencing. *Exp Parasitol* 2009; 121: 352-361.
20. Fraga J, Montalvo AM, De Doncker S, Dujardin JC, Van der Auwera G. Phylogeny of *Leishmania* species based on the heat-shock protein 70 gene. *Infect Genet Evol* 2010; 10: 238-245.
21. Garcia L, Kindt A, Bermudez H, Llanos-Cuentas A, De Doncker S, Arevalo J, Wilber Quispe Tintaya K, Dujardin JC. Culture-independent species typing of neotropical *Leishmania* for clinical validation of a PCR-based assay targeting heat shock protein 70 genes. *J Clin Microbiol* 2004; 42: 2294-2297.
22. Fotouhi-Ardakani R, Dabiri S, Ajdari S, Alimohammadian MH, Alaei-Novin E, Taleshi N, Parvizi P. Assessment of nuclear and mitochondrial genes in precise identification and analysis of genetic polymorphisms for the evaluation of *Leishmania* parasites. *Infect Genet Evol* 2016; 46: 33-41.
23. Hajjaran H, Mohebbali M, Mamishi S, Vasigheh E, Oshaghi MA, Naddaf SR, Teimouri A, Edrissian GH, Zarei Z. Molecular identification and polymorphism determination of cutaneous and visceral leishmaniasis agents isolated from human and animal hosts in Iran. *Biomed Res Int* 2013; 789326.
24. Mirahmadi H, Salimi Khorashad A, Sohrabnabad A, Heydarian P, Bizhani N. Species identification and molecular typing of *Leishmania* spp. using targeting HSP70 gene in suspected patients of cutaneous leishmaniasis from Sistan and Baluchestan Province, Southeast Iran. *Iran J Parasitol* 2016; 11: 489-498.
25. Hall TA. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 1999; 41: 95-98.
26. Tamura K, Stecher G, Peterson D, Filipinski A, Kumar S. MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* 2013; 30: 2725-2729.
27. Hurvich CM, Tsai CL. Regression and time series model selection in small samples. *Biometrika* 1989; 76: 297-307.
28. Posada D. jModelTest: phylogenetic model averaging. *Mol Biol Evol* 2008; 25: 1253-1256.
29. Excoffier L, Lischer HE. Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Mol Ecol Resour* 2010; 10: 564-567.
30. Tajima F. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* 1989; 123: 585-595.
31. Fu YX. Statistical tests of neutrality of mutations against population growth, hitchhiking and background selection. *Genetics* 1997; 147: 915-925.
32. Cupolillo E, Medina-Acosta E, Noyes H, Momen H, Grimaldi G Jr. A revised classification for *Leishmania* and *Endotrypanum*. *Parasitol Today* 2000; 16: 142-143.
33. Lainson R. *Evolution, Classification and Geographical Distribution*. Cambridge, Massachusetts, USA. Academic Press. 1987.
34. Rioux JA, Lanotte G, Serres E, Pratlong F, Bastien P, Perieres J. *Taxonomy of Leishmania*. Use of isoenzymes. Suggestions for a new classification. *Ann Parasitol Hum Comp* 1990; 65: 111-125.
35. Van der Auwera G, Fraga J, Montalvo AM, Dujardin JC. *Leishmania* taxonomy up for promotion? *Trends Parasitol* 2011; 27: 49-50.
36. da Silva LA, de Sousa CdoS, da Graça GC, Porrozzi R, Cupolillo E. Sequence analysis and PCR-RFLP profiling of the hsp70 gene as a valuable tool for identifying *Leishmania* species associated with human leishmaniasis in Brazil. *Infect Genet Evol* 2010; 10: 77-83.