

Research Article



CrossMark

Open Access

파풍망 종류에 따른 키위의 발아, 개화 및 과실 특성

곽용범¹, 김홍림¹, 이목희¹, 이한철¹, 곽연식², 이용복^{2*}

¹

²

Budbreak, Floral Bud and Fruit Characteristics of Kiwifruit as Affected by Various Windbreaks

Yong-Bum Kwack¹, Hong Lim Kim¹, Mockhee Lee¹, Han-Cheol Rhee¹, Youn-Sig Kwak² and Yong Bok Lee^{2*}
(¹Namhae Branch, National Institute of Horticultural and Herbal Science, Rural Development Administration, Namhae 52430, Korea, ²Institute of Agriculture & Life Science, Gyeongsang National University, Jinju 52828, Korea)

Received: 23 September 2017/ Revised: 6 September 2017/ Accepted: 15 September 2017

Copyright © 2017 The Korean Society of Environmental Agriculture

This is an Open-Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Yong Bok Lee

<http://orcid.org/0000-0002-7651-4556>

Yong-Bum Kwack

<http://orcid.org/0000-0003-2506-7693>

Abstract

BACKGROUND: Kiwifruit growers build their vineyards using many windbreaks to protect their kiwifruit vines from defoliation injury by strong winds such as typhoon. In this study, we have compared fruit quality, budbreak rate and floral bud as affected by windbreaks. And also we surveyed several microclimate indices of kiwifruit orchard depending on the covering materials of arch-type windbreaks.

METHODS AND RESULTS: Five different windbreak materials including polyethylene film (PE), blue- and white-colored nets were tested in pipe-framed arch-type kiwifruit vineyards as the covering materials. Photosynthetically active radiation (PAR), annual mean temperature (AMT) and chill unit (CU) as well as fruit quality were compared among the covering materials. In all treatments, annual PAR was more than $400 \mu\text{mol m}^{-2} \text{s}^{-1}$, in which kiwifruit leaf could reach its maximum photosynthesis, since the leaves were emerged. Annual mean temperature was greater in 0.1 mm-PE covering as much as $1-2^\circ\text{C}$ than other windbreaks. In CU calculated by three different models, all windbreaks showed more

than 1400 CU that is fully fulfilled CU for kiwifruit rest completion. There were no difference in budbreak rate among the covering materials. Fruit weight was heavier in 0.1 mm-PE and white-net (4 mm) than other windbreaks.

CONCLUSION: Regardless of the windbreak materials, the PAR quantity was enough for kiwifruit photosynthesis. And CU for kiwifruit rest completion was fully achieved in all treatments. However, with respect to fruit weight, quantity of PAR, and AMT, etc., It is highly recommended for kiwifruit growers to choose 0.1 mm-PE and white-net (4 mm) as for their windbreaks materials.

Key words: Chill unit, Fruit quality, Kiwifruit, Microclimate, Windbreaks

서론

1,300 ha가 (, 2016).

1 3 (, 2011)

가

*Corresponding author: Yong Bok Lee

Phone: +82-55-772-1967; Fax: +82-55-772-1969;

E-mail: yblee@gnu.ac.kr

Table 1. Physical features of cover materials of arch-type windbreaks used in this experiment

Windbreaks	Average mesh size (mm)	Net porosity (%)
Blue net (2 mm)	1.36×2.39	49.3
Blue net (4 mm)	5.21×3.23	72.7
White net (2 mm)	1.73×2.29	65.6
White net (4 mm)	5.56×3.42	82.1

²Net porosity was calculated by mesh area and number per 100 cm².

가 (Tanhora, 1988; Kwack *et al.*, 2012, 2013, 2015), (Choi *et al.*, 2002) 가

과실희성 및 익년 발아, 개화 조사
2012 2013 20
가
2013 2014 가
(bud) (sprouted bud)
1 가
(PR-32 α, Atago, Bellevue, USA)
(TitroLine Easy, Schott, Germany)

재료 및 방법

시험 품종 및 방풍시설

31 (Hayward) 5 m, 4.3 m
1.7 m
0.1 mm (PE)
2010 12 (Open field)
△△ 2 mm (Blue 2 mm) 4 mm (Blue 4 mm) 2 mm (White 2 mm) 4 mm (White 4 mm)
(Table 1). (%) 100 cm² 5 m×6 m 6

통계분석

2 SAS 9.2, Enterprise 4.3 (SAS Institute, Inc., Cary, N.C., USA) (*P*≤0.05)

결 과

파풍망의 특성 및 과원 내 미세기상

가
가 PE mesh 가
(Table 1). 2 mm mesh 가 3.25, 3.96 mm², 16.82, 19.01 mm² 2 mm 4 mm 49.3% 2 mm 4 mm 72.7% 65.6% 82.1% mesh PAR Fig 1 PE가

미세기상 조사

400~700 nm (PAR) 가 WatchDog 2475 (Spectrum technology, USA) 30 cm 30 (chill unit, CU) Sunley (2006) 3가




Fig. 1. PAR and mean temperature of kiwifruit orchard during seven months from April through October in 2012 and 2013 depending on covering materials of arch-type windbreaks. Bars labeled with same letter are not significantly different by the Duncan's multiple range test ($p=0.05$). Vertical bars indicate standard errors of the means.

Table 2. Several microclimate indices under different covering materials of arch-type windbreaks during seven months from April through October

Year	Windbreaks	PAR (μMOL m ⁻² s ⁻¹)	Temperature (°C)
2012	Open field	638.0	23.1
	PE	497.1	24.3
	Blue net (2 mm)	488.4	23.0
	Blue net (4 mm)	498.4	23.6
	White net (2 mm)	530.3	23.4
	White net (4 mm)	613.9	23.0
2013	Open field	691.7	19.5
	PE	557.2	21.9
	Blue net (2 mm)	563.7	19.4
	Blue net (4 mm)	617.0	20.0
	White net (2 mm)	677.9	20.2
	White net (4 mm)	701.0	19.9

^zMean separation within columns by the Duncan's multiple range test at $p=0.05$.

(Fig. 1). PAR

mesh 가
PAR (Table 2). PAR 2 mm 4 mm 2012 488.4, 498.4 μmol m⁻² s⁻¹, 2013 563.7, 617.0 μmol m⁻² s⁻¹, 2 mm 4 mm 2012 530.3,




Fig. 2. Chilling unit of kiwifruit orchard calculated by three different models. It is sum of five months from November through March in the year 2012/2013 and 2013/2014 under different covering materials of arch-type windbreaks. Bars labeled with 'ns' are not significantly different by the Duncan's multiple range test ($p=0.05$). Vertical bars indicate standard errors of the means.

Table 3. Fruit characteristics of 31-year-old 'Hayward' kiwifruit grown under different covering materials of arch-type windbreaks

Year	Windbreaks	Fruit weight (g)	Soluble solids content (Brix)	Titratable acidity (%)
2012	PE	92.5	13.5	0.8
	Blue net (2 mm)	90.8	13.5	0.8
	Blue net (4 mm)	88.2	13.9	0.9
	White net (2 mm)	97.9	12.7	0.8
	White net (4 mm)	93.8	13.3	0.7
2013	Open field	96.5	12.8	1.1
	PE	118.2	14.3	1.1
	Blue net (2 mm)	109.6	13.5	1.2
	Blue net (4 mm)	99.1	12.8	1.1
	White net (2 mm)	101.7	13.5	1.1
	White net (4 mm)	120.2	13.5	1.0

^zMean separation within columns by the Duncan's multiple range test at $p=0.05$.

613.9 μmol m⁻² s⁻¹, 2013 677.9, 701.0 μmol m⁻² s⁻¹ mesh (PAR)

가 가 가
(chill unit, CU)
37가 (Sunley *et al.*, 2006)
mesh

(Fig. 2).

파풍망에 따른 과실특성 가

Table 4. Characteristics of budbreak and flowering of 31-year-old 'Hayward' kiwifruit grown under different covering materials of arch-type windbreaks

Year	Windbreaks	Budbreak (%)	Floral bud (%)	Flower (No/floral shoot)
2013	Open field	67.5 a ^z	40.7 c	2.2 c
	PE	66.0 a	84.0 a	3.6 a
	Blue net (2 mm)	65.4 a	61.9 b	3.0 b
	Blue net (4 mm)	65.8 a	65.3 b	3.2 a
	White net (2 mm)	66.9 a	71.4 b	3.2 a
	White net (4 mm)	68.3 a	71.8 b	3.3 a
2014	Open field	66.7 a	89.1 a	4.1 a
	PE	66.2 a	85.8 a	3.6 b
	Blue net (2 mm)	60.6 a	80.6 a	3.4 b
	Blue net (4 mm)	65.3 a	75.6 a	3.3 b
	White net (2 mm)	71.6 a	79.6 a	3.2 b
	White net (4 mm)	66.3 a	84.4 a	3.1 b

^zMean separation within columns by the Duncan's multiple range test at $p=0.05$.

(g FW), (Brix), (%)
 (Table 3).
 가, PAR 가
 4 mm 1~2°C
 0.1 mm PE 가 2013 가
 2012 4 mm 13.9 Brix, 2013 0.1 mm
 PE 14.3 Brix 가
 mm 4

파풍망에 따른 익년 발아 및 개화

. 2013 2014
 (bud) 가 (Table 4).
 가 가
 (floral bud) 2014
 가 . 2013
 40.7% 가 PE 84.0%
 가 . 2013 2014
 mesh
 2013

고 찰

(Norton, 1988),
 (Judd and McAneney, 1984; Jarvis, 1985)

(Briassoulis *et al.*, 2007). mesh
 (Gravina *et al.*,
 2011), 2 mm
 , 4 mm mesh
 9.4~16.3%
 가
 , 5 10~11 μmol
 $\text{CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ (Piller and Meekings, 1997),
 (PAR) 375 $\mu\text{mol m}^{-2} \text{ s}^{-1}$
 가 (Buwalda *et al.*, 1991).
 mesh 가 PAR가 가
 mesh PAR가
 PAR
 400 $\mu\text{mol m}^{-2} \text{ s}^{-1}$
 (Fig. 1; Table 2).
 PE 가
 1~2°C (Fig. 1; Table 2). Norton
 (1988)
 Williams (1970) 1°C
 가
 2013 PE (Table 3)
 4 mm PE 가
 PAR
 (Table 2)
 (Sunley *et al.*, 2006).
 (Weinberger, 1950, 1954; Oukabli *et al.*, 2003). 가
 (Sunley *et al.*, 2006).
 950~1000
 (4°C) (Brundell, 1976; Lionakis
 and Schwabe, 1984). Caldwell(1989)
 950
 가
 1150
 2012~2014
 11 3 5 1 3
 가 (Sunley *et al.*, 2006)
 (Fig. 2). PE

1400 CU 가
 (Fig. 2).
 가 (Table 4), 2013
 가 2012
 (Kwack *et al.*, 2013, 2014, 2015)
 PE
 1~2°C (Table 2)
 0.1 mm PE (PAR)
 (2 mm, 4 mm)
 가
 가 PE 가
 가 PAR, PE 4 mm
요 약
 5 4.3 m
 (PAR)
 4~10 400 $\mu\text{mol m}^{-2} \text{s}^{-1}$
 PAR PE
 1~2°C 가 (CU)
 1400 CU
 가 PE 4 mm
 가 PAR
 PE
 4 mm 가

References

Briassoulis, D., Mistriotis, A., & Eleftherakis, D. (2007). Mechanical behaviour and properties of agricultural nets-Part I: Testing methods for agricultural nets. *Polymer Testing*, 26(6), 822-832.

Brundell, D. J. (1976). The effect of chilling on the termination of rest and flower bud development of the chinese gooseberry. *Scientia Horticulturae*, 4(2), 175-182.

Buwalda, J. G., Meekings, J. S., & Smith G. S. (1991). Seasonal changes in photosynthetic capacity of leaves of kiwifruit (*Actinidia deliciosa*) vines. *Physiologia Plantarum*, 83(1), 93-98.

Caldwell, J. (1989). Kiwifruit performance in South Carolina and effect of winter chilling. *Clemson University, Clemson South Carolina*, 10, 127-129.

Choi, S. T., Kang, S. M., Park, D. S., Song, W. D., & Seo, G. K. (2002). Thinning effect on fruit characteristics and reserve accumulation of persimmon trees defoliated in early autumn. *Horticulture, Environment, and Biotechnology*, 43(6), 660-665.

Gravina, A., Cataldo, J., Gambetta, G., Pardo, E., & Fornero, C. (2011). Relation of peel damage in citrus fruit to wind climate in orchard and its control. *Scientia Horticulturae*, 129(1), 45-51.

Jarvis, P. P. G. (1985). Coupling of transpiration to the atmosphere in horticultural crops: the omega factor. In *I International Symposium on Water Relations in Fruit Crops*, 171, 187-206.

Judd, M. J., & McAneney, K. J. (1984). Water use by tamarillos (*Cyphomandra betacea*) within a sheltered orchard environment. *Agricultural and Forest Meteorology*, 32(1), 31-40.

Kwack, Y. B., Kim, H. L., Choi, Y. H., Lee, J. H., Kim, J. G., & Lee, Y. B. (2012). Fruit quality and fruit locule air hole of kiwifruit (*Actinidia deliciosa* cv. Hayward) affected by early defoliation, *Korean Journal of Environmental Agriculture*, 31(3), 229-234.

Kwack, Y. B., Kim, H. L., Chae, W. B., Lee, J. H., Lee, E. H., Kim, J. G., & Lee, Y. B. (2013). Regrowth of buds and flower bud formation in kiwifruit as affected by early defoliation, *Korean Journal of Environmental Agriculture*, 32(3), 201-206.

Kwack, Y. B., Kim, H. L., Chae, W. B., Kim, S. H., Lee, Y. B., & Kim, J. G. (2014). Carbohydrate reserves of non-fruiting young kiwifruit vines as affected by early artificial defoliation, *Horticulture, Environment, and Biotechnology*, 55(6), 462-470.

Notes

The author declare no conflict of interest.

Acknowledgement

This study was funded by a research program (PJ012046) of Rural Development Administration (RDA), Republic of Korea.

- Kwack, Y. B., Kang, S. M., Kim, H. L., Kim, M. J., Kim, S. C., & Lee, Y. B. (2015). Regrowth of axillary buds the current season and early growth and development the following year in fruiting young kiwifruit as affected by early defoliation, *Korean Journal of Environmental Agriculture*, 34(4), 288-293.
- Lionakis, S., & Schwabe, W. W. (1984). Bud dormancy in the kiwifruit, *Actinidia chinensis* Planch. *Annals of Botany*, 54(4), 467-484.
- Norton, R. L. (1988). Windbreaks: benefits to orchard and vineyard crops. *Agriculture, Ecosystems & Environment*, 22-23, 205-213.
- Oukabli, A., Bartolini, S., & VITI, R. (2003). Anatomical and morphological study of apple (*Malus domestica* Borkh.) flower buds growing under inadequate winter chilling. *Journal of Horticultural Science & Biotechnology*, 78(4), 580-585.
- Piller, G. J., & Meekings, J. S. (1997). The acquisition and utilization of carbon in early spring by kiwifruit shoots. *Annals of Botany*, 79(5), 573-581.
- Sunley, R. J., Atkinson, C. J., & Jones, H. G. (2006). Chill unit models and recent changes in the occurrence of winter chill and spring frost in the united kingdom. *Journal of Horticultural Science & Biotechnology*, 81(6), 949-958.
- Tanhora, K. (1988). *Encyclopedia of kiwifruit*, pp. 150-152, Yaki Inc., Japan.
- Weinberger, J. H. (1950). Chilling requirement of peach varieties. *Proceedings of the American Society for Horticultural Science*, 56, 122-128.
- Weinberger, J. H. (1954). Effects of high temperatures during the breaking of the rest of 'Sullivan Elberta' peach buds. *Proceedings of the American Society for Horticultural Science*, 63, 157-162.
- Williams, R. R. (1970). Factors affecting pollination in fruit trees. *Physiology of Tree Crops*. Luckwill, L. C., Cutting, C. V. (eds.). pp. 193-207, Academic Press, London.