

ARTICLE

Characteristics of Kwark Cheese Supplemented with *Bifidobacterium longum* KACC 91563

Minyu Song, Won Seo Park, Jayeon Yoo, Gi-Sung Han, Bu-Min Kim, Pil-Nam Seong, Mi-Hwa Oh, Kyung-Woon Kim¹, and Jun-Sang Ham*

Animal Products Research and Development Division, National Institute of Animal Science, RDA, Wanju 55365, Korea

¹Animal Biotechnology Division, National Institute of Animal Science, RDA, Wanju 55365, Korea



Received March 20, 2017
Revised September 21, 2017
Accepted October 8, 2017

*Corresponding author

Jun-Sang Ham

Animal Products Research and Development Division, National Institute of Animal Science, RDA, Wanju, 55365, Korea

Tel: +82-63-238-7366

Fax: +82-63-238-7397

E-mail: hamjs@korea.kr

Copyright © Korean Society for Food Science of Animal Resources

This is an open access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

The effect of addition of the probiotic *Bifidobacterium longum* KACC 91563 on the chemical and sensory properties of Kwark cheese produced using CHN-11 as a cheese starter were investigated. The addition of *B. longum* KACC 91563 to Kwark cheese did not change the composition or pH value of the cheese, compared with control. *B. longum* KACC 91563 survived at a level of 7.58 Log CFU/g and did not have any negative effect on survival of the cheese starter. A sensory panel commented that the addition of *B. longum* KACC 91563 made Kwark cheese more desirable to consumers, and that the probiotic supplementation had no effect on perceived taste. Thus, *B. longum* KACC 91563 can be used for inclusion of probiotic bacteria in cheese.

Keywords *Bifidobacterium longum*, probiotic, kwark cheese, sensory property

Introduction

The use of probiotic lactic acid bacteria (LAB) is a current topic of interest and a growing trend in the dairy industry. Probiotic bacteria are primarily used to manufacture dairy products. As reported by several authors, cheese is an excellent medium for addition of these probiotics. However, individual probiotic strains should be evaluated to determine whether they alter the sensory characteristics of a cheese, and to determine the effects of cheese production and storage on survival of the probiotic cells (Grattepanche *et al.*, 2008; Vinderola *et al.*, 2009; Yerlikaya and Özer, 2014). Cheese has been shown to be a good medium for transfer of probiotics into the intestine, as the cheese creates a buffer against the highly acidic conditions in the gastrointestinal tract (GIT) and thus creates a favorable environment for bacterial survival during gastric transit (Karimi *et al.*, 2012a; Karimi *et al.*, 2012b; Ortakci *et al.*, 2012). Supplementation of cheeses with probiotic LAB adds value and provides potential health benefits (Gomes *et al.*, 2011; Minervini *et al.*, 2012). Intake of cheese supplemented with probiotic bacteria has been associated with a variety of health-promoting benefits, such as immune system improvement, oral and gut health effects in the elderly, prevention of food allergies, and strengthening of intestinal immunity (Albenzio *et al.*, 2013a; Albenzio *et al.*, 2013b; Hatakka *et al.*, 2007; Ibrahim *et al.*, 2010; Lollo *et al.*, 2012; McFarland, 2000;

Medici *et al.*, 2004; Modzelewska-Kapituła *et al.*, 2010). In a previous study, we isolated probiotics from fecal samples of healthy Korean neonates. We have used one of the bacteria isolated in the previous study in this study, *Bifidobacterium longum* KACC 91563, a subspecies of *B. longum*, as it is a well-known probiotic strain that exhibits positive host effects (Shanahan, 2010). In addition, *B. longum* KACC 91563 produces family 5 extracellular solute-binding protein (ESBP), which not only reduces food allergies (Kim *et al.*, 2016), and also exhibits antioxidant activity (Chang *et al.*, 2013), and capacity for production of antihypertensive peptides (Ha *et al.*, 2015) by degrading milk proteins. Kwark, also known as quark or quarg, is a natural, soft, white, and un-ripened variety of fresh cheese ($\geq 50\%$ moisture) originating from Central Europe, where it is generally manufactured from cow milk only. These fresh cheeses appear to be ideally suited for use in delivery of probiotic organisms. Because they are stored at refrigeration temperatures, prolonged periods of ripening are not necessary (Heller *et al.*, 2003). Kwark is generally made from acid milk gels that are concentrated after fermentation with lactic cultures to \sim pH 4.6. Kwark is snowy white in color, with a subtle taste similar to sour cream, but a soft texture similar to cottage cheese. The health-enhancing properties of Kwark cheese can be improved by incorporation of functional probiotic bacteria. Several probiotics, which are well established in terms of their positive health effects, have been used in various dairy foods including Kwark cheese (Kadiya *et al.*, 2014; Kelly and O’Kennedy, 2001; Kosikowski, 1982; Lake *et al.*, 2005). However, the combined use of functional probiotic bacteria with the Kwark cheese starter has seldom been reported. Therefore, in this study, Kwark cheese was manufactured with commercial starter and *B. longum* KACC 91563, and its chemical and sensory properties, as well as the survival of the probiotic bacteria, were evaluated.

Materials and Methods

Materials

A cheese starter culture consisting of freeze dried CHN-11 (Chr. Hansen, Denmark) and *B. longum* KACC 91563 was used. Raw fresh cow’s milk was obtained from the National Institute of Animal Science.

Kwark cheese making

Kwark cheese was manufactured using the method described by Davis (1976), with some modifications, as shown

in Fig. 1. Kwark cheese was made using 10 L of pasteurized milk (63°C, 30 min) that was then cooled to 33°C using a cheese vat. The milk was inoculated with starter culture (0.002% CHN-11, v/v), and then the same amount of *B. longum* KACC 91563 (approximately 10^5 - 10^6 CFU/g) was separately added to the milk. The control was produced with starter culture alone. Rennet (100 μ L/10 mL) was added and mixed thoroughly. The cheese was incubated at 33°C until it reached pH 4.8. The resulting curd was cut into 2-cm cubes with cheese knives. The whey was removed, and the curds were cooled at low pressure, and then stored at 4°C.

Proximate composition

Moisture, protein, fat, and salt contents were analyzed using a Food Scan (Food Scan™ Lab 78810, Foss Tecator Co. Ltd., Denmark).

Chemical characteristics

The pH of the samples was measured using a pH-meter (CH/SevenEasy S20K, Mettler-Toledo, Switzerland). Acetic acid, d-lactic acid, l-lactic acid and lactose were analyzed using a Automated Chemistry Analyzer (Thermo Scientific, Finland).

Microbiological analyses

Samples were serially diluted and then used for viable plate counts. Viable counts of lactic acid bacteria in the cheese samples were determined using de Man, Rogosa, and Sharpe agar (MRS; Difco, USA). Viable counts of bifidobacteria in cheese samples were determined using *Bifidobacterium* Selective Medium (BSM; TOS supplemented with mupirocin, Thitaram *et al.*, 2005). Kwark cheese (10 g) was homogenized for 1 min in a sterile stomacher bag with 90 mL of sterile distilled water using a stomacher (Bagmixer® 400W, Interscience, France) for 1 min at high speed to obtain a slurry for the first dilution, and subsequent serial dilutions were made in diluent before spread plating on BSM and MRS plates. The BSM plates were incubated for 48 h at 37°C under anaerobic conditions (GasPak™ EZ Anaerobe Container System, Dickinson and Company, USA). The MRS agar plates were incubated under aerobic conditions for 24-48 h at 37°C. Colony forming units (CFU) per gram were counted per plate.

Sensory evaluation of Kwark cheese

Semi-trained panelists (n=10) evaluated and analyzed

samples of Kwark cheese with and without *B. longum* KACC 91563. The panelists used were chosen among the members of Animal Products Development Division, and based on their previous experiences in sensory evaluation of dairy products. The panelists were asked to score the samples for color, flavor, texture, taste, and overall acceptance using the following hedonic 9-point scale: like extremely (9), like very much (8), like moderately (7), like slightly (6), like moderately (5), neither like nor dislike (4), dislike moderately (3), dislike very much (2), dislike extremely (1) (Jaclyn *et al.*, 2014).

Statistical analyses

Data were analyzed using the Statistical Analysis System program (version 9.2) (SAS, 2010). Means were compared by analysis of variance (ANOVA) followed by Duncan's multiple range test and the difference after 10 d storage were compared by a Student's t-test. Significance of differences was defined at the 5% level ($p < 0.05$). All of the experiments were performed twice in duplicate ($n = 4$).

Results and Discussion

Chemical characteristics of Kwark cheese

The proximate composition of Kwark cheese with and without addition of probiotic bacteria is presented in Table 1. Control was made with using commercial starter but, treatment was made with commercial starter and *B. longum* KACC 91563. The average moisture content of Kwark

cheese ranged between 68.27 and 67.10% and no differences were observed between the treatment and control. The protein and fat content of treatment was lower than in control. The salt content of treatment was $0.70 \pm 0.08\%$, and showed no difference from that of control. This finding was in agreement with the results reported by Gursoy *et al.* (2014). The addition of *B. longum* KACC 91563 did not change the cheese composition compared with control. Thus, the treatment and control showed no significant differences in moisture, protein, fat, or salt contents.

As shown in Table 2, pH and chemical composition of Kwark cheese with and without addition of probiotic bacteria are examined during the storage. pH of treatment was lower than that of control. Magdoub *et al.* (2005) investigated that the decrease of pH may be due to the convert to residual lactose in cheese to lactic acid and free fatty acid which had developed in the cheese. After 10 d, pH of treatment and control cheese was increased from 4.36 to 4.54, and 4.52 to 4.63, respectively. The level of acetic acid and d-lactic acid of treatment was higher than that of control. Treatment and control were increased during storage days in acetic acid and d-lactic acid. The control and treatment revealed the level of l-lactic acid increasing over 10 d. Lactose of treatment was higher than that of control. However, it was seen that the lactose of all the samples has decreased during storage days. Lactose of fresh cheese like quark was 2 to 4% (Park, 2003). Thus, the results obtained that pH of Kwark cheese with commercial starter and *B. longum* KACC 91563 seemed

Table 1. Proximate composition of Kwark cheese supplemented with commercial starter and *Bifidobacterium longum* KACC 91563

Kwark cheese	Moisture (%)	Protein (%)	Fat (%)	Salt (%)
C ¹	66.27±4.73	12.46±1.89	17.50±2.43	0.65±0.02
T ²	67.10±3.08	12.10±1.77	16.79±1.29	0.70±0.08

¹Control, Kwark cheese added with commercial starter; ²Treatment, Kwark cheese supplemented with commercial starter and *B. longum* KACC 91563

Data are expressed as mean±standard deviation ($n = 4$).

Values in the same column are not significantly different ($p > 0.05$).

Table 2. Chemical analysis of Kwark cheese supplemented with commercial starter and *Bifidobacterium longum* KACC 91563

Kwark cheese	Storage days	pH	Acetic acid (g/l)	D-Lactic Acid (g/l)	L-Lactic Acid (g/l)	Lactose (g/l)
C ¹	0	4.52±0.21	0.60±0.00	0.18±0.04	6.79±0.68	21.57±0.29
	10	4.63±0.06	0.84±0.06*	0.31±0.02	7.20±0.36	20.12±0.98
T ²	0	4.36±0.07	0.64±0.03	0.22±0.02	6.69±0.26	22.17±0.38
	10	4.54±0.01	0.91±0.00*	0.32±0.02*	7.21±0.30	21.43±0.76

¹Control, Kwark cheese added with commercial starter; ²Treatment, Kwark cheese supplemented with commercial starter and *B. longum* KACC 91563

Data are expressed as mean±standard deviation ($n = 4$).

*Values in the same group are significantly different by t-test ($p < 0.05$).

to be reduced due to more the production of lactic acid and other organic acids than that of control.

Microbial characteristics of Kwark cheese

The growth of CHN-11 and *B. longum* KACC 91563, based on viable cell counts in Kwark cheese, are shown in Table 3. According to Table 3, the number of lactic acid bacteria and *B. longum* KACC 91563 of cells increased about 2 and 1 Log CFU/g immediately after inoculation, respectively. In treatment, the number of lactic acid bacteria was lower than in the control, but the difference was not significant ($p>0.05$). Bifidobacterial counts on selective agar plates incubated under anaerobic conditions showed successful incorporation into Kwark cheese at a level of 7.58 Log CFU/g (Table 3). Thus, Kwark cheese with and without addition of probiotics showed no significant difference ($p>0.05$) viable cell counts. It should be noted that different *Bifidobacterium* species will exhibit different survivability or have different impacts on the sensory attributes of dairy products because bifidobacteria species differ in their nutrient requirements, growth characteristics, and metabolic activity.

Several factors must be considered when adding probiotics to fermented foods such as cheese. Mainly, the probiotics must be present at high viable cell counts at the time of consumption to achieve the desired benefits (Gomes *et al.*, 1995). For maximal benefit, a probiotic dairy

product should contain at least 10^6 - 10^7 CFU/g probiotic bacteria at the time of consumption, and should be consumed regularly at a quantity of higher than 100 g per day (Boylston *et al.*, 2004; Gomes and Malcata, 1999; Matijević *et al.*, 2009; Medici *et al.*, 2004). According to these criteria, daily consumption of 10 g of Kwark cheese supplemented with *B. longum* KACC 91563 (containing 10^7 CFU/g) would meet the minimum probiotic bacteria requirements.

Sensory properties of Kwark cheese

The results of the sensory evaluation of the Kwark cheese samples are shown in Table 4. In sensory properties, flavor, texture, taste and overall acceptance were decreased generally after 10 d. The results seemed to be due to loss of freshness of the cheese. The individual effect of supplementation of Kwark cheese with or without *B. longum* KACC 91563 on taste and overall acceptance was statistically significant during storage days ($p<0.05$). Mahmudi *et al.* (2012) found that supplementation of Iranian white cheese with *B. animalis* and *Lactobacillus rhamnosus* had no significant effect on the texture and flavor of the cheese. These results are consistent with the findings of previous studies by Gursoy and Kinik (2010) and Zomorodi *et al.* (2010). In addition, cheese supplemented with bifidobacteria shows higher levels of acetic acids compared with controls (Ong *et al.*, 2007); however, the

Table 3. Viable cell counts in Kwark cheese supplemented with commercial starter and *Bifidobacterium longum* KACC 91563

		Lactic acid bacteria	<i>Bifidobacterium longum</i> KACC 91563
Initial inoculation (CFU/g)	C ¹	5.96±0.16	-
	T ²	5.59±0.09	6.50±0.05
Whey off (CFU/g)	C ¹	7.78±0.08	-
	T ²	7.73±0.14	7.14±0.39
Kwark cheese (CFU/g)	C ¹	8.26±0.44	-
	T ²	7.74±0.04	7.58 ± 0.05

¹Control, Kwark cheese added with commercial starter; ²Treatment, Kwark cheese supplemented with commercial starter and *B. longum* KACC 91563

Data are expressed as mean±standard deviation (n=4).

Values are not significantly different ($p>0.05$).

Table 4. Sensory evaluation of Kwark cheese supplemented with commercial starter and *Bifidobacterium longum* KACC 91563

Kwark cheese	Storage days	Color	Flavor	Texture	Taste	Overall acceptance
C ¹	0	7.58±0.10	6.05±0.59	5.93±0.21	5.75±0.64*	6.15±0.87*
	10	7.65±0.21	5.40±1.13	5.40±0.85	4.45±1.34	4.75±1.48
T ²	0	7.68±0.21	6.28±0.57	5.98±0.26	5.83±0.25*	6.30±0.42*
	10	7.60±0.14	5.55±0.21	5.35±0.49	4.85±0.21	5.30±0.00

¹Control, Kwark cheese added with commercial starter; ²Treatment, Kwark cheese supplemented with commercial starter and *B. longum* KACC 91563; panel=10

Data are expressed as mean±standard deviation (n=4).

*Values in the same group are significantly different by t-test ($p<0.05$).

organoleptic properties were unchanged (Gobbetti *et al.*, 1998). The inclusion of some probiotic bacteria in dairy foods, such as cheese, does not markedly change the sensory profile of the food (Champagne *et al.*, 2005; Cruz *et al.*, 2009). Escobar *et al.* (2012) suggested that the probiotic supplementation of Panela cheese had no perceived effect. In addition, Buriti *et al.* (2005) reported that the addition of *L. acidophilus* to Minas fresh cheese had no effect on flavor compared with a control. In agreement with these results, our study showed that cheese supplemented with probiotic bifidobacteria attained equal or greater acceptance in sensory evaluation compared with control, and the addition of *B. longum* KACC 91563 to Kwark cheese did not create any sensorial defects.

Conclusions

In this study, we produced Kwark cheese supplemented with *B. longum* KACC 91563 to investigate the effects on the chemical and sensory characteristics of the cheese. The compositional analysis showed that any differences were not significant ($p > 0.05$). The chemical analysis showed that pH of Kwark cheese with commercial starter and *B. longum* KACC 91563 was lower than that of control. In addition, no significant differences ($p > 0.05$) in lactic acid bacterial counts were detected between Kwark cheese with and without addition of probiotics. Kwark cheese supplemented with *B. longum* KACC 91563, which has the ability to alleviate food allergies, retained a viable cell count $> 10^7$ CFU/g of bifidobacteria. Thus, daily consumption of 10 g of Kwark cheese would meet the minimum probiotic requirement. Kwark cheese supplemented with *B. longum* KACC 91563 was preferred over the control, but addition of probiotics did not significantly alter the color, flavor, texture, taste, or overall acceptance of the Kwark cheese. Based on these findings, addition of *B. longum* KACC 91563 improved product quality without significant negative effects on the characteristics of the Kwark cheese. Therefore, Kwark cheese supplemented with *B. longum* KACC 91563 shows promise for use as a probiotic or functional cheese against food allergies.

Acknowledgements

This study was supported by 2017-Postdoctoral Fellowship Program of National Institute of Animal Science (Project No. PJ01196001), Rural Development Administration, Republic of Korea.

References

- Albenzio, M., Santillo, A., Caroprese, M., Braghieri, A., Sevi, A., and Napolitano, F. (2013a) Composition and sensory profiling of probiotic Scamorza ewe milk cheese. *J. Dairy Sci.* **96**, 2792-2800.
- Albenzio, M., Santillo, A., Caroprese, M., Ruggieri, D., Napolitano, F., and Sevi, A. (2013b) Physicochemical properties of Scamorza ewe milk cheese manufactured with different probiotic cultures. *J. Dairy Sci.* **96**, 2781-2791.
- Boylston, T. D., Vinderola, C. G., Ghoddusi, H. B., and Reinheimer, J. A. (2004) Incorporation of bifidobacteria into cheeses: challenges and rewards. *Int. Dairy J.* **14**, 375-387.
- Buriti, F. C. A., da Rocha, J. S., and Saad, S. M. I. (2005) Incorporation of *Lactobacillus acidophilus* in Minas fresh cheese and its implications for textural and sensorial properties during storage. *Int. Dairy J.* **15**, 1279-1288.
- Champagne, C. P., Gardner, N. J., and Roy, D. (2005) Challenges in the addition of probiotic cultures to foods. *Crit. Rev. Food Sci. Nutr.* **45**, 61-84.
- Chang, O. K., Seol, K. H., Jeong, S. G., Oh, M. H., Park, B. Y., Perin, C., and Ham, J. S. (2013) Casein hydrolysis by *Bifidobacterium longum* KACC 91563 and antioxidant activities of peptides derived therefrom. *J. Dairy Sci.* **96**, 5544-5555.
- Cruz, A. G., Buriti, F. C. A., De Souza, C. H. B., Favia, J. A. F., and Saad, S. M. I. (2009) Probiotic cheese: Health benefits, technological and stability aspects. *Trends Food Sci. Tech.* **20**, 344-354.
- Davis, J. G. (1976) Cheese, volume (III) manufacturing methods. Churchill Livingstone, London, pp. 757-760.
- Escobar, M. C., Van Tassell, M. L., Martínez-Bustos, F., Singh, M., Castaño-Tostado, E., Amaya-Llano, S. L., and Miller, M. J. (2012) Characterization of a Panela cheese with added probiotics and fava bean starch. *J. Dairy Sci.* **95**, 2779-2787.
- Gobbetti, M., Corsetti, A., Smacchi, E., Zocchetti, A., and de Angelis, M. (1998) Production of Crescenza cheese by incorporation of bifidobacteria. *J. Dairy Sci.* **81**, 37-47.
- Gomes, A. A., Braga, S. P., Cruz, A. G., Cadena, R. S., Lollo, P. C. B., Carvalho, C., Amaya-Farfán, J., Faria, J. A. F., and Bolini, H. M. A. (2011) Effect of the inoculation level of *Lactobacillus acidophilus* in probiotic cheese on the physicochemical features and sensory performance compared with commercial cheeses. *J. Dairy Sci.* **94**, 4777-4786.
- Gomes, A. M. P. and Malcata, F. X. (1999) *Bifidobacterium* spp. and *Lactobacillus acidophilus*: biological, biochemical, technological and therapeutic properties relevant for use as probiotics. *Trends Food Sci. Technol.* **10**, 139-157.
- Gomes, A. M. P., Malcata F. X., Klaver, F. A. M., and Grande, H. J. (1995) Incorporation and survival of *Bifidobacterium* sp. strain Bo and *Lactobacillus acidophilus* strain Ki in a cheese product. *Neth. Milk Dairy J.* **49**, 71-95.
- Grattepanche, F., Miescher-Schwenninger, S., Meile, L., and Lacroix, C. (2008) Recent developments in cheese cultures with protective and probiotic functionalities. *Dairy Sci. Technol.* **88**, 421-444.

- Gursoy, O., Gokce, R., Con, A. H., and Kinik, O. (2014) Survival of *Bifidobacterium longum* and its effect on physicochemical properties and sensorial attributes of white brined cheese. *Int. J. Food Sci. Nutr.* **65**, 816-820.
- Gursoy, O. and Kinik, O. (2010) Incorporation of adjunct cultures of *Enterococcus faecium*, *Lactobacillus paracasei* ssp. *paracasei* and *Bifidobacterium bifidum* into white cheese. *J. Food Agric. Environ.* **8**, 107-112.
- Ha, G. E., Chang, O. K., Jo, S. M., Han, G. S., Park, B. Y., Ham, J. S., and Jeong, S. G. (2015) Identification of antihypertensive peptides derived from low molecular weight casein hydrolysates generated during fermentation by *Bifidobacterium longum* KACC 91563. *Korean J. Food Sci. An.* **35**, 738-747.
- Hatakka, K., Ahola, A. J., Yli-Knuutila, H., Richardson, M., Poussa, T., Meurman, J. H., and Korpela, R. (2007) Probiotics reduce the prevalence of oral *Candida* in the elderly-A randomized controlled trial. *J. Dent. Res.* **86**, 125-130.
- Heller, K. J., Bockelmann, W., Schrezenmeir, J., and deVrese, M. (2003) Cheese and its potential as a probiotic food. In E. R. Farnworth (ed), Handbook of fermented functional foods. CRC Press, BR, pp. 203-225.
- Ibrahim, F., Ruvio, S., Granlund, L., Salminen, S., Viitanen, M., and Ouwehand, A. C. (2010) Probiotics and immunosenescence: Cheese as a carrier. *FEMS Immunol. Med. Microbiol.* **59**, 53-59.
- Jaclyn, J. K., Charles, A. S., Lorenzo, A. P., Derek, J. S., and Linda, M. B. (2014) Comparison of the hedonic general labeled magnitude scale with the hedonic 9-point scale. *J. Food Sci.* **79**, S238-245.
- Kadiya, K. S., Kanawjia, S. K., and Solanki, A. K. (2014) Survival of free and encapsulated probiotic bacteria and their effect on the sensory properties of Quarg cheese. *IJFF.* **3**, 61-76.
- Karimi, R., Mortazavian, A. M., and Karimi, M. (2012a) Incorporation of *Lactobacillus casei* in Iranian ultrafiltered Feta cheese made by partial replacement of NaCl with KCl. *J. Dairy Sci.* **95**, 4209-4222.
- Karimi, R., Sohrabvandi, S., and Mortazavian, A. M. (2012b) Sensory characteristics of probiotic cheese. *Compr. Rev. Food Sci. Food Saf.* **11**, 437-452.
- Kelly, P. M. and O'Kennedy, B. T. (2001) The effect of casein/whey protein ratio and minerals on the rheology of fresh cheese gels using a model system. *Int. Dairy J.* **11**, 525-532.
- Kim, J. H., Jeun, E. J., Hong, C. P., Kim, S. H., Jang, M. S., Lee, E. J., Moon, S. J., Yun, C. H., Im, S. H., Jeong, S. G., Park, B. Y., Kim, K. T., Seoh, J. Y., Kim, Y. K., Oh, S. J., Ham, J. S., Yang, B. G., and Jang, M. H. (2016) Extracellular vesicle-derived protein from *Bifidobacterium longum* alleviates food allergy through mast cell suppression. *J. Allergy Clin. Immunol.* **137**, 507-516.
- Kosikowski, F. (1982) Cheese and fermented milk foods. 2th ed, Brooktondale, NY, pp. 144-155.
- Lake, L., Hudson, A., Cressey, P., and Gilbert, S. (2005) Risk Profile: *Listeria monocytogenes* in soft cheeses. Client report FW0382 of New Zealand Food Safety Authority.
- Lollo, P. C., Cruz, A. G., Morato, P. N., Moura, C. S., Carvalho-Silva, L. B., Oliveira, C. A., Faria, J. A., and Amaya-Farfan, J. (2012) Probiotic cheese attenuates exercise-induced immune suppression in Wistar rats. *J. Dairy Sci.* **16**, 317-321.
- Magdoub, M. N., Osman, S. H. G., and El-Kenawy, M. M. (2005) Effect of different starter cultures on composition and microbiological quality of Ain shams cheese. *Egyptian J. Applied Sci.* **10**, 132-141.
- Mahmoudi, M., Asl, A. K., and Zomorodi, S. (2012) The influence of probiotic bacteria on the properties of Iranian white cheese. *Int. J. Dairy Technol.* **65**, 561-567.
- Matijević, B., Božanić, R., and Tratnik, L. (2009) The influence of lactulose on growth and survival of probiotic bacteria *Lactobacillus acidophilus* La-5 and *Bifidobacterium animalis* subsp. *lactis* BB-12 in reconstituted sweet whey. *Mljekarstvo.* **59**, 20-27.
- McFarland, L. (2000) A review of evidences of health claims for biotherapeutic agents. *Microb. Ecol. Health Dis.* **12**, 65-76.
- Medici, M., Vinderola, C. G., and Perdigon, G. (2004) Gut mucosal immunomodulation by probiotic fresh cheese. *Int. Dairy J.* **14**, 611-618.
- Minervini, F., Siragusa, S., Faccia, M., Dal Bello, F., Gobetti, M., and De Angelis, M. (2012) Manufacture of Fior di Latte cheese by incorporation of probiotic lactobacilli. *J. Dairy Sci.* **95**, 508-520.
- Modzelewska-Kapituła, M., Kobukowski, J. A. N., and Kłębukowska, L. (2010) The influence of feeding diets containing white cheese, produced with prebiotics and the potentially probiotic *Lactobacillus plantarum* strain, on the gastrointestinal microflora of rats. *Czech J. Food Sci.* **28**, 139-145.
- Ong, L., Henriksson, A., and Shah, N. P. (2007) Chemical analysis and sensory evaluation of Cheddar cheese produced with *Lactobacillus acidophilus*, *Lb. casei*, *Lb. paracasei* or *Bifidobacterium* sp. *Int. Dairy J.* **17**, 937-945.
- Ortakci, F., Broadbent, J. R., Mcmanus, W. R., and Mchahon, D. J. (2012) Survival of microencapsulated probiotic *Lactobacillus paracasei* LBC-1e during manufacture of Mozzarella cheese and simulated gastric digestion. *J. Dairy Sci.* **95**, 6274-6281.
- Park, S. Y. (2003) Milk production and processing. Yu Han Mun Hwa sa, Seoul, Korea, p. 285. (in Korean)
- SAS. (2010) SAS/STAT Software for PC. Release 9.2, SAS Institute Inc., Cary, NC, USA.
- Shanahan, F. (2010) Probiotics in perspective. *Gastro.* **139**, 1808-1812.
- Thitaram, S. N., Siragusa, G. R., and Hinton Jr., A. (2005) *Bifidobacterium*-selective isolation and enumeration from chicken caeca by a modified oligosaccharide antibiotic-selective agar. *Lett. Appl. Microbiol.* **41**, 355-360.
- Vinderola, G., Prosello, W., Molinari, F., Ghiberto, D., and Reinheimer, J. (2009) Growth of *Lactobacillus paracasei* A13 in Argentinian probiotic cheese and its impact on the characteristics of the product. *Int. J. Food Microbiol.* **135**, 171-174.
- Yerlikaya, O. and Özer, E. (2014) Production of fresh white cheese using co-culture with *Streptococcus thermophilus*. *Food Sci. Technol.* **34**, 471-477.

Zomorodi, S., Khosrowshahi Asl, A., Razavirohani, S. M., and Miraghaei, S. (2010) Survival of *Lactobacillus casei*, *Lactobacillus plantarum* and *Bifidobacterium bifidum* and their eff-

ect on composition of Iranian white cheese produced by ultra-filtration technique. *Int. J. Dairy Technol.* **63**, 1-8.