

Prevalence and Intensity of *Clonorchis sinensis* Metacercariae in Freshwater Fish from Wicheon Stream in Gunwi-gun, Gyeongsangbuk-do, Korea

Woon-Mok Sohn^{1*}, Byoung-Kuk Na¹, Shin-Hyeong Cho², Jung-Won Ju², Dong-Chul Son³

¹Department of Parasitology and Tropical Medicine, and Institute of Health Sciences, Gyeongsang National University College of Medicine, Jinju 52727, Korea; ²Division of Vectors and Parasitic Diseases, Centers for Disease Control and Prevention, Osong 28159, Korea; ³Infectious Disease Research Section, Gyeongsangbuk-do Institute of Health and Environment, Yeongcheon 38874, Korea

Abstract: The infection status of *Clonorchis sinensis* metacercariae (CsMc) was examined in freshwater fish from a highly prevalent site, Wicheon (a branch of Nakdong-gang), which is located in Gunwi-gun, Gyeongsangbuk-do, the Republic of Korea. Total 1,162 fish in 32 species were examined by the artificial digestion method through 6 years. CsMc were detected in 720 (67.5%) out of 1,067 fish (26 spp.) and their density was 610 per fish infected. In the susceptible gobioninid fish group, i.e., *Pungtungia herzi*, *Squalidus gracilis majimae*, *Squalidus japonicus coreanus*, *Sarcocheilichthys variegatus wakiyae* and *Pseudorasbora parva*, all of 323 fish were infected with an average of 1,310 CsMc. Total 23 (95.8%) gobioninid fish, i.e., *Pseudogobio esocinus*, *Abbottina springeri*, *Hemibarbus longirostris*, *Microphysogobio koreensis*, and *Microphysogobio jeoni*, were infected with 127 CsMc in average. In the acheilognathinid fish (bitterlings) group, the prevalence was 77.0%, and the density was 50 CsMc per fish infected. In the rasborinid fish (chubs) group, i.e., *Zacco platypus*, *Zacco temminckii*, *Zacco koreanus*, and *Opsariichthys uncirostris amurensis*, 147 (36.5%) out of 403 fish examined were infected with 15 CsMc in average. The susceptibility indices of CsMc were 412 in the overall positive fish group, 1,310 in the gobioninid group-1, 122 in the gobioninid group-2, 38.5 in the acheilognathinid group, and 5.5 in the rasborinid fish group. Conclusively, it was confirmed that CsMc are highly prevalent in fish from Wicheon, and their infection tendency varied according to the subfamily groups in Cyprinidae fish hosts.

Key words: *Clonorchis sinensis*, metacercaria, susceptibility index, gobioninae, acheilognathinae, rasborinae, cyprinidae fish host, Wicheon

INTRODUCTION

Nowadays *Clonorchis sinensis* infection is the most endemic parasitic disease in the Republic of Korea (= Korea). The prevalence of this endemic disease was 1.9%, and about 932,540 residents are estimated to be infected, which is the highest value among the prevalences in the nationwide survey on helminthic infections in Korea [1]. The endemicity of clonorchiasis has maintained at relatively high levels in riverside areas of Korea [2-8]. Especially, in 1981, high prevalences were reported from the riverside residents in 7 major rivers, i.e., Nakdong-gang (gang= River) (40.2%), Yeongsan-gang (30.8%), Seomjin-gang (17.3%), Tamjin-gang (15.9%), Han-gang (15.7%),

Geum-gang (12.0%), and Mangyeong-gang (8.0%) [2]. Also in 2006 and 2007, the egg positive rates of *C. sinensis* were surveyed from riverside residents in major 5 rivers of Korea [5,7]. Recently, Jeong et al. [8] reported the prevalence of clonorchiasis from the residents of 5 major rivers as Nakdong-gang (11.7%), Seomjin-gang (9.9%), Geum-gang (6.5%), Yeongsan-gang (3.1%), and Han-gang (1.0%).

On the other hand, many Korean workers surveyed on freshwater fish, the infection sources of clonorchiasis, collected from various endemic riverside areas to estimate the endemicities of clonorchiasis. The results revealed that more than 49 fish species in 7 families have been reported as the second intermediate hosts of *C. sinensis* [9-18]. Especially, Kim et al. [12] examined 677 freshwater fish (21 spp.) collected from 34 localities to know the infection status of *C. sinensis* metacercariae (CsMc). In addition, Cho et al. [14] surveyed the infection status of CsMc in freshwater fish from 3 wide regions, which are tentatively divided by the latitudinal levels of the Korean peninsula. They examined total 136 freshwater fish (16 spp.) from

•Received 27 November 2017, revised 29 December 2017, accepted 3 January 2018.

*Corresponding author (wmsohn@gnu.ac.kr)

© 2018, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Wicheon, the same area in this study, in 2008. Later, in 2014, Cho et al. [16] investigated the prevalence of zoonotic trematode metacercariae in freshwater fish from Gangwon-do (Province), Korea. Recently, Sohn et al. [17] surveyed the infection status of digenetic trematode metacercariae including *C. sinensis* in freshwater fish from the water systems of Hantan-gang and Imjin-gang in northern regions of Korea. Sohn et al. [18] also investigated the infection status with *C. sinensis* metacercariae in fish from water systems of Seomjin-gang.

Wicheon is one of the branch streams of Nakdong-gang, which arises from a mountainous area of Goro-myeon (myeon = township) in Gunwi-gun (gun = county), flows via Gunwi-gun and Euseong-gun, and unites with the main stream at Sangju-si (si = city), Gyeongsangbuk-do. The water ecosystem of this stream is more or less healthy but the ecological conditions for fish is not so good [19]. However, it has been known that freshwater fish such as *Pungtungia herzi*, *Squalidus gracilis majimae*, *S. japonicus coreanus*, and *Pseudogobio esocinus* from this stream were heavily infected with CsMc [14]. Therefore, for a period of 7 years, we focused on the infection status of CsMc in fish from a highly prevalent site of Wicheon, Gyeongsangbuk-do, Korea, and analyzed the infection status of CsMc according to the subfamily groups of Cyprinidae fish hosts of *C. sinensis*.

MATERIALS AND METHODS

Fish collection site and freshwater fish examined

We collected total 1,162 freshwater fish in 32 species in Wicheon (a stream of Nakdong-gang), which is located in Woobo-myeon, Gunwi-gun (latitude: 38.43268; longitude: 127.4375), Gyeongsangbuk-do, Korea. The numbers and species of fish examined in each year were as follows:

In 2011, total 105 freshwater fish in 13 species (no. of fish examined), i.e., *Zacco platypus* (49), *Squalidus japonicus coreanus* (10), *Zacco temminckii* (10), *Squalidus gracilis majimae* (8), *Pungtungia herzi* (5), *Acheilognathus majusculus* (5), *Opsariichthys uncirostris amurensis* (5), *Carassius auratus* (3), *Pseudogobio esocinus* (3), *Acanthorhodeus macropterus* (3), *Sarcocheilichthys variegatus wakiyae* (2), and *Acheilognathus koreensis* (1), *Cobitis sinensis* (1), were examined. In 2013, total 107 freshwater fish in 12 species, i.e., *P. herzi* (20), *Z. platypus* (20), *Acheilognathus yamatsutae* (16), *Z. temminckii* (15), *A. koreensis* (10), *S. gracilis majimae* (9), *S. variegatus wakiyae* (6), *Coreoperca herzi* (5), *Abbottina springeri* (2), *C. sinensis* (2), *C. auratus* (1), and *Lepomis macrochirus* (1), were examined.

In 2014, total 338 freshwater fish in 24 species, i.e., *Z. platypus* (60), *Z. temminckii* (50), *P. herzi* (47), *A. koreensis* (29), *A. yamatsutae* (25), *S. japonicus coreanus* (14), *C. auratus* (13), *S. gracilis majimae* (13), *Acheilognathus rhombeus* (11), *S. variegatus wakiyae* (11), *A. majusculus* (10), *Micropterus salmoides* (8), *C. sinensis* (8), *C. herzi* (7), *P. esocinus* (7), *Misgurnus anguillicaudatus* (6), *Odontobutis platycephala* (4), *Pseudorasbora parva* (4), *L. macrochirus* (4), *Cobitis lutheri* (3), *Acheilognathus lanceolatus* (2), *Microphysogobio jeoni* (1), and *Microphysogobio koeensis* (1), were examined. In 2015, total 245 freshwater fish in 19 species, i.e., *Z. platypus* (50), *Z. temminckii* (49), *P. herzi* (41), *A. koreensis* (29), *A. yamatsutae* (16), *S. japonicus coreanus* (15), *S. gracilis majimae* (11), *C. auratus* (10), *A. rhombeus* (7), *A. springeri* (4), *S. variegatus wakiyae* (3), *O. uncirostris amurensis* (3), *P. parva* (1), *Rhynchocypris oxycephalus* (1), *A. macropterus* (1), *P. esocinus* (1), *C. herzi* (1), *M. koeensis* (1), and *A. lanceolatus* (1), were examined.

In 2016, total 168 freshwater fish in 17 species, i.e., *A. koreensis* (35), *Z. platypus* (30), *P. herzi* (30), *A. rhombeus* (15), *A. yamatsutae* (15), *Zacco koreanus* (14), *O. platycephala* (6), *C. auratus* (5), *S. japonicus coreanus* (4), *A. lanceolatus* (3), *C. herzi* (3), *P. esocinus* (2), *S. gracilis majimae* (2), *Hemibarbus longirostris* (1), *M. anguillicaudatus* (1), *Siniperca scherzeri* (1), and *M. koreensis* (1), were examined. In 2017, total 199 freshwater fish in 17 species, i.e., *Z. platypus* (50), *S. japonicus coreanus* (31), *A. yamatsutae* (27), *P. herzi* (26), *A. koreensis* (26), *Z. temminckii* (10), *P. parva* (7), *Acanthorhodeus gracilis* (5), *S. variegatus wakiyae* (3), *C. herzi* (2), *C. auratus* (2), *Z. koreanus* (2), *A. rhombeus* (2), *S. scherzeri* (2), *M. salmoides* (2), *C. lutheri* (1), and *Pseudobagrus fulvidraco* (1), were examined.

Examination methods

All collected fish with ice were transferred to the laboratory of the Department of Parasitology and Tropical Medicine, Gyeongsang National University College of Medicine, Jinju, Korea. After identification of fish species, they were individually ground with a mortar or grinder. Each ground fish meat was mixed with artificial gastric juice, and the mixture was incubated at 36°C for 2 hr. The digested material was filtered with 1 × 1 mm of mesh, and washed with 0.85% saline until the supernatant became clear. The sediment was carefully examined under a stereomicroscope. The metacercariae of *C. sinensis* were separately collected by the general morphological feature [13,15], and they were counted to get hold of the infection rate and density by fish species. The susceptibility indices of

Table 1. Infection status of *Clonorchis sinensis* metacercariae in fish from Wicheon (a stream of Nakdong-gang) in Gunwi-gun, Gyeong-sangbuk-do

Year and fish species	No. of fish examined	No. (%) of fish infected	No. of metacercariae detected		
			Total	Range	Average
2011					
<i>Pungtungia herzi</i>	5	5 (100)	5,216	340-2,420	1,043
<i>Squalidus japonicus coreanus</i>	10	10 (100)	818	13-295	82
<i>Squalidus gracilis majimae</i>	8	8 (100)	4,960	9-1,035	620
<i>Sarcocheilichthys variegatus</i>	2	2 (100)	1,901	2-12	951
<i>Pseudogobio esocinus</i>	3	3 (100)	16	-	5
<i>Acheilognathus majusculus</i>	5	1 (20.0)	1	2	1
<i>Acanthorhodeus macropterus</i>	3	3 (100)	91	87	30
<i>Acheilognathus koreensis</i>	1	1 (100)	9	-	9
<i>Zacco platypus</i>	49	18 (59.3)	129	1-32	7
Subtotal	86	51 (59.3)	13,141	1-2,420	258
2013					
<i>Pungtungia herzi</i>	20	20 (100)	22,790	27-11,290	1,140
<i>Squalidus gracilis majimae</i>	9	9 (100)	8,687	33-2,720	965
<i>Sarcocheilichthys variegatus</i>	6	6 (100)	2,545	26-1,245	424
<i>Abbottina springeri</i>	2	2 (100)	95	44-51	48
<i>Acheilognathus koreensis</i>	10	10 (100)	58	1-14	6
<i>Acheilognathus yamatsutae</i>	16	6 (37.5)	23	1-10	4
<i>Zacco platypus</i>	20	8 (40.0)	129	1-70	16
<i>Zacco temminckii</i>	15	1 (6.7)	2	-	2
Subtotal	98	62 (63.3)	34,329	1-11,290	554
2014					
<i>Pungtungia herzi</i>	47	47 (100)	116,805	6-31,250	2,485
<i>Squalidus japonicus coreanus</i>	14	14 (100)	25,291	140-8,460	1,807
<i>Squalidus gracilis majimae</i>	13	13 (100)	20,285	132-6,130	1,560
<i>Sarcocheilichthys variegatus</i>	11	11 (100)	8,936	13-2,730	812
<i>Pseudorasbora parva</i>	4	4 (100)	2,911	483-1,180	728
<i>Microphysogobio jeoni</i>	1	1 (100)	365	-	365
<i>Acheilognathus koreensis</i>	29	29 (100)	1,099	1-184	38
<i>Microphysogobio koeensis</i>	1	1 (100)	12	-	12
<i>Acheilognathus yamatsutae</i>	25	22 (88.0)	560	1-136	26
<i>Acheilognathus rhombeus</i>	11	9 (81.8)	782	1-290	87
<i>Acheilognathus majusculus</i>	10	2 (20.0)	3	1-2	2
<i>Acheilognathus lanceolatus</i>	2	1 (50.0)	283	-	283
<i>Zacco platypus</i>	60	40 (66.7)	447	1-66	11
<i>Zacco temminckii</i>	50	1 (2.0)	1	-	1
<i>Carassius auratus</i>	13	1 (7.7)	1	-	1
<i>Micropterus salmoides</i>	8	1 (12.5)	1	-	1
<i>Coreoperca herzi</i>	7	1 (14.3)	3	-	3
<i>Misgurnus anguillicaudatus</i>	6	2 (33.3)	2	-	1
Subtotal	319	206 (64.6)	178,007	1-31,250	864
2015					
<i>Pungtungia herzi</i>	41	41 (100)	53,753	24-7,750	1,311
<i>Squalidus japonicus coreanus</i>	15	15 (100)	21,164	359-2,670	1,411
<i>Squalidus gracilis majimae</i>	11	11 (100)	34,255	146-7,680	3,114
<i>Sarcocheilichthys variegatus</i>	3	3 (100)	1,632	157-1,318	544
<i>Pseudorasbora parva</i>	1	1 (100)	1,538	-	1,538
<i>Pseudogobio esocinus</i>	1	1 (100)	1,325	-	1,325
<i>Abbottina springeri</i>	4	4 (100)	155	10-80	39
<i>Microphysogobio koeensis</i>	1	1 (100)	139	-	139
<i>Acheilognathus koreensis</i>	29	29 (100)	2,454	4-678	85
<i>Acheilognathus yamatsutae</i>	16	4 (25.0)	38	1-28	10
<i>Acheilognathus rhombeus</i>	7	7 (100)	1,081	70-295	154
<i>Acheilognathus lanceolatus</i>	1	1 (100)	245	-	245
<i>Acanthorhodeus macropterus</i>	1	1 (100)	79	-	79
<i>Zacco platypus</i>	50	33 (66.0)	965	1-420	29
<i>Zacco temminckii</i>	49	4 (8.2)	7	1-4	2
<i>Opsariichthys uncirostris</i>	3	2 (66.7)	2	-	1
<i>Carassius auratus</i>	10	2 (20.0)	2	-	1
<i>Rhynchocypris oxycephalus</i>	1	1 (100)	28	-	28
Subtotal	244	161 (66.0)	118,862	1-7,750	738

(Continued to the next page)

Table 1. Continued

Year and fish species	No. of fish examined	No. (%) of fish infected	No. of metacercariae detected		
			Total	Range	Average
2016					
<i>Pungtungia herzi</i>	30	30 (100)	36,686	28-9,870	1,223
<i>Squalidus japonicus coreanus</i>	4	4 (100)	3,603	210-1,480	901
<i>Squalidus gracilis majimae</i>	2	2 (100)	3,323	653-2,670	1,662
<i>Pseudogobio esocinus</i>	2	2 (100)	540	36-504	270
<i>Microphysogobio koreensis</i>	1	1 (100)	40	-	40
<i>Hemibarbus longirostris</i>	1	1 (100)	13	-	13
<i>Acheilognathus koreensis</i>	35	34 (97.1)	871	2-287	26
<i>Acheilognathus rhombeus</i>	15	15 (100)	3,006	64-329	200
<i>Acheilognathus yamatsutae</i>	15	8 (53.3)	25	1-9	3
<i>Acheilognathus lanceolatus</i>	3	3 (100)	327	4-255	109
<i>Zacco platypus</i>	30	19 (63.3)	192	1-30	10
<i>Siniperca scherzeri</i>	1	1 (100)	6	-	6
Subtotal	139	120 (86.3)	48,632	1-9,870	405
2017					
<i>Pungtungia herzi</i>	26	26 (100)	26,736	22-8,950	1,028
<i>Squalidus japonicus coreanus</i>	31	31 (100)	13,161	4-2,750	425
<i>Sarcocheilichthys variegatus</i>	3	3 (100)	759	156-352	253
<i>Pseudorasbora parva</i>	7	7 (100)	5,260	365-2,225	751
<i>Acheilognathus yamatsutae</i>	27	15 (55.6)	45	1-2,030	7
<i>Acheilognathus koreensis</i>	26	19 (73.1)	129	1-30	5
<i>Acanthorhodeus gracilis</i>	5	5 (100)	26	2-12	1
<i>Acheilognathus rhombeus</i>	2	1 (50.0)	1	-	16
<i>Zacco platypus</i>	50	21 (42.0)	345	1-205	1
<i>Coreoperca herzi</i>	2	1 (50.0)	1	-	5
<i>Micropterus salmoides</i>	2	1 (50.0)	5	-	387
Subtotal	181	120 (66.3)	46,468	1-8,950	
Total	1,067	720 (67.5)	439,439	1-31,250	610

CsMc were calculated by the formula, prevalence/100 × mean metacercarial density per fish infected.

Subfamily groups in Cyprinidae fish hosts of *C. sinensis*

The gobioninid (Gobioninae) group-1 (323 fish) included *P. herzi* (169), *S. gracilis majimae* (43), *S. japonicus coreanus* (74), *S. variegatus wakiyae* (25), and *P. parva* (12). The gobioninid group-2 (24 fish) included *P. esocinus* (13), *A. springeri* (6), *M. koreensis* (3), *M. jeoni* (1), and *H. longirostris* (1). The acheilognathinid (Acheilognathinae) group (296 fish) included *A. koreensis* (130), *A. yamatsutae* (99), *A. rhombeus* (35), *A. majusculus* (15), *A. macropterus* (4), *A. lanceolatus* (6), and *A. gracilis* (7). The rasborinid (Rasborinae) group (403 fish) included *Z. platypus* (259), *Z. temminckii* (136), and *O. uncirostris amurensis* (8).

RESULTS

Infection status of CsMc in overall examined fish

CsMc were detected in 720 (62.0%) out of 1,162 fish in 32 species examined, and their average density was 610 per fish infected. The infection status by fish species and surveyed years

is shown in Table 1.

Infection status of CsMc in gobioninid fish group-1

CsMc were detected in all (100%) of 323 fish examined, and their average density was 1,310 per fish infected. The densities were highest in 2014 (1,958), followed by 2015 (1,582), 2016 (1,211), 2013 (972), 2017 (685), and 2011 (516) (Table 2). The infection status of CsMc by fish species, i.e., *P. herzi*, *S. gracilis majimae*, *S. japonicus coreanus*, *S. variegatus wakiyae* and *P. parva*, is revealed in Table 3.

Infection status of CsMc in gobioninid fish group-2

CsMc were detected in 23 (95.8%) out of 24 fish examined, and their average density was 127 per fish infected. The infection status by fish species, i.e., *P. esocinus*, *A. springeri*, *M. koreensis*, *M. jeoni* and *H. longirostris*, is shown in Table 4.

Infection status of CsMc in acheilognathinid fish group

CsMc were detected in 228 (77.0%) out of 296 fish examined, and their average density was 50 per fish infected. The densities were highest in 2015 (93), followed by 2016 (70),

Table 2. Infection status of *Clonorchis sinensis* metacercariae in susceptible gobionid fish from Wicheon in Gunwi-gun, Gyeongsangbuk-do

Year examined	No. of fish examined	No. (%) of fish infected	No. of metacercariae detected		
			Total	Range	Average
2011	25	25 (100)	12,895	9-2,420	516
2013	35	35 (100)	34,022	26-11,290	972
2014	89	89 (100)	174,228	6-31,250	1,958
2015	71	71 (100)	112,342	24-7,750	1,582
2016	36	36 (100)	43,612	28-9,870	1,211
2017	67	67 (100)	45,916	4-8,950	685
Total	323	323 (100)	423,015	4-31,250	1,310

Table 3. Infection status of *Clonorchis sinensis* metacercariae by fish species in the gobionid group-1

Species of fish	No. of fish examined	No. (%) of fish infected	No. of metacercariae detected		
			Total	Range	Average
<i>Pungtungia herzi</i>	169	169 (100)	261,986	6-31,250	1,550
<i>Squalidus japonicus coreanus</i>	74	74 (100)	64,037	4-8,460	865
<i>Squalidus gracilis majimae</i>	43	43 (100)	71,510	9-7,680	1,663
<i>Sarcocheilichthys variegatus</i>	25	25 (100)	15,773	13-2,730	631
<i>Pseudorasbora parva</i>	12	12 (100)	9,709	365-2,225	809
Total	323	323 (100)	423,015	4-31,250	1,310

Table 4. Infection status of *Clonorchis sinensis* metacercariae by fish species in the gobionid group-2

Species of fish	No. of fish examined	No. (%) of fish infected	No. of metacercariae detected		
			Total	Range	Average
<i>Pseudogobio esocinus</i>	13	12 (92.3)	2,101	1-1,325	175
<i>Abbottina springeri</i>	6	6 (100)	250	10-80	42
<i>Hemibarbus longirostris</i>	1	1 (100)	13	-	13
<i>Microphysogobio koreensis</i>	3	3 (100)	191	12-139	64
<i>Microphysogobio jeoni</i>	1	1 (100)	365	-	365
Total	24	23 (95.8)	2,920	1-1,325	127

Table 5. Infection status of *Clonorchis sinensis* metacercariae in acheilognathinid fish from Wicheon in Gunwi-gun, Gyeongsangbuk-do

Year examined	No. of fish examined	No. (%) of fish infected	No. of metacercariae detected		
			Total	Range	Average
2011	9	5 (55.6)	101	1-87	20
2013	26	16 (61.5)	81	1-14	5
2014	77	63 (81.8)	2,727	1-290	43
2015	54	42 (77.8)	3,897	1-678	93
2016	68	60 (88.2)	4,229	1-329	70
2017	60	40 (66.7)	201	1-30	5
Total	296	228 (77.0)	11,372	1-678	50

2014 (43), 2011 (20), 2013 (5), and 2017 (5) (Table 5). The infection status of CsMc by fish species, i.e., *A. koreensis*, *A. yamatsutae*, *A. rhombeus*, *A. majusculus*, *A. macroptera*, *A. lanceolatus* and *A. gracilis*, is revealed in Table 6.

Infection status of CsMc in rasborinid fish group

CsMc were detected in 147 (36.5%) out of 403 fish examined, and their average density was 15 per fish infected. The densities were highest in 2015 (25), followed by 2017 (16), 2013 (15), 2014 (11), 2016 (10), and 2011 (7) (Table 7). The

Table 6. Infection status of *Clonorchis sinensis* metacercariae by fish species in the acheilognathinid group

Species of fish	No. of fish examined	No. (%) of fish infected	No. of metacercariae detected		
			Total	Range	Average
<i>Acheilognathus koreensis</i>	130	122 (93.9)	4,620	1-678	38
<i>Acheilognathus yamatsutae</i>	99	55 (55.6)	691	1-136	13
<i>Acheilognathus rhombeus</i>	35	32 (91.4)	4,870	1-295	152
<i>Acheilognathus majusculus</i>	15	3 (20.0)	4	1-2	1.3
<i>Acanthorhodeus macropterus</i>	4	4 (100)	170	2-87	43
<i>Acheilognathus lanceolatus</i>	6	5 (83.3)	855	4-283	171
<i>Acanthorhodeus gracilis</i>	7	7 (100)	162	2-82	23
Total	296	228 (77.0)	11,372	1-678	50

Table 7. Infection status of *Clonorchis sinensis* metacercariae in rasborinid fish from Wicheon in Gunwi-gun, Gyeongsangbuk-do

Year examined	No. of fish examined	No. (%) of fish infected	No. of metacercariae detected		
			Total	Range	Average
2011	64	18 (28.1)	129	1-32	7
2013	35	9 (25.7)	131	1-70	15
2014	110	41 (37.3)	448	1-66	11
2015	102	39 (38.2)	974	1-420	25
2016	44	19 (43.2)	192	1-30	10
2017	62	21 (33.9)	345	1-205	16
Total	403	147 (36.5)	2,219	1-420	15

Table 8. Infection status of *Clonorchis sinensis* metacercariae by fish species in the rasborinid group

Species of fish	No. of fish examined	No. (%) of fish infected	No. of metacercariae detected		
			Total	Range	Average
<i>Zacco platypus</i>	259	139 (53.7)	2,207	1-420	16
<i>Zacco temminckii</i>	136	6 (4.4)	10	1-4	2
<i>Opsariichthys uncirostris</i>	8	2 (25.0)	2	-	1
Total	403	147 (36.5)	2,219	1-420	15

Table 9. Susceptibility index of *Clonorchis sinensis* metacercariae by surveyed year and fish groups

Group	Susceptibility index ^a						
	2011	2013	2014	2015	2016	2017	Total
Overall positive fish group	153	351	558	487	350	257	412
Gobioninid group-1	516	972	1,958	1,582	1,211	685	1,310
Acheilognathinid group	11.2	3.1	35.2	72.4	61.7	3.3	38.5
Rasborinid group	2	3.9	4.1	9.6	4.3	5.4	5.5

^aPrevalence/100 × mean metacercarial density per fish infected.

infection status of CsMc by fish species, i.e., *Z. platypus*, *Z. temminckii* and *O. uncirostris amurensis*, is shown in Table 8.

Susceptibility index of CsMc by fish groups

The susceptibility indices of CsMc were 412 in the overall positive fish group; 1,310 in the gobioninid group-1, 122 in the gobioninid group-2, 38.5 in the acheilognathinid, and 5.5

in the rasborinid fish group (Table 9).

DISCUSSION

By the present study, it was reconfirmed that CsMc are highly prevalent in fish from Wicheon, Gyeongsangbuk-do, Korea. Moreover, the infection status showed a certain tendency ac-

cording to the subfamily groups of fish hosts, i.e., Gobioninae, Acheilognathinae, and Rasborinae, among the Cyprinidae. The prevalences were 100%, 79.7%, and 35.5%, and metacercarial densities were 1,310, 50, and 15 per fish infected in 3 fish groups, respectively. These findings also suggested that the endemicity of CsMc is closely related to the fish group, and this host-parasite relationship may originate from a long-lasting evolutionary process.

The fish sampling is one of the important factors in metacercarial surveys for the epidemiology of fishborne zoonotic trematode infections. Total 1,162 fish in 32 species were collected through 6 years (2011 and 2013-2017) in the same site of Woobo-myeon in Gunwi-gun. Among them, the pale chub (259 *Z. platypus*: 22.2%) was the most dominant fish species like Sohn et al. [18] in Seomjin-gang, followed by the striped shinner (169 *P. herzi*: 14.5%), dark chub and/or Korean chub (136 *Z. temminckii* and/or *Z. koreanus*: 11.7%), oily bitterling (130 *A. koreensis*: 11.2%), Korean striped bitterling (99 *A. yamatsutae*: 8.5%), and short barbell gudgeon (74 *S. japonicus coreanus*: 6.4%). The number of fish examined was 867 (74.4%) in major 6 species and 298 (25.6%) in the remaining 26 species. The disproportion of the fish number examined was dependent upon the ecological conditions of each fish species, and moreover, we collected all fish only by the casting net in the daytime. If we used together with other methods for fish catching like a netting, we could have collected more various species of fish, including nocturnal ones.

Total 49 fish species (in 34 genera 7 families) have been reported as the second intermediate hosts of *C. sinensis* in Korea [9-18]. In the present study, CsMc were found in 26 fish species, i.e., *Abbottina springeri*, *Acanthorhodeus macropterus*, *A. gracilis*, *Acheilognathus koreensis*, *A. lanceolatus*, *A. majusculus*, *A. rhombeus*, *A. yamatsutae*, *Carassius auratus*, *Coreoperca herzi*, *Hemibarbus longirostris*, *Micropterus salmoides*, *Microphysogobio koeensis*, *M. jeoni*, *Misgurnus anguillicaudatus*, *Opsariichthys uncirostris*, *Pseudogobio esocinus*, *Pungtungia herzi*, *Pseudorasbora parva*, *Rhynchocypris oxycephalus*, *Sarcocheilichthys variegates wakiyae*, *Siniperca scherzeri*, *Squalidus gracilis majimae*, *S. japonicus coreanus*, *Zacco platypus*, and *Z. temminckii*. Among them, 2 fish species, i.e., *M. salmoides* (Centrarchidae) and *M. anguillicaudatus* (Cobitidae), are newly added in the list of the second intermediate hosts of *C. sinensis* in Korea [9-18]. Accordingly, total 51 fish species in 9 families are listed as the second intermediate hosts of *C. sinensis* in Korea.

The large mouth bass, *Micropterus salmoides*, was imported

from USA as an edible fish species in 1973. However, this fish is widely spread in the water systems of the whole country, and then became notorious as an agitating fish species of the ecosystem in Korea [20]. In this study, we examined total 10 *M. salmoides* and detected 1 and 5 CsMc from 2 fish. Although the number of fish examined and metacercariae detected were not so many, the fact, infection with CsMc in foreign fish species, is highly important. Thus, we should pay attention to the fish species from foreign countries in the metacercarial survey for the fishborne zoonotic trematodes.

Fish species, i.e., Mandarin fish (*S. scherzeri*), Korean aucha perch (*C. herzi*), common carp (*Cyprinus carpio*), and crucian carp (*C. auratus*), edible in raw conditions practically act as the infection source of clonorchiasis in Korea. Fortunately, these fish species are less prevalent with CsMc. Total 3 (8.8%) out of 34 *C. auratus* were infected with total 3 CsMc, 2 (11.1%) *C. herzi* was infected with a total of 4 CsMc, and only 1 (33.3%) *S. scherzeri* was infected with 6 CsMc, respectively, even in such a highly endemic area, Wicheon. In a survey of the fish from Seomjin-gang, no CsMc was detected in 4 fish species, i.e., *C. herzi* (57), *C. auratus* (42), *S. scherzeri* (11), and *C. carpio* (2), edible in raw conditions [18].

The striped shinner, *P. herzi*, is appropriate to be an index fish species to check the endemicity of *C. sinensis* infection. This fish species broadly lives in the water systems of rivers in Korea and is highly susceptible to CsMc. In the present study, all of 169 *P. herzi* examined were infected with an average of 1,550 CsMc per fish. The prevalences of CsMc in this fish species were 80.4%, 72.5%, and 30.6%, and metacercarial densities were 15.2, 46.1, and 175.9 per fish infected in the upper, middle, and lower reaches of Seomjin-gang, respectively [18]. The positive rates of CsMc were 6.2%, 31.8%, and 92.9%, and densities were 2.8, 214, and 409 per fish infected in *P. herzi* from the northern, middle, and southern regions of the Korean peninsula, respectively [14]. Especially, in this fish species from the streams of Nakdong-gang, i.e., Wicheon in Gunwi-gun, Banbyuncheon in Yeongyang-gun, Gyeongsangbuk-do, and Yangcheon in Sancheong-gun, Gyeongsangnam-do, the endemicities of CsMc were much higher like in this study [14].

Conclusively, it has been confirmed that the endemicity of CsMc is much higher in fish from Wicheon, and a strong tendency was shown in the prevalence and intensity of CsMc according to the subfamily groups of fish, i.e., Gobioninae, Acheilognathinae, and Rasborinae, in the family Cyprinidae fish hosts. Among the highly susceptible gobioninid fish, *P.*

herzi is recommended as a useful index fish to check the endemicity of clonorchiasis in a certain area of Korea.

ACKNOWLEDGMENTS

This study was supported by an anti-communicable diseases control program, 2014-E54002-00 (Investigation of fish-borne parasites and acquisition of their biological resources in the southern regions of Korea) and 2015-E54001-00 (Investigation of fish-borne parasites and acquisition of their biological resources in the eastern regions of Korea) of National Institute of Health (NIH), Korea Centers for Disease Control and Prevention (KCDCP), Korea. We thank Jung-A Kim and Hee-Joo Kim (Department of Parasitology and Tropical Medicine, Gyeongsang National University College of Medicine, Jinju, Korea), for their help in the examination of fish.

CONFLICT OF INTEREST

The authors have no conflicts of interest concerning the work reported in this paper.

REFERENCES

1. Korea Centers for Disease Control and Prevention; Korea National Institute of Health. National survey of the prevalence of intestinal parasitic infections in Korea, 2012. The 8th Report. Osong, Chungcheongbuk-do, Korea. 2013.
2. Seo BS, Lee SH, Cho SY, Chai JY, Hong ST, Han IS, Sohn JS, Cho BH, Ahn SR, Lee SK, Chung SC, Kang KS, Shim HS, Hwang IS. An epidemiologic study on clonorchiasis and metagonimiasis in riverside areas in Korea. *Korean J Parasitol* 1981; 19: 137-150.
3. Rim HJ. The current pathobiology and chemotherapy of clonorchiasis. *Korean J Parasitol* 1986; 24 (suppl): 1-141.
4. Hong ST, Hong SJ. *Clonorchis sinensis* and clonorchiasis in Korea. Food-Borne Helminthiasis in Asia. *Asian Parasitology* 2005; 1: 35-56.
5. Cho SH, Lee KY, Lee BC, Cho PY, Cheun HI, Hong ST, Sohn WM, Kim TS. Prevalence of clonorchiasis in southern endemic areas of Korea in 2006. *Korean J Parasitol* 2008; 46: 133-137.
6. Kim TS, Cho SH, Huh S, Kong Y, Sohn WM, Hwang SS, Chai JY, Lee SH, Park YK, Oh DK, Lee JK. A nationwide survey on the prevalence of intestinal parasitic infections in the Republic of Korea, 2004. *Korean J Parasitol* 2009; 47: 37-47.
7. Kim HK, Cheun HI, Chung BS, Lee KY, Kim TS, Lee SE, Lee WJ, Cho SH. Prevalence of *Clonorchis sinensis* infections along the five major rivers in Republic of Korea, 2007. *Osong Public Health Res Perspect* 2010; 1: 43-49.
8. Jeong YI, Shin HE, Lee SE, Cheun HI, Ju JW, Kim JY, Park MY, Cho SH. Prevalence of *Clonorchis sinensis* infection among residents along 5 major rivers in the Republic of Korea. *Korean J Parasitol* 2016; 54: 215-219.
9. Kong HH, Choi BR, Moon HC, Choi DW. Larval digenetic trematodes from freshwater fish in river Miryang, Korea. *Jpn J Parasitol* 1995; 44: 112-118.
10. Sohn WM, Choi YS. Infection status with trematode metacercariae in the freshwater fish from Chunamchosuchi (pond), Uichang-gun, Kyongsangnam-do, Korea. *Korean J Parasitol* 1997; 35: 165-170 (in Korean).
11. Joo JY, Chung MS, Kim SJ, Kang CM. Changing patterns of *Clonorchis sinensis* infections in Kyongbuk, Korea. *Korean J Parasitol* 1997; 35: 155-164.
12. Kim EM, Kim JL, Choi SY, Kim JW, Kim S, Choi MH, Bae YM, Lee SH, Hong ST. Infection status of freshwater fish with metacercariae of *Clonorchis sinensis* in Korea. *Korean J Parasitol* 2008; 46: 247-251.
13. Sohn WM. Fish-borne zoonotic trematode metacercariae in the Republic of Korea. *Korean J Parasitol* 2009; 47 (suppl): 103-113.
14. Cho SH, Sohn WM, Na BK, Kim TS, Kong Y, Eom K, Seok WS, Lee T. Prevalence of *Clonorchis sinensis* metacercariae in freshwater fish from three latitudinal regions of the Korean Peninsula. *Korean J Parasitol* 2011; 49: 385-398.
15. Sohn WM. Invertebrate Founa of Korea Vol. 6, No. 1. Trematodes. Incheon, Korea: The National Institute of Biological Resources; 2013. pp 1-125.
16. Cho SH, Lee WJ, Kim TS, Seok WS, Lee TJ, Jeong KJ, Na BK, Sohn WM. Prevalence of zoonotic trematode metacercariae in freshwater fish from Gangwon-do, Korea. *Korean J Parasitol* 2014; 52: 399-412.
17. Sohn WM, Na BK, Cho SH, Lee SW, Choi SB, Seok WS. Trematode metacercariae in freshwater fish from water systems of Hantangang and Imjingang in Republic of Korea. *Korean J Parasitol* 2015; 53: 289-298.
18. Sohn WM, Na BK, Cho SH, Park MY, Kim CH, Hwang MA, No KW, Yoon KB, Lim HC. Prevalence of *Clonorchis sinensis* metacercariae in fish from water systems of Seomjin-gang (River). *Korean J Parasitol* 2017; 55: 305-312.
19. Wicheon in Gyeongbuk Province (South Korea) in Wikipedia - The free encyclopedia: <http://en.wikipedia.org>.
20. Kim IS, Kang EJ. Colored Fish of Korea. Seoul, Korea. Academy Publishing Company. 1993, pp 1-477 (in Korean).