

APPLICATIONS OF THE COUPLED FIXED POINT THEOREM TO THE NONLINEAR MATRIX EQUATIONS

SEJONG KIM AND HOSOO LEE

ABSTRACT. In this article we consider certain types of nonlinear matrix equations including the stochastic rational Riccati equation and show the existence and uniqueness of the positive definite solution by using Bhaskar-Lakshmikantham's coupled fixed point theorem.

1. Introduction

The fixed point theory has been studied widely in nonlinear analysis, and its results have been developed in a variety of areas in mathematics and engineering. In recent years, there has been a lot of interest in establishing fixed point theorems on ordered metric spaces with a contractive condition which holds for all points that are related by partial ordering. This trend was initiated by Ran and Reurings in [15] where they extended the Banach contraction principle in partially ordered sets with some applications to matrix equations.

Bhaskar and Lakshmikantham [3] introduced the notions of a mixed monotone mapping and a coupled fixed point, and proved some coupled fixed point theorems for mixed mappings in ordered metric spaces. Afterwards, Lakshmikantham and Ćirić [10] established coupled coincidence and coupled fixed point theorems. Many different kinds of coupled fixed point theorems with applications have been developed; see the literatures [2, 7, 10, 13]. In this article we mainly focus on solving the certain nonlinear matrix equations using the coupled fixed point theorem.

In [1] Berzig, Duan, and Samet have studied the positive definite solution to the nonlinear matrix equation

$$(1.1) \quad X = Q - A^* X^{-1} A + B^* X^{-1} B,$$

where Q is an $n \times n$ positive definite Hermitian matrix, A, B are arbitrary $n \times n$ matrices. This is a special stochastic rational Riccati equation arisen in stochastic control theory. Also, the special cases of (1.1) such as $X = Q - A^* X^{-1} A$ and $X = Q + B^* X^{-1} B$ have been studied (see [8, 14]).

Received December 22, 2016; Revised May 23, 2017; Accepted July 17, 2017.

2010 *Mathematics Subject Classification.* 47H10, 15B48.

Key words and phrases. mixed monotone property, coupled fixed point theorem.

Furthermore, many researchers have considered the following nonlinear matrix equation generalizing the equations

$$(1.2) \quad X - \sum_{i=1}^m A_i^* X^{\delta_i} A_i = Q,$$

where $0 < |\delta_i| < 1$ and A_i 's are arbitrary $n \times n$ matrices for all $i = 1, \dots, m$, and suggested numerical methods for finding a solution [5, 6, 12]. Huang, Huang, Tsai [9] and Duan, Liao, Tang [5] have shown the existence and uniqueness of the positive definite solution of the equation (1.2) by using Hilbert's projective metric and fixed point theorems for monotone and mixed monotone operators, respectively. Moreover, Lim [11] has shown it by proving that the map $F(X) = Q + \sum_{i=1}^m A_i^* X^{\delta_i} A_i$ is a strict contraction for the Thompson metric with the contraction coefficient less than or equal to $\delta := \max\{|\delta_i|\}_{i=1}^m$.

In this article, we investigate the matrix equation generalized the equations (1.1) and (1.2)

$$(1.3) \quad X = Q \pm \sum_{i=1}^m A_i X^{\pm p_i} A_i^* \mp \sum_{j=1}^n B_j X^{\pm q_j} B_j^*,$$

where $p_i, q_j \in (0, 1]$, and A_i 's and B_j 's are arbitrary $n \times n$ matrices for all $i = 1, \dots, m$ and $j = 1, \dots, n$. We show the existence and uniqueness of the positive definite solution to the nonlinear matrix equation (1.3) using Bhaskar-Lakshmikantham's coupled fixed point theorem. We also discuss some simple cases of the equation (1.3) including two equations (1.1) and (1.2).

2. Preliminaries

Throughout this paper, we denote by $\mathcal{H}(n)$ the set of all $n \times n$ Hermitian matrices. For $A, B \in \mathcal{H}(n)$, $A \leq B$ ($A < B$) means that $B - A$ is positive semidefinite (positive definite, respectively). Moreover, $X \in [A, B]$ means that $A \leq X \leq B$, and $X \in [A, \infty)$ means that $X \geq A$. We denote by $\|A\|$ and $\|A\|_{\text{tr}}$ the spectral norm and trace norm, respectively, that is,

$$\|A\| = \max\{\sigma_j(A) : j = 1, \dots, n\},$$

$$\|A\|_{\text{tr}} = \sum_{j=1}^n \sigma_j(A),$$

where $\sigma_j(A)$, $j = 1, \dots, n$ are the singular values of A .

The following lemmas will be useful later.

Lemma 2.1. *Let $A, B \in \mathcal{H}(n)$ such that $A \geq O$ and $B \geq O$. Then*

$$0 \leq \text{tr}(AB) \leq \|A\| \text{tr}(B).$$

Lemma 2.2 ([4]). *If $A, B > O$ with $A \leq B$, then $A^t \leq B^t$ for any $0 \leq t \leq 1$ and $A^{-1} \geq B^{-1}$.*

Lemma 2.3 ([4]). *Let P and Q be positive definite matrices of the same order with $P, Q \geq aI$, where $a > 0$. Then for every unitarily invariant norm $\|\cdot\|$ and $0 < t \leq 1$*

$$\begin{aligned} \|P^t - Q^t\| &\leq ta^{t-1}\|P - Q\|, \\ \|P^{-t} - Q^{-t}\| &\leq ta^{-(t+1)}\|P - Q\|. \end{aligned}$$

Let (X, \preceq) be a partially ordered set and $F : X \times X \rightarrow X$ be a given mapping. We say that F has the *mixed monotone property* if for any $x, y \in X$

$$\begin{aligned} x_1, x_2 \in X, \quad x_1 \preceq x_2 &\Rightarrow F(x_1, y) \preceq F(x_2, y), \\ y_1, y_2 \in X, \quad y_1 \preceq y_2 &\Rightarrow F(x, y_1) \succeq F(x, y_2). \end{aligned}$$

We say that (x, y) is a *coupled fixed point* of F if $x = F(x, y)$ and $y = F(y, x)$.

The following fixed point theorem is the key of the main result.

Theorem 2.4 ([3]). *Let (X, \preceq) be a partially ordered set endowed with a metric d such that (X, d) is complete. Let $F : X \times X \rightarrow X$ be a continuous mapping having the mixed monotone property on X . Assume that there exists a $\delta \in [0, 1)$ such that*

$$d(F(x, y), F(u, v)) \leq \frac{\delta}{2}[d(x, u) + d(y, v)]$$

for any $(x, y), (u, v) \in X \times X$ with $x \succeq u$ and $y \preceq v$. We suppose that there exist $x_0, y_0 \in X$ such that $x_0 \preceq F(x_0, y_0)$ and $y_0 \succeq F(y_0, x_0)$. Then

- (a) F has a coupled fixed point $(\bar{x}, \bar{y}) \in X \times X$; and
- (b) the sequences $\{x_n\}$ and $\{y_n\}$ defined by $x_{n+1} = F(x_n, y_n)$ and $y_{n+1} = F(y_n, x_n)$ converge to \bar{x} and \bar{y} , respectively.

In addition, suppose that every pair of elements has a lower bound and an upper bound, then

- (c) F has a coupled fixed point $(\bar{x}, \bar{y}) \in X \times X$;
- (d) $\bar{x} = \bar{y}$; and
- (e) we have the following estimate

$$\max\{d(x_n, \bar{x}), d(y_n, \bar{x})\} \leq \frac{\delta^n}{2(1-\delta)}[d(F(x_0, y_0), x_0) + d(F(y_0, x_0), y_0)].$$

3. Solving $X = Q + \sum_{i=1}^m A_i X^{p_i} A_i^* - \sum_{j=1}^n B_j X^{q_j} B_j^*$

In this section, we consider the following matrix equation

$$(3.4) \quad X = Q + \sum_{i=1}^m A_i X^{p_i} A_i^* - \sum_{j=1}^n B_j X^{q_j} B_j^*,$$

where $p_i, q_j \in (0, 1]$.

For $a, b > 0$ the following assumptions are considered:

1. $aI + b_M \sum_{j=1}^n B_j B_j^* \leq Q \leq bI - b_M \sum_{i=1}^m A_i A_i^*$,
2. $\sum_{i=1}^m \|A_i A_i^*\| < \frac{a}{2a_M}$, $\sum_{j=1}^n \|B_j B_j^*\| < \frac{a}{2a_M}$,

where

$$a_M := \max\{a^{p_i}, a^{q_j} : 1 \leq i \leq m, 1 \leq j \leq n\},$$

$$b_M := \max\{b^{p_i}, b^{q_j} : 1 \leq i \leq m, 1 \leq j \leq n\}.$$

Theorem 3.1. *Under the assumptions 1 and 2, we have*

- (I) *Equation (3.4) has a unique solution $\bar{X} \in [aI, bI]$, and*
- (II) *the sequences $\{X_k\}$ and $\{Y_k\}$ defined by*

$$\begin{cases} X_0 = aI, \\ X_{k+1} = Q + \sum_{i=1}^m A_i X_k^{p_i} A_i^* - \sum_{j=1}^n B_j Y_k^{q_j} B_j^*, \\ Y_0 = bI, \\ Y_{k+1} = Q + \sum_{i=1}^m A_i Y_k^{p_i} A_i^* - \sum_{j=1}^n B_j X_k^{q_j} B_j^*, \end{cases}$$

converge to \bar{X} , that is,

$$\lim_{k \rightarrow \infty} \|X_k - \bar{X}\|_{\text{tr}} = \lim_{k \rightarrow \infty} \|Y_k - \bar{X}\|_{\text{tr}} = 0.$$

Furthermore, the error estimation is given by

$$\max\{\|X_k - \bar{X}\|_{\text{tr}}, \|Y_k - \bar{X}\|_{\text{tr}}\} \leq \frac{\delta^k}{1 - \delta} \max\{\|X_1 - X_0\|_{\text{tr}}, \|Y_1 - Y_0\|_{\text{tr}}\},$$

where $0 < \delta < 1$.

Proof. For $X, Y \in \mathcal{H}(n)$ let

$$F(X, Y) = Q + \sum_{i=1}^m A_i X^{p_i} A_i^* - \sum_{j=1}^n B_j Y^{q_j} B_j^*.$$

By Lemma 2.2 F is a continuous mapping having the mixed monotone property.

We claim that $F([aI, bI]^2) \subseteq [aI, bI]$. Let $X, Y \in [aI, bI]$, that is, $aI \leq X, Y \leq bI$. This implies by assumption 1 that

$$F(X, Y) \leq Q + \sum_{i=1}^m A_i X^{p_i} A_i^* \leq Q + \sum_{i=1}^m b^{p_i} A_i A_i^* \leq Q + b_M \sum_{i=1}^m A_i A_i^* \leq bI,$$

and

$$F(X, Y) \geq Q - \sum_{j=1}^n B_j Y^{q_j} B_j^* \geq Q - \sum_{j=1}^n b^{q_j} B_j B_j^* \geq Q - b_M \sum_{j=1}^n B_j B_j^* \geq aI.$$

Let $X, Y, U, V \in [aI, bI]$ such that $X \geq U$ and $Y \leq V$. Then

$$\begin{aligned}
 & \|F(X, Y) - F(U, V)\|_{\text{tr}} \\
 &= \left\| \sum_{i=1}^m A_i(X^{p_i} - U^{p_i})A_i^* + \sum_{j=1}^n B_j(V^{q_j} - Y^{q_j})B_j^* \right\|_{\text{tr}} \\
 &\leq \sum_{i=1}^m \|A_i(X^{p_i} - U^{p_i})A_i^*\|_{\text{tr}} + \sum_{j=1}^n \|B_j(V^{q_j} - Y^{q_j})B_j^*\|_{\text{tr}} \\
 &\leq \sum_{i=1}^m \|A_i A_i^*\| \|X^{p_i} - U^{p_i}\|_{\text{tr}} + \sum_{j=1}^n \|B_j B_j^*\| \|V^{q_j} - Y^{q_j}\|_{\text{tr}} \\
 &\leq \left(\sum_{i=1}^m p_i a^{p_i-1} \|A_i A_i^*\| \right) \|X - U\|_{\text{tr}} + \left(\sum_{j=1}^n q_j a^{q_j-1} \|B_j B_j^*\| \right) \|V - Y\|_{\text{tr}} \\
 &\leq \frac{a_M}{a} \left[\left(\sum_{i=1}^m \|A_i A_i^*\| \right) \|X - U\|_{\text{tr}} + \left(\sum_{j=1}^n \|B_j B_j^*\| \right) \|V - Y\|_{\text{tr}} \right].
 \end{aligned}$$

The first inequality follows from the triangle inequality, the second from Lemma 2.1, the third from Lemma 2.3, and the last from the following inequality: for any $t \in \{p_i, q_j\}$

$$0 < ta^{t-1} \leq \frac{a^t}{a} \leq \frac{a_M}{a}.$$

This implies that

$$\|F(X, Y) - F(U, V)\|_{\text{tr}} \leq \frac{\delta}{2} [\|X - U\|_{\text{tr}} + \|V - Y\|_{\text{tr}}],$$

where

$$\delta = \frac{2a_M}{a} \max \left\{ \sum_{i=1}^m \|A_i A_i^*\|, \sum_{j=1}^n \|B_j B_j^*\| \right\}.$$

From condition 2 we can easily see that $0 \leq \delta < 1$.

Taking $X_0 = aI$ and $Y_0 = bI$ we can show from condition 1 that $X_0 \leq F(X_0, Y_0)$ and $Y_0 \geq F(Y_0, X_0)$. On the other hand, for every $X, Y \in \mathcal{H}(n)$ there exist a greatest lower bound and a least upper bound. Thus, (I) and (II) follow immediately from Theorem 2.4, and \bar{X} is the unique solution to Equation (3.4) in $[aI, bI]$. \square

The following results are immediate consequences of Theorem 3.1.

Corollary 3.2. Consider the matrix equation (3.4) with $Q = I$. Suppose that

$$\sum_{i=1}^m \|A_i A_i^*\| \leq \min \left\{ \frac{b-1}{b_M}, \frac{a}{2a_M} \right\} \quad \text{and} \quad \sum_{j=1}^n \|B_j B_j^*\| \leq \min \left\{ \frac{1-a}{b_M}, \frac{a}{2a_M} \right\}.$$

Then items (I) and (II) of Theorem 3.1 hold.

Corollary 3.3. Consider the matrix equation (3.4) with unitary matrices A_i and B_j for all i and j . Suppose that

1. $(a + nb_M)I \leq Q \leq (b - mb_M)I$, and
2. $a_M < \min\left\{\frac{a}{2m}, \frac{a}{2n}\right\}$.

Then items (I) and (II) of Theorem 3.1 hold.

The following is the case when $p_i = q_j = t \in (0, 1]$ for all i, j , and $A_i = B_j = O$ for all i, j except one term.

Corollary 3.4. Consider the matrix equation

$$(3.5) \quad X = Q + AX^t A^* - BX^t B^*,$$

where $t \in (0, 1]$. Suppose that

1. $aI + b^t BB^* \leq Q \leq bI - b^t AA^*$, and
2. $\|AA^*\| < \frac{a^{1-t}}{2}$, $\|BB^*\| < \frac{a^{1-t}}{2}$.

Then items (I) and (II) of Theorem 3.1 hold.

$$4. \text{ Solving } X = Q - \sum_{i=1}^m A_i X^{-p_i} A_i^* + \sum_{j=1}^n B_j X^{-q_j} B_j^*$$

In this section, we consider the following matrix equation

$$(4.6) \quad X = Q - \sum_{i=1}^m A_i X^{-p_i} A_i^* + \sum_{j=1}^n B_j X^{-q_j} B_j^*,$$

where $p_i, q_j \in (0, 1]$.

For $a, b > 0$ the following assumptions are considered:

1. $aI + \frac{1}{a_m} \sum_{i=1}^m A_i A_i^* \leq Q \leq bI - \frac{1}{a_m} \sum_{j=1}^n B_j B_j^*$,
2. $\sum_{i=1}^m \|A_i A_i^*\| < \frac{a_m a}{2}$, $\sum_{j=1}^n \|B_j B_j^*\| < \frac{a_m a}{2}$,

where

$$a_m := \min\{a^{p_i}, a^{q_j} : 1 \leq i \leq m, 1 \leq j \leq n\}.$$

Theorem 4.1. Under the assumptions 1 and 2,

- (I) Equation (4.6) has a unique solution $\hat{X} \in [aI, \infty)$, and
- (II) the sequences $\{X_k\}$ and $\{Y_k\}$ defined by

$$\begin{cases} X_0 = aI, \\ X_{k+1} = Q - \sum_{i=1}^m A_i X_k^{-p_i} A_i^* + \sum_{j=1}^n B_j Y_k^{-q_j} B_j^*, \end{cases}$$

$$\begin{cases} Y_0 = bI, \\ Y_{k+1} = Q - \sum_{i=1}^m A_i Y_k^{-p_i} A_i^* + \sum_{j=1}^n B_j X_k^{-q_j} B_j^*, \end{cases}$$

converge to \hat{X} , that is,

$$\lim_{k \rightarrow \infty} \|X_k - \hat{X}\|_{\text{tr}} = \lim_{k \rightarrow \infty} \|Y_k - \hat{X}\|_{\text{tr}} = 0.$$

Furthermore, the error estimation is given by

$$\max\{\|X_k - \hat{X}\|_{\text{tr}}, \|Y_k - \hat{X}\|_{\text{tr}}\} \leq \frac{\delta^k}{1 - \delta} \max\{\|X_1 - X_0\|_{\text{tr}}, \|Y_1 - Y_0\|_{\text{tr}}\},$$

where $0 < \delta < 1$.

Proof. For $X, Y \in \mathcal{H}(n)$ let

$$G(X, Y) = Q - \sum_{i=1}^m A_i X^{-p_i} A_i^* + \sum_{j=1}^n B_j Y^{-q_j} B_j^*.$$

By Lemma 2.2, G is a continuous mapping having the mixed monotone property.

We claim that $G([aI, \infty)^2) \subseteq [aI, \infty)$. Let $X, Y \in [aI, \infty)$, that is, $X, Y \geq aI$. This implies by assumption 1 that

$$G(X, Y) \geq Q - \sum_{i=1}^m A_i X^{-p_i} A_i^* \geq Q - \sum_{i=1}^m a^{-p_i} A_i A_i^* \geq Q - a_m^{-1} \sum_{i=1}^m A_i A_i^* \geq aI.$$

Let $X, Y, U, V \in [aI, \infty)$ such that $X \geq U$ and $Y \leq V$. Then

$$\begin{aligned} & \|G(X, Y) - G(U, V)\|_{\text{tr}} \\ &= \left\| \sum_{i=1}^m A_i (U^{-p_i} - X^{-p_i}) A_i^* + \sum_{j=1}^n B_j (Y^{-q_j} - V^{-q_j}) B_j^* \right\|_{\text{tr}} \\ &\leq \sum_{i=1}^m \|A_i (U^{-p_i} - X^{-p_i}) A_i^*\|_{\text{tr}} + \sum_{j=1}^n \|B_j (Y^{-q_j} - V^{-q_j}) B_j^*\|_{\text{tr}} \\ &\leq \sum_{i=1}^m \|A_i A_i^*\| \|U^{-p_i} - X^{-p_i}\|_{\text{tr}} + \sum_{j=1}^n \|B_j B_j^*\| \|Y^{-q_j} - V^{-q_j}\|_{\text{tr}} \\ &\leq \left(\sum_{i=1}^m p_i a^{-p_i-1} \|A_i A_i^*\| \right) \|U - X\|_{\text{tr}} + \left(\sum_{j=1}^n q_j a^{-q_j-1} \|B_j B_j^*\| \right) \|Y - V\|_{\text{tr}} \\ &\leq \frac{1}{a_m a} \left[\left(\sum_{i=1}^m \|A_i A_i^*\| \right) \|U - X\|_{\text{tr}} + \left(\sum_{j=1}^n \|B_j B_j^*\| \right) \|Y - V\|_{\text{tr}} \right]. \end{aligned}$$

The last follows from the following inequality: for any $t \in \{p_i, q_j\}$

$$0 < ta^{-t-1} \leq \frac{1}{a^t a} \leq \frac{1}{a_m a}.$$

This implies that

$$\|G(X, Y) - G(U, V)\|_{\text{tr}} \leq \frac{\delta}{2} [\|X - U\|_{\text{tr}} + \|V - Y\|_{\text{tr}}],$$

where

$$\delta = \frac{2}{a_m a} \max \left\{ \sum_{i=1}^m \|A_i A_i^*\|, \sum_{j=1}^n \|B_j B_j^*\| \right\}.$$

From condition 2 we can easily see that $0 \leq \delta < 1$.

Taking $X_0 = aI$ and $Y_0 = bI$ we can show from condition 1 that $X_0 \leq G(X_0, Y_0)$ and $Y_0 \geq G(Y_0, X_0)$. On the other hand, for every $X, Y \in \mathcal{H}(n)$ there exist a greatest lower bound and a least upper bound. Thus, (I) and (II) follow immediately from Theorem 2.4, and \hat{X} is the unique solution to Equation (4.6) in $[aI, \infty)$. \square

The following results are immediate consequences of Theorem 4.1.

Corollary 4.2. *Consider the matrix equation (4.6) with $Q = I$. Suppose that*

1. $\sum_{i=1}^m A_i A_i^* \leq a_m(1-a)I$, $\sum_{j=1}^n B_j B_j^* \leq a_m(b-1)I$, and
2. $\sum_{i=1}^m \|A_i A_i^*\| < \frac{a_m a}{2}$, $\sum_{j=1}^n \|B_j B_j^*\| < \frac{a_m a}{2}$.

Then items (I) and (II) of Theorem 4.1 hold.

Corollary 4.3. *Consider the matrix equation (4.6) with unitary matrices A_i and B_j for all i and j . Suppose that*

1. $\left(a + \frac{m}{a_m}\right)I \leq Q \leq \left(b - \frac{n}{a_m}\right)I$, and
2. $a_m > \max\left\{\frac{2m}{a}, \frac{2n}{a}\right\}$.

Then items (I) and (II) of Theorem 4.1 hold.

The following is the case when $p_i = q_j = t \in (0, 1]$ for all i, j , and $A_i = B_j = O$ for all i, j except one term.

Corollary 4.4. *Consider the matrix equation*

$$(4.7) \quad X = Q - AX^{-t}A^* + BX^{-t}B^*,$$

where $t \in (0, 1]$. Suppose that

1. $aI + a^{-t}AA^* \leq Q \leq bI - a^{-t}BB^*$, and
2. $\|AA^*\| < \frac{a^{1+t}}{2}$, $\|BB^*\| < \frac{a^{1+t}}{2}$.

Then items (I) and (II) of Theorem 4.1 hold.

Acknowledgement. The work of S. Kim was supported by Basic Science Research Program through the National Research Foundation of Korea funded by the Ministry of Science, ICT and Future Planning (NRF-2015R1C1A1A020364 07). The work of H. Lee was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF-2015R1D1A1A0105 9900) funded by the Ministry of Education.

References

- [1] M. Berzig, X. Duan, and B. Samet, *Positive definite solution of the matrix equation $X = Q - A^*X^{-1}A + B^*X^{-1}B$ via Bhaskar-Lakshmikantham fixed point theorem*, Math. Sci. **6** (2012), Art. 27, 6 pp.
- [2] M. Berzig and B. Samet, *An extension of coupled fixed point's concept in higher dimension and applications*, Comput. Math. Appl. **63** (2012), no. 8, 1319–1334.
- [3] T. Bhaskar and V. Lakshmikantham, *Fixed point theorems in partially ordered metric spaces and applications*, Nonlinear Anal. **65** (2006), no. 7, 1379–1393.
- [4] R. Bhatia, *Matrix Analysis*, Graduate Texts in Mathematics, Vol. 169, Springer-Verlag, New York, 1997.
- [5] X. Duan, A. Liao, and B. Tang, *On the nonlinear matrix equation $X - \sum_{i=1}^m A_i^* X^{\delta_i} A_i = Q$* , Linear Algebra Appl. **429** (2008), no. 1, 110–121.
- [6] A. Ferrante and B. Levy, *Hermitian solutions of the equation $X = Q + N^*XN$* , Linear Algebra Appl. **247** (1996), 359–373.
- [7] J. Harjani, B. López, and K. Sadarangani, *Fixed point theorems for mixed monotone operators and applications to integral equations*, Nonlinear Anal. **74** (2011), no. 5, 1749–1760.
- [8] V. Hasanov, *Positive definite solutions of the matrix equations $X \pm A^*X^{-q}A = Q$* , Linear Algebra Appl. **404** (2005), 166–182.
- [9] M. Huang, C. Huang, and T. Tsai, *Applications of Hilbert's projective metric to a class of positive nonlinear operators*, Linear Algebra Appl. **413** (2006), no. 1, 202–211.
- [10] V. Lakshmikantham and L. Ćirić, *Coupled fixed point theorems for nonlinear contractions in partially ordered metric spaces*, Nonlinear Anal. **70** (2009), no. 12, 4341–4349.
- [11] Y. Lim, *Solving the nonlinear matrix equation $X = Q + \sum_{i=1}^m M_i X^{\delta_i} M_i^*$ via a contraction principle*, Linear Algebra Appl. **430** (2009), no. 4, 1380–1383.
- [12] X. G. Liu and H. Gao, *On the positive definite solutions of the matrix equation $X^s \pm A^T X^{-t} A = I_n$* , Linear Algebra Appl. **368** (2003), 83–97.
- [13] N. V. Luong and N. X. Thuan, *Coupled fixed points in partially ordered metric spaces and application*, Nonlinear Anal. **74** (2011), no. 3, 983–992.
- [14] B. Meini, *Efficient computation of the extreme solutions of $X + A^*X^{-1}A = Q$ and $X - A^*X^{-1}A = Q$* , Math. Comput. **71** (2001), 1189–1204.
- [15] A. C. M. Ran and M. C. B. Reurings, *A fixed point theorem in partially ordered sets and some applications to matrix equations*, Proc. Amer. Math. Soc. **132** (2004), no. 5, 1435–1443.

SEJONG KIM
 DEPARTMENT OF MATHEMATICS
 CHUNGBUK NATIONAL UNIVERSITY
 CHEONGJU 28644, KOREA
 E-mail address: skim@chungbuk.ac.kr

HOSOO LEE
DEPARTMENT OF MATHEMATICS
SUNGKYUNKWAN UNIVERSITY
SUWON 16419, KOREA
E-mail address: `hosoollee@skku.edu`