

## CORRIGENDUM

29권 3호(p. 349-358)의 “복부비만과 위험음주가 성인의 고혈압에 미치는 영향” 이은숙 중 p. 349의 Abstract과 p. 353 Table 1, p.354의 Table 2~4의 오류를 정정합니다.

## 1) p. 349 - Abstract - 5~8줄

**Before correction**

The odds ratio of hypertension for risk drinkers with abdominal obesity was 4.81 compared to non-risk drinkers with normal waist circumference, whereas the odds ratios of hypertension for risk drinkers with normal waist circumference and non-risk drinkers with abdominal obesity were 1.58 and 2.37 respectively.

**After correction**

The odds ratio of hypertension for risk drinkers with abdominal obesity was 4.81 compared to non-risk drinkers with normal waist circumference, whereas the odds ratios of hypertension for risk drinkers with normal waist circumference and non-risk drinkers with abdominal obesity were 1.58 and 2.37 respectively.

## 2) p. 353 - Table 1. Characteristics 열의 12줄

**Before correction**

Waist circumference

**After correction**

Waist circumference

## 3) p. 354 - Table 2

**Before correction****Table 2.** Odds Ratio for Hypertension according to Waist Circumstance

(N=13,885)

Variable	Categories	Model 1 <sup>†</sup>		Model 2 <sup>‡</sup>	
		OR (95% CI)	p	OR (95% CI)	p
Waist circumference	< 90 (men)/85 (women) cm	1		1	
	≥ 90 (men)/85 (women) cm	2.57 (2.33~2.83)	<.001	2.18 (1.95~2.43)	<.001

OR=odds ratio; CI=confidence interval; <sup>†</sup> Adjusted for age, sex; <sup>‡</sup> Adjusted for age, sex, education, marital status, occupation, smoking, diabetes mellitus, hyperlipidemia, frequency of drinking, typical drinking quantity.

**After correction****Table 2.** Odds Ratio for Hypertension according to Waist Circumference

(N=13,885)

Variable	Categories	Model 1 <sup>†</sup>		Model 2 <sup>‡</sup>	
		OR (95% CI)	p	OR (95% CI)	p
Waist circumference	< 90 (men)/85 (women) cm	1		1	
	≥ 90 (men)/85 (women) cm	2.57 (2.33~2.83)	<.001	2.18 (1.95~2.43)	<.001

OR=odds ratio; CI=confidence interval; <sup>†</sup> Adjusted for age, sex; <sup>‡</sup> Adjusted for age, sex, education, marital status, occupation, smoking, diabetes mellitus, hyperlipidemia, frequency of drinking, typical drinking quantity.

## 4) p. 354 - Table 3. 각주

**Before correction**

WC (waist circumference)

**After correction**

WC (waist circumference)

## 5) p. 354 - Table 4. 각주

**Before correction**

WC (waist circumference)

**After correction**

WC (waist circumference)

DOI of original article <https://doi.org/10.12799/jkachn.2018.29.3.349>

Corresponding author: **Lee, Eun Sook**

Department of Nursing, Gyeongnam National University of Science and Technology, 33 Dongjin-ro, Jinju 52725, Korea.

Tel: +82-55-751-3655, Fax: +82-55-751-3659, E-mail: eslee5335@gntech.ac.kr