

Effects of *Rubus coreanus* byproducts on intestinal microbiota and the immune modulation

Da Yoon Yu^{1,a}, Sung Hak Kim^{2,a}, Jeong A Kim¹, In Sung Kim¹, Yang Soo Moon³, Sang Suk Lee⁴, Hwa Chun Park⁵, Jong Hyun Jung⁵, Yi Hyung Chung⁶, Dae Keun Shin⁷, Ki Chang Nam⁴, In Soon Choi⁸, and Kwang Keun Cho^{1,*}

* **Corresponding Author:** Kwang Keun Cho
Tel: +82-55-751-3286, **Fax:** +82-55-751-3689,
E-mail: chotwo2@daum.net

¹ Department of Animal Resources Technology, Gyeongsang National University of Science and Technology, Jinju 52725, Korea

² Department of Animal Science, Chonnam National University, Gwangju 61186, Korea

³ Department of Animal Science & Biotechnology, Gyeongsang National University of Science and Technology, Jinju 52725, Korea

⁴ Department of Animal Science and Technology, Sunchon National University, Suncheon 57922, Korea

⁵ Dasan Genetics, Namwon 55716, Korea

⁶ Jeonbuk Institute for Bioindustry, Jeonju 54810, Korea

⁷ Berry & Biofood Research Institute, Gochang 56417, Korea

⁸ Department of Life Science, Silla University, Busan 46958, Korea

^a These authors contributed equally to this study.

ORCID

Da Yoon Yu

<https://orcid.org/0000-0001-5415-2294>

Sung Hak Kim

<https://orcid.org/0000-0003-4882-8600>

Jeong A Kim

<https://orcid.org/0000-0002-1993-3316>

In Sung Kim

<https://orcid.org/0000-0001-9127-0732>

Yang Soo Moon

<https://orcid.org/0000-0001-9858-1779>

Sang Suk Lee

<https://orcid.org/0000-0003-1540-7041>

Hwa Chun Park

<https://orcid.org/0000-0001-8987-7789>

Jong Hyun Jung

<https://orcid.org/0000-0003-3667-7710>

Yi Hyung Chung

<https://orcid.org/0000-0001-5470-8360>

Dae Keun Shin

<https://orcid.org/0000-0003-3977-0208>

Ki Chang Nam

<https://orcid.org/0000-0002-2432-3045>

In Soon Choi

<https://orcid.org/0000-0001-8803-9142>

Kwang Keun Cho

<https://orcid.org/0000-0001-8834-5369>

Submitted Oct 2, 2017; Revised Nov 26, 2017;

Accepted Jan 4, 2018

Objective: Although the efficacy of *Rubus coreanus* (RC) byproducts as a feed additive has been recognized, its effects on intestinal microorganisms and the immune system are still unknown.

Methods: Six-week-old male rats were treated with 0.5% RC (T1), 1.0% RC (T2), and 1.5% RC (T3) for 4 weeks.

Results: We found that treatment with RC byproducts significantly increased the daily gain of body weight and feed intake. Treg-cell differentiation was enhanced in the mesenteric lymph nodes and spleen from the rats fed with RC byproducts. Illumina sequencing showed that bacteria in the phylum *Firmicutes* decreased and while those in the phylum *Bacteroidetes* increased in RC-treated groups. Particularly, the pathogenic microorganisms in the family *Peptococcaceae* decreased, and the non-pathogenic families *Lachnospiraceae* and *S24-7* increased. Quantitative polymerase chain reaction analysis showed that the RC byproducts increased the lactic acid bacteria *Bifidobacterium* spp., *Oscillospira* spp., *Leuconostoc citreum*, and *Weissella cibaria* in a concentration-dependent manner.

Conclusion: RC byproducts may be effective in immunomodulation by affecting intestinal microorganisms.

Keywords: *Rubus coreanus*; Th1/Th2 Balance; Galectin-9; Illumina Sequencing; Intestinal Microbiota; Immune Modulation

INTRODUCTION

Antibiotics have been used as feed additives for the purpose of preventing disease, promoting growth, and improving feed efficiency in animals. As antibiotics can remain in livestock products such as meat, eggs, and milk causing antibiotic resistance in humans, their use is increasingly banned across countries. Recently, the number of antibiotics used as feed additives has been reduced from 53 to 23, and among the antibiotics used in 25 feed additives, the tetracycline series and seven generic antibiotics have been banned [1]. Due to these antibiotic restrictions in feed additives, various natural resource materials have been developed.

Rubus coreanus (RC) is a deciduous shrub belonging to the *Rosaceae* family and generally contains 87.0 g of water, 1.6 g of protein, 1.7 g of fat, 1.3 g of glucose, 2.9 g of fiber and 0.5 g of ash in 100 g as well as Ca, Mg, Na and Fe and vitamins such as vitamin A, vitamin B1, and vitamin C [2]. It has been reported that organic acids such as gallic acid, polyphenols such as quercetin, kaempferol, and tannins in RC have various anticancer, antioxidant, cholesterol-lowering and anti-obesity effects [3]. In addition, ellagitannins and ellagic acid, which possess anti-cancer, anti-diabetic and anti-inflammatory effects, are very poorly ab-

sorbed by the gastrointestinal system and the unabsorbed compounds are metabolized to urolithin by intestinal microorganisms [4]. As the activity of useful ingredients in RC has been reported, various functional products are attracting attention. The byproducts account for approximately 20% of the RC juice extracts, but are mostly being discarded [5]. The hot water extract of RC byproducts contains 8.9% of organic acid, 69.9% of neutral sugar, 19.1% of pectin and 5.2% of anthocyanin [6].

Although studies on the polyphenol content [7] of RC byproducts using ethanol concentration have been reported, studies using RC byproducts have not been performed yet. Therefore, this study was conducted to investigate the effects of RC byproducts on intestinal microbes and the immune system in white rats. This may prove useful for the production of functional products using RC.

MATERIALS AND METHODS

Feed and experimental animal feeding experiment

The RC byproducts used in this experiment were obtained by generating a RC extract, drying the remaining byproducts (peel, seed) in a 50°C dry oven for 24 hours and then grinding them into a powder. Six-week-old male Sprague-Dawley rats were purchased from Samtako (Osan, Korea). A total of 6 rats were used in each treatment group, which were further divided into six repetitions consisting of one rat per repetition. The rats were subjected to a basal diet (AIN-76, Feed Lab, Guri, Korea) (Control group), a basal diet mixed with 0.5% dried RC byproduct powder (T1 group), a basal diet mixed with 1.0% dried RC byproduct powder (T2 group), and a basal diet mixed with 1.5% dried RC byproduct powder (T3 group). The animal housing was maintained at a temperature of 22°C ±3°C, relative humidity of 55%±5%, and light and dark cycles of 12 hours each. The rats underwent a 7-day acclimatization period before subjecting them to the aforementioned diets for 4 weeks, and the feed and drinking water were provided *ad libitum*. The weight and feed intake of each rat were measured once weekly throughout the experimental period. All procedures were approved by the Institutional Animal Care and Use Committees at Gyeongnam National University of Science and Technology (No. 2016-10).

DNA extraction and Illumina sequencing

All animals were sacrificed by diethyl ether and fecal samples of the rats were collected and stored at -80°C until genomic DNA (gDNA) extraction. DNA was extracted using Fecal DNA MiniPrep kits (Zymo Research, Irvine, CA, USA), and the sequence was analyzed using Illumina MiSeq (Chunlab Inc., Seoul, Korea) [8].

Quantitative polymerase chain reaction

As mentioned above, DNA was extracted from rat fecal samples. Quantitative polymerase chain reaction (qPCR) was performed using Rotor-Gene SYBR Green PCR Kits and Rotor-Gene Q (Qiagen, Hilden, Germany). The PCR cycling conditions were as follows: 40 cycles of 95°C for 10 s, 56°C for 30 s, 72°C for 10 s. Universal primer was used as the reference gene (Table 1).

Separation and analysis of RNA from rat spleen and mesenteric lymph nodes and RT-PCR

To analyze the immunomodulatory effects of the RC byproducts, rats were sacrificed and mesenteric lymph nodes (MLNs) and spleen were separated. Trizol was added to MLN and spleens and they were homogenized using SilentCrusher M (Heidolph, Schwabach, Germany). RNA was extracted and stored at -20°C until cDNA synthesis using reverse transcription-PCR (RT-PCR) kits (TaKaRa Co., Tokyo, Japan) according to the manufacturer's instructions. RT-PCR (GeneAMP PCR System 9700; Thermo Fisher Scientific Inc., Waltham, MA, USA) cycling conditions are as follows: initial denaturation step at 50°C for 30 min, 94°C for 2 min, followed by 35 cycles of 94°C for 30 s, 55°C for 30 s, and 72°C for 59 s, and a final extension step at 72°C for 2 min. Glyceraldehyde-3-phosphate dehydrogenase was used as the reference gene (Table 2).

Table 1. Quantitative real-time polymerase chain reaction primer design

Target group/specificity		Primer sequence
Universal	Forward	GTGSTGCAYGGYYGTCGTCA
	Reverse	ACGTCRTCCMCNCCTTCTCTC
<i>Bacteroides</i> spp.	Forward	GAAGGTCCCCACATTG
	Reverse	CGCKACTTGGCTGGTTTCAG
<i>Roseburia</i> spp. and <i>E. rectale</i>	Forward	GCGGTRCGGCAAGTCTGA
	Reverse	CCTCCGACTCTAGTMCAGC
<i>Faecalibacterium prausnitzii</i>	Forward	GGAGGAAGAAGTCTTCGG
	Reverse	AATCCGCCTACTCTGCACT
Cluster IV <i>Ruminococcus</i> spp.	Forward	GGCGGCYTRCTGGGCTTT
	Reverse	CCAGGTGGATWACTTATTGTGTTAA
<i>Bifidobacterium</i> spp.	Forward	TCCGCTCYGGTGTGAAAG
	Reverse	GGTGTCTTCCGATATCTACA
Methanogens	Forward	GGATTAGATACCCSGGTAGT
	Reverse	GTTGARTCCAATTAACCGCA
<i>Oscillospira</i> spp.	Forward	ACGGTACCCCTTGAAATAAGCC
	Reverse	TCCCGCACACCTAGTATTG
<i>Leuconostoc mesenteroid</i>	Forward	TGATGCATAGCCGAGTTGAG
	Reverse	GAAAGCCTTCATCACACACG
<i>Leuconostoc citreum</i>	Forward	GGAAACAGATGCTAATACCGAATA
	Reverse	TTTACCCACCAACTAATAATG
<i>Weissella cibaria</i>	Forward	GGGAAACCTACTCTTAGCA
	Reverse	GGACCATCTCTTAGTGATAGCA
<i>Weissella koreensis</i>	Forward	GGGCTACACACGTGTACAA
	Reverse	GATTCCGACTTCGTGTAGGC
<i>Lactobacillus sakei</i>	Forward	CCATGTGTAGCGGTGAAATG
	Reverse	ATCCTGTTGTACCCATGC

Table 2. Reverse transcriptase polymerase chain reaction primer design

Target group/ specificity	Primer sequence
TGF-β	Forward GAAGGCAGAGTTCAGGGTCTT
	Reverse GGTTCTGTCTTTGTGGTGAA
IL-10	Forward ATAAGTGCACCACTTCCCA
	Reverse TCATTTCCGATAAGGCTTGG
IFN-γ	Forward TCAAGTGGCATAGATGTGGAAGAA
	Reverse TGGCTCTGCAGGATTTTCATG
IL-17A	Forward TTCATCTGTGTCTCTGATGCT
	Reverse TTGACCTTCACATTCTGGAG
IL-4	Forward ACAGGAGAAGGGACGCCAT
	Reverse GAAGCCCTACAGACGAGCTCA
Foxp3	Forward CCCATCCCAGGAGTCTTG
	Reverse CCATGACTAGGGCACTGTA
GATA3	Forward CATTACCCTATCCGCCCTATG
	Reverse CACACACTCCCTGCCTTCTGT
T-bet	Forward TCAACCAGCACCAGACAGAG
	Reverse AAACATCCTGTAATGGCTTGTG
RORγt	Forward TTCACCCCACTCCACTG
	Reverse TGCAAGGGATCACTTCAATTT

TGF-β, transforming growth factor beta; IL-10, interleukin-10; IFN-γ, interferon-γ; Foxp3, forkhead box P3; GATA3, GATA binding protein 3; T-bet, T-box expressed in T cells; RORγt, Rar-related orphan receptor gamma T.

Statistical analysis

All results are shown as means±standard deviation and the variance of the results obtained through repeated experiments was analyzed using SPSS 12.0 (SPSS Inc., Chicago, IL, USA). The significance of the results was tested using Duncan’s multiple range test at a level of p<0.05.

RESULTS

Body weight, daily gain, daily feed intake and feed efficiency in rats

Rats in the T3 RC byproduct-fed group showed the greatest increase in both average daily gain and average daily feed intake (Table 3). The treatments with T1, T2, and T3 caused a marked feed efficiency when compared to the control group.

Table 3. Rat fed *Rubus coreanus* byproducts weight gain, daily feed intake, and feed efficiency

	C ¹⁾	T1	T2	T3
Initial body weight (g)	220.34±0.53	220.05±0.79	220.21±0.48	219.90±0.70
Finished body weight (g)	308.45±0.17 ^a	307.38±0.22 ^a	308.18±0.88 ^a	311.35±0.97 ^b
Average daily gain (g)	3.15±0.02 ^a	3.12±0.02 ^a	3.14±0.04 ^a	3.27±0.04 ^b
Average daily feed intake (g)	20.49±0.68 ^c	17.93±0.39 ^a	18.67±0.19 ^b	19.19±0.40 ^b
Feed efficiency	0.15±0.005 ^a	0.17±0.004 ^b	0.17±0.003 ^b	0.17±0.003 ^b

SD, standard deviation; ADG, average daily gain; ADFI, average daily feed intake.

Data represent means±standard deviation of 6 replicates.

¹⁾ C, control; T1, 0.5% *Rubus coreanus* byproducts; T2, 1.0% *Rubus coreanus* byproducts; T3, 1.5% *Rubus coreanus* byproducts.

ADG, ADFI and feed efficiency increased when *Rubus coreanus* byproducts were fed as compared to control.

^{a-c} Means are significantly different within the same row (p<0.05).

mRNA expression of galectin-9 in MLN and spleen

Galectin-9 is known to induce a Treg-mediated immune response [9]. We performed RT-PCR analysis to investigate whether RC byproducts can increase the mRNA expression of galectin-9 in MLNs and spleens of the rats. mRNA expression of galectin-9 was significantly increased in MLNs as well as spleens from the RC-treated (T1, T2, and T3) rats compared to that in the control group. In addition, the treatment with T1 caused the highest increase in the mRNA expression of galectin-9 in MLN (Figure 1).

Th1/Th2 balance and the development of Treg cells in MLN and spleen

In order to investigate whether T cell polarization is influenced by RC, we evaluated the expression of the transcription factors T-box expressed in T cells (T-bet) for Th1, GATA binding protein 3 (GATA 3) for Th2, RAR-related orphan receptor gamma T (RORγT) for Th17 and forkhead box P3 (Foxp3) for Treg cells in the MLN and spleen. In the MLN, the expression of T-bet and Foxp3 were increased, while the expression of GATA-3 and RORγT were decreased in the T1, T2, and T3 treatment groups compared to that in the control group (Figures 2A-2D). Th1/Th2 and Treg/Th2 ratios were significantly increased in a dose-dependent manner in rats fed with T1, T2, and T3, inducing a Th1 and Treg cell-mediated immunity response (Figures 2E, 2G). However, the Treg/Th1 ratio was decreased and the Treg/(Th1+Th2) ratio was not changed (Figures 2E-2H). Consistent with the results in the MLN, T-bet and Foxp3 expression were increased in the T1, T2, and T3 treatment groups (Figures 3A, 3D), whereas GATA-3 and RORγT expression were decreased in the spleen compared to that in the control group (Figures 3B, 3C). The ratios of Th1/Th2 and Treg/Th1, and Treg/Th2 were increased to induce Th1 and Treg cell-mediated immunity (Figures 3E-3G). The ratio of Treg/(Th1+Th2) was also increased by T1, T2, and T3 treatment (Figure 3H). Therefore, these results indicate that RC byproducts can improve rat immunity by enhancing Treg cells in the MLN and spleen.




Figure 1. Rat fed *Rubus coreanus* byproducts show enhanced galectin-9 development in mesenteric lymph nodes (MLN) and spleen. C, control; T1, 0.5% *Rubus coreanus* byproducts; T2, 1.0% *Rubus coreanus* byproducts; T3, 1.5% *Rubus coreanus* byproducts. The expression of galectin-9 in MLN was higher in the group fed with *Rubus coreanus* byproducts than in control, and highest in the group of 0.5% *Rubus coreanus* byproducts. Galectin-9 expression in the spleen was also higher in the group fed with *Rubus coreanus* byproducts than in control. a-c Means are significantly different within the same row ($p < 0.05$). Data represent means \pm standard deviation of 6 replicates.




Figure 2. Rat fed *Rubus coreanus* byproducts show enhanced Treg-cell development in mesenteric lymph nodes (MLN). C, control; T1, 0.5% *Rubus coreanus* byproducts; T2, 1.0% *Rubus coreanus* byproducts; T3, 1.5% *Rubus coreanus* byproducts. T-cell polarization in MLN was evaluated by analyzing the expression of T-bet (Th1, A), GATA binding protein 3 (GATA-3) (Th2, B), RAR-related orphan receptor gamma T (RORγT) (Th17, C) and forkhead box P3 (Foxp3) (Treg, D). The expression of Foxp3 tended to increase in rat fed *Rubus coreanus* byproducts, while GATA-3 and RORγT expression was decreased. Expression of T-bet was not changed. Rat fed *Rubus coreanus* byproducts showed an increased Th1/Th2 (E) and Treg/Th2 ratio (F), but the Treg/Th1 ratio decreased (G). The ratio between Treg cells and effector Th1 and Th2 cells (H) was not changed. ^{a-c} Means are significantly different within the same row ($p < 0.05$). Data represent means \pm standard deviation of 6 replicates.




Figure 3. Rat fed *Rubus coreanus* byproducts show enhanced Treg-cell development in spleen. C, Control; T1, 0.5% *Rubus coreanus* byproducts; T2, 1.0% *Rubus coreanus* byproducts; T3, 1.5% *Rubus coreanus* byproducts. T-cell polarization in spleen was evaluated by analyzing the expression of T-box expressed in T cells (T-bet) (Th1, A), GATA binding protein 3 (GATA-3) (Th2, B), RAR-related orphan receptor gamma T (RORγT) (Th17, C) and forkhead box P3 (Foxp3) (Treg, D). The expression of Foxp3 tended to increased in rat fed *Rubus coreanus* byproducts, while GATA-3 and RORγT expression was decreased. Expression of T-bet was no changed. Rat fed *Rubus coreanus* byproducts showed an increased Th1/Th2 (E) and Treg/Th2 ratio (F), but the Treg/Th1 ratio decreased (G). The ratio between Treg cells and effector Th1 and Th2 cells (H) was increased. ^{a-c} Means are significantly different within the same row (p<0.05). Data represent means±standard deviation of 6 replicates.

Intestinal microbiological environment in rats

When rats were supplied with T1, T2, and T3, *Firmicutes* (F) levels in the intestinal microflora were markedly decreased while *Bacteroidetes* (B) levels were increased in a dose-dependent manner (Figure 4). The F/B ratio was decreased depending on the concentration of T1, T2, and T3. Particularly, *Clostridia* from the phylum *Firmicutes* and pathogenic microorganisms belonging to the *Peptococcaceae* decreased by the consumption of T1, T2, and T3, and the non-pathogenic *Lachnospiraceae* and *Ruminococcaceae* microorganisms increased (Table 4). Among the class *Bacteroidia*, which occupies most of the *Bacteroidetes*, the number of microbes belonging to the non-

pathogenic *S24-7* and *Rikenellaceae* family increased. These results show that microorganisms promoting intestinal health were increased in rats fed with RC byproducts.

Microorganisms showing obesity, anti-obesity, and butyrate-producing traits

Rats fed with T1, T2, and T3 byproducts showed a decrease in anti-obesity microorganisms, while an increase in obesity- and butyrate-producing microorganisms occurred in a concentration-dependent manner (Figure 5). Therefore, these data suggest that RC byproducts can improve intestinal health by increasing the number of butyrate-producing microorganisms.




Figure 4. Effects of dietary supplementation of *Rubus coreanus* byproducts on diversity of phyla of Intestinal microbiota in rats. C, control; T1, 0.5% *Rubus coreanus* byproducts; T2, 1.0% *Rubus coreanus* byproducts; T3, 1.5% *Rubus coreanus* byproducts. Duncan’s multiple range test was used to show the *Firmicutes* and *Bacteroidetes* contents and F/B ratios of each sample at the phylum level. F/B ratio was reduced by *Rubus coreanus* byproducts. ^{a-c} Means are significantly different within the same row ($p < 0.05$). Data represent means \pm standard deviation of 6 replicates.

Intestinal lactic acid bacteria and intestinal microflora

The qPCR analysis showed that RC byproducts induced 12 intestinal probiotics and microorganisms compared to that in the control group. Particularly, *Bifidobacterium* spp., *Leuconostoc citreum*, *Weissella cibaria*, *Oscillospira* spp. showed the most prominent increase (Figure 6).

DISCUSSION

The RC byproducts account for approximately 20% of the bok-bunja juice extract, but are mostly being discarded [5]. There

have been some studies on the general components and poly-phenol contents of RC byproducts [6,7]. To observe the effects of these byproducts on the intestinal microorganism environment and immune system of experimental animals, MLN and spleen mRNA was purified and analyzed for the composition of intestinal microorganisms via pyro-sequencing, and by investigating the mechanisms of immune function regulation. The expression of galectin-9 in the MLNs and spleens of rats fed 0.5%, 1.0%, and 1.5% RC was higher than that of the control group. This implies that RC byproducts can activate galectin-9 in the body. Previous studies have shown that

Table 4. The mean of the relative abundance of dominant phyla, classes and familia in fecal sample obtained from rat

Items ²⁾	Treatments ¹⁾			
	C	T1	T2	T3
	n = 6 mean% (SD)	n = 6 mean% (SD)	n = 6 mean% (SD)	n = 6 mean% (SD)
Firmicutes	85.79 (3.68) ^c	80.90 (2.97) ^b	76.45 (2.77) ^a	74.96 (2.56) ^a
Clostridia	68.49 (3.19) ^b	65.01 (1.71) ^a	63.13 (2.80) ^a	65.05 (2.08) ^a
<i>Lachnospiraceae</i>	39.74 (2.21) ^a	40.50 (2.78) ^a	42.10 (2.16) ^{ab}	44.93 (2.24) ^b
<i>Ruminococcaceae</i>	19.32 (4.18)	19.84 (2.06)	19.59 (2.80)	18.40 (1.78)
<i>Peptococcaceae</i>	3.19 (0.19) ^c	1.31 (0.15) ^b	0.16 (0.07) ^a	0.13 (0.04) ^a
Bacilli	15.86 (2.04) ^c	13.11 (1.80) ^b	10.95 (1.23) ^a	8.90 (1.96) ^a
<i>Lactobacillaceae</i>	15.37 (2.81) ^c	12.71 (2.76) ^{bc}	10.50 (2.92) ^{ab}	8.60 (1.41) ^a
Bacteroidetes	11.23 (2.50) ^a	17.14 (2.05) ^b	21.95 (2.76) ^c	23.77 (3.07) ^c
Bacteroidia	10.84 (1.25) ^a	16.82 (2.50) ^b	21.74 (1.91) ^c	23.38 (1.98) ^c
<i>S24-7_f</i>	7.33 (1.02) ^a	8.98 (1.71) ^a	11.32 (2.17) ^b	14.95 (1.59) ^c
<i>Prevotellaceae</i>	1.28 (0.74) ^a	6.08 (1.90) ^c	6.68 (1.77) ^c	3.40 (1.00) ^b
<i>Rikenellaceae</i>	1.68 (0.57) ^a	1.16 (0.52) ^a	2.88 (1.48) ^b	4.29 (1.08) ^c

SD, standard deviation.

Data represent means \pm SD of 6 replicates.

¹⁾ C, control; T1, 0.5% *Rubus coreanus* byproducts; T2, 1.0% *Rubus coreanus* byproducts; T3, 1.5% *Rubus coreanus* byproducts.

²⁾ Statistical tests of over- or under-representation of bacterial lineages among at each sample were made at the phylum (bold), class (bold) and family levels using Duncan’s multiple range test. Rat fed *Rubus coreanus* byproducts show *Firmicutes* decreased and *Bacteroidetes* increased.

^{a-c} Means are significantly different within the same row ($p < 0.05$).




Figure 5. Rat fed *Rubus coreanus* byproducts show relatively changed intestinal microorganisms. C, control; T1, 0.5% *Rubus coreanus* byproducts; T2, 1.0% *Rubus coreanus* byproducts; T3, 1.5% *Rubus coreanus* byproducts. The relative abundance of obesity microorganism (*Roseburia*, *Eubacterium_g8*, *Pseudoflavonifractor*, *Ruminococcaceae_uc*, *Ruminococcus*, *JN713389_g*, *Eubacterium_g23*, *AB185816_g*, *Caproiciproducens*, *Ruminococcus_g2*, *JQ084194_g*, *Subdoligranulum*, *Oscillibacter*, *Butyricicoccus*, *EF602946_g*, *Allobaculum*, *Erysipelotrichaceae_uc*, *Coprobacillus*, *Faecalibaculum*, *EF445272_g*; (A) and antiobesity microorganism (*Bacteroides*, *Parabacteroides*, *Prevotella*, *Prevotellaceae_uc*, *Alistipes*, *AB239481_g*, *Christensenellaceae_uc*, *HM124260_g*, *GU302849_g*, *Christensenella*, *Bacteroides_g1*; (B) tended to increase in rat fed *Rubus coreanus* byproducts, while butyric acid microorganism (*Eubacterium_g17*, *Eubacterium_g4*, *Roseburia*, *Eubacterium_g6*, *Eubacterium_g20*, *Eubacterium_g21*, *Eubacterium_g8*, *Pseudoflavonifractor*, *Ruminococcaceae_uc*, *Ruminococcus*, *JN713389_g*, *Eubacterium_g23*, *AB185816_g*, *Caproiciproducens*, *Ruminococcus_g2*, *JQ084194_g*, *Subdoligranulum*, *Oscillibacter*, *Anaerotruncus*, *Butyricicoccus*, *EF602946_g*, *Phascolarctobacterium*, *Clostridium_g21*, *AB606237_g*, *FJ880395_g*, *Blautia*, *JPZU_g*, *AJ576336_g*, *AB606254_g*, *HM123979_g*, *Coprococcus_g2*, *Ruminococcus_g4*, *GU324393_g*, *Murimonas*, *EU509117_g*, *Marvinbryantia*; (C) relative abundance was decreased. Data represent means±standard deviation of 6 replicates.

galectin-9 induces Treg cell immune responses and neutralizes immunoglobulin E antibodies [10]. In addition, RC byproducts activated the mRNA expression of galectin-9 in the MLN and spleen with increased T-bet and Foxp3 as a reflection of Th1 and Treg cells, respectively. T-bet and Foxp3 could bind to multiple distal sites at immune regulatory genes for T cell fate determination as transcription factors [11]. This is consistent with the results of Kivit et al [12] who showed that galectin-9 induces Th1 and Treg cell development and increases interferon- γ and interleukin-10 secretion to suppress the allergic response and regulate immunity [12]. In addition, galectin-9, which induces apoptosis of T cells, inhibited the activation level of Treg/Th1 immune factors. Therefore, the RC byproducts activate the expression of galectin-9, and the activated galectin-9 promotes Th 1 and Treg immune responses

and thus induces immunomodulation.

The mammalian intestinal microbiome is composed of hundreds of millions of microorganisms and could affect host metabolism and diseases such as obesity, arteriosclerosis, and type 2 diabetes. In addition to the host's metabolism, these microbes also provide many advantages, including the development of the immune system and maintenance of the epithelial barrier [13]. *Firmicutes* and *Bacteroidetes* are the predominant phyla living in the intestine of humans and most mammals, and *Proteobacteria*, *Actinobacteria*, and *Tenericutes* together constitute other phyla [14]. *Bacteroidetes* [15], which contribute to defense against potential pathogens in the intestine, are increased by RC byproducts in the diet. The increase in *Bacteroidetes* is associated with a decrease in metabolic syndrome indicators such as high cholesterol and blood pressure




Figure 6. Rat fed *Rubus coreanus* byproducts show increased intestinal lactobacilli and microorganisms by quantitative polymerase chain reaction (qPCR). C, control; T1, 0.5% *Rubus coreanus* byproducts; T2, 1.0% *Rubus coreanus* byproducts; T3, 1.5% *Rubus coreanus* byproducts. The expression of *Bifidobacterium* spp., *Leuconostoc citreum*, *Weissella cibaria* and *Oscillospira* spp. tended to increased in rat fed *Rubus coreanus* byproducts. ^{a-c} Means are significantly different within the same row ($p < 0.05$). Data represent means \pm standard deviation of 6 replicates.

[16]. Patients with type 1 diabetes or hypertension have increased intestinal levels of *Firmicutes*, which in turn increases the ratio of *Firmicutes/Bacteroidetes* [17]. Rats treated with RC byproducts showed a relative decrease of *Firmicutes*, and the ratio of *Firmicutes/Bacteroidetes* decreased in a concentration-dependent manner. The family *Peptococcaceae* in the *Clostridia* class consists of pathogenic microorganisms showing a positive correlation with CCL5, an inflammation-inducing cytokine, and decreased in response to RC treatment [18]. *Lachnospiraceae* and *Ruminococcaceae* of class *Clostridia*, the levels of which increase when RC byproducts are given, are microorganisms associated with metabolic health [19]. It is also known that RC produces intestinal anti-inflammatory compounds and butyrate which has immunomodulating properties [20]. The increased levels of microbes in the S24-7 family in the *Bacteroidia* class act as a protective factor for diabetes [21], while *Rikenellaceae* alter the intestinal environment and exhibit antitumor effects [22]. The decrease in pathogenic microorganisms and the increase in non-pathogenic microorganisms suggest that intestinal health is enhanced by RC byproducts

in rats. The bioavailability of ellagitannins and ellagic acid in RC byproducts is very low. These molecules are converted by intestinal microorganisms to urolithin, which is much better absorbed in the body [23]. Urolithin production is more abundant in *Firmicutes*, especially in *Clostridium leptum*, than in *Bacteroidetes* [24]. Urolithin is a microflora metabolite of ellagic acid derivatives [25], which enhances anti-proliferative, anti-inflammatory, and vascular endothelial function in cancer cell models [26]. Dietary fibers, such as RC byproducts, are fermented by microorganisms in the large intestine to produce short-chain fatty acids containing butyric acid [27]. Butyrate-producing microorganisms belong to the *Firmicutes*, and among them, *Ruminococcus*, *Roseburia*, and *Eubacterium* are prominent examples [28]. *Bifidobacterium* spp. is an important microorganism with immunomodulatory and anticancer activity [29], and its levels increased when bifidobacterial byproducts were ingested. In rats fed with RC byproducts, levels of butyrate-producing microorganisms increased, leading to improved intestinal health. In conclusion, the consumption of RC byproducts increased the expression of Th1 and Treg

cell transcription factors and galectin-9 in this study. In addition, the intestinal environment was changed by decreasing *Firmicutes*, increasing *Bacteroidetes*, and decreasing the *Firmicutes/Bacteroidetes* ratio. This suggests that bokbunja byproducts improve immune function and promote bowel health.

CONFLICT OF INTEREST

We certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript. Park HC, Jung JH are employees of Dasan Genetics, and Shin DK is employee of Berry & Biofood Research Institute.

ACKNOWLEDGMENTS

This research was supported by the Economic Cooperation Industrial Development Project (No. R0005730), Ministry of Trade, Industry, and Energy, Korea.

REFERENCES

- Jang YD, Oh HK, Piao LG, et al. Evaluation of probiotics as an alternative to antibiotic on growth performance, nutrient digestibility, occurrence of diarrhea and immune response in weaning pigs. *Anim Feed Sci Technol* 2009;51:25-32.
- Sung KC. A Study on the pharmaceutical and chemical characteristics and analysis of natural Bokbunja extract. *J Korean Oil Chem Soc* 2012;29:402-11.
- Jang TS, Yang JC, Lim SY, Kim BA. Antioxidant and antihemolytic activity of ethanol extracts of *Rubus coreanus* Miquel. *J Korean Oil Chem Soc* 2014;31:130-5.
- Espín JC, Larrosa M, García-Conesa MT, Tomás-Barberán F. Biological significance of urolithins, the gut microbial ellagic acid-derived metabolites: the evidence so far. *Evid Based Complement Alternat Med* 2013;2013:Article ID 270418.
- Ku CS, Mun SP. Optimization of the extraction of anthocyanin from Bokbunja (*Rubus coreanus* Miq.) marc produced during traditional wine processing and characterization of the extracts. *Bioresour Technol* 2008;99:8325-30.
- Ku CS, Mun SP. Antioxidant activities of ethanol extracts from seeds in fresh Bokbunja (*Rubus coreanus* Miq.) and wine processing waste. *Bioresour Technol* 2008;99:4503-9.
- Jeong YJ, Seo JH. Conditions for pigment extraction from Bokbunja (*Rubus coreanus* Miquel) byproducts. *Korean J Food Preserv* 2009;16:400-4.
- Jeon YS, Park SC, Lim J, Chun J, Kim BS. Improved pipeline for reducing erroneous identification by 16S rRNA sequences using the Illumina MiSeq platform. *J Microbiol* 2015;53:60-9.
- Rabinovich GA, Toscano MA. Turning 'sweet' on immunity: galectin-glycan interactions in immune tolerance and inflammation. *Nat Rev Immunol* 2009;9:338-52.
- Niki T, Tsutsui S, Hirose S, et al. Galectin-9 is a high affinity IgE-binding lectin with anti-allergic effect by blocking IgE-antigen complex formation. *J Biol Chem* 2009;284:32344-52.
- Kanhere A, Hertweck A, Bhatia U, et al. T-bet and GATA3 orchestrate Th1 and Th2 differentiation through lineage-specific targeting of distal regulatory elements. *Nat Commun* 2012;3:1268.
- de Kivit S, Saeland E, Kraneveld AD, et al. Galectin-9 induced by dietary synbiotics is involved in suppression of allergic symptoms in mice and humans. *Allergy* 2012;67:343-52.
- Wells JM, Rossi O, Meijerink M, van Baarlen P. Epithelial crosstalk at the microbiota-mucosal interface. *Proc Natl Acad Sci USA* 2011;108(Suppl 1):4607-14.
- Yang T, Owen JL, Lightfoot YL, Kladder MP, Mohamadzadeh M. Microbiota impact on the epigenetic regulation of colorectal cancer. *Trends Mole Med* 2013;19:714-25.
- Wexler HM. Bacteroides: the good, the bad, and the nitty-gritty. *Clin Microbiol Rev* 2007;20:593-621.
- Henning SM, Yang J, Shao P, et al. Health benefit of vegetable/fruit juice-based diet: Role of microbiome. *J Sci Rep* 2017;7:2167.
- Pellegrini S, Sordi V, Bolla AM, et al. Duodenal mucosa of patients with type 1 diabetes shows distinctive inflammatory profile and microbiota. *J Clin Endocrinol Metab* 2017;102:1468-77.
- Houlden A, Goldrick M, Brough D, et al. Brain injury induces specific changes in the caecal microbiota of mice via altered autonomic activity and mucoprotein production. *Brain Behav Immun* 2016;57:10-20.
- Morowitz MJ, Di Caro V, Pang D, et al. Dietary supplementation with nonfermentable fiber alters the gut microbiota and confers protection in murine models of sepsis. *Crit Care Med* 2017;45:e516-e523.
- Aktas B, De Wolfe TJ, Tandee K, et al. The effect of *Lactobacillus casei* 32G on the mouse cecum microbiota and innate immune response is dose and time dependent. *PLOS ONE* 2015;10:e0145784.
- Krych Ł, Nielsen DS, Hansen AK, Hansen CH. Gut microbial markers are associated with diabetes onset, regulatory imbalance, and IFN- γ level in NOD mice. *Gut Microbes* 2015;6:101-9.
- Wu M, Wu Y, Deng B, et al. Isoliquiritigenin decreases the incidence of colitis-associated colorectal cancer by modulating the intestinal microbiota. *Oncotarget* 2016;7:85318-31.
- Espín JC, Larrosa M, García-Conesa MT, Tomás-Barberán F. Biological significance of urolithins, the gut microbial ellagic acid-derived metabolites: the evidence so far. *Evid Based Complement Alternat Med* 2013;2013:Article ID 270418.
- García-Villalba R, Beltrán D, Espín JC, Selma MV, Tomás-Barberán FA. Time course production of urolithins from ellagic acid by human gut microbiota. *J Agric Food Chem* 2013;61:8797-806.

25. Tang L, Mo Y, Li Y, et al. Urolithin A alleviates myocardial ischemia/reperfusion injury via PI3K/Akt pathway. *Biochem Biophys Res Commun* 2017;486:774-80.
26. Selma MV, González-Sarrías A, Salas-Salvadó J, et al. The gut microbiota metabolism of pomegranate or walnut ellagitannins yields two urolithin-metabotypes that correlate with cardiometabolic risk biomarkers: Comparison between normoweight, overweight-obesity and metabolic syndrome. *Clin Nutr* 2017; S0261-5614(17):30103-6.
27. Duncan SH, Louis P, Flint HJ. Lactate-utilizing bacteria, isolated from human feces, that produce butyrate as a major fermentation product. *Appl Environ Microbiol* 2004;70:5810-7.
28. Wu M, Wu Y, Deng B, et al. Isoliquiritigenin decreases the incidence of colitis-associated colorectal cancer by modulating the intestinal microbiota. *Oncotarget* 2016;7:85318-31.
29. Vuotto C, Longo F, Donelli G. Probiotics to counteract biofilm-associated infections: promising and conflicting data. *Int J Oral Sci* 2014;6:189-94.