



ARTICLE

Effects of Calcium Powder Mixtures and Binding Ingredients as Substitutes for Synthetic Phosphate on the Quality Properties of Ground Pork Products

Min Guk Cho and Jong Youn Jeong*

School of Food Biotechnology & Nutrition, Kyungsung University, Busan 48434, Korea

OPEN ACCESS

Received October 2, 2018
Revised October 17, 2018
Accepted October 18, 2018

*Corresponding author : Jong Youn Jeong
School of Food Biotechnology & Nutrition,
Kyungsung University, Busan 48434, Korea
Tel: +82-51-663-4711
Fax: +82-51-622-4986
E-mail: jeongjy@ks.ac.kr

*ORCID
Min Guk Cho
0000-0002-3205-5837
Jong Youn Jeong
0000-0001-5284-4510

Abstract This study aimed to investigate the combined effect of using natural calcium mixtures and various binding ingredients as replacers for synthetic phosphate in ground pork products. We performed seven treatments: control (0.3% phosphate blend), treatment 1 (0.5% natural calcium mixtures [NCM, which comprised 0.2% oyster shell calcium and 0.3% egg shell calcium powder] and 0.25% egg white powder), treatment 2 (0.5% NCM and 0.25% whey protein concentrate), treatment 3 (0.5% NCM and 0.25% concentrated soybean protein), treatment 4 (0.5% NCM and 0.25% isolated soybean protein), treatment 5 (0.5% NCM and 0.25% carrageenan), and treatment 6 (0.5% NCM and 0.25% collagen powder). All the treatment mixtures had higher pH and lower cooking loss than the control, which was treated with phosphate. We found that NCM and binding ingredients had no negative effects on the moisture content, lightness, and yellowness of the cooked ground pork products. Treatments 3 and 4 showed significantly lower CIE a^* values than the control. Treatments 2 and 6 improved the textural properties of the products. In conclusion, the combination of NCM with whey protein concentrate or collagen powder could be suitable for producing phosphate-free meat products.

Keywords ground pork products, oyster shell powder, egg shell powder, phosphate replacer, ingredients

Introduction

Phosphates are one of the most widely used food additives in poultry and meat products because of their multifunctional properties (Petracci et al., 2013; Sebranek, 2009, Sebranek, 2015). Phosphates have multiple functional abilities such as improving water-holding capacity, increasing pH, chelating on metal ions, improving the freeze-thaw drip and cooking stability, and emulsifying stability. These abilities can improve yield, texture, and sensory properties, as well as extend the shelf life of meat products (Long et al., 2011; McKee and Alvarado, 2004; Sebranek, 2015; Thorainsdottir et al., 2004; Xiong, 2005; Young and Lyon, 1997). Despite these advantages, the use of phosphates in the meat industry is under severe scrutiny due to increasing interest in

healthier foods and the poor consumer perception of phosphates owing to health risks associated with synthetic phosphate (Petracci et al., 2013). However, a direct substitute for phosphates has not been discovered so far (Sindelar, 2015), as it is difficult to cover their multifunctional properties.

Many previous studies have attempted to improve the functionalities of meat products by using various functional ingredients, such as SavorPhos (blends of citrus flour and sodium carbonate) (Casco et al., 2013), dried plums (Jarvis et al., 2012; Jarvis et al., 2015), functional carbohydrates, including guar gum, carrageenan, alginic acid, and chitosan (Park et al., 2008), isolated soy protein (Rhee, 1994), collagen products (Prestes et al., 2013; Schilling et al., 2003), whey protein isolates (Youssef and Barbut, 2011), and modified tapioca starch, wheat bran, and sodium citrate (Ruusunen et al., 2003). Recently, new approaches for identifying phosphate alternatives have been attempted to satisfy the market demand for meat products with natural and clean labels. Casco et al. (2013) reported that marinated chicken breasts with SavorPos (blends of citrus flour and sodium carbonate) had higher injection yield, lower cooking loss, and higher tenderness than those with a commercial phosphate blend. Egg shell calcium and oyster shell calcium are obtained from by-products generated during the processing of egg and oyster products, and widely used as calcium supplements because of their low cost and sustainability (Jang et al., 2010; Kim et al., 2015; Shin and Kim, 1997). These natural calcium powders are generally prepared by pulverization after washing with water and drying or calcining at about 800°C to 1,000°C. Thus, the calcium powders from natural sources have high calcium contents and their main compound is calcium carbonate, which can be distinguished from synthetic calcium products such as calcium carbonate, calcium lactate, and calcium citrate (Kim et al., 2015; Lee and Park, 2002; Shin and Kim, 1997). In previous studies, the utilization of egg shell calcium and oyster calcium has been investigated in meat products to replace a synthetic phosphate. Park (2011b) attempted to evaluate the effects of oyster calcium, as a natural source, and whey protein concentrate on the functional properties of emulsion pork sausages. They found that pork sausages with 0.3% or 0.5% oyster calcium powder had significantly higher water holding capacity and hardness and chewiness values; they suggested that oyster calcium powder could be a suitable substitute for phosphate in the presence of 0.5% whey protein, providing pork sausages of quality similar to those with 0.3% phosphate. Another study compared the efficacy of 4 types of natural calcium (eggshell calcium, oyster shell calcium, marine algae calcium, and milk calcium) as substitutes for synthetic phosphates in meat products (Bae et al., 2017). It was suggested that eggshell calcium could be an ideal phosphate replacer for minimizing cooking loss, likely because of an increase in pH of the products, and oyster shell calcium appeared to be the second-best candidate for a phosphate substitute. In order to determine the optimal ratio of oyster shell and egg shell calcium, Cho et al. (2017) compared the effects of using oyster shell and egg shell calcium (alone and in combination) on the qualities of cooked pork products. They concluded that a combination of 0.2% oyster shell calcium and 0.3% egg shell calcium could replace synthetic phosphate in pork products, but textural properties of the final product were less desirable than the control with synthetic phosphate. Further research has been required if a combination of natural calcium mixtures and food ingredients could be even more effective for improving the quality properties of cooked ground pork products.

Therefore, we investigated the effects of different combinations of premixed natural calcium powders and various binding ingredients on the quality properties of cooked pork products, with the aim of developing high-quality phosphate-free meat products.

Materials and Methods

Preparation and processing of pork products

Raw pork ham (*M. biceps femoris*, *M. semitendinosus*, and *M. semimembranosus*) used in this study was purchased from a local meat processor (Pukyung Pig Farmers Livestock Cooperatives, Korea) at 24–48 h postmortem. After removing subcutaneous and intermuscular fat and visible connective tissues from the fresh ham, lean pork meat was cut in squares of 4–5 cm, vacuum-packaged in nylon/PE film bags, and stored in a freezer at -18°C until processing. Pork back fat was prepared in the same manner. The frozen pork ham and back fat were stored at 2°C – 3°C for 24–36 h before processing. The thawed raw materials were ground in a chopper (TC-22 Elegant plus, Tre Spade, Italy) with an 8-mm plate and then again ground with a 3-mm plate. A phosphate blend (mixtures of sodium tripolyphosphate, tetrasodium pyrophosphate, and sodium hexametaphosphate) (Polymix-CS, SDBNI Co. Ltd., Korea) was used as the synthetic phosphate. The egg shell calcium (egg shell calcium 50, Essentron Co. Ltd., Korea) and oyster shell calcium (oyster shell calcium 40, JK Biochem Co. Ltd., Korea) were purchased from commercial suppliers. The egg shell calcium and oyster calcium powders tested in this study contained 39.6% and 51.6% calcium, respectively. As binding ingredients, egg white powder (Egg white, TOF Co. Ltd., Korea), whey protein concentrate (WPC, TOF Co. Ltd., Korea), concentrated soybean protein (CSP, TOF Co. Ltd., Korea), isolated soybean protein (ISP, Yujin Co. Ltd., Korea), carrageenan (carrageenan WG-5, MSC Co. Ltd.), and collagen powder (collagen powder, Ikjin Co. Ltd., Korea) were also prepared. The formulation of the pork meat products with the calcium powder mixtures and binding ingredients is presented in Table 1.

The control was processed by mixing ground pork meat and back fat with 1.5% NaCl, 0.3% phosphate blend, and 15% ice water in a mixer (KitchenAid 5K5SS, Whirlpool Corp., USA) for 5 min (Fig. 1). For treatments, 1.5% NaCl and 15% ice water were incorporated into the ground pork meat and back fat; samples were then produced by combining 0.5% natural calcium mixtures (NCM, comprising 0.2% oyster shell calcium and 0.3% egg shell calcium powder) with 0.25% egg white

Table 1. Formulations for ground pork meat products with natural calcium mixtures and various binding ingredients

Materials (%)	Treatments ¹⁾						
	Control	Treatment 1	Treatment 2	Treatment 3	Treatment 4	Treatment 5	Treatment 6
Ground pork	68.2	66.75	66.75	66.75	66.75	66.75	66.75
Back fat	15.0	15.00	15.00	15.00	15.00	15.00	15.00
Ice	15.0	15.00	15.00	15.00	15.00	15.00	15.00
Salt	1.5	1.50	1.50	1.50	1.50	1.50	1.50
Phosphate blend	0.3	-	-	-	-	-	-
Oyster shell calcium powder	-	0.20	0.20	0.20	0.20	0.20	0.20
Egg shell calcium powder	-	0.30	0.30	0.30	0.30	0.30	0.30
Egg white powder	-	0.25	-	-	-	-	-
Whey protein concentrate	-	-	0.25	-	-	-	-
Concentrated soybean protein	-	-	-	0.25	-	-	-
Isolated soybean protein	-	-	-	-	0.25	-	-
Carrageenan	-	-	-	-	-	0.25	-
Collagen powder	-	-	-	-	-	-	0.25
Total	100.0	100.00	100.00	100.00	100.00	100.00	100.00

¹⁾ Control, 0.3% phosphate blend added; Treatment 1, 0.5% natural calcium mixtures (NCM; 0.2% oyster shell calcium powder and 0.3% egg shell calcium powder) and 0.25% egg white powder added; Treatment 2, 0.5% NCM and 0.25% whey protein concentrate added; Treatment 3, 0.5% NCM and 0.25% concentrated soybean protein added; Treatment 4, 0.5% NCM and 0.25% isolated soybean protein added; Treatment 5, 0.5% NCM and 0.25% carrageenan added; Treatment 6, 0.5% NCM and 0.25% collagen powder added.




Fig. 1. Schematic diagram of manufacturing process for ground pork meat products with natural calcium powders and various binding ingredients.

powder (treatment 1), whey protein concentrate (treatment 2), concentrated soybean protein (treatment 3), isolated soybean protein (treatment 4), carrageenan (treatment 5), or collagen powder (treatment 6). The meat mixtures from each batch were stuffed into conical tubes (50 g each) using a stuffer (MOD.5/V Deluxe, Tre Spade, Italy). The stuffed tubes were centrifuged at $2,000\times g$ for 10 min in a centrifuge (FELTA5, Hanil Scientific Inc., Korea) to remove air pockets. The tubes were then transferred into a 90°C water bath (CB60L, Dong Won Scientific System, Korea) and cooked to an internal temperature of 75°C . Changes in temperature were monitored by inserting a K-type thermocouple attached to a digital thermometer (TES-1384, Ketch Scientific Instrument Co., Ltd., Taiwan) into the center of the samples. After cooking, the samples were immediately cooled on ice slurry for 20 min and placed at 2°C – 3°C overnight, prior to analyzing dependent variables. All experiments were performed in triplicate.

Determination of pH values

Five grams of the samples was weighed before and after cooking and homogenized with 20 mL of distilled water for 1 min in a homogenizer (DI 25 basic, IKA[®]-Werke GmbH & Co. KG, Germany). The pH was measured using a pH meter (Fisher Scientific Accumet AB150, Thermo Fisher Scientific, Inc., Singapore) equipped with a glass electrode.

Determination of moisture content

Moisture content of the cooked products was measured using the drying method (AOAC, 2007). The samples were dried using a 105°C dry oven (CB60L, Dongwon Scientific System, Korea) for 12 h, and the difference between the weight of each sample before and after drying was calculated.

Determination of cooking loss

As previously described, the weight of each sample in the conical tube was measured prior to cooking and then again after cooking and cooling to determine the cooking loss using the following equation:

$$\text{Cooking loss (\%)} = \frac{\text{Weight before cooking} - \text{Weight after cooking}}{\text{Weight before cooking}} \times 100$$

Instrumental color measurement

Commission Internationale de l'Eclairage (CIE) measurements were used to determine lightness (CIE L*), redness (CIE a*), and yellowness (CIE b*). CIE L*a*b* values were measured on freshly cut surfaces of the cooked samples using a colorimeter (Chroma Meter CR-400, illuminant C; Konica Minolta Sensing Inc., Japan). Two slices from each cooked sample were cut in half (parallel to the longitudinal axis), and three CIE L*a*b* readings per slice were taken immediately after cutting. The colorimeter was standardized against a white calibration plate (CIE L* = +94.90, a* = -0.39, and b* = +3.88).

Texture profile analysis

After cutting the samples to a width of 2.5 cm perpendicular to the longitudinal axis, their hardness, springiness, cohesiveness, gumminess, and chewiness were measured using a texture analyzer (TA-XT2i, Stable Micro System, UK) equipped with a 50-mm-diameter aluminum cylinder (Bourne, 1978). The cross-head speed for measurement was 5 mm/s.

Statistical analysis

All experiments were performed in triplicate. Statistical analysis of the results was performed using the generalized linear model (GLM) procedure of the SAS package program (SAS, 2012). When the analysis of variance showed a significant difference between the means of different samples, the significance of the differences was further analyzed by Duncan's multiple range test ($p < 0.05$).

Results and Discussion

pH values, moisture content, and cooking loss

The pH of meat products can affect their water holding capacity. The incorporation of alkaline phosphates and their blends in meat products can increase the pH, causing unequal positive and negative charges in meat proteins and resulting in higher water holding capacity (Glorieux et al., 2017; Lamkey, 1998; Long et al., 2011). The pH values of ground pork products treated with calcium powder mixtures and various binding ingredients before and after cooking are presented in Table 2. Before cooking, the treatments had higher ($p < 0.05$) pH than the control, although no differences in pH were observed

Table 2. pH values, moisture contents, and cooking loss of ground pork meat products with natural calcium mixtures and various binding ingredients

Traits	Treatments ¹⁾						
	Control (+)	Treatment 1	Treatment 2	Treatment 3	Treatment 4	Treatment 5	Treatment 6
pH (uncooked)	6.37±0.02 ^B	7.47±0.05 ^A	7.53±0.04 ^A	7.56±0.05 ^A	7.80±0.24 ^A	7.68±0.00 ^A	7.82±0.34 ^A
pH (cooked)	6.65±0.01 ^E	7.63±0.01 ^C	7.56±0.01 ^D	7.57±0.01 ^D	7.63±0.01 ^C	7.69±0.01 ^B	7.83±0.02 ^A
Moisture content (%)	68.87±0.09 ^A	69.05±0.19 ^A	69.23±0.24 ^A	68.91±0.04 ^A	69.00±0.13 ^A	69.41±0.07 ^A	69.06±0.10 ^A
Cooking loss (%)	2.79±0.11 ^A	1.66±0.07 ^{CD}	1.41±0.11 ^{DE}	1.36±0.10 ^{DE}	2.21±0.10 ^B	1.80±0.12 ^C	1.15±0.09 ^E

All values are presented as means±standard error of triplicates.

^{A-E} Means within the same row with different superscript letters are significantly different ($p < 0.05$).

¹⁾ Control, 0.3% phosphate blend added; Treatment 1, 0.5% natural calcium mixtures (NCM; 0.2% oyster shell calcium powder and 0.3% egg shell calcium powder) and 0.25% egg white powder added; Treatment 2, 0.5% NCM and 0.25% whey protein concentrate added; Treatment 3, 0.5% NCM and 0.25% concentrated soybean protein added; Treatment 4, 0.5% NCM and 0.25% isolated soybean protein added; Treatment 5, 0.5% NCM and 0.25% carrageenan added; Treatment 6, 0.5% NCM and 0.25% collagen powder added.

($p > 0.05$) among the pork products treated with the different combinations of NCM and binding ingredients. The increase in pH of the treatments was probably due to the NCM added. Cho et al. (2017) had found that the incorporation of egg shell calcium and oyster shell calcium powders to ground pork products can increase the pH. They had also reported that egg shell calcium could more significantly increase the pH than oyster calcium. After cooking, the pH of the control was lower ($p < 0.05$) than that of all the treatments. Among the treatments, treatment 6 had the highest pH, whereas treatments 2 and 3 had the lowest. The pH of treatments 1 and 4 were in between that of the others. According to Prestes et al. (2013), the pH of collagen fiber powder and collagen fiber were 7.78 and 8.05, respectively. Therefore, the high pH of treatment 6 could be attributed to the combined effect of calcium powders and collagen powder. Generally, higher pH in meat results in more water retention (Sebranek, 2009, Sebranek, 2015; Trout and Schmidt, 1983).

Based on the observed pH of the cooked products, it is expected that water retention ability would vary in the different formulations. However, all treatments had similar ($p > 0.05$) moisture contents, compared to the control (Table 2). It has been shown that phosphates and salt in meat products increase the meat protein solubility and ionic strength, thereby contributing to water retention ability (Petracci et al., 2013; Sebranek, 2015; Trout and Schmidt, 1983). Therefore, this result indicated that the absence of synthetic phosphate had no detrimental effect on moisture content, which might have been due to the variation in pH caused by the addition of natural calcium powder and binding ingredients.

The cooking loss of ground pork products treated with calcium powder mixtures and various binding ingredients are shown in Table 2. Interestingly, the cooking loss ranged from 1.15%–2.21% in products with calcium powder mixtures and binding ingredients, which was lower ($p < 0.05$) than that observed in the control (2.79%). In particular, treatments 2, 3, and 6 had the lowest ($p < 0.05$) cooking loss, indicating that calcium powder mixtures coupled with whey protein concentrate, concentrated soybean protein, or collagen powder could be potential phosphate replacers for industrial applications. Bae et al. (2017), after testing eggshell calcium, oyster shell calcium, marine algae calcium, and milk calcium in ground pork products, suggested that eggshell calcium could be an ideal phosphate replacer for minimizing cooking loss. Park et al. (2008) tested guar gum, κ -carrageenan, alginic acid, and chitosan to identify a suitable substitute for inorganic polyphosphate in the pork model system. They reported that pork samples treated with guar gum and κ -carrageenan showed lower cooking loss than those treated with polyphosphate.

Instrumental color

Table 3 shows the instrumental color values of ground pork products treated with calcium powder mixtures and various

Table 3. Instrumental color values of ground pork meat products formulated with natural calcium mixtures and various binding ingredients

Traits	Treatments ¹⁾						
	Control (+)	Treatment 1	Treatment 2	Treatment 3	Treatment 4	Treatment 5	Treatment 6
CIE L*	69.02±0.22 ^A	69.69±0.29 ^A	69.44±0.18 ^A	69.14±0.29 ^A	69.48±0.28 ^A	69.46±0.12 ^A	69.18±0.06 ^A
CIE a*	5.95±0.11 ^A	5.54±0.14 ^{AB}	5.55±0.12 ^{AB}	5.21±0.19 ^B	5.21±0.16 ^B	5.39±0.40 ^{AB}	5.44±0.07 ^{AB}
CIE b*	8.18±0.09 ^A	8.21±0.11 ^A	7.98±0.24 ^A	8.31±0.24 ^A	8.29±0.10 ^A	8.25±0.35 ^A	8.49±0.09 ^A

All values are presented as means±standard error of triplicates.

^{A,B} Means within the same row with different superscript letters are significantly different ($p < 0.05$).

¹⁾ Control, 0.3% phosphate blend added; Treatment 1, 0.5% natural calcium mixtures (NCM; 0.2% oyster shell calcium powder and 0.3% egg shell calcium powder) and 0.25% egg white powder added; Treatment 2, 0.5% NCM and 0.25% whey protein concentrate added; Treatment 3, 0.5% NCM and 0.25% concentrated soybean protein added; Treatment 4, 0.5% NCM and 0.25% isolated soybean protein added; Treatment 5, 0.5% NCM and 0.25% carrageenan added; Treatment 6, 0.5% NCM and 0.25% collagen powder added.

binding ingredients. We found no differences ($p > 0.05$) in CIE L* values between the control and the treatments. In addition, all treatments with NCM and binding ingredients had similar ($p > 0.05$) CIE L* values. CIE a* values of treatments 1, 2, 5, and 6 were not different ($p > 0.05$) from the control, but those of treatments 3 and 4 were lower ($p < 0.05$) than that of the control. Generally, cooked meat products with higher pH are darker and redder in color (Barbut, 2009; Casco et al., 2013; Trout, 1989). However, treatment with 0.5% NCM and 0.25% binding ingredients had no effect on the CIE L* value in this study. The lighter color of the binding ingredients likely contributed to lighter products by reducing the effect of pH (Hughes et al., 1998; Youssef and Barbut, 2011). The low CIE a* values in treatments 3 and 4 could be attributed to the inherent color of the soybean products used in this study. This result is consistent with that of Park (2011a), who reported that redness decreased with the addition of concentrated soybean protein in cooked pork patties. We found no differences ($p > 0.05$) in CIE b* values between the control and treatments. These findings are consistent with the results of Cho et al. (2017), who demonstrated that cooked pork products treated with a combination of oyster shell calcium and egg shell calcium powder had similar CIE b* values.

Textural properties

Table 4 shows the textural properties of ground pork products treated with calcium powder mixtures and various binding ingredients. Among the samples tested in this study, the control had hardness similar ($p > 0.05$) to treatments 2 and 6. Choi

Table 4. Textural properties of ground pork meat products formulated with natural calcium mixtures and various binding ingredients

Traits	Treatments ¹⁾						
	Control (+)	Treatment 1	Treatment 2	Treatment 3	Treatment 4	Treatment 5	Treatment 6
Hardness (kg)	5.74±0.17 ^A	4.82±0.15 ^C	5.33±0.16 ^{ABC}	4.89±0.23 ^C	4.99±0.22 ^C	5.04±0.22 ^{BC}	5.59±0.17 ^{AB}
Springiness	0.93±0.00 ^A	0.93±0.00 ^A	0.93±0.00 ^A	0.94±0.00 ^A	0.93±0.00 ^A	0.93±0.00 ^A	0.93±0.00 ^A
Cohesiveness	0.73±0.00 ^A	0.74±0.00 ^A	0.73±0.00 ^A	0.73±0.00 ^A	0.74±0.00 ^A	0.74±0.00 ^A	0.74±0.00 ^A
Gumminess (kg)	4.18±0.13 ^A	3.55±0.12 ^B	3.87±0.11 ^{AB}	3.59±0.16 ^B	3.69±0.15 ^B	3.75±0.15 ^B	4.16±0.12 ^A
Chewiness (kg)	3.91±0.12 ^A	3.30±0.10 ^C	3.60±0.11 ^C	3.36±0.15 ^C	3.44±0.14 ^C	3.48±0.14 ^{BC}	3.86±0.11 ^{AB}

All values are presented as means±standard error of triplicates.

^{A-C} Means within the same row with different superscript letters are significantly different ($p < 0.05$).

¹⁾ Control, 0.3% phosphate blend added; Treatment 1, 0.5% natural calcium mixtures (NCM; 0.2% oyster shell calcium powder and 0.3% egg shell calcium powder) and 0.25% egg white powder added; Treatment 2, 0.5% NCM and 0.25% whey protein concentrate added; Treatment 3, 0.5% NCM and 0.25% concentrated soybean protein added; Treatment 4, 0.5% NCM and 0.25% isolated soybean protein added; Treatment 5, 0.5% NCM and 0.25% carrageenan added; Treatment 6, 0.5% NCM and 0.25% collagen powder added.

et al. (2014) obtained similar results by adding 0.5% oyster calcium and 0.5% whey protein to restructured pork ham, and Meullenet et al. (1994) reported similar results for chicken frankfurter treated with collagen fibers. The hardness of treatments 1, 3, 4, and 5 was lower ($p < 0.05$) than that of the control. However, springiness and cohesiveness were not different between the control and treatments (Table 4). Cho et al. (2017) reported that springiness and cohesiveness of pork products treated with 0.5% oyster shell calcium or a combination of 0.3% oyster shell calcium and 0.2% eggshell calcium powder did not differ significantly from those of the control treated with synthetic phosphate blend, which is consistent with our findings. The control had higher ($p < 0.05$) gumminess than all the treatments except treatments 2 and 6, which had a similar ($p > 0.05$) gumminess as the control. Chewiness was higher ($p < 0.05$) in the control than in treatments 1, 2, 3, 4, and 5, but no differences ($p > 0.05$) in chewiness were found between the control and treatment 6. Previous studies have reported higher gumminess and chewiness in meat products with phosphate. Bae et al. (2017) reported that hardness, gumminess, and chewiness were significantly higher in the control samples with synthetic phosphate than in samples treated with natural calcium powders. Cho et al. (2017) found that gumminess and chewiness were the highest in pork products containing a phosphate blend and that a combination of 0.2% oyster shell and 0.3% egg shell calcium improved gumminess and chewiness, which may complement the textural properties of pork products. However, in this study, the combined use of calcium powders and whey protein concentrate or collagen powder improved the textural properties of pork products (Table 4). Lee et al. (2011) had reported that the addition of oyster shell powder (0.15%, 0.3%, and 0.5%) and 0.5% whey protein to emulsion sausages significantly increased their hardness, springiness, and chewiness, compared with sausages with or without 0.3% sodium tripolyphosphate, which was partially consistent with the results of our study.

Conclusions

In this study, NCM and various non-meat ingredients were added to ground pork products to evaluate potential substitutes for synthetic phosphate. NCM and the tested ingredients increased the water holding capacity (lowering the cooking loss), possibly by increasing the pH of the pork products. In addition, no negative effects were observed regarding the moisture content and instrumental color of the final products, although treatments 3 and 4, which contained soybean protein, had lower CIE a^* values than the control. Compared to the control (containing synthetic phosphate blend), treatments 2 and 6 had a notable effect on the textural properties of the ground pork products. Therefore, the combination of NCM and whey protein concentrate or collagen powder could be ideal for producing phosphate-free meat products. Further research is required for determining the optimal proportions of the combinations for industrial application.

Conflicts of Interest

The authors declare no potential conflict of interest.

Acknowledgments

This research was supported by Kyungsoo University Research Grants in 2017.

References

AOAC. 2007. Official methods of analysis. 18th ed. Association of Official Analytical Chemists. Washington, DC, USA. p

931.

- Bae SM, Cho MG, Jeong JY. 2017. Effects of various calcium powders as replacers for synthetic phosphate on the quality properties of ground pork meat products. *Korean J Food Sci An* 37:456-463.
- Barbut S. 2009. Pale, soft, and exudative poultry meat—Reviewing ways to manage at the processing plant. *Poult Sci* 88:1506-1512.
- Bourne MC. 1978. Texture profile analysis. *Food Technol* 32:62-66, 72.
- Casco G, Veluz GA, Alvarado CZ. 2013. SavorPhos as an all-natural phosphate replacer in water- and oil-based marinades for rotisserie birds and boneless-skinless breast. *Poult Sci* 82:3236-3243.
- Cho MG, Bae SM, Jeong JY. 2017. Egg shell and oyster shell powder as alternatives for synthetic phosphate: Effects on the quality of cooked ground pork products. *Korean J Food Sci An* 37:571-578.
- Choi JS, Lee HJ, Jin SK, Lee HJ, Choi YI. 2014. Effect of oyster shell calcium powder on the quality of restructured pork ham. *Korean J Food Sci An* 34:372-377.
- Glorieux S, Goemaere O, Steen L, Fraeye I. 2017. Phosphate reduction in emulsified meat products: Impact of phosphate type and dosage on quality characteristics. *Food Technol Biotechnol* 55:390-397.
- Hughes E, Mullen AM, Troy DJ. 1998. Effect of fat level, tapioca starch and whey protein on frankfurters formulated with 5% and 12% fat. *Meat Sci* 48:169-180.
- Jang SY, Jeong YJ, Kwon TK, Seo JH. 2010. Effects of water-soluble calcium supplements made from eggshells and oyster shells on the calcium metabolism of growing rats. *J Food Sci Nutr* 15:78-82.
- Jarvis N, Clement AR, O'Bryan CA, Babu D, Crandall PG, Owens CM, Meullenet JF, Ricke SC. 2012. Dried plum products as a substitute for phosphate in chicken marinade. *J Food Sci* 77:S253-S257.
- Jarvis N, O'Bryan C, Ricke SC, Crandall PG. 2015. The functionality of plum ingredients in meat products: A review. *Meat Sci* 102:41-48.
- Kim JS, Jung NY, Jang SJ, Lee HJ, Park SH, Kim MJ, Heu MS. 2015. Characteristics of the shells and calcined powders from the butter clam *Saxidomus purpuratus* and littleneck clam *Ruditapes philippinarum* as a natural calcium resource. *Korean J Fish Aquat Sci* 48:168-177.
- Lamkey JW. 1998. Non-meat ingredients for meat processing. 51st Annual Reciprocal Meat Conference, Storrs, CT, USA. pp 48-52.
- Lee JJ, Park SH, Choi JS, Kim JH, Lee SH, Choi SH, Choi YI, Jung DS. 2011. Effect of oyster shell powder on quality properties and storage stability of emulsion-type pork sausages. *Korean J Food Sci An* 31:469-476.
- Lee SK, Park JH. 2002. Studies of egg-shell calcium (I). The effects of elution condition of egg-shell calcium on elution quantity and ionization rate. *J Food Hyg Saf* 17:173-187.
- Long NHBS, Gál R, Buňka F. 2011. Use of phosphate in meat products. *Afr J Biotechnol* 10:19874-19882.
- McKee S, Alvarado C. 2004. Marination and brining of poultry products: Value added. 57th Annual Reciprocal Meat Conference, Lexington, KY, USA, pp 129-135.
- Meullenet JF, Chang HC, Carpenter JA, Resurreccion AVA. 1994. Textural properties of chicken Frankfurters with added collagen fibers. *J Food Sci* 59:729-733.
- Park JW. 2011a. A study on the quality property of pork patties supplemented by concentrated soy protein. M.S. thesis, Chung-Ang Univ., Anseong, Korea.
- Park KS, Choi YI, Lee SH, Kim CH, Auh JH. 2008. Application of functional carbohydrates as a substitute for inorganic

- polyphosphate in pork meat processing. *Korean J Food Sci Technol* 40:118-121.
- Park SH. 2011b. Effects of natural calcium powder addition on functional properties and storage characteristics of emulsion-type sausages. M.S. thesis, Chungbuk National Univ., Cheongju, Korea.
- Petracci M, Bianchi M, Mudalal S, Cavani C. 2013. Functional ingredients for poultry meat products. *Trends Food Sci Technol* 33:27-39.
- Prestes RC, Graboski A, Roman SS, Kempka AP, Toniazzo G, Demiate IM, Luccino MD. 2013. Effects of the addition of collagen and degree of comminution in the quality of chicken ham. *J Appl Poult Res* 22:885-903.
- Rhee KC. 1994. Functionality of soy proteins. In *Protein functionality in food systems*. Hettiarachchy NS, Ziegler GR (ed). Marcel Dekker, New York, NY, USA. pp 311-324.
- Ruusunen M, Vainionpää J, Puolanne E, Lyly M, Lähteenmäki L, Niemistö M, Ahvenainen R. 2003. Physical and sensory properties of low-salt phosphate-free frankfurters composed with various ingredients. *Meat Sci* 63:9-16.
- SAS. 2012. SAS/STAT® Software for PC. Release 9.4 version, SAS Institute Inc., Cary, NC, USA.
- Schilling MW, Mink LE, Gochenour PS, Marriott NG, Alvarado CZ. 2003. Utilization of pork collagen for functionality improvement of boneless cured ham manufactured from pale, soft, and exudative pork. *Meat Sci* 65:547-553.
- Sebranek JG. 2009. Basic curing ingredients. In *Ingredients in meat products*. Tarté R (ed). Springer Science+Business Media LLC, New York, NY, USA. pp 1-24.
- Sebranek JG. 2015. An overview of functional non-meat ingredients in meat processing: The current toolbox. 68th Annual Reciprocal Meat Conference, Lincoln, NE, USA. pp 42-46.
- Shin HS, Kim KH. 1997. Preparation of calcium powder from eggshell and use of organic acids for enhancement of calcium ionization. *Appl Biol Chem* 40:531-535.
- Sindelar JJ. 2015. Impact of removing functional non-meat ingredients in processed meat products and exploring possible alternatives. 68th Annual Reciprocal Meat Conference, Lincoln, NE, USA. pp 48-51.
- Thorainsdottir KA, Gudmundsdottir G, Arason S, Thorkelsson G, Kristbergsson K. 2004. Effects of added salt phosphate, and proteins on the chemical and physicochemical characteristics of frozen cod (*Gadus morhua*) fillets. *J Food Sci* 69:SNQ144-SNQ152.
- Trout GR, Schmidt GR. 1983. Utilization of phosphates in meat products. 36th Annual Reciprocal Meat Conference, Fargo, ND, USA. pp 24-27.
- Trout GR. 1989. Variation in myoglobin denaturation and color of cooked beef, pork, and turkey meat as influenced by pH, sodium chloride, sodium tripolyphosphate, and cooking temperature. *J Food Sci* 54:536-540.
- Xiong Y. 2005. Role of myofibrillar proteins in water-binding in brine enhanced meats. *Food Res Int* 38:281-287.
- Young LL, Lyon CE. 1997. Effect of postchill aging and sodium tripolyphosphate on moisture binding properties, color, and Warner-Bratzler shear values of chicken breast meat. *Poult Sci* 76:1587-1590.
- Youssef MK, Barbut S. 2011. Effects of two types of soy protein isolates, native and preheated whey protein isolates on emulsified meat batters prepared at different protein levels. *Meat Sci* 87:54-60.