



Is Osteoporosis a Common Comorbidity in Different Chronic Airway Diseases?

Seung Hyeun Lee, M.D., Ph.D.

Division of Respiratory and Critical Care Medicine, Department of Internal Medicine, Kyung Hee University Hospital, Kyung Hee University School of Medicine, Seoul, Korea

Although a recent genome-wide association study to prove “Dutch hypothesis,” meaning that both asthma and chronic obstructive pulmonary disease (COPD) share a genetic background, has failed to show a common genetic component between the two¹, considerable number of patients have clinical features of both asthma and COPD²⁻⁴. Some patients with COPD may have eosinophilic airway inflammation or bronchodilator response and some asthmatics may have neutrophilic airway inflammation or fixed airflow limitation. Such condition has been termed asthma-COPD overlap syndrome. However, the use of “syndrome” in asthma-COPD overlap (ACO) is now being challenged because such overlap includes several different phenotypes caused by various underlying mechanisms rather than a single disease entity⁵.

Nevertheless, recognition and diagnosis of ACO are important as ACO has different clinical features compared to asthma or COPD alone. More frequent exacerbation, poorer quality of life, more rapid decline in lung function, and more utilization of healthcare resources have been reported in ACO than in traditional chronic airway diseases⁶⁻⁸. Epidemiologic studies have reported that the prevalence of ACO ranges from 15% to 55% depending on criteria used by different investigators^{9,10}. Recent studies have reported a relatively consistent prevalence rate; approximately 15% of patients with COPD are

categorized as ACO both in a large Spanish cohort and COPD Gene consortium^{11,12}. It has been shown that the prevalence of ACO is increased with age.²

Less is known about comorbidities of ACO. In this issue of *Tuberculosis and Respiratory Diseases*, Oh et al.¹³ reported results of a retrospective cross-sectional study of 321 patients. They investigated the prevalence of osteoporosis or bone mineral density (BMD) in patients with asthma, COPD, and ACO¹³. Although they failed to demonstrate different prevalence of osteoporosis among three chronic airway diseases, they found that patients with ACO had significantly lower BMD compared to patients with asthma or COPD alone. In addition, they showed that the use of inhaled corticosteroid (ICS) was not associated with low BMD in their population by multivariate analysis¹³.

Regarding comorbidity in ACO, van Boven et al.¹⁴ have reported comprehensive results about its pattern and clinical impact on hospitalization risk using a large Spanish cohort. They showed that allergic rhinitis (odds ratio [OR], 1.81), anxiety (OR, 1.18), and gastroesophageal reflux disease (OR, 1.18) were more prevalent in patients with ACO compared to those in patients with COPD. Moreover, osteoporosis was also more frequent in ACO (OR, 1.14). Interestingly, they reported that chronic kidney disease (OR, 0.79) and ischemic heart disease (OR, 0.88) were less frequent in ACO while cardiovascular diseases were associated with hospitalization risk of patients with ACO¹⁴. Using 2009 Korean National Health Insurance database, Rhee et al.³ have also reported that a variety of comorbidities including diabetes, hypertension, ischemic heart disease, depression, and osteoporosis are more frequent in patients with ACO compared to those in patients with COPD without asthma. Although the study of Oh et al.¹³ could not find significant higher prevalence of osteoporosis in patients with ACO, their results suggest that lower BMD in patients with ACO compared to patients with asthma or COPD alone could be a characteristic of the overlap, consistent with results of two aforementioned studies^{3,14}.

Interestingly, in this particular study, the use of ICS was not associated with low BMD. Although the association between ICS use and prevalence of osteoporosis remains controversial,

Address for correspondence: Seung Hyeun Lee, M.D., Ph.D.

Division of Respiratory and Critical Care Medicine, Department of Internal Medicine, Kyung Hee University Hospital, Kyung Hee University School of Medicine, 23 Kyungheedaero, Dongdaemun-gu, Seoul 02447, Korea

Phone: 82-2-958-8511, **Fax:** 82-2-968-1848

E-mail: humanmd04@hanmail.net

Received: Oct. 25, 2017

Revised: Nov. 1, 2017

Accepted: Nov. 6, 2017

Published online: Dec. 13, 2017

©It is identical to the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>).



Copyright © 2018
The Korean Academy of Tuberculosis and Respiratory Diseases.
All rights reserved.

evidences suggest that using ICS for treating chronic airway diseases may not be harmful or even beneficial from the viewpoint of osteoporosis. Two systematic reviews have demonstrated that long-term ICS use does not have a significant impact on bone mineral density^{15,16}. Moreover, a recent study has reported that the incidence of osteoporosis in ICS users is significantly lower than that in non-ICS users (hazard ratio, 0.73) in female patients with COPD¹⁷. In that study, higher ICS dose is associated with lower risk of osteoporosis (p for trend=0.0023) after adjusting for age, income, and medications¹⁷. How can we explain these phenomena? First, ICS can reduce systemic inflammation which might be associated with increased osteoporosis in chronic airway diseases^{18,19}. Evidences support that pro-inflammatory cytokines including tumor necrosis factor- α and interleukin-6 are associated with increased risk of osteoporosis^{7,20}. Second, regular use of ICS can prevent osteoporosis by preventing acute exacerbation of COPD characterized by surge of both pulmonary and systemic inflammation. A previous study has demonstrated that acute exacerbation of COPD is independently associated with progression of osteoporosis²¹. Third, improved symptoms, lung function, and quality of life by regular use of ICS may prevent immobility-associated bone loss via increased physical activity.

Although a relatively small number of ACO patients were included, this particular study provides insight into comorbidity among different chronic airway diseases¹³. Future investigations should encompass the impact of treatment including ICS on pro-inflammatory cytokines or bone turnover markers and on the development or progression of osteoporosis as well as influence of osteoporosis to the risk of fracture and quality of life in patients with ACO using a large population.

ACO is still an area of unknown. Its pathophysiology, biomarker, clinical characteristics should be elucidated. Identifying ACO is important so that appropriate treatment can be selected. Unfortunately, there is scant information on the optimal treatment for ACO because most previous large clinical trials on asthma or COPD have excluded ACO cases. Although somewhat late, efforts to find potentially beneficial new drugs, ranging from small kinase blocker to monoclonal antibodies for this overlap, are now ongoing²². Recent positive results on mepolizumab, a monoclonal antibody against interleukin-5 for patients with eosinophilic COPD, are shedding light on those efforts²³.

Conflicts of Interest

No potential conflict of interest relevant to this article was reported.

References

- Smolonska J, Koppelman GH, Wijmenga C, Vonk JM, Zanen P, Bruinenberg M, et al. Common genes underlying asthma and COPD? Genome-wide analysis on the Dutch hypothesis. *Eur Respir J* 2014;44:860-72.
- Pleasant RA, Ohar JA, Croft JB, Liu Y, Kraft M, Mannino DM, et al. Chronic obstructive pulmonary disease and asthma-patient characteristics and health impairment. *COPD* 2014;11:256-66.
- Rhee CK, Yoon HK, Yoo KH, Kim YS, Lee SW, Park YB, et al. Medical utilization and cost in patients with overlap syndrome of chronic obstructive pulmonary disease and asthma. *COPD* 2014;11:163-70.
- Sin DD. Asthma-COPD overlap syndrome: what we know and what we don't. *Tuberc Respir Dis* 2017;80:11-20.
- GOLD 2017 Global strategy for the diagnosis, management and prevention of COPD [Internet]. Gaithersburg: Global Initiative for Chronic Obstructive Lung Disease; 2017 [cited 2017 Nov 11]. Available from: <http://goldcopd.org/gold-2017-global-strategy-diagnosis-management-prevention-copd/>.
- Alshabanat A, Zafari Z, Albanyan O, Dairi M, FitzGerald JM. Asthma and COPD overlap syndrome (ACOS): a systematic review and meta analysis. *PLoS One* 2015;10:e0136065.
- Liang B, Feng Y. The association of low bone mineral density with systemic inflammation in clinically stable COPD. *Endocrine* 2012;42:190-5.
- Kang HK, Kim K, Lee H, Jeong BH, Koh WJ, Park HY. COPD assessment test score and serum C-reactive protein levels in stable COPD patients. *Int J Chron Obstruct Pulmon Dis* 2016;11:3137-43.
- Kauppi P, Kupiainen H, Lindqvist A, Tammilehto L, Kilpelainen M, Kinnula VL, et al. Overlap syndrome of asthma and COPD predicts low quality of life. *J Asthma* 2011;48:279-85.
- Weatherall M, Travers J, Shirtcliffe PM, Marsh SE, Williams MV, Nowitz MR, et al. Distinct clinical phenotypes of airways disease defined by cluster analysis. *Eur Respir J* 2009;34:812-8.
- Gibson PG, Simpson JL. The overlap syndrome of asthma and COPD: what are its features and how important is it? *Thorax* 2009;64:728-35.
- Almagro P, Cabrera FJ, Diez J, Boixeda R, Alonso Ortiz MB, Murio C, et al. Comorbidities and short-term prognosis in patients hospitalized for acute exacerbation of COPD: the EPOC en Servicios de medicina interna (ESMI) study. *Chest* 2012;142:1126-33.
- Oh JY, Lee YS, Min KH, Lee SY, Shim JJ, Kang KH, et al. Osteoporosis in patients with asthma-chronic obstructive pulmonary disease overlap syndrome. *Tuberc Respir Dis* 2018;81:73-9.
- van Boven JF, Roman-Rodriguez M, Palmer JF, Toledo-Pons N, Cosio BG, Soriano JB. Comorbidity, pattern, and im-

- pact of asthma-COPD overlap syndrome in real life. *Chest* 2016;149:1011-20.
15. Halpern MT, Schmier JK, Van Kerkhove MD, Watkins M, Kalberg CJ. Impact of long-term inhaled corticosteroid therapy on bone mineral density: results of a meta-analysis. *Ann Allergy Asthma Immunol* 2004;92:201-7.
 16. Jones A, Fay JK, Burr M, Stone M, Hood K, Roberts G. Inhaled corticosteroid effects on bone metabolism in asthma and mild chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2002;(1):CD003537.
 17. Liu SF, Kuo HC, Liu GH, Ho SC, Chang HC, Huang HT, et al. Inhaled corticosteroids can reduce osteoporosis in female patients with COPD. *Int J Chron Obstruct Pulmon Dis* 2016;11:1607-14.
 18. Sin DD, Lacy P, York E, Man SF. Effects of fluticasone on systemic markers of inflammation in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2004;170:760-5.
 19. Ozol D, Aysan T, Solak ZA, Mogulkoc N, Veral A, Sebik F. The effect of inhaled corticosteroids on bronchoalveolar lavage cells and IL-8 levels in stable COPD patients. *Respir Med* 2005;99:1494-500.
 20. Omoigui S. The Interleukin-6 inflammation pathway from cholesterol to aging: role of statins, bisphosphonates and plant polyphenols in aging and age-related diseases. *Immun Ageing* 2007;4:1.
 21. Kiyokawa H, Muro S, Oguma T, Sato S, Tanabe N, Takahashi T, et al. Impact of COPD exacerbations on osteoporosis assessed by chest CT scan. *COPD* 2012;9:235-42.
 22. Barnes PJ. Therapeutic approaches to asthma-chronic obstructive pulmonary disease overlap syndromes. *J Allergy Clin Immunol* 2015;136:531-45.
 23. Gauderman WJ, Urman R, Avol E, Berhane K, McConnell R, Rappaport E, et al. Association of improved air quality with lung development in children. *N Engl J Med* 2015;372:905-13.