

Influence of dietary nonstructural carbohydrate concentration on growth performance and carcass characteristics of Holstein steers

Daniel Ramos-Aviña¹, Alejandro Plascencia^{1,*}, and Richard Zinn²

* **Corresponding Author:** Alejandro Plascencia
Tel: +52-686-5636906 (111), **Fax:** +52-686-5636907,
E-mail: aplas_99@yahoo.com

¹ Instituto de Investigaciones en Ciencias Veterinarias,
Universidad Autónoma de Baja California, Mexicali
21386, México

² Department of Animal Science, University of
California, Davis, CA 95616, USA

ORCID

Daniel Ramos-Aviña
<https://orcid.org/0000-0001-7495-5449>
Alejandro Plascencia
<https://orcid.org/0000-0003-2151-5372>
Richard Zinn
<https://orcid.org/0000-0003-3014-3814>

Submitted Jun 1, 2017; Revised Aug 31, 2017;
Accepted Oct 23, 2017

Objective: Since very little information exists about the topic; in this experiment we compare, in a long-term finishing program, the growth-performance responses and carcass characteristics of Holstein steers where non-structural carbohydrate concentration of the diet is reduced from 64% to 51% (dry matter basis).

Methods: Sixty Holstein steer calves (129±2.2 kg) were blocked by initial weight into five groups and randomly assigned within weight groupings to 10 pens. Calves were fed with a steam-flaked corn-based finishing diets containing 51% higher fiber (HF) or 64% lower fiber (LF) nonstructural carbohydrates. Non-structural carbohydrates concentrations were manipulated substituting dried distiller grain with solubles and alfalfa hay for flaked corn. Cattle were weighed every 112 days and at the end of the experiment (day 308) when the cattle were harvested and carcass characteristics were evaluated.

Results: Steers fed the HF diet showed improvement (8.8%) in average daily gain (ADG) during the initial 112-d period. This effect was followed by a numerical trend for greater ADG throughout the remainder of the study so that overall ADG tended to be greater (4.9%, $p = 0.06$) for the HF than for LF. There were no treatment effects on dry matter intake. Gain efficiency and estimated dietary net energy (NE) were greater 8.3% and 5.2%, respectively for HF during the initial 112-d period. Overall (308-d) gain efficiency and estimated dietary NE were similar for both dietary treatments. However, due to differences in tabular dietary NE, the ratio of observed:expected dietary NE tended to be greater (4.1%, $p = 0.06$) for the HF vs LF diet. There were no treatment effects on carcass characteristics except for a tendency toward a slightly greater (0.5%, $p = 0.09$) estimated carcass yield.

Conclusion: Reducing the non-structural carbohydrate concentration of a conventional steam-flaked corn-based growing finishing diet for Holstein steers can effectively enhance growth performance, particularly during the early growing and late finishing phases.

Keywords: Holstein; Finishing; Diet Energy Density; Performance; Carcass

INTRODUCTION

Holstein steers calves grow faster than beef steers at comparable weights [1], capable of doubling their daily live weight during the first 90 to 100 d in the feedlot [2,3]. To fully express this growth potential, receiving and growing diets for the calf-fed Holsteins are energy dense [4,5]. Due to their light initial weight upon entering the feedlot (~130 kg) and heavy mature final weight (>575 kg), the duration of the growing-finishing period of calf-fed Holsteins is protracted, exceeding 300 days [6,7]. Protracted feeding of energy-dense diets (characteristically high in starch and low in forage), increases individual risk of digestive upset and overall growth-performance [8]. This effect is magnified in Holsteins due to their genetically enhanced gut capacity, potentiating wider swings in voluntary feed intake and associated incidence of metabolic death, bloat, and liver abscess than observed in conventional beef breeds [6]. Conventional steam-flaked corn-based growing finishing diets fed to Holstein

steers in the Southwestern USA are high in readily fermentable non-structural carbohydrate (64% to 66% dry matter [DM] basis). Decreasing the level of non-structural carbohydrate may reduce the risk of digestive dysfunctions. However, the alternative of increasing dietary forage and reducing dietary starch in growing finishing diets may, itself, limit energy intake and hence, increase the risk for decreased growth-performance [9,10]. To our knowledge, very little, if any information has been reported that evaluates the impact of feeding decreased levels of non-fiber carbohydrate across the entirety of the growing finishing phases on feedlot performance of calf-fed Holstein steers. In this study we compare the growth-performance responses and carcass characteristics of calf-fed Holstein steers where non-structural carbohydrate concentration of the diet is reduced from 64% to 51% (DM basis).

MATERIALS AND METHODS

All procedures involving animal care and management were in accordance with and approved by the University of California, Davis, Animal Use and Care Committee.

Animals, feeding, treatments, and carcass measurements

Sixty Holstein steer calves (129 ± 2.2 kg initial weight) were utilized to evaluate the effects of dietary forage level on growth performance, dietary energetics, and carcass characteristics. The number of pen replicates (5) and animals (30) within treatments are enough to determine statistical differences on growth performance of feedlot cattle. Indeed, four pens replicates would have been sufficient. Based on power analysis and standard deviation for measure, we had a power of 0.955 for detecting a 5% difference in gain efficiency. Upon arrival at the University of California Desert Research Center (El Centro, CA, USA), steer calves were vaccinated against IBR, BVD (type 1 and 2), PI3, BRSV (Cattle Master Gold FP 5 L5, Zoetis, New York, NY, USA), clostridia (Ultrabac 8, Zoetis, New York, NY, USA), treated against internal and external parasites (Dectomax, Zoetis, New York, NY, USA), injected with 1,500 IU vitamin E (as d-alpha-tocopherol) 500,000 IU vitamin A (as retinyl-palmitate) and 50,000 IU vitamin D₃ (Vital E-AD, Stuart Products, Bedford, TX, USA), and 300 mg tulathromycin (Draxxin, Zoetis, USA). Steers calves were blocked by initial shrunk (off truck) weight into five groups and randomly assigned within weight groupings to 10 pens (six steers per pen). Pens were 62 m² with 25 m² overhead shade, automatic waterers and 2.4 m fence-line feed bunks. Treatments consisted in a steam-flaked corn-based finishing diets containing 51% higher fiber (HF) vs 64% lower fiber (LF) nonstructural carbohydrates (Table 1). The total nonstructural carbohydrate concentration of the diets was manipulated using dried distiller grain plus solubles and forage (alfalfa hay). Accordingly, nonstructural carbohydrates and neutral deter-

Table 1. Composition of experimental diets

Item	Nonstructural carbohydrates (%)	
	51	64
Diet designation	Higher fiber (HF)	Lower fiber (LF)
Ingredient composition (% DM)		
Alfalfa hay	14.00	6.00
Sudangrass hay	6.00	6.00
Steam-flaked corn	54.65	76.23
Dried distiller grain with solubles	15.00	
Yellow grease	2.00	2.00
Cane molasses	6.00	6.00
Limestone	1.00	1.27
Urea	0.50	1.30
Trace mineral salt ¹⁾	0.40	0.40
Magnesium oxide	0.15	0.15
Dicalcium phosphate	0.30	0.65
Nutrient composition, DM basis ²⁾		
Net energy (Mcal/kg)		
Maintenance	2.11	2.19
Gain	1.45	1.53
Crude protein (%)	14.6	12.9
Rumen undegradable protein (%)	36.3	34.5
Rumen degradable protein (%)	63.7	65.5
Nonstructural carbohydrates (%)	51.3	64.2
Neutral detergent fiber (%)	21.7	13.3
Ether extract (%)	6.39	5.54
Calcium (%)	0.81	0.80
Phosphorous (%)	0.41	0.40

DM, dry matter.

¹⁾ Trace mineral salt contained: CoSO₄, 0.068%; CuSO₄, 1.04%; FeSO₄, 3.57%; ZnO, 0.75% MnSO₄, 1.07%; KI, 0.052%; NaCl, 93.4%.

²⁾ Based on tabular values for individual feed ingredients [22].

gent fiber (NDF) concentrations were 51.3% and 64.2%, and 21.7% and 13.3% to HF and LF, respectively. Steers were allowed *ad libitum* access to feed and water. Fresh feed was provided twice daily at 06:00 and 14:00 h, offering approximately 30% of daily consumption in the morning feeding and the remainder in the afternoon feeding. Feed and refusal samples were collected daily for DM analysis, which involved oven drying the samples at 105°C until no further weight loss occurred (method 930.15 [11]). The experiment lasted 308 days.

Hot carcass weights (HCW) were obtained at time of slaughter. After carcasses chilled for 24 h, the following measurements were obtained: Longissimus muscle (LM) area (cm²) by direct grid reading of the muscle at the 12th rib; subcutaneous fat (cm) over the LM at the 12th rib taken at a location 3/4 the lateral length from the chine bone end (adjusted by eye for unusual fat distribution); kidney, pelvic and heart fat (KPH) as a percentage of HCW; marbling score ([12] using 3.0 as minimum slight, 4.0 as minimum small, 5.0 as minimum modest, 6.0 as minimum moderate, etc.), and estimated retail yield of boneless, closely trimmed retail cuts from the round, loin, rib and chuck (% of HCW) = 52.56 – 1.95 × subcutaneous

fat – 1.06×KPH + 0.106×LM area – 0.018×HCW [13].

Calculations

Energy gain (EG, Mcal/d) was calculated by the equation: $EG = 0.0557 \times \text{shrunk body weight (SBW)}^{0.75} \times \text{average daily gain (ADG)}^{1.097}$; where EG is the daily deposited energy and SBW is the equivalent of body weight×0.96 [14]. Maintenance energy (EM, Mcal/d) was calculated by the equation: $EM = 0.084 \text{ SBW}^{0.75}$ [15]. From the derived estimates of energy required for maintenance and gain, the NE_m and NE_g values of the diet were obtained using the quadratic formula: $x = (-b - \sqrt{b^2 - 4ac}) / 2c$, where $x = \text{diet NE}_m$, Mcal/kg, $a = -0.41EM$, $b = 0.877EM +$

0.41 dry matter intake (DMI)+EG, and $c = -0.877DMI$, and $NE_g = 0.877 \text{ NE}_m - 0.41$ [16].

Statistical analyses: The trial was analyzed as a randomized complete block design using the pen as experimental unit [17] according to the following statistical model: $Y_{ij} = \mu + B_i + T_j + \epsilon_{ij}$, where μ is the common experimental effect, B_i represents initial weight block effect, T_j represent dietary treatment effect, and ϵ_{ij} represents the residual error. Performance data were determined considering initial shrunk weight groupings for blocks. In determination of ADG, interim and final weights were reduced 4% to account for digestive tract fill. The effects of energy level in the diet were tested using orthogonal contrast. Significant difference was considered when the p-value was ≤ 0.05 and tendencies when $p > 0.05 \leq 0.10$. Data were analyzed using the MIXED procedure of SAS [17].

Table 2. Effects dietary nonstructural carbohydrate concentration on growth performance and dietary net energy of Holstein steers on long-term feeding

Items	Nonstructural carbohydrates (%)		SEM	p value
	51	64		
Pen replicates	5	5		
Body weight (kg) ¹⁾				
Initial	127.5	131.2	0.99	0.06
112	280.0	270.3	2.89	0.08
224	511.6	497.4	3.65	0.06
Final	591.9	573.8	5.89	0.09
ADG (kg/d)				
1-112 days	1.36	1.24	0.02	0.03
112-224 days	1.68	1.63	0.32	0.38
224-308 days	1.46	1.42	0.01	0.16
1-308 days	1.50	1.43	0.01	0.06
DMI (kg/d)				
1-112 days	5.66	5.62	0.09	0.81
112-224 days	9.72	9.18	0.23	0.19
224-308 days	11.89	11.34	0.32	0.30
1-308 days	8.83	8.48	0.16	0.20
Gain to feed (kg/kg)				
1-112 days	0.240	0.220	0.003	0.01
112-224 days	0.173	0.178	0.002	0.24
224-308 days	0.123	0.126	0.002	0.54
1-308 days	0.170	0.169	0.002	0.69
Dietary NE (Mcal/kg)				
1-112 days				
NE _m	1.93	1.83	0.02	0.02
NE _g	1.29	1.20	0.01	0.02
112-224 days				
NE _m	2.03	2.05	0.02	0.55
NE _g	1.37	1.39	0.02	0.55
224-308 days				
NE _m	1.97	1.99	0.04	0.82
NE _g	1.32	1.33	0.03	0.82
1-308 days				
NE _m	2.01	2.00	0.02	0.77
NE _g	1.35	1.35	0.01	0.77

SEM, standard error of the mean; ADG, average daily gain; DMI, dry matter intake; NE, net energy; NE_m, net energy of maintenance; NE_g, net energy of gain.

¹⁾ Initial weight is the shrunk off truck arrival weight. Interim and final weights were reduced 4% to account for digestive tract fill.

RESULTS AND DISCUSSION

Treatment effects on growth performance, dietary net energy (NE) and carcass characteristics are shown in Table 2, 3. Steers fed HF diet showed marked improvement (8.8%, $p = 0.03$) in ADG during the initial 112-d period. This effect was followed by a numerical trend for greater ADG throughout the remainder of the study so that overall ADG tended to be greater (4.9%, $p = 0.06$) for the HF than for LF.

The relationship between observed and projected [18] incremental ADG during the course of the trial is shown in Figure 1. Overall projected ADG for the 308-d feeding period was 1.43 kg. This value agrees with observed overall ADG (1.43 kg) for steers fed the LF diet. Thus, the 5% greater ADG of steers on the HF diet was not due to a poorer than expected performance of steers on the LF diet. Rather instead, feeding the HF diet resulted in a greater than generally anticipated overall ADG for implanted calf-fed Holstein steers across the weight range in consideration. This effect is consistent with

Table 3. Effects of dietary nonstructural carbohydrate concentration on carcass characteristics of calf-fed Holstein steers

Item	Nonstructural carbohydrates (%)		SEM	p value
	51	64		
Carcass weight (kg)	366.4	355.2	3.6	0.10
Dressing (%)	61.6	61.5	0.19	0.79
LM area (cm ²)	79.9	80.3	1.4	0.85
Fat thickness (cm)	0.84	0.77	0.04	0.26
KPH fat (%)	2.27	2.28	0.09	0.94
Yield grade (%) ¹⁾	51.96	52.22	0.086	0.09
Marbling score ²⁾	4.08	4.61	0.25	0.22

SEM, standard error of the mean; LM, longissimus muscle; KPH, kidney, pelvic and heart fat as a percentage of carcass weight.

¹⁾ Percentage of closely trimmed, mostly boneless, retail product from the round, loin, rib, and chuck.

²⁾ Coded: minimum slight = 3, minimum small = 4.




Figure 1. Treatment effects on average daily weight gain (ADG). ADG-LF, lower fiber diet; ADG-HF, higher fiber diet; ADG-PRD, generalized projected ADG based on Torrentera et al [18].

the hypothesis of this study; particularly apparent during the initial and final 112 d of the 308-d feeding period (Figure 1). However, it's important to clarify that the substitution of steam-flaked corn with alfalfa and dry distillers grains with solubles (DDGS) increased crude protein concentration of the HF diet (14.6% vs 12.9% CP) versus that of the LF diet. Hence, a portion of the increase in ADG during the initial growing phase may be attributable to the great protein level of the HF diet. The latter is especially important, as lysine is a limiting amino acid during this period [3,19]. According to the diet formulation (Table 1) the metabolizable lysine for steers fed a LF diet during the first 112-d of the experiment was slightly lower of the requirements indicated by NRC ([19], Level 1).

With respect to forage level, Defoor et al [20] likewise observed that increasing dietary NDF in practical feedlot diets enhanced weight gain, presumably through reduction in day-to-day occurrence of subclinical digestive dysfunctions. Erratic patterns of intake have been associated with digestive disorders [21,22]. In the present study, variation in DMI (data not shown) was 23% lower in HF group (coefficient of variation = 5.34%) than in the LF group (coefficient of variation = 7.32%). Additionally, the risk of subclinical acidosis may be assessed in function of the non-structural carbohydrate:NDF ratio of the HF and LF diets (2.3 and 4.8, respectively; Table 1). Based on NRC [19], the predicted (Level 1 approach) average ruminal pH for steers fed the HF and LF diets are 5.82 and 5.69, respectively.

Notwithstanding differences in dietary NE (Table 1), treatment differences in DMI were not appreciable ($p > 0.10$). Indeed, gain efficiency (ADG/DMI) and estimated dietary NE were greater (8.3% and 5.2%, respectively, $p \leq 0.02$) for HF vs LF

during the initial 112-d period. Overall (308-d) gain efficiency and estimated dietary NE were similar ($p \geq 0.69$) for both dietary treatments. However, due to differences in tabular dietary NE, the ratio of observed:expected dietary NE tended to be greater (4.1%, $p = 0.06$) for the HF vs LF diet.

Consistent with the tendency for increased ADG, carcass weight also tended to be greater (11.2 kg, $p = 0.10$) for HF- vs LF-fed steers. Otherwise, there were no treatment effects ($p > 0.10$) on carcass characteristics. Although there was a tendency for a slightly greater (0.5%, $p = 0.09$) estimated carcass yield.

CONCLUSION

It is concluded that reducing the non-structural carbohydrate concentration of a conventional steam-flaked corn-based growing finishing diet for calf-fed Holstein steers can effectively enhance growth performance, particularly during the early growing and late finishing phases.

CONFLICT OF INTEREST

We certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript.

REFERENCES

1. Thooney ML. Growth, feed efficiency and variation of individually fed Angus, Polled Hereford and Holstein steers. *J Anim Sci* 1987;65:1-8.
2. Smith JL, Wilson LL, Swanson DL. Implant sequence effects

- in intact male Holstein veal calves: live and slaughter traits. *J Anim Sci* 1999;77:3125-32.
3. Zinn RA, Calderon JF, Corona L, et al. Phase feeding strategies to meet metabolizable amino acid requirements of calf-fed Holstein steers. *Prof Anim Sci* 2007;23:333-9.
 4. Stock RA, Laudert SB, Stroup WW, et al. Effect of monensin and tylosin combination on feed intake variation of feedlot steers. *J Anim Sci* 1995;73:39-44.
 5. Ware RA, Zinn RA. Influence of forage source and NDF level on growth performance in feedlot cattle. *Proc West Sect Am Soc Anim Sci* 2004;55:637-41.
 6. Duff GC, McMurphy CP. Feeding Holstein steers from start to finish. *Vet Clin North Am Food Anim Pract* 2007; 23:281-97.
 7. Walter LAJ, McEvers TJ, May ND, et al. The effects of days on feed and zilpaterol hydrochloride supplementation on feeding behavior and live growth performance of Holstein steers. *J Anim Sci* 2016;94:2139-50.
 8. Cheng KJ, McCallister TA, Popp JD, et al. A review of bloat in feedlot cattle. *J Anim Sci* 1998;76:299-308.
 9. Zinn RA, Plascencia A, Barajas R. Interaction of forage level and monensin in diets for feedlot cattle on growth performance and digestive function. *J Anim Sci* 1994;72:2209-15.
 10. Calderon-Cortes JF, Zinn RA. Influence of dietary forage level and forage coarseness of grind on growth performance and digestive function in feedlot steers. *J Anim Sci* 1996;74:2310-6.
 11. AOAC. Official methods of analysis. 17th ed. Association of Official Analytical Chemists. Gaithersburg, MD, USA: AOAC International; 2000. p. 69.
 12. USDA. United States Standards for Grading of Carcass Beef. Washington, DC, USA: Agricultural Marketing Service, USDA; 1997.
 13. Murphey CE, Hallett DK, Tyler WE, Pierce JC. Estimating yields of retail cuts from beef carcasses. In *Proceeding 62nd Meet American Society Animal Production*; Chicago, IL, USA. 1960. p. 1-12.
 14. Committee on Nutrient Requirement of Beef Cattle, National Research Council. Nutrient requirements of beef cattle. 6th rev ed. Washington, DC, USA: National Academy Press; 1984.
 15. Garrett W. Energetic efficiency of beef and dairy steers. *J Anim Sci* 1971;31:452-6.
 16. Zinn RA, Shen Y. An evaluation of ruminally degradable intake protein and metabolizable amino acid requirements of feedlot calves. *J Anim Sci* 1998;76:1280-9.
 17. SAS. User's guide: statistics version SAS/STAT 9. 6th ed. Cary, NC, USA: SAS Inst., Inc.; 2000.
 18. Torrentera NG, Barreras A, Plascencia A, Zinn RA. Delay implant strategy in calf-fed Holstein steers: growth performance, growth rate and carcass characteristics. *J Appl Anim Res* 2017;45:454-9.
 19. Committee on Nutrient Requirement of Beef Cattle, National Research Council. Nutrient requirements of beef cattle. 7th rev ed. Washington, DC, USA: National Academy Press; 2000.
 20. Defoor PJ, Galyean ML, Saylor GB, Nunnery GA, Parsons CH. Effects of roughage source and concentration on intake and performance by finishing heifers. *J Anim Sci* 2002;80:1395-404.
 21. Stock RA, Laudert SB, Stroup WW, et al. Effect of monensin and tylosin combination on feed intake variation of feedlot steers. *J Anim Sci* 1995;73:39-44.
 22. Vasconcelos JT, Galyean ML. ASAS Centennial Paper: contributions in the Journal of Animal Science to understanding cattle metabolic and digestive disorders. *J Anim Sci* 2008;86: 1711-21.