

Immunomodulatory effects of phytochemicals in chickens and pigs — A review

C. M. Huang¹ and T. T. Lee^{1,2,*}

* **Corresponding Author:** T. T. Lee
Tel: +886-4-2287-0613, **Fax:** +886-4-2286-0265,
E-mail: tlee@dragon.nchu.edu.tw

¹ Department of Animal Science, National Chung Hsing University, Taichung 402, Taiwan

² The iEGG and Animal Biotechnology Center, National Chung Hsing University, Taichung 402, Taiwan

ORCID

C. M. Huang

<https://orcid.org/0000-0001-7692-5542>

T. T. Lee

<https://orcid.org/0000-0002-0063-5714>

Submitted Sept 4, 2017; Revised Oct 17, 2017;
Accepted Nov 17, 2017

Abstract: Environmental stressors like pathogens and toxins may depress the animal immune system through invasion of the gastrointestinal tract (GIT) tract, where they may impair performance and production, as well as lead to increased mortality rates. Therefore, protection of the GIT tract and improving animal health are top priorities in animal production. Being natural-sourced materials, phytochemicals are potential feed additives possessing multiple functions, including: anti-inflammatory, anti-fungal, anti-viral and antioxidative properties. This paper focuses on immunity-related physiological parameters regulated by phytochemicals, such as carvacrol, cinnamaldehyde, curcumin, and thymol; many studies have proven that these phytochemicals can improve animal performance and production. On the molecular level, the impact of inflammatory gene expression on underlying mechanisms was also examined, as were the effects of environmental stimuli and phytochemicals in initiating nuclear factor kappa B and mitogen-activated protein kinases signaling pathways and improving health conditions.

Keywords: Phytochemicals; Phytochemicals; Immune Response; Inflammatory Cytokines; Domestic Animals

INTRODUCTION

Pathogens like viruses, bacteria, fungi and other environmental stimuli are latent stressors which may damage animal tissues and cells. Excessive stressors may increase the concentration of reactive oxygen species (ROS), causing lipid peroxidation and oxidative damage to cellular membranes. The lack of sufficient antioxidants to eliminate ROS will lead to oxidative damage and trigger inflammation [1-3]. When inflammation occurs, cells have to recruit other cells to local sites by secreting more inflammatory cytokines and chemokines; this is a necessary process for the host to fight against infection and to repair tissue. However, prolonged inflammation may lead to unnecessary energy expenditure. As such, the alleviation and prevention of over-inflammation and the return of immune status to normal condition will benefit animal production.

As the first line of defense, the gastrointestinal tract (GIT) has a large surface area where food degradation and absorption take place; pathogens and toxins may also gain access to the body through the GIT by destroying the mucus layer [4]. Accordingly, protection of the GIT against the incursion of pathogens and toxins is of vital importance. Common toxins, such as aflatoxin and lipopolysaccharide (LPS), may cause great damage to intestinal cells, induce inflammation and immune response, and impair production [5,6]. Toxic airborne gases, like ammonia, may harm the respiratory system of animals, resulting in depressed performance [7,8]. Exogenous stressors may exacerbate the pathogenic invasion in the gut of animals, indicated by decreased bodyweight and feed intake, infiltration of heterophils in

the intestine and increased *Salmonella* numbers in the spleens of heat stressed chickens after *Salmonella* infection [9]. Coccidiosis, a parasitic disease in birds infected mainly with *Eimeria* spp., is a major problem in the poultry industries worldwide, resulting in enormous annual losses [10]. These stressors impair mucosal immune responses to other pathogens and cause intestinal damage and poor growth performance [6,11]. Therefore, many different feed additives, like antibiotics, probiotics, prebiotics and phytochemicals, have been added to animal diets to improve the hosts' capacity to eliminate the negative influence of environmental stressors and improve growth performance [12-17].

Antibiotics had been widely used in the livestock industry, but adverse effects, including drug resistance and reduced beneficial endogenous bacteria counts, led to subsequent prohibition against antibiotic supplementation by the European Union in 2006. Many other governments also began to restrict antibiotic supplementation, prompting a great deal of new research into phytochemicals as alternative feed additives. It was observed that supplementation of phytochemicals had positive effects on the pancreatic and intestinal enzyme secretion in broilers [18,19], and improved dry matter and crude protein digestibility in pigs [20]. Although inconsistent results of feed intake suggested that phytochemicals might not improve feed palatability, feeding phytochemicals did promote nutrient absorption and feed conversion ratio (FCR) in poultry [21]. Many authors have pointed out that phytochemicals in animal diet improved the immune response and the protective capacity of the GIT against exogenous stressors [22-26]; however, the mechanisms by which phytochemicals act on host cells were still unclear. In this review, we summarize the known relevant mechanisms in mammals and poultry, and suggest the potential immunomodulatory effects of phytochemicals on chickens and pigs.

EFFECTS OF PHYTOCHEMICALS ON IMMUNE RESPONSE

Phytochemicals, the active constituents in phytochemicals, can be roughly divided into two groups based on the different synthetic pathways in plants: terpenes and terpenoids, or aromatic and aliphatic compounds [27]. Some phenolic compounds and terpenes have been reported to exhibit anti-inflammatory activity [28,29]. However, as all the phytochemicals were extracted under different conditions or by using different parts of the plant, it is difficult to evaluate their effect. Phytochemicals have traditionally been used in powder form, but antagonizing or additive reactions between compounds in a single phytochemical may result, as inconsistent results of immunoregulatory action between treatments of phytochemical and its individual component were found [30,31]. It's difficult to study the possible interactions between compounds in phytochemicals, not to

mention the diversity of phytochemicals and phytochemicals. As such, the complexity of the immunoregulatory actions of phytochemicals limits the progress of related research. Nevertheless, phytochemicals and phytochemicals still remain among the most promising feed additives thanks to their positive effect on regulating the immune response in chickens and pigs. Here we review the effects of some popularly applied phytochemicals on the immune response in animals, including *Echinacea purpurea* L. (EP), oregano, cinnamon, *Curcuma longa*, and thyme (Table 1).

Echinacea purpurea L.

In our previous studies, EP extract, the herbal medicine normally used to prevent respiratory infections in humans, was evaluated for its antioxidant activity in broilers [32-35]. The results demonstrated that EP ethanol extracts had no major influence on the viability of the mononuclear cells in peripheral blood [32]. Dietary supplementation with dried EP improved both the FCR and meat quality [33,34]. It was also reported that 5.0 g/kg EP supplementation in broilers could increase the antibody titer against sheep red blood cells (SRBCs) and Newcastle disease virus (NDV) [35]. The EP, which was fed to laying hens and fattening pigs as pressed juice preserved with ethanol, still showed immunomodulatory effects, as indicated by the increased count of lymphocytes and NDV antibodies in the blood [36]. The 0.1% supplementation of EP aqueous extract in the diet enhanced cutaneous basophil hypersensitivity and antibody response against SRBCs in broilers [37]. However, no significant differences were observed in sows fed 1.2% and 3.6% EP cobs during gestation, in sows fed 0.5% and 1.5% cobs during lactation, or in piglets fed 1.8% cobs compared to the control group, while the FCR and erysipelas antibodies were improved in grower-finisher pigs supplemented with 1.5% cobs or 4 to 6 mL pressed juice per day [38]. Accordingly, EP supplementation may promote host immunity in broilers and fattening pigs, but determining its immunomodulatory effect and effective dosage in sows and piglets needs more research.

Oregano

Oregano belongs to the Labiatae family. Oregano is the popular name for a variety of widely distributed plant species, including *Origanum vulgare* L. and *O. onites* L., and is characterized as containing great amounts of carvacrol as the main source of its fragrance. Another plant called oregano is *Coridothymus capitatus* (L.) Reichenb. fil. (Spanish oregano) [39]; it has been widely studied as a feed additive in animals. Mohiti-Asli and Ghanaatparast-Rashti [40] reported that dietary supplementation with 300 ppm of oregano essential oil (OEO) improved the European production efficiency factor in broilers, which was calculated according to body weight, livability, FCR and trial duration; 500 ppm OEO had no positive effect, while

Table 1. Effect of five phytochemicals, *Echinacea purpurea* L., oregano, cinnamon, turmeric, and thyme, on the immune responses of chickens and pigs

Phytochemical materials	Species of experimental animals	Supplemented form	Effective dosages	Bioactive components	Enhanced performance/production	Immune responses	References
<i>Echinacea purpurea</i> L.	Broilers	Aerial part powder	5 and 10 g/kg diet		AWG, FI, and FCR	Higher antibody titers against SRBC and NDV	Landy et al [35]
	Laying hens and pigs	Pressed juice	0.25 mL/kg BW ^{0.75}		No effect	Higher count of lymphocytes and NDV antibodies in the blood	Böhmer et al [36]
	Broilers	Aqueous extract	1 mL/L drinking water		No effect	Higher antibody titer against SRBC	Rahimi et al [37]
	Grower/finishing pigs	Cobs or pressed juice	1.5% cobs in the diet or 4-6 mL juice per day		FCR	Higher erysipelas antibody	Maass [38]
Oregano	Broilers	<i>Essential oil</i>	300 ppm in the diet		AWG and FI	Higher secondary antibody titer and IgG titer, lowered H/L ratio	Mohiti-Asli and Ghanaat-parast-Rashti [40]
	Broilers	<i>Essential oil</i>	50 and 100 ppm in drinking water	Carvacrol and thymol	BW	Higher antibody titer against NDV and avian influenza virus	Galal et al [41]
	Sows Broilers	<i>Essential oil</i> <i>Essential oil</i>	250 ppm in the diet 60-200 ppm in the diet	Carvacrol and thymol Carvacrol and thymol	Energy intake AWG and FCR	Higher thymus lymphocytes Higher total antibody and IgG antibody, lowered H/L ratio	Ariza-Nieto et al [43] Hashemipour et al [44]
Cinnamon	Broilers	Powder	4 and 8 g/kg in the diet		FCR	Higher lymphocytes proportion	Najafi and Taherpour [51]
	Broilers	Oil extract	100 and 200 ppm in the diet		WG, FI, and FCR	Lowered H/L ratio, higher white blood cells	Al-Kassie [52]
	Broilers	Powder	5 g/L in the drinking water		Negative effect on BW	Higher antibody titer against NDV	Sadeghi et al [53]
Turmeric (<i>Curcuma longa</i>)	Broilers	Rhizome powder	2.5, 5, and 7.5 g/kg diet		Not detected	Higher IgA, IgG, and IgM levels, and decreased ratio of monocytes	Emadi and Kermanshahi [55]
	Broilers	Rhizome powder	2 g/kg diet		AWG and FCR	Higher total secondary antibody titer, and decreased H/L ratio	Akhavan-Salamat and Ghasemi [56]
	Broilers	Dried rhizome extract	35 mg/kg diet		BW	Higher antibody level against <i>Eimeria</i> microneme protein	Kim et al [23]
Thyme	Broilers	Oil extract	100 and 200 ppm in the diet		AWG, FI, and FCR	Lowered H/L ratio, higher white blood cells	Al-Kassie [52]
	Broilers	<i>Essential oil</i>	0.1% in the diet		Not detected	Higher white blood cells and decreased H/L ratio	Khaksar et al [60]
	Broilers	<i>Essential oil</i>	0.02% in the diet		FCR	Higher primary antibody titer	Najafi and Toriki [64]
	Broilers	Powder	2, 5, and 8 g/kg diet	Essential oils: thymol, γ -terpinene, P-cymene, carvacrol Phenolic acids: salicylic acid, ellagic acid, benzoic acid, chlorogenic acid, caffeic acid, gallic acid, vanillic acid, cinnamic acid Flavonoids: peridin, rosmarinic, quercitrin, narerigin, rutin, quercetin, narenginin, kaempferol, epicatechin, luteolin	BW, AWG, and FI	Higher levels of lymphocytes, white blood cells, and IgG	Hassan and Awad [119]

BW, body weight; AWG, average weight gain; FI, feed intake; FCR, feed conversion ratio; SRBC, sheep red blood cell; NDV, Newcastle disease virus; Ig, immunoglobulin; H/L ratio, heterophil to lymphocyte ratio.

300 ppm OEO in the diet produced higher secondary total antibody titer against SRBCs; in addition, the immunoglobulin G (IgG) titer was higher than it was in the control group. Another study reported that higher doses (500 and 1,000 ppm) of OEO supplementation led to positive immune responses in NDV- and avian influenza-vaccinated broilers [41]. Serum heterophil counts and the heterophil to lymphocyte ratio (H/L ratio), both considered stress indicators [42], were decreased in birds fed 300 ppm OEO. On day 14 of lactation, higher thymus lymphocyte concentrations were observed in sows' milk after 250 ppm OEO was added to their diet, indicating the immunostimulatory effect of OEO [43]. Other studies also

reported that 300 ppm OEO or 300 ppm of a mixture of carvacrol and thymol could improve the growth performance or the immune response, as indicated by lowered excreta oocyst counts of coccidia [24], and increased hypersensitivity response, total antibody titer, and H/L ratio [44]. However, the growth promoting effects of OEO were inconsistent [45], probably because the immunomodulatory actions of OEO could not improve growth performance in a less harsh experimental environment when compared to practical rearing conditions.

Cinnamon

The essential oils from some *Cinnamomum* species contain

great amounts of trans-cinnamaldehyde (*C. burmannii*), while other species do not (*C. tamala*, *C. pauciflorum*, and *C. zeylanicum*) [46,47]. Dietary supplementation with cinnamon leaves or extracted oil have been demonstrated to improve growth performance in broilers [48,49], and decrease coliform bacteria in the jejunum and large intestine [50]. Najafi and Taherpour [51] reported that the inclusion of 0.4% and 0.8% cinnamon in broiler diets improved the FCR and increased the concentration of hemoglobin and the proportion of lymphocytes in the blood. Dosages of 100 and 200 ppm essential cinnamon oil improved FCR and body weight gain from 1 to 6 weeks, reduced the H/L ratio and increased the level of white blood cells in broilers [52]. Conversely, positively modulated immune responses and impaired growth performance were detected in 21-day-old broilers that were given 5.0 g/L cinnamon in their drinking water [53]. This discrepancy may be because the majority of studies supplied cinnamon to animals via their diet versus their drinking water. Furthermore, the variety of species used in the different experiments make it difficult to determine the effective dosage of cinnamon or cinnamon oil, or to make valid comparisons across studies, especially if the content of the major bioactive compounds was not measured.

Turmeric (*Curcuma longa*)

There are many species in the genus *Curcuma*, among which *C. longa* (turmeric) is the most studied. The major phytochemicals in the rhizome of *C. longa* vary dramatically with the environment and location. Li et al [54] concluded that curcumin, demethoxycurcumin and bisdemethoxycurcumin are the main compounds in the rhizomes, powders and extract products, whereas the concentrations of ar-turmerone, α -turmerone, and β -turmerone are related to the quality of the turmeric oils and oleoresin products. Most studies evaluated this phytochemical in the form of turmeric rhizome powder (TRP). TRP inclusion in the diet significantly increased blood IgA, IgG, and IgM levels, and decreased the ratio of monocytes in 42-day-old broilers challenged with SRBCs [55]. Similarly, 0.2% TRP supplementation decreased the H/L ratio and increased the total secondary antibody titer against SRBCs in broilers undergoing heat stress treatment [56]. Moreover, the addition of 0.33% and 1.0% TRP, which contained 1.16% curcumin, improved the FCR in broilers during the entire feeding period [57]. Inclusions of 0.33%, 0.66%, and 1.0% TRP significantly reduced abdominal fat content and serum triglyceride concentration [57]. Therefore, dietary curcumin could regulate immune responses in animals; it has the potential to enhance growth performance.

Thyme

Thyme is the popular name for *Thymus* spp.; thymol, carvacrol, p-cymene and γ -terpinene are the major active components

in the essential oil [58,59]. Dietary supplementation of thyme was in the form of essential oil or leaf powder. It has been shown that supplementation with thyme essential oil improved cutaneous basophil hypersensitivity to phytohaemagglutinin-P (PHA-P) and reduced the H/L ratio. It also produced a more balanced intestinal microbiology, as indicated by higher *Lactobacillus* and *Bifidobacterium* counts and lower *Escherichia coli* (*E. coli*) levels in the ileum [60]. Previous studies suggested that heat stress may impair the cell-mediated immune response to PHA-P in chicks [61], highlighting the benefit of feeding thyme essential oil to birds. The H/L ratio is a stress indicator, as mentioned above. A reduced H/L ratio demonstrates the enhanced immunoregulatory capacity of the host. Intestinal microbiota plays an important role in the immune system of chickens [62]. Environmental stressors may change the intestinal microbiota and increase the susceptibility to colonization by pathogens in the intestine [63]. It was also reported that thyme essential oil improved bodyweight gain and FCR in broilers [52,64]. As such, thyme has potential as a feed additive in poultry production.

INTERACTIONS BETWEEN PHYTOCHEMICALS AND THE MAPK- AND NF κ B-REGULATED PATHWAYS IN CHICKENS AND PIGS

NF κ B and MAPKs in immune responses

Innate immunity, also known as non-specific immunity, includes activation and the recruitment of innate leukocytes like macrophages, thereby enhancing phagocytosis. As pathogens invade, innate leukocytes identify and engulf them while expressing cytokines, thereby initiating the immune response and leading to an inflammatory reaction [65]. Adaptive immunity is a specific immune system process initiated by the presentation of antigens. Pathogens and exogenous stimuli are identified and eliminated by B cells and T cells. Among these, helper T cells are induced by effector cytokines to differentiate into cell types Th1 or Th2, and subsequently deliver signals to macrophages or B cells, respectively [66]. Therefore, homeostasis is maintained through the mutual actions of the innate and adaptive immune systems.

In the innate immune system, toll-like receptors (TLRs) are the most studied pattern-recognition receptors in poultry and livestock [67-69]. As TLRs are stimulated, many signaling pathways are activated simultaneously through adaptor proteins such as myeloid differentiation primary response 88 (MyD88) and toll-IL-1-resistance domain-containing adaptor, inducing interferon β [70]. The Src kinase plays a vital role in the activation of mitogen-activated protein kinases (MAPKs) and nuclear factor kappa B (NF κ B), the two major pathways that modulate immune responses [71-73]. MAPKs are a group of protein kinases that regulate cellular activities. Among

MAPKs, p38 protein kinase, extracellular signal-regulated protein kinase (ERK1/2), and c-Jun N-terminal kinase have been reported to be associated with the regulation of inflammation and immunity in response to environmental stress [74-76]. The NF κ B-mediated pathway involves many physical reactions and is extremely important in the modulation of immunity, inflammation and apoptosis [71]. It is well known that NF κ B is a dimer consisting of different subunits [71]. Under normal physical conditions, the dimer binds to the inhibitor of kappa B (I κ B) and presents as an inactive form in the cytosol. When the cell is activated by exogenous stimuli, NF κ B is liberated from its I κ B-inhibited form. The inhibitor of kappa B kinase (IKK) phosphorylates the linked I κ B, leading to the ubiquitination of phosphorylated I κ B, and the subsequent identification and degradation by the 26S proteasome [77]. Disconnected NF κ B enters the nucleus, binding to a specific region of DNA, and promotes the expression of downstream genes. Translated proteins are secreted into the extracellular space to modulate the immune system and inflammatory response [78]. Depending on the strength, timing and site of stimulation, different downstream products are generated and exert diverse functions, including cellular growth, differentiation, tissue repair and synthesis of tissue collagen. Therefore, studies on NF κ B signaling include its onset and therapies for many diseases as well as the apoptosis of cancer cells [79]. Pro-inflammatory cytokines, ROS and mitogens generated by bacterial or viral infection, chemical and biological tissue damage or other stressors, induce the activation of NF κ B signaling [80,81], followed by the expression of more inflammatory cytokines and chemokines. Interleukin 1 (IL-1), IL-6, and tumor necrosis factor alpha (TNF- α) are the three most investigated pro-inflammatory cytokines that can be expressed by monocytes and macrophages after invading pathogens are identified. These cytokines mediate metabolic changes, which enhance the immune response and disease resistance, thereby stunting growth performance [82,83]. TNF- α may be secreted by activated macrophages and lead to chemotaxis of monocytes [84]. Expressions of IL-1 β , IL-6 and TNF- α are highly associated with the immune status of poultry and livestock [85].

Nitric oxide (NO) and prostaglandin E₂ (PGE₂) are important inflammatory mediators that modulate the development of inflammation [86,87]. After NF κ B is activated, NF κ B-mediated signaling molecules upregulate the expression of the downstream gene COX-2, synthesizing cyclooxygenase 2, which converts arachidonic acid into PGE₂. Inducible nitric oxide synthase (iNOS) is also activated, which subsequently increases NO production [88]. The enhanced production of histamine and PGE₂ in response to stressors not only induces phagocytosis of macrophages, but also affects the production of cytokines by T cells, macrophages and dendritic cells [87,89]. These cytokines include IL-1 and IL-6, which are transported

through the circulatory system to the brain, activating the hypothalamic-pituitary-adrenal (HPA) axis [90]. In response to acute inflammation, the HPA axis is stimulated to secrete more glucocorticoids, which have been thought to elevate the H/L ratio in birds and the neutrophil to lymphocyte ratio in mammals. Glucocorticoids also induce heterophils (neutrophils) to express pro-inflammatory cytokines, including IL-1 β , IL-6, IL-10, IL-12 α , and IL-18 [91]. Increased heterophils (neutrophils) and decreased lymphocytes in the circulatory system represent the development of the innate response in the host as inflammation is triggered. Elevated H/L ratio is considered a stress marker. Exogenous substances are eliminated through acute inflammation in organisms. However, under stressful conditions, the suppressed immune system is unable to cope, leading to chronic inflammation [92]. It has been proven that chronic inflammation is associated with bacterial disease and heat stress in animals [93,94]. In the previous section, we reviewed that phytochemicals, such as oregano, cinnamon, turmeric, and thyme, are able to modulate humoral responses in chickens and pigs. Their major components have been reported to regulate NF κ B and/or MAPKs signaling pathways in rodents [95-97]. Therefore, phytochemicals that modulate the immune response and regulate the expression of inflammatory mediators in chickens and pigs were introduced (Figure 1); they are listed below.

Carvacrol

Carvacrol (2-methyl-5-(1-methylethyl)-phenol), the major active component in oregano and thyme essential oils, is a phenolic monoterpene. In mice, carvacrol reduced local levels of IL-1 β and PGE₂, and mRNA levels of COX-2 and IL-1 β [98]. The mRNA and protein levels of TNF- α , IL-6, Inos, and COX-2 were also downregulated in D-galactosamine-induced hepatotoxic rats after treatment with carvacrol. The suppression of pro-inflammatory cytokine production may be attributed to the inhibition of TLR2- and TLR4-mediated signaling [99]. Zou et al [100] reported that OEO, which contains 81.92% carvacrol, lowered the protein level of MAPKs and NF κ B, and decreased the expression of the pro-inflammatory cytokines IL-1 β , IL-6, interferon- γ (IFN- γ), TNF- α and monocyte chemoattractant protein-1 in the jejunum of pigs, proving its anti-inflammatory effects. Moreover, the mRNA levels of occludin and zonula occludens-1, two epithelial tight junction proteins, were increased, demonstrating that carvacrol may improve the integrity of intestinal epithelia. Reduced expression of IL-6 and LPS-induced TNF- α factor were observed in *Eimeria* oocyst-challenged broilers supplemented with essential oils containing 81.89% carvacrol [101]. It is speculated that carvacrol may suppress TLRs and/or MAPKs and NF κ B signaling pathways in animals; however, more studies are needed to elucidate its mechanisms in detail.




Figure 1. The possible immunomodulatory action of carvacrol, cinnamaldehyde, curcumin, and thymol regulating expression of pro-inflammatory mediators and enzymes involved in the nuclear factor kappa B (NFκB) and mitogen-activated protein kinases (MAPKs) signaling pathways, inflammation, and immune response in chickens and pigs.

Cinnamaldehyde

In vitro studies have demonstrated that both *Cinnamomum cassia* oil and cinnamaldehyde, the major bioactive compound in cinnamon, inhibited mRNA expression of pro-inflammatory cytokines, including IL-1 β , IL-6, and TNF- α , and increased the production of anti-inflammatory cytokine IL-10 in LPS-activated J774A.1 cells. Subsequently, the expression of iNOS and COX-2 was decreased, as was NO production [102]. Youn et al [95] reported that cinnamaldehyde could suppress LPS-induced activation of NFκB and interferon regulatory factor 3, resulting in the decreased expression of the TLR4 target genes *COX-2* and *IFN- β* . In broilers, an analysis of global gene expression profiles indicated that the immunomodulatory effects of dietary cinnamaldehyde were related to antigen presentation, humoral immune response and inflammatory disease [25]. *In vitro* tests showed that chicken spleen lymphocytes treated with cinnamaldehyde (25 to 400 ng/mL) significantly promoted cell proliferation compared to the control group. The dosages of 1.2 to 5.0 μ g/mL of cinnamaldehyde stimulated macrophages to produce more NO, while 0.6 to 2.5 μ g/mL inhibited the growth of chicken tumor cells; 10 and 100 μ g/mL of cinnamaldehyde lowered the viability of *Eimeria tenella* [103]. Conversely, *in vivo* test results showed that dietary supplementation with 14.4 mg/kg cinnamaldehyde increased mRNA expression of the pro-inflammatory cytokines IL-1 β , IL-6, IL-15 and IFN- γ . Cinnamaldehyde

supplementation also improved the body weight gain of *E. acervulina* or *E. maxima*-infected chickens [103]. It should be noted that the dosage of cinnamaldehyde used in this study was very low, and treatment with low concentrations of cinnamaldehyde (0.01 to 0.5 μ g/mL) increased NFκB activation in LPS-stimulated human cells *in vitro* [104]. Further experiments are needed to prove whether cinnamaldehyde has a dose-dependent immunoregulatory effect on birds.

Curcumin

Curcumin (diferuloylmethane), one of the major components in turmeric, has been extensively researched. Previous studies have shown that curcumin involved in the p38 MAPK signaling pathway in LPS-stimulated RAW 264.7 macrophages [96], produced similar results in dendritic cells [105], and in the colonic tissue of rats with intracolonic injuries [106]. Additionally, Lubbad et al [107] reported that curcumin could suppress inflammation through the reduction of protein levels of TLR4, MyD88, and NFκB. MyD88 is normally recruited after TLR4 activation, and leads to the recruitment of more downstream molecules, as well as the activation of NFκB [108]. Acute carbon tetrachloride-induced liver damage was also alleviated by reducing the protein production of TNF- α and IL-1 β in rats with curcumin treatment [109]. In poultry, the mRNA expression of IL-1 β , IL-6, IL-12, IL-18 and tumor necrosis factor superfamily 15 were enhanced in chicken

macrophages treated *in vitro* with organic extract of turmeric [110]. The mRNA expression of IL-6 in the liver of broilers was increased by aflatoxin B1 treatment, but alleviated by turmeric powder supplementation [111]. The results of microarray hybridization demonstrated that the gene expression of myeloperoxidase, cluster of differentiation 28 and lactotransferrin are associated with the inflammatory response in broilers [112-114], which was downregulated by treatment with organic extract of turmeric [23]. A series of studies was conducted to evaluate the effect of turmeric oleoresin, which contained 98% curcuminoids, on the immune response of swine [26,76, 115,116]. The supplementation significantly reduced serum IL-1 β and TNF- α levels in porcine reproductive and respiratory syndrome virus-challenged pigs [26] and counteracted the effects of *E. coli*, as indicated by the markedly altered mRNA expression of IL-1 β , mucin 2, COX-2 and p38 MAPK in ileal mucosa [76] and serum TNF- α concentration [115]. According to the above reports, in animals, curcumin mainly modulated the p38 MAPK signaling pathway and regulated downstream pro-inflammatory mediators, such as IL-1 β , IL-6, and TNF- α .

Thymol

As the major compound in the essential oils of thyme and oregano, thymol is the isomer of carvacrol; thymol has been shown to possess anti-inflammatory properties; it suppresses the maturation of dendritic cells and activation of T cell proliferation *in vitro* [117]. Mature dendritic cells play an important role in the promotion of the immune response. Additionally, thymol inhibited the phosphorylation of NF κ B and MAPKs, and lowered the production of IL-6, TNF- α , iNOS, and COX-2 in LPS-stimulated mouse mammary epithelial cells [97]. It also reduced the production of NO and hydrogen peroxide in LPS-stimulated cells [118]. In the jejunum of broilers, supplementation with thyme powder, which contains the essential oils thymol (50.48%), γ -terpinene (11.03%), P-cymene (9.77%), and carvacrol (4.30%), significantly decreased transcriptional factor NF κ B. Pro-inflammatory cytokines such as IL-6, IFN- γ , and TNF- α were also downregulated in broilers following treatment with thyme powder [119]. This is consistent with previously mentioned results, suggesting that the anti-inflammatory effects of thyme make it suitable for application in animal production.

CONCLUSION

As it is inevitable for animals to be faced with environmental stressors like heat, disease and microbial toxins, the defense system of the host plays a vital role in maintaining health as well as growth performance and production. According to the above reports, the immune response improved when animal diets were supplemented with cinnamon, oregano, turmeric or thyme products. The stress response was alleviated since

the NF κ B and MAPKs signaling pathways were suppressed, and the expression of anti-inflammatory cytokines was increased. As a result, the blood levels of non-specific immune cells such as heterophils (neutrophils) were lowered, and lymphocyte and antibody production were promoted to defend against invading pathogens. Essentially, the less energy expended by chickens and pigs on the non-specific immune system, the more energy that is available for growth and production. Conclusively, phytochemicals did modulate NF κ B and/or MAPKs signaling pathways in chickens and pigs, but the detailed mechanisms, including expression of upstream molecules involving NF κ B and MAPKs signaling as well as the relationship between antioxidant and anti-inflammation, are still under investigation.

CONFLICT OF INTEREST

We certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript.

ACKNOWLEDGMENTS

The authors would like to express the sincere gratitude to the Ministry of Science and Technology (MOST 106-2313-B-005-048) and the iEGG and Animal Biotechnology Center for financially supporting the study to Prof. Tzu-Tai Lee.

REFERENCES

1. Lee MT, Lin WC, Yu B, Lee TT. Antioxidant capacity of phytochemicals and their potential effects on oxidative status in animals — A review. Asian-Australas J Anim Sci 2017;30: 299-308.
2. Blackwell TS, Blackwell TR, Holden EP, Christman BW, Christman JW. *In vivo* antioxidant treatment suppresses nuclear factor-kappa B activation and neutrophilic lung inflammation. J Immunol 1996;157:1630-7.
3. Cuzzocrea S, Riley DP, Caputi AP, Salvemini D. Antioxidant therapy: a new pharmacological approach in shock, inflammation, and ischemia/reperfusion injury. Pharmacol Rev 2001; 53:135-59.
4. Tarabova L, Makova Z, Piesova E, Szaboova R, Faixova Z. Intestinal mucus layer and mucins (a review). Folia Vet 2016; 60:21-5.
5. Arce C, Rami' rez-Boo M, Lucena C, Garrido JJ. Innate immune activation of swine intestinal epithelial cell lines (IPEC-J2 and IPI-2I) in response to LPS from *Salmonella typhimurium*. Comp Immunol Microbiol Infect Dis 2010;33:161-74.
6. Yunus AW, Razzazi-Fazeli E, Bohm J. Aflatoxin B1 in Affecting broiler's performance, immunity, and gastrointestinal tract: a review of history and contemporary issues. Toxins 2011;3:

- 566-90.
7. Miles DM, Branton SL, Lott BD. Atmospheric ammonia is detrimental to the performance of modern commercial broilers. *Poult Sci* 2004;83:1650-4.
 8. Drummond JG, Curtis SE, Simon J, Norton HW. Effect of aerial ammonia on growth and health of young pigs. *J Anim Sci* 1980;50:1085-9.
 9. Quinteiro-Filho WM, Gomes AVS, Pinheiro ML, et al. Heat stress impairs performance and induces intestinal inflammation in broiler chickens infected with *Salmonella enteritidis*. *Avian Pathol* 2012;41:5:421-7.
 10. Williams RB. A compartmentalised model for the estimation of the cost of coccidiosis to the world's chicken production industry. *Int J Parasitol* 1999;29:1209-29.
 11. McLamb BL, Gibson AJ, Overman EL, Stahl C, Moeser AJ. Early weaning stress in pigs impairs innate mucosal immune responses to enterotoxigenic *E. coli* challenge and exacerbates intestinal injury and clinical disease. *PLoS One* 2013;8:e59838.
 12. Lin WC, Lee MT, Lo CT, Chang SC, Lee TT. Effects of dietary supplementation of *Trichoderma pseudokoningii* fermented enzyme powder on growth performance, intestinal morphology, microflora and serum antioxidative status in broiler chickens. *Ital J Anim Sci* 2018;17:153-64.
 13. Lin WC, Lee MT, Chang YL, et al. Effects of mulberry leaves on production performance and the potential modulation of antioxidative status in laying hens. *Poult Sci* 2017;96:1191-203.
 14. Teng PY, Chung CH, Chao YP, et al. Effects of *Bacillus amyloliquefaciens* and *Saccharomyces cerevisiae* as directed-fed microbials on performance, intestinal microflora, and intestinal morphology in broiler chickens. *J Poult Sci* 2017;54:134-41.
 15. Lin MJ, Chang SC, Jea YS, et al. *In vitro* antioxidant capability and performance assessment of White Roman goose supplemented with dried *Toona sinensis*. *J Appl Anim Res* 2016;44:359-402.
 16. Lai LP, Lee MT, Chen J, Yu B, Lee TT. Effects of co-fermented *Pleurotus eryngii* stalk residues and soybean hulls by *Aureobasidium pullulans* on performance and intestinal morphology in broiler chickens. *Poult Sci* 2015;94:2959-69.
 17. Lin MJ, Chang SC, Jea YS, Chen WS, Lee TT. Effects of dietary garlic scape meal on the growth and meat characteristics of geese. *Br Poult Sci* 2015;56:716-22.
 18. Jang IS, Ko YH, Yang HY, et al. Influence of essential oil components on growth performance and the functional activity of the pancreas and small intestine in broiler chickens. *Asian-Australas J Anim Sci* 2004;17:394-400.
 19. Jamroz D, Wiliczkiwicz A, Wiertelcki T, Orda J, Skorupińska J. Use of active substances of plant origin in chicken diets based on maize and locally grown cereals. *Br Poult Sci* 2005;46:485-93.
 20. Li P, Piao X, Ru Y, et al. Effects of adding essential oil to the diet of weaned pigs on performance, nutrient utilization, immune response and intestinal health. *Asian-Australas J Anim Sci* 2012;25:1617-26.
 21. Fascina VB, Sartori JR, Gonzales E, et al. Phytogetic additives and organic acids in broiler chicken diets. *R Bras Zootec* 2012;41:2189-97.
 22. Kim DK, Lillehoj HS, Lee SH, et al. Immune effects of dietary anethole on *Eimeria acervulina* infection. *Poult Sci* 2013;92:2625-34.
 23. Kim DK, Lillehoj HS, Lee SH, et al. Dietary *Curcuma longa* enhances resistance against *Eimeria maxima* and *Eimeria tenella* infections in chickens. *Poult Sci* 2013;92:2635-43.
 24. Alp M, Midilli M, Kocabağlı N, et al. The effects of dietary oregano essential oil on live performance, carcass yield, serum immunoglobulin G level, and oocyst count in broilers. *J Appl Poult Res* 2012;21:630-6.
 25. Lillehoj HS, Kim DK, Bravo DM, Lee SH. Effects of dietary plant-derived phytonutrients on the genome-wide profiles and coccidiosis resistance in the broiler chickens. *BMC Proc* 2011;5(Suppl 4):S34.
 26. Liu Y, Che TM, Song M, et al. Dietary plant extracts improve immune responses and growth efficiency of pigs experimentally infected with porcine reproductive and respiratory syndrome virus. *J Anim Sci* 2013;91:5668-79.
 27. Bakkali F, Averbeck S, Averbeck D, Idaomar M. Biological effects of essential oils - A review. *Food Chem Toxicol* 2008;46:446-75.
 28. Fang SC, Hsu CL, Yen GC. Anti-inflammatory effects of phenolic compounds isolated from the fruits of *Artocarpus heterophyllus*. *J Agric Food Chem* 2008;56:4463-8.
 29. Siani AC, Souza MC, Henriques MG, Ramos MF. Anti-inflammatory activity of essential oils from *Syzygium cumini* and *Psidium guajava*. *Pharm Biol* 2013;51:881-7.
 30. Lin SJ, Chen CS, Ho CC, et al. *In vitro* anti-microbial and *in vivo* cytokines modulation effects of different prepared Chinese herbal medicines. *Food Chem Toxicol* 2006;44:2078-85.
 31. Lin SS, Lu TM, Chao PC, et al. *In vivo* cytokine modulatory effects of cinnamaldehyde, the major constituent of leaf essential oil from *Cinnamomum osmophloeum* Kaneh. *Phytother Res* 2011;25:1511-8.
 32. Lee TT, Chen CL, Shieh ZH, Lin JC, Yu B. Study on antioxidant activity of *Echinacea purpurea* L. extracts and its impact on cell viability. *Afr J Biotechnol* 2009;8:5097-105.
 33. Lee TT, Chen CL, Wang CC, Yu B. Growth performance and antioxidant capacity of broilers supplemented with *Echinacea purpurea* L. in the diet. *J Appl Poult Res* 2012;21:484-91.
 34. Lee TT, Ciou JY, Chen CL, Yu B. Effect of *Echinacea purpurea* L. on oxidative status and meat quality in Arbor Acres broilers. *J Sci Food Agric* 2013;93:166-72.
 35. Landy N, Ghalamkari Gh, Toghyani M, Moattar F. The effects of *Echinacea purpurea* L. (purple coneflower) as an antibiotic growth promoter substitution on performance, carcass characteristics and humoral immune response in broiler chickens. *J Med Plants Res* 2011;5:2332-8.

36. Böhmer BM, Salisch H, Paulicks BR, Roth FX. *Echinacea purpurea* as a potential immunostimulatory feed additive in laying hens and fattening pigs by intermittent application. *Livest Sci* 2009;122:81-5.
37. Rahimi S, Zadeh ZT, Torshizi MAK, Omidbaigi R, Rokni H. Effect of the three herbal extracts on growth performance, immune system, blood factors and intestinal selected bacterial population in broiler chickens. *J Agric Sci Tech* 2011;13:527-39.
38. Maass N, Bauer J, Paulicks BR, Böhmer BM, Roth-Maier DA. Efficiency of *Echinacea purpurea* on performance and immune status in pigs. *J Anim Physiol Anim Nutr* 2005;89:244-52.
39. Lagouri V, Blekas G, Tsimidou M, Kokkini S, Boskou D. Composition and antioxidant activity of essential oils from Oregano plants grown wild in Greece. *Z Lebensm Unters Forsch* 1993;197:20-3.
40. Mohiti-Asli M, Ghanaatparast-Rashti M. Comparison of the effect of two phytochemicals on growth performance and immune response of broilers. *J Appl Anim Res* 2017;45:603-8.
41. Galal AAE, El-Araby IE, Hassanin O, El-Said Omar A. Positive impact of oregano essential oil on growth performance, humoral immune responses and chicken interferon alpha signalling pathway in broilers. *Adv Anim Vet Sci* 2016;4:57-65.
42. Gross WB, Siegel HS. Evaluation of the heterophil/lymphocyte ratio as a measure of stress in chickens. *Avian Dis* 1983;27:972-9.
43. Ariza-Nieto C, Bandrick M, Baidoo SK, et al. Effect of dietary supplementation of oregano essential oils to sows on colostrum and milk composition, growth pattern and immune status of suckling pigs. *J Anim Sci* 2011;89:1079-89.
44. Hashemipour H, Kermanshahi H, Golian A, Veldkamp T. Effect of thymol and carvacrol feed supplementation on performance, antioxidant enzyme activities, fatty acid composition, digestive enzyme activities, and immune response in broiler chickens. *Poult Sci* 2013;92:2059-69.
45. Karimi A, Yan F, Coto C, et al. Effects of level and source of oregano leaf in starter diets for broiler chicks. *J Appl Poult Res* 2010;19:137-45.
46. Wang R, Wang R, Yang B. Extraction of essential oils from five cinnamon leaves and identification of their volatile compound compositions. *Innov Food Sci Emerg Technol* 2009;10:289-92.
47. Schmidt E, Jirovetz L, Buchbauer G, et al. Composition and antioxidant Activities of the essential oil of cinnamon (*Cinnamomum zeylanicum* Blume) leaves from Sri Lanka. *J Essent Oil Bear Pl* 2006;9:170-82.
48. Shirzadegan K. Reactions of modern broiler chickens to administration of cinnamon powder in the diet. *Iran J Appl Anim Sci* 2014;4:367-71.
49. Sang-Oh P, Chae-Min R, Byung-Sung P, Jong H. The meat quality and growth performance in broiler chickens fed diet with cinnamon powder. *J Environ Biol* 2013;34:127-33.
50. Al-Kassie GAM. The effect of thyme and cinnamon on the microbial balance in gastro intestinal tract on broiler chicks. *Int J Poult Sci* 2010;9:495-8.
51. Najafi S, Taherpour K. Effects of dietary ginger (*Zingiber Officinale*), cinnamon (*Cinnamomum*), synbiotic and antibiotic supplementation on performance of broilers. *J Anim Sci Adv* 2014;4:658-67.
52. Al-Kassie GAM. Influence of two plant extracts derived from thyme and cinnamon on broiler performance. *Pakistan Vet J* 2009;29:169-73.
53. Sadeghi GH, Karimi A, Padidar Jahromi SH, Azizi T, Daneshmand A. Effects of cinnamon, thyme and turmeric infusions on the performance and immune response in of 1- to 21-day-old male broilers. *Rev Bras Cienc Avic* 2012;14:15-20.
54. Li S, Yuan W, Deng G, et al. Chemical composition and product quality control of turmeric (*Curcuma longa* L.). *Pharm Crop* 2011;2:28-54.
55. Emadi M, Kermanshahi H. Effect of turmeric rhizome powder on immunity responses of broiler chickens. *J Anim Vet Adv* 2007;6:833-6.
56. Akhavan-Salamat H, Ghasemi HA. Alleviation of chronic heat stress in broilers by dietary supplementation of betaine and turmeric rhizome powder: dynamics of performance, leukocyte profile, humoral immunity, and antioxidant status. *Trop Anim Health Prod* 2016;48:181-8.
57. Nouzarian R, Tabeidian SA, Toghiani M, Ghalamkari G, Toghiani M. Effect of turmeric powder on performance, carcass traits, humoral immune responses, and serum metabolites in broiler chickens. *J Anim Feed Sci* 2011;20:389-400.
58. Hudaib M, Speroni E, Pietra AMD, Cavrini V. GC/MS evaluation of thyme (*Thymus vulgaris* L.) oil composition and variations during the vegetative cycle. *J Pharm Biomed Anal* 2002;29:691-700.
59. Senatore F. Influence of Harvesting Time on yield and composition of the essential oil of a thyme (*Thymus pulegioides* L.) growing wild in Campania (Southern Italy). *J Agric Food Chem* 1996;44:1327-32.
60. Khaksar V, Golian A, Kermanshahi H. Immune response and ileal microflora in broilers fed wheat-based diet with or without enzyme Endofeed W and supplementation of thyme essential oil or probiotic PrimaLac®. *Afr J Biotechnol* 2012;11:14716-23.
61. Jahanian R, Rasouli E. Dietary chromium methionine supplementation could alleviate immunosuppressive effects of heat stress in broiler chicks. *J Anim Sci* 2015;93:3355-63.
62. Brisbin JT, Gong J, Sharif S. Interactions between commensal bacteria and the gut-associated immune system of the chicken. *Anim Health Res Rev* 2008;9:101-10.
63. Burkholder KM, Thompson KL, Einstein ME, Applegate TJ, Patterson JA. Influence of stressors on normal intestinal microbiota, intestinal morphology, and susceptibility to *Salmonella enteritidis* colonization in broilers. *Poult Sci* 2008;87:1734-41.
64. Najafi P, Torki M. Performance, blood metabolites and im-

- munocompetence of broiler chicks fed diets included essential oils of medicinal herbs. *J Anim Vet Adv* 2010;9:1164-8.
65. Akira S, Uematsu S, Takeuchi O. Pathogen recognition and innate immunity. *Cell* 2006;124:783-801.
66. Medzhitov R, Janeway Jr CA. Innate immunity: impact on the adaptive immune response. *Curr Opin Immunol* 1997;9:4-9.
67. Li H, Zhang L, Chen L, et al. *Lactobacillus acidophilus* alleviates the inflammatory response to enterotoxigenic *Escherichia coli* K88 via inhibition of the NF- κ B and p38 mitogen-activated protein kinase signaling pathways in piglets. *BMC Microbiol* 2016;16:273.
68. MacKinnon KM, He H, Nerren JR, et al. Expression profile of toll-like receptors within the gastrointestinal tract of 2-day-old *Salmonella enteritidis*-infected broiler chickens. *Vet Microbiol* 2009;137:313-9.
69. Ramasamy KT, Verma P, Reddy MR. Toll-like receptors gene expression in the gastrointestinal tract of *Salmonella* serovar Pullorum-infected broiler chicken. *Appl Biochem Biotechnol* 2014;173:356-64.
70. Zhu J, Mohan C. Toll-like receptor signaling pathways--therapeutic opportunities. *Mediators Inflamm* 2010;2010:781235.
71. Li Q, Verma IM. NF-kappaB regulation in the immune system. *Nat Rev Immunol* 2002;2:725-34.
72. Byeon SE, Yi YS, Oh J, et al. The role of Src kinase in macrophage-mediated inflammatory responses. *Mediators Inflamm* 2012;2012:Article ID 512926.
73. Yu T, Yi YS, Yang Y, et al. The pivotal role of TBK1 in inflammatory responses mediated by macrophages. *Mediators Inflamm* 2012;2012:Article ID 979105.
74. Raingeaud J, Gupta S, Rogers JS, et al. Pro-inflammatory cytokines and environmental stress cause p38 mitogen-activated protein kinase activation by dual phosphorylation on tyrosine and threonine. *J Biol Chem* 1995;270:7420-6.
75. Genovese KJ, He H, Lowry VK, Kogut MH. Comparison of MAP and tyrosine kinase signaling in heterophils from commercial and wild-type turkeys. *Dev Comp Immunol* 2007;31:927-33.
76. Liu Y, Song M, Che TM, et al. Dietary plant extracts modulate gene expression profiles in ileal mucosa of weaned pigs after an *Escherichia coli* infection. *J Anim Sci* 2014;92:2050-62.
77. Coux O, Tanaka K, Goldberg AL. Structure and functions of the 20S and 26S proteasomes. *Annu Rev Biochem* 1996;65:801-47.
78. Taofiq O, Martins A, Barreiro MF, Ferreira ICFR. Anti-inflammatory potential of mushroom extracts and isolated metabolites. *Trends Food Sci Technol* 2016;50:193-210.
79. Ravi R, Bedi GC, Engstrom LW, et al. Regulation of death receptor expression and TRAIL/Apo2L-induced apoptosis by NF-kappaB. *Nat Cell Biol* 2001;3:409-16.
80. Gessner DK, Fiesel A, Most E, et al. Supplementation of a grape seed and grape marc meal extract decreases activities of the oxidative stress-responsive transcription factors NF- κ B and Nrf2 in the duodenal mucosa of pigs. *Acta Vet Scand* 2013;55:18.
81. Zhao FQ, Zhang ZW, Wang C, et al. The role of heat shock proteins in inflammatory injury induced by cold stress in chicken hearts. *Cell Stress Chaperones* 2013;18:773-83.
82. Klasing KC, Johnstone BJ. Monokines in growth and development. *Poult Sci* 1991;70:1781-9.
83. Johnson RW. Inhibition of growth by pro-inflammatory cytokines: an integrated view. *J Anim Sci* 1997;75:1244-55.
84. Koch AE, Kunkel SL, Harlow LA, et al. Enhanced production of monocyte chemoattractant protein-1 in rheumatoid arthritis. *J Clin Invest* 1992;90:772-9.
85. Ghareeb K, Awad WA, Soodoi C, et al. Effects of feed contaminant deoxynivalenol on plasma cytokines and mRNA expression of immune genes in the intestine of broiler chickens. *PLoS One* 2013;8:e71492.
86. Guzik TJ, Korbut R, Adamek-Guzik T. Nitric oxide and superoxide in inflammation and immune regulation. *J Physiol Pharmacol* 2003;54:469-87.
87. Harizi H, Gualde N. Pivotal role of PGE2 and IL-10 in the cross-regulation of dendritic cell-derived inflammatory mediators. *Cell Mol Immunol* 2006;3:271-7.
88. Hseu YC, Wu FY, Wu JJ, et al. Anti-inflammatory potential of *Antrodia Camphorata* through inhibition of iNOS, COX-2 and cytokines via the NF-kappaB pathway. *Int Immunopharmacol* 2005;5:1914-25.
89. Dy M, Schneider E. Histamine-cytokine connection in immunity and hematopoiesis. *Cytokine Growth Factor Rev* 2004;15:393-410.
90. Besedovsky HO, del Rey A, Klusman I, et al. Cytokines as modulators of the hypothalamus-pituitary-adrenal axis. *J Steroid Biochem Mol Biol* 1991;40:613-8.
91. Shini S, Shini A, Kaiser P. Cytokine and chemokine gene expression profiles in heterophils from chickens treated with corticosterone. *Stress* 2010;13:185-94.
92. Shini S, Huff GR, Shini A, Kaiser P. Understanding stress-induced immunosuppression: exploration of cytokine and chemokine gene profiles in chicken peripheral leukocytes. *Poult Sci* 2010;89:841-51.
93. Kabir SML. Avian colibacillosis and salmonellosis: a closer look at epidemiology, pathogenesis, diagnosis, control and public health concerns. *Int J Environ Res Public Health* 2010;7:89-114.
94. Jin Y, Hu Y, Han D, Wang M. Chronic Heat stress weakened the innate immunity and increased the virulence of highly pathogenic avian influenza virus H5N in mice. *J Biomed Biotechnol* 2011;2011:367846.
95. Youn HS, Lee JK, Choi YJ, et al. Cinnamaldehyde suppresses toll-like receptor 4 activation mediated through the inhibition of receptor oligomerization. *Biochem Pharmacol* 2008;75:494-502.
96. Guimarães MR, Leite FRM, Spolidorio LC, Kirkwood KL, Rossa

- C. Curcumin abrogates LPS-induced proinflammatory cytokines in RAW 264.7 macrophages. Evidence for novel mechanisms involving SOCS-1, -3 and p38 MAPK. *Arch Oral Biol* 2013;58:1309-17.
97. Liang D, Li F, Fu Y, et al. Thymol inhibits LPS-stimulated inflammatory response via down-regulation of NF- κ B and MAPK signaling pathways in mouse mammary epithelial cells. *Inflammation* 2014;37:214-22.
 98. Lima Mda S, Quintans-Júnior LJ, de Santana WA, et al. Anti-inflammatory effects of carvacrol: evidence for a key role of interleukin-10. *Eur J Pharmacol* 2013;699:112-7.
 99. Cho S, Choi Y, Park S, Park T. Carvacrol prevents diet-induced obesity by modulating gene expressions involved in adipogenesis and inflammation in mice fed with high-fat diet. *J Nutr Biochem* 2012;23:192-201.
 100. Zou Y, Xiang Q, Wang J, Peng J, Wei H. Oregano essential oil improves intestinal morphology and expression of tight junction proteins associated with modulation of selected intestinal bacteria and immune status in a pig model. *Biomed Res Int* 2016;2016:Article ID5436738.
 101. Lu H, Adedokun SA, Adeola L, Ajuwon KM. Anti-inflammatory effects of non-antibiotic alternatives in coccidia challenged broiler chickens. *J Poult Sci* 2014;51:14-21.
 102. Pannee C, Chandhane I, Wacharee L. Antiinflammatory effects of essential oil from the leaves of *Cinnamomum cassia* and cinnamaldehyde on lipopolysaccharide-stimulated J774A.1 cells. *J Adv Pharm Technol Res* 2014;5:164-70.
 103. Lee SH, Lillehoj HS, Jang SI. Cinnamaldehyde enhances *in vitro* parameters of immunity and reduces *in vivo* infection against avian coccidiosis. *Br J Nutr* 2011;106:862-9.
 104. Roth-Walter F, Moskovskich A, Gomez-Casado C, et al. Immune suppressive effect of cinnamaldehyde due to inhibition of proliferation and induction of apoptosis in immune cells: implications in cancer. *PLoS One* 2014;9:e108402.
 105. Kim GY, Kim KH, Lee SH, et al. Curcumin inhibits immunostimulatory function of dendritic cells: MAPKs and translocation of NF-kappa B as potential targets. *J Immunol* 2005;174:8116-24.
 106. Camacho-Barquero L, Villegas I, Sánchez-Calvo JM, et al. Curcumin, a *Curcuma longa* constituent, acts on MAPK p38 pathway modulating COX-2 and iNOS expression in chronic experimental colitis. *Int Immunopharmacol* 2007;7:333-42.
 107. Lubbad A, Oriowo MA, Khan I. Curcumin attenuates inflammation through inhibition of TLR-4 receptor in experimental colitis. *Mol Cell Biochem* 2009;322:127-35.
 108. Karnati HK, Pasupuleti SR, Kandi R, et al. TLR-4 signalling pathway: MyD88 independent pathway up-regulation in chicken breeds upon LPS treatment. *Vet Res Commun* 2015;39:73-8.
 109. Reyes-Gordillo K, Segovia J, Shibayama M, et al. Curcumin protects against acute liver damage in the rat by inhibiting NF-kappaB, proinflammatory cytokines production and oxidative stress. *Biochim Biophys Acta* 2007;1770:989-96.
 110. Lee SH, Lillehoj HS, Hong YH, et al. *In vitro* effects of plant and mushroom extracts on immunological function of chicken lymphocytes and macrophages. *Br Poult Sci* 2010;51:213-21.
 111. Yarru LP, Settivari RS, Gowda NK, et al. Effects of turmeric (*Curcuma longa*) on the expression of hepatic genes associated with biotransformation, antioxidant, and immune systems in broiler chicks fed aflatoxin. *Poult Sci* 2009;88:2620-7.
 112. Che Z, Liu Y, Wang H, et al. The protective effects of different mycotoxin adsorbents against blood and liver pathological changes induced by mold-contaminated feed in broilers. *Asian-Australas J Anim Sci* 2011;24:250-7.
 113. Malek M, Hasenstein JR, Lamont SJ. Analysis of chicken TLR4, CD28, MIF, MD-2, and LITAF genes in a *Salmonella enteritidis* resource population. *Poult Sci* 2004;83:544-9.
 114. Gilbert ER, Cox CM, Williams PM, et al. *Eimeria* species and genetic background influence the serum protein profile of broilers with coccidiosis. *PLoS One* 2011;6:e14636.
 115. Liu Y, Song M, Che TM, et al. Dietary plant extracts alleviate diarrhea and alter immune responses of weaned pigs experimentally infected with a pathogenic *Escherichia coli*. *J Anim Sci* 2013;91:5294-306.
 116. Liu Y, Song M, Che TM, et al. Effects of capsicum oleoresin, garlic botanical, and turmeric oleoresin on gene expression profile of ileal mucosa in weaned pigs. *J Anim Sci* 2014;92:3426-40.
 117. Amirghofran Z, Ahmadi H, Karimi MH, et al. *In vitro* inhibitory effects of thymol and carvacrol on dendritic cell activation and function. *Pharm Biol* 2016;54:1125-32.
 118. Kavooosi G, Teixeira da Silva JA, Saharkhiz, MJ. Inhibitory effects of *Zataria multiflora* essential oil and its main components on nitric oxide and hydrogen peroxide production in lipopolysaccharide-stimulated macrophages. *J Pharm Pharmacol* 2012;64:1491-500.
 119. Hassan FAM, Awad A. Impact of thyme powder (*Thymus vulgaris* L.) supplementation on gene expression profiles of cytokines and economic efficiency of broiler diets. *Environ Sci Pollut Res Int* 2017;24:15816-26.