



# Depolarizing Effectors of Bradykinin Signaling in Nociceptor Excitation in Pain Perception

Seung-In Choi and Sun Wook Hwang\*

Department of Biomedical Sciences and Department of Physiology, Korea University College of Medicine, Seoul 02841, Republic of Korea

## Abstract

Inflammation is one of the main causes of pathologic pain. Knowledge of the molecular links between inflammatory signals and pain-mediating neuronal signals is essential for understanding the mechanisms behind pain exacerbation. Some inflammatory mediators directly modulate the excitability of pain-mediating neurons by contacting the receptor molecules expressed in those neurons. For decades, many discoveries have accumulated regarding intraneuronal signals from receptor activation through electrical depolarization for bradykinin, a major inflammatory mediator that is able to both excite and sensitize pain-mediating nociceptor neurons. Here, we focus on the final effectors of depolarization, the neuronal ion channels, whose functionalities are specifically affected by bradykinin stimulation. Particular G-protein coupled signaling cascades specialized for each specific depolarizer ion channels are summarized. Some of these ion channels not only serve as downstream effectors but also play critical roles in relaying specific pain modalities such as thermal or mechanical pain. Accordingly, specific pain phenotypes altered by bradykinin stimulation are also discussed. Some members of the effector ion channels are both activated and sensitized by bradykinin-induced neuronal signaling, while others only sensitized or inhibited, which are also introduced. The present overview of the effect of bradykinin on nociceptor neuronal excitability at the molecular level may contribute to better understanding of an important aspect of inflammatory pain and help future design of further research on the components involved and pain modulating strategies.

**Key Words:** Bradykinin, Pain, Nociceptor neuron, Ion channel, Depolarization

## INTRODUCTION

Inflammation is frequently accompanied by pain, where several inflammatory pain mediators generated from inflamed tissues have been known to contribute to this pain induction, e.g., bradykinin, nerve growth factors, prostaglandins, and a group of cytokines (Patapoutian *et al.*, 2009). These mediators stimulate the primary nociceptor neurons innervating inflamed locations. The resultant firing of electrical signals is then transmitted to the brain, leading to the perception of pain. Acquiring information on the nature of the stimulatory mechanisms may help to improve therapeutic pain control strategies, and the relevant approaches at cellular and molecular levels have been conducted for many years. Rarely found regarding the properties for other mediators, bradykinin is able to induce action potential firing of the nociceptors as well as to sensitize those to other stimulations. The mechanisms seem to involve various ion channels that function as the final effec-

tors of excitatory outcomes. Although important frames for the molecular signaling that support the mechanisms were built in late 20th century, the molecular identities and detailed properties of most of the ionotropic players were reported during the 21st century.

As early as the 1950s, the hypothesis that bradykinin mediates pain via nociceptor excitation started to be confirmed in various experimental settings with *in vitro* and *in vivo* animal models, as well as human subjects. Administration of bradykinin to human skin and muscle clearly elicited pain perception (Armstrong *et al.*, 1957; Whalley *et al.*, 1987; Manning *et al.*, 1991; Kindgen-Milles *et al.*, 1994; Babenko *et al.*, 1999). Injections to the skin, vascular areas, and the peritoneal cavity caused nocifensive reflexes in model animals including mice, rats, cats, rabbits, dogs, and monkeys (Kumazawa and Mizumura, 1976; Steranka *et al.*, 1988; Walter *et al.*, 1989; Khan *et al.*, 1992; Hong and Abbott, 1994; Griesbacher *et al.*, 1998; Katanosaka *et al.*, 2008). Fiber recordings revealed that

**Open Access** <https://doi.org/10.4062/biomolther.2017.127>

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.


Received Jun 17, 2017 Revised Oct 13, 2017 Accepted Oct 24, 2017

Published Online Jan 30, 2018

\*Corresponding Author

E-mail: sunhwang@korea.ac.kr

Tel: +82-2-2286-1204, Fax: +82-2-925-5492



**Fig. 1.** Summary of the roles of important effector ion channels which account for bradykinin-induced excitation of pain-mediating nociceptors. AA, arachidonic acid; AC, adenylate cyclase; AKAP, A kinase anchoring protein; ANO1, anoctamin 1; B1R, bradykinin receptor B1; B2R, bradykinin receptor B2; BK, bradykinin; cAMP, 3',5'-cyclic adenosine monophosphate; COX, cyclooxygenase; DAG, diacylglycerol; EP/IP, prostaglandin E2 receptor and prostaglandin I2 receptor; HPETE, hydroperoxyeicosatetraenoic acid; IKCa, Ca<sup>2+</sup>-activated K<sup>+</sup> channels; IP<sub>3</sub>, inositol 1,4,5-trisphosphate; KCNQ, voltage-gated K<sup>+</sup> channel subfamily KCNQ; LOX, lipoxygenase; PG, prostaglandin; PIP<sub>2</sub>, phosphatidylinositol 4,5-bisphosphate; PKA, protein kinase A; PKC, protein kinase C; PLA2, phospholipase A2; TRPA1, transient receptor potential ankyrin subtype 1; TRPV1, transient receptor potential vanilloid subtype 1.

the nociceptor depolarization initiated those painful outcomes (Juan and Lembeck, 1974; Chahl and Iggo, 1977; Dray *et al.*, 1992; Soukhova-O'Hare *et al.*, 2006), in which models utilizing testis-spermatoc nerve and skin-saphenous nerve preparations have greatly contributed to the provision of fundamental information on bradykinin-controlling sensory modalities and phases, nociceptor categorizing, and signaling participants (Beck and Handwerker, 1974; Kumazawa and Mizumura, 1976). As a result, it is now firmly known that the polymodal nociceptors comprising the unmyelinated C and thinly myelinated A $\delta$  neurons are responsible for bradykinin-induced pain, that the B2 receptor is more constitutively responsible for bradykinin detection than the B1 receptor, and that both discharging of action potentials and lowering of its threshold can be caused by bradykinin action (Mizumura *et al.*, 2009).

Following this, the molecular evidence has kept being corroborated regarding bradykinin receptor-mediated signals, using extended technologies such as culture platforms, molecular biology, genetics, and the patch clamp. Bradykinin acts on the B1 and B2 receptors that are among the metabotropic G protein-coupled receptors (GPCRs) expressed at the surface membrane (Burgess *et al.*, 1989; McGuirk *et al.*, 1989; McGehee and Oxford, 1991; Dray *et al.*, 1992; McGuirk and Dolphin, 1992). The majority of the downstream information was obtained from B2 studies, but as for many molecular processes, both receptors have been shown to share similar mechanisms of action (Petho and Reeh, 2012). Commonly, Gq/11-mediated phospholipase C (PLC) and Gi/o-mediated phospholipase A2 (PLA2) activation lead to diverse cellular effects. In nociceptor neurons, several depolarizing effectors are activated or positively regulated (sensitized) through such signaling, which are critical steps necessary for action potential firing or threshold lowering. Here we summarize the identities of the depolarizing molecules and bradykinin-related mechanisms for activation and sensitization.

## TRANSIENT RECEPTOR POTENTIAL VANILLOID SUBTYPE 1 ION CHANNEL

Transient Receptor Potential Vanilloid subtype 1 ion channel (TRPV1) functions as a receptor and a cation channel in nociceptor sensory neurons. Sensitive to noxious temperature ranges (>43°C), protons (pH <5.5), and pungent chemicals (e.g., capsaicin), TRPV1 responds by opening its pore. Cation influx through TRPV1 depolarizes the nociceptor membrane, discharging action potentials when the membrane voltage reaches its firing threshold. Other mechanisms for activation and activity modulation have been revealed, and bradykinin has been shown to be tightly linked.

### Bradykinin-induced activation of TRPV1 via arachidonic acid metabolism

TRPV1-mediated action potential spike generation upon bradykinin exposure has successfully been repeated in the primary sensory afferents from various sources, including cutaneous nociceptors, cardiac afferents, jejunal afferents, and tracheobronchial afferents (Fig. 1) (Carr *et al.*, 2003; Pan and Chen, 2004; Rong *et al.*, 2004; Lee *et al.*, 2005a). Research efforts have been put into seeking the link between bradykinin-initiated G protein signaling and depolarizing effector functions. Increased production of arachidonic acid by bradykinin and its further metabolism has been considered an important candidate for the signaling (Thayer *et al.*, 1988; Burgess *et al.*, 1989; Gammon *et al.*, 1989). Not only in neurons but also in other tissues, Gi/o mediated arachidonic acid liberation through bilayer digestion of PLA2 activated by bradykinin has been proposed to be involved (Burch and Axelrod, 1987; Gammon *et al.*, 1989; Yanaga *et al.*, 1991). The resultant excitation and sensitization of the nociceptor has also been demonstrated (Taiwo *et al.*, 1990; Ferreira *et al.*, 2004). The role of members of the lipoxygenase (LOX) in furthering arachidonic acid

metabolism has been raised for the immediate depolarization caused by bradykinin (McGuirk and Dolphin, 1992; Shin *et al.*, 2002; Carr *et al.*, 2003; Wu and Pan, 2007). Formation of 12-LOX products like 12(S)-hydroxyeicosatetraenoic acid (12(S)-HETE) was also confirmed in nociceptor neurons and skin (Wang *et al.*, 1999; Shin *et al.*, 2002; Tang *et al.*, 2005).

Production of arachidonic acid can result from PLC metabolism, as well as from PLA2-mediated one. Diacylglycerol (DAG) produced via PLC action may further be converted into arachidonic acid by DAG lipase and monoacylglycerol lipase (Gammon *et al.*, 1989; Allen *et al.*, 1992; Dray *et al.*, 1992). Attenuation of locally injected bradykinin-elicited acute nocifensive behaviors that reflect immediate sensory action potential discharge is achieved via PLC inhibitors, as well as PLA2 and 5-LOX inhibitors (Ferreira *et al.*, 2004).

LOX metabolites have been shown to activate TRPV1 (Hwang *et al.*, 2000). A number of HETEs, hydroperoxyeicosatetraenoic acids (HPETEs), and leukotrienes displayed this activity, where 12(S)-HPETE has the greatest potency. A later study from the same group reported the B2 receptor-PLA2-LOX axis for the signaling cascade for TRPV1 activation (Shin *et al.*, 2002). The same signaling paradigm was successfully repeated by a pharmacological approach for visceral afferents by using multiple TRPV1 blockers and LOX inhibitors (Carr *et al.*, 2003). LOX product pools that have TRPV1 activating potential and the neuronal outcomes other than nociception for example memory, are both being currently extended (Gibson *et al.*, 2008; Gregus *et al.*, 2012). Although the binding pocket for LOX-produced ligands or TRPV1 is less conclusively determined, their structural similarity with capsaicin as predicted by 3-dimensional simulation suggests that the capsaicin binding site, the S4-S5 intracellular linker region of TRPV1 protein is a strong candidate (Ikeda *et al.*, 2001b; Gao *et al.*, 2016).

### Bradykinin-induced activation of TRPV1 via protein kinase C

Gq/11-activated PLC metabolizes the membrane bilayer phosphatidylinositol 4,5-bisphosphate (PIP2) into DAG and inositol 1,4,5-trisphosphate (IP3). The increased level of these two products upon bradykinin stimulation has been reported in nociceptor dorsal root ganglionic (DRG) neurons, indicating the activation of Gq/11-PLC axis by bradykinin (Thayer *et al.*, 1988; Burgess *et al.*, 1989; Gammon *et al.*, 1989; Kano *et al.*, 1994; Tang *et al.*, 2004). Both of the two messengers contribute to protein kinase C (PKC) activation.

Evidence has been accumulating that PKC activation upon bradykinin exposure leads to an immediate excitation in studies with various platforms using ectopically receptor expressing cells (Crandall *et al.*, 2002), cultured sensory neurons (Rang and Ritchie, 1988; Burgess *et al.*, 1989; McGehee and Oxford, 1991; McGuirk and Dolphin, 1992), afferent nerve fibers (Mizumura *et al.*, 1997; Guo *et al.*, 1998, 1999), spinal cord-tail preparations (Dray *et al.*, 1988, 1992), or animals with nocifensive behaviors (Ferreira *et al.*, 2004). Suppression of excitatory responses by pharmacological inhibition of PKC and mimicking of depolarization when exposed to PKC-activating phorbol esters support the finding. The excitatory effect appears to be caused by the increased permeability of the neuronal membrane to both Na<sup>+</sup> and K<sup>+</sup> ions, indicating that nonselective cation channels are probably a final effector for this bradykinin-induced PKC action (Rang and Ritchie, 1988; Burgess *et al.*, 1989; McGehee and Oxford, 1991).

Further research revealed that TRPV1 participates as the ion-permeating effector and suggested that bradykinin leads to TRPV1 activation by PKC phosphorylation and that PKC $\epsilon$  plays a crucial role (Premkumar and Ahern, 2000; Vellani *et al.*, 2004). When strongly depolarized, sensory afferents secrete peptidergic transmitters, specifically Substance P and calcitonin gene-related peptide, which is augmented by PKC activation and reduced by PKC downregulation (Barber and Vasko, 1996). Two serine residues located in the intracellular S2-S3 linker and N-terminal end of TRPV1 protein were identified as PKC $\epsilon$  phosphorylation targets (Numazaki *et al.*, 2002; Bhave *et al.*, 2003; Choi *et al.*, 2014). Although neuronal excitation by PKC has readily been repeated, ectopic expression systems have often shown less effective induction of the inward current but relatively prominent sensitization of the TRPV1 response to other stimulation, which remains to be quantitatively dissected.

### Bradykinin-induced sensitization of TRPV1 activity

In comparison with an acute excitatory action, constantly sensitized nociception caused by a mediator may more broadly explain pathologic pain mechanisms. Since TRPV1 is the major heat sensing molecule, heat hyperalgesia induced by bradykinin, which has long been studied in pain research, may putatively involve changes in TRPV1 activity. Therefore, here we provide an overview of the role of bradykinin in pathology-induced heat hyperalgesia and then discuss the evidence supporting the possible participation of TRPV1 in this type of bradykinin-exacerbated thermal pain. Different from acute nociception where data were produced mostly in B2 receptor setting, the focus may include both B1 and B2-mediated mechanisms underlying pathology-induced chronic nociception, since roles for inducible B1 may emerge in certain disease states. A number of specific pathologies may even show pronounced dependence on B1 function. Nonetheless, both receptors likely share the intracellular signaling mechanisms for effector sensitization.

**B1 receptor-dependent pathologic pain:** Since the 1980s, B2 receptor involvement has been extensively demonstrated in relatively short-term inflammation models primed with an adjuvant carrageenan or other mediator treatments (Costello and Hargreaves, 1989; Ferreira *et al.*, 1993b; Ikeda *et al.*, 2001a). On the other hand, B1 receptor seems to be more tightly involved in heat hyperalgesia in relatively chronic inflammatory pain models such as the complete Freund's adjuvant (CFA)-induced inflammation model. While B2 knockout mice failed to show any difference in comparison with wild types, either B1 knockouts or B1 antagonism leads to reduced heat hyperalgesia (Rupniak *et al.*, 1997; Ferreira *et al.*, 2001; Porreca *et al.*, 2006). Because of the ignorable difference in CFA-induced edema between wild types and B1 knockouts, B1 is thought to be involved in heightened neuronal excitability rather than inflammation itself (Ferreira *et al.*, 2001). In diabetic neuropathy models, B1 knockouts are resistant to development of the heat hyperalgesia, and treatment with a B1 antagonist was effective in preventing heat hyperalgesia in naïve animals (Gabra and Sirois, 2002, 2003a, 2003b; Gabra *et al.*, 2005a, 2005b). In a brachial plexus avulsion model, B1 knockouts but not B2 knockouts have shown prolonged resistance to heat hyperalgesia (Quintão *et al.*, 2008). Pharmacological studies on ultraviolet (UV) irradiation models have also shown B1 dominance (Perkins and Kelly, 1993; Perkins

*et al.*, 1993; Gougat *et al.*, 2004). Both the peptidergic antagonist des-Arg<sup>9</sup>,Leu<sup>8</sup>-bradykinin and a synthetic B1 antagonist SSR240612 commonly prevented UV-induced heat hyperalgesia, whereas the effect of HOE 140, a B2 antagonist, was largely limited. The hyperalgesia was further aggravated by a relatively selective B1 agonist des-Arg<sup>9</sup>-bradykinin and reversed only by the B1 antagonist.

**B1 & B2 receptor-dependent pathologic pain:** In neuropathic pain models, both B1 and B2 receptor-mediated mechanisms are generally important (Levy and Zochodne, 2000; Yamaguchi-Sase *et al.*, 2003; Ferreira *et al.*, 2005; Petcu *et al.*, 2008; Luiz *et al.*, 2010). In the models of chronic constriction injury, infraorbital nerve constriction injury, and partial sciatic nerve ligation, selective pharmacological antagonism of either of the receptor types was effective against the putatively TRPV1-mediated heat hyperalgesia, as well as cold hyperalgesia and mechanical allodynia. Heat hyperalgesia occurring in a rat plantar incision model was once shown to be unrelated to bradykinin-mediated mechanisms (Leonard *et al.*, 2004). Later, a contradictory result that the heat hyperalgesia was partially reversed by treatment with either B1 or B2 receptor antagonist was obtained in a different laboratory (Füredi *et al.*, 2010). In the same model, treatment with an LOX inhibitor or a TRPV1 antagonist was also effective. Interestingly, in the same study, heat injury-evoked heat hyperalgesia was attenuated only by B2 antagonist treatment.

**Bradykinin-induced heat hypersensitivity:** Injection of bradykinin itself has also been shown to augment heat pain sensitivity in humans, monkeys, and rats (Manning *et al.*, 1991; Khan *et al.*, 1992; Schuligoi *et al.*, 1994; Griesbacher *et al.*, 1998). It is commonly likely that the heat sensitivity was left-shifted with lowered heat threshold by bradykinin injection. There are several different points when speculating possible mechanisms that could explain direct excitation and sensitization. Direct nociception in response to bradykinin generally undergoes strong tachyphylaxis, but such sensitization appears to be relatively persistent in time scale. In-depth analyses at the cellular or molecular levels that are mentioned below have shown that the sensitizing effect sometimes occurs in the absence of direct excitation (Beck and Handwerker, 1974; Kumazawa *et al.*, 1991; Khan *et al.*, 1992). Nonetheless, nociceptors that more readily fire upon bradykinin exposure appeared to tend to be more sensitized in heat responsiveness (Kumazawa *et al.*, 1991; Liang *et al.*, 2001). Common PKC-centered machinery is hypothesized to be responsible for both excitation and sensitization, which still requires further careful dissection to understand how those differentiated outcomes are realized.

**The sensitizing action of bradykinin on nociceptors:** After feline nociceptors were once demonstrated to be sensitized by acute bradykinin exposure of their termini in terms of heat-evoked spike discharges in an *in vivo* model, many similar *in vitro* or *ex vivo* results were produced, again for example, in rodent skin-saphenous nerve and canine testis-spermatic nerve models (Beck and Handwerker, 1974; Lang *et al.*, 1990; Kumazawa *et al.*, 1991). As shown in the *in vivo* experiments mentioned above, the potency and efficacy of heat-induced electrical responses were increased by bradykinin stimulation of the relevant receptive fields, which was mainly observed in unmyelinated C- or thinly myelinated A $\delta$  nociceptors with polymodality (Kumazawa *et al.*, 1991; Koltzenburg *et al.*, 1992; Haake *et al.*, 1996; Liang *et al.*, 2001). Such facilitation

occurred at lower doses than needed for bradykinin-evoked excitation, and moreover, subpopulations of nociceptors that were without bradykinin- or heat-evoked excitation in a naive stage became sensitive to heat by bradykinin exposure (Kumazawa *et al.*, 1991; Liang *et al.*, 2001). The observed population enlargement is unlikely to be due to an elevated expression of TRPV1 at the surface membrane as this failed to be demonstrated in a more recent study (Camprubi-Robles *et al.*, 2009). Although the experiment did not manipulate heat, research revealed that the capsaicin responses in trachea-innervating vagal C-fibers was sensitized by bradykinin, underlying cough exacerbation upon bradykinin accumulation as an adverse effect of treatment with angiotensin converting enzyme inhibitors for hypertension (Fox *et al.*, 1996). B2 receptor participation was confirmed in the models above.

**TRPV1 as a principal actuator for bradykinin-induced heat sensitization:** As mentioned above, PKC activation is involved in TRPV1 activation and sensitization. Electrophysiological recordings of canine testis-spermatic nerve preparations raised a role for PKC in the bradykinin-induced sensitization of the heat responses (Mizumura *et al.*, 1997). PKC $\epsilon$  phosphorylation initiated by bradykinin was proposed to sensitize the native heat-activated cation channels of cultured nociceptor neurons (Cesare and McNaughton, 1996; Cesare *et al.*, 1999). This was successfully repeated in TRPV1 experiments after its genetic identification and the temperature threshold for TRPV1 activation was lowered by PKC phosphorylation (Vellani *et al.*, 2001; Sugiura *et al.*, 2002). Not only to heat but also to other activators such as protons and capsaicin, TRPV1 responses were sensitized by PKC phosphorylation in several different experimental models (Stucky *et al.*, 1998; Crandall *et al.*, 2002; Lee *et al.*, 2005b; Camprubi-Robles *et al.*, 2009). However, it remains to be elucidated if inducible B1 receptor may utilize the same pathway.

**Molecular mechanisms for TRPV1 sensitization by PKC phosphorylation:** TRPV1 protein contains a number of target amino acid residues for phosphorylation by various protein kinases. The phosphorylation of these residues largely contributes to the facilitation of TRPV1 activity but it is likely that bradykinin mainly utilizes PKC for its TRPV1 sensitization according to an *in vitro* analysis of phosphorylated proteins (Lee *et al.*, 2005b). PKC has been shown to directly phosphorylate two TRPV1 serine residues that are located in the first intracellular linker region between the S2 and S3 transmembrane domains, and in the C-terminal (Numazaki *et al.*, 2002; Bhave *et al.*, 2003; Wang *et al.*, 2015). Mutant TRPV1 that was missing these target sequences were tolerant in terms of sensitization upon bradykinin treatment. Interestingly, an adaptor protein seems to be critical to access to the target residues by PKC. Members of A kinase anchoring proteins (AKAPs) are able to modulate intracellular signaling by recruiting diverse kinase and phosphatase enzymes (Fischer and McNaughton, 2014). The activity of some of ion channels is known to be controlled by this modulation when these proteins form a complex, the best known example being the interaction of TRPV1 with AKAP79/150 (AKAP79 indicates the human homolog of the rodent AKAP150) (Zhang *et al.*, 2008). Specifically, AKAP79/150 bound to the C-terminal of TRPV1 protein has been shown to greatly facilitate PKC action by physically recruiting it upon bradykinin stimulation.

**Other proposed mechanisms for bradykinin-induced TRPV1 sensitization:** As mentioned above, pharmacological inhibition

or genetic downregulation studies have demonstrated that PKC may be a predominant factor required for bradykinin-induced TRPV1 sensitization. In addition, different molecular mechanisms also appear to contribute to this sensitization under bradykinin-mediated pathway. PLC is an upstream enzyme of PKC, which consumes bilayer PIP2 to produce IP3 and DAG. PIP2 seems to bind to TRPV1 protein in its resting state, tonically inhibiting TRPV1 activity. In response to bradykinin stimulation however, PIP2 is consumed by Gq-mediated PLC, thereby disinhibiting TRPV1. Responses to capsaicin and protons have been shown to be augmented by this mechanism, although sensitization to heat by bradykinin remains to be explored (Chuang *et al.*, 2001; Prescott and Julius, 2003). As arachidonic acid metabolizing enzymes, LOXs generate a number of endogenous TRPV1 activators. Interestingly, 12-LOX inhibition was also effective at erasing bradykinin-induced heat sensitization of TRPV1 (Shin *et al.*, 2002). 12(S)-HPETE or 12(S)-HETE, its metabolites may therefore not only activate TRPV1, but also facilitate the heat response, which remains to be experimentally confirmed.

Complicated results were obtained from research exploring the role of a different group of arachidonic acid metabolizing enzymes, cyclooxygenases (COXs), as mentioned in the *Unsolved problems* section. COX activation undoubtedly contributes as a downstream from bradykinin signaling to nociceptor sensitization and excitation but it remains under debate where among nociceptors and surrounding cells COXs plays the predominant role. Whatever the major source of prostaglandins is, the activation of neuronal G-protein-coupled receptors, including EP1, EP4, and IP, by secreted prostaglandins causes activation of not only PKC, but also protein kinase A (PKA) that also phosphorylates and sensitizes TRPV1 (Moriyama *et al.*, 2005).

### Unsolved problems

While the major heat sensor TRPV1 has been massively studied regarding bradykinin-induced nociception, the contribution of thermosensitive cousins of TRPV1 in the same superfamily, including TRPV2 and TRPV3, have largely been unexplored. Interestingly, another polymodal TRPV member, TRPV4, which is sensitive to warm temperatures and mechanical stretches, has very recently been shown to mediate at least mechanical hyperalgesia of a mouse under paclitaxel-induced peripheral neuropathy via bradykinin-B1/B2-PKC $\epsilon$  axis (Costa *et al.*, 2018). In addition, for bradykinin-induced sensitization, effector components other than such direct temperature sensors are possibly involved. For example, knockouts of a tetrodotoxin-insensitive voltage-gated Na<sup>+</sup> channel, Nav1.9, have displayed a blunted phenotype in bradykinin-induced heat hyperalgesia (Amaya *et al.*, 2006).

Still, it remains as theoretical predictions how exactly PKC underlies both activation and sensitization of TRPV1. It is reasonable to hypothesize that these differential outcomes might result from phosphorylation of different serine residues, which possibly contributes to channel gating in different manners. Alternatively, at the point of AKAP79/150 action, the differential roles of PKC could be diverged. Although it seems to be limited to a particular tissue like cutaneous areas, the transcellular mechanism involving prostaglandins may exclusively be engaged in sensitization.

The central molecular mechanisms for TRPV1 activation and sensitization have firmly been shown to engage voltage-

dependence (Voets *et al.*, 2004). The relevant stimuli, including heat, capsaicin, protons, endogenous ligands, phosphorylations, etc., appear to converge into the leftward shift of TRPV1 voltage-dependence. In this regard, given multiple stimuli may be additive or synergistic for enhancing TRPV1 voltage sensitivity, which can be seen as one stimulus facilitates the response to others (Vyklícký *et al.*, 1999). Accordingly, bradykinin-induced phosphorylation may left-shift the effect of heat on TRPV1 voltage-dependence, leading to augmented firing of the nociceptors upon heat stimulation. An extreme shift may enable TRPV1 activation by ambient temperatures, which can be seen as bradykinin directly excites the neurons. Since TRPV1 is known to essentially undergo Ca<sup>2+</sup>-induced desensitization to itself, Reeh and colleagues have suggested that, prior to desensitization, bradykinin may induce short-term direct firing, and that the relatively blunted shift of TRPV1 sensitivity may look as if its lowered heat threshold during desensitized state (Reeh and Pethö, 2000; Liang *et al.*, 2001). A newly found mechanism unrelated to voltage dependence or even to other signal transductions mentioned above has recently been proposed. Exocytic trafficking of TRPV1-containing vesicle may selectively contribute to the sensitization of peptidifergic nociceptors, which awaits replication (Mathivanan *et al.*, 2016).

The major tissue type where bradykinin induces COX-dependent prostaglandin secretion remains elusive. While nociceptor neurons has been raised as a crucial source of prostaglandins in the pharmacological inhibition of COXs and labeling of COX expression (Mizumura *et al.*, 1987; Kumazawa *et al.*, 1991; Dray *et al.*, 1992; Rueff and Dray, 1993; Vasko *et al.*, 1994; Weinreich *et al.*, 1995; Maubach and Grundy, 1999; Jenkins *et al.*, 2003; Oshita *et al.*, 2005; Inoue *et al.*, 2006; Tang *et al.*, 2006; Jackson *et al.*, 2007), other studies have failed to corroborate this finding and have instead suggested surrounding tissues innervated by neuronal termini (Lembeck and Juan, 1974; Lembeck *et al.*, 1976; Juan, 1977; Franco-Cereceda, 1989; McGuirk and Dolphin, 1992; Fox *et al.*, 1993; Sauer *et al.*, 1998; Kajekar *et al.*, 1999; Sauer *et al.*, 2000; Pethö *et al.*, 2001; Shin *et al.*, 2002; Ferreira *et al.*, 2004). Possibly, COXs in non-neuronal cells may be of more importance during the initial stages of bradykinin action and a relatively long term exposure (~hours or longer) is needed for the induction of neuronal expression of COXs (Oshita *et al.*, 2005). However, the relative importance of COX-1 and COX-2 needs to be fully assessed (Jackson *et al.*, 2007; Mayer *et al.*, 2007). In addition, many lines of pharmacological evidence for COX participation include the reduction in bradykinin-evoked immediate excitation of nociceptors by COX inhibition. On the other hand, the protein kinase-mediated molecular mechanisms of bradykinin action mentioned above only explain sensitized heat responses.

### TRANSIENT RECEPTOR POTENTIAL ANKYRIN SUBTYPE 1 ION CHANNEL

Transient Receptor Potential ankyrin subtype 1 (TRPA1) is a comparably important TRP channel in nociception with regards to polymodality. The opening of TRPA1 depolarizes polymodal nociceptors in response to temperatures <17°C, mechanical stretches, and reactive irritants (e.g., mustard oil, cinnamaldehyde, air pollutants, prostaglandins with  $\alpha,\beta$ -

carbonyl carbon, etc.) (Bang and Hwang, 2009). Inflammatory pain mediators such as bradykinin also appear to positively modulate TRPA1 activity, leading to pain exacerbation.

### Bradykinin-induced activation of TRPA1 via arachidonic acid metabolism

In an early study where cinnamaldehyde was first found as a specific agonist for TRPA1, bradykinin also displayed an ability to activate TRPA1 via intracellular signaling. In a heterologous expression system co-transfected with DNAs encoding B2 receptor and TRPA1, immediate TRPA1-specific responses occurred upon bradykinin perfusion, as measured by TRPA1-mediated electrical currents and  $Ca^{2+}$  influx (Bardell *et al.*, 2004). Perfusions of a membrane-permeable DAG analog and an arachidonic acid analog also replicated this response, indicating that the bradykinin pathway may utilize PLC (perhaps together with DAG lipase) for TRPA1 activation and possibly PLA2. Although further downstream signaling has not been thoroughly explored, it is also possible that other substances more metabolized from arachidonic acid can activate TRPA1. For example, a number of prostaglandins (PGs) have also been shown to activate TRPA1 (Andersson *et al.*, 2008; Materazzi *et al.*, 2008). The PGs include 15-deoxy- $\Delta$ 12, 14-PGJ2,  $\Delta$ 12-PGJ2, PGA1, PGA2, and 8-iso PGA2, all of which contain a reactive carbon that can covalently bind to reactive serine or cysteine residues in TRPA1 protein in the same manner that mustard oil and cinnamaldehyde interact (Hinman *et al.*, 2006; Macpherson *et al.*, 2007). Because the PGs are non-enzymatically generated from COX products such as PGH2 and PGE2, the bradykinin-mediated COX activation mentioned above may be linked to depolarization resulting from TRPA1 activation. Whatever the strongest contributor among the metabolites is, bradykinin appears to depolarize nociceptor neurons not only via TRPV1 but also via TRPA1, which was confirmed in TRPA1 knockout studies through action potential firing and nocifensive behaviors (Bautista *et al.*, 2006; Kwan *et al.*, 2006). TRPA1 knockouts have also exhibited reduced hypersensitivity in response to bradykinin (Bautista *et al.*, 2006; Kwan *et al.*, 2006).

### Bradykinin-induced sensitization of TRPA1 activity

**Molecular mechanisms for TRPA1 sensitization by bradykinin:** Not only activation, but also sensitization of TRPA1 when exposed to bradykinin occurs in nociceptor neurons (Fig. 1). The same research group has suggested that there exist two parallel signaling pathways for bradykinin-induced TRPA1 sensitization, which were the PLC and PKC pathways (Dai *et al.*, 2007; Wang *et al.*, 2008). However, this awaits further confirmation because of some discrepancies. The Gq/11-mediated PLC pathway was raised first (Dai *et al.*, 2007). Without further requirement of downstream signaling such as PKC activation, bilayer PIP2 consumption has been demonstrated to disinhibit TRPA1, which seems to adequately explain enhanced TRPA1 activity observed when exposed to a known specific agonist for TRPA1. This study proposed that the membrane PIP2 intrinsically masks the channel's activity in the resting state, which was confirmed by an independent study showing that the addition of intracellular PIP2 inhibits TRPA1 opening (Kim *et al.*, 2008). Two other studies have shown the opposite effect, where TRPA1 is directly activated by PIP2 (Akopian *et al.*, 2007; Karashima *et al.*, 2008), while another group failed to show this activation (Kim and Cavanaugh,

2007). TRPV1 has once been demonstrated to be either positively or negatively modulated by the presence of PIP2, which may depend on the extent of channel activation, which is not shown yet to be the case for TRPA1 modulation (Lukacs *et al.*, 2007).

Another proposed mechanism for TRPA1 sensitization by bradykinin is via the PKA. As mentioned above, TRPV1 can be sensitized in a similar manner, but PKA action appears to take a relatively long time (~10 minutes) and requires PG synthesis as an upstream signal. However, fast sensitization of TRPA1 was shown to be dependent on Gs-mediated adenylate cyclase activity and subsequent PKA activation but unlikely with PG production. Such Gs-mediated signaling by bradykinin stimulation has been reported to occur in different cell types (Stevens *et al.*, 1994; Liebmann *et al.*, 1996; Bae *et al.*, 2003). TRPA1, as well as TRPV1, needs further repetition in this regard.

**Evidence from nociceptors and animals:** Formalin and mustard oil are TRPA1-selective activators that were used as experimental stimulants for nociceptor excitation in the pain research field prior to their relationship with TRPA1 being discovered. Acute nocifensive behaviors are typically evoked by intraplantar administration of either of formalin or mustard oil, and were shown to be significantly facilitated by injections in the same location of bradykinin itself or bradykinin receptor specific agonists (De Campos *et al.*, 1998; Wang *et al.*, 2008). In addition to these chemical-specific modalities, TRPA1 seems to be involved in noxiously mechanical ones to an extent due to its intrinsic mechanosensitivity (Kwan *et al.*, 2006; Petrus *et al.*, 2007; Brierley *et al.*, 2009; Kwan *et al.*, 2009; Yu and Ouyang, 2009). Nociceptor firing in response to mechanical stimuli was significantly diminished in TRPA1-deficient mice or by pharmacological antagonism (Brierley *et al.*, 2005; Brierley *et al.*, 2009; Yu and Ouyang, 2009).

Therefore, it is worth speculating the relationship between TRPA1 and the molecular mechanisms underlying bradykinin-elicited mechanical hypersensitivities that have been proposed from behavioral studies. Protein kinase G (PKG) has been relatively unexplored with regards to TRPA1 modulation, and PKG inhibition has been shown to reduce bradykinin-induced mechanical hyperalgesia (Nakamura *et al.*, 1996). The same study actually suggested that the nitric oxide synthase (NOS)-guanylate cyclase (GC)-PKG cascade mediates the mechanical hypersensitivity. NOS is possibly activated by PLC-IP3-mobilized  $Ca^{2+}$ . However, NO itself is known to react with TRPA1 protein and seemed to be inadequate to lead to hyperalgesia despite the heightened level of NO, indicating that further signal amplification via subsequent GC and PKG activation may be required.

Other studies have raised the role of the PLA2-COX pathway in the development of bradykinin-induced mechanical hyperalgesia (Taiwo and Levine, 1988; Taiwo *et al.*, 1990). COX induction by bradykinin may require a transcellular process in the sensitized heat responses mentioned above. In a multitude of studies on this mechanical hypersensitivity, details particularly including complicated upstreams for the COX mechanism have been described (Ferreira *et al.*, 1993a; Ferreira *et al.*, 1993b; Poole *et al.*, 1999; Cunha *et al.*, 2007). Collectively, the causality seems to involve a transcellular mechanism and to be that the sequential secretions of TNF- $\alpha$ , other cytokines including interleukin-1 $\beta$  (IL-1 $\beta$ ), IL-6, and IL-8, and then finally prostaglandins and sympathetic amines. It appears that the

major sources of TNF- $\alpha$  and sympathetic amines may be vicinal macrophages and sympathetic nerve termini respectively, indicating that the bradykinin-induced mechanical hyperalgesia may also involve transcellular processes. Seeing as surgical sympathectomy alleviated mechanical, but not heat hyperalgesia, the downstream sympathetic amines from the postganglionic neurons may only control the peripheral mechanical function of nociceptors (Koltzenburg *et al.*, 1992; Meyer *et al.*, 1992; Schuligoi *et al.*, 1994).

Interestingly, the occurrence of mechanical hypersensitivity appears to depend on the location of bradykinin accumulation (Manning *et al.*, 1991; Khan *et al.*, 1992; Khasar *et al.*, 1993). This may again indicate that not only the changes in the functionality of nociceptors but also transcellular interactions where specific cellular components that additionally participate are important. In accordance with a study showing that mechanical hyperalgesia was induced by intradermal injections of bradykinin or PGE<sub>2</sub>, but not readily by subcutaneous treatments, later studies using a diverse range of nociceptor markers demonstrated that nociceptor termini are differentially distributed in terms of the depth of the skin layer, and that a more superficial subpopulation may supposedly be responsible for mechanical hyperalgesia (Khasar *et al.*, 1993; Zylka *et al.*, 2005; Cavanaugh *et al.*, 2009). Interestingly, it has recently been demonstrated that TRPA1 in the central terminal of nociceptors also contribute to the development of mechanical hyperalgesia by participating in B1 receptor-dependent spinal neuroinflammation where intracellular and transcellular mechanisms may operate in a similar manner as mentioned above (Meotti *et al.*, 2017).

**TRPA1 contributes to bradykinin-induced heat hyperalgesia:** Although TRPA1 is not intrinsically sensitive to hot temperatures, TRPA1 knockouts have exhibited a blunted phenotype in bradykinin-induced heat hyperalgesia when compared to wild type littermates (Bautista *et al.*, 2006). In the same study, however, CFA-induced heat hyperalgesia was not affected by TRPA1 deletion. TRPA1 may only mildly influence TRPV1-based heat sensation. Intracellular Ca<sup>2+</sup> elevated first by TRPV1 opening in response to heat was once proposed to link TRPV1 activation to the subsequent TRPA1 activation. However a current theory is that a part of TRPV1 and TRPA1 proteins may be physically coupled to form a sensory complex located on the surface of the nociceptor terminal (Staruschenko *et al.*, 2010; Fischer *et al.*, 2014; Weng *et al.*, 2015).

**Difference between TRPA1 and PIEZO2:** Piezo-type mechanosensitive ion channel component 2 (PIEZO2) is a recently discovered cation channel that has been shown to be a sensor responsible for innocuous touch and proprioception by displaying rapidly-inactivating feature with a low mechanical threshold and by being expressed in a medium to large diameter non-nociceptive population of sensory neurons, whereas TRPA1 represents a slowly-inactivating component with a high mechanical threshold. Interestingly, one study has shown that a fraction of the nociceptive subpopulation expresses PIEZO2 that can be functionally modified in response to bradykinin signaling (Dubin *et al.*, 2012). Unlike TRPA1, PIEZO2 was only sensitized but not activated by bradykinin exposure. Its mechanosensitivity and response efficacy were both heightened with a moderate threshold shift. PKA and PKC are involved in the enhancement in mechanosensitivity, but surprisingly, nothing upstream of PKC activation related to Gq/11- or Gi/o-coupled PLC activation was associated.

## Ca<sup>2+</sup>-ACTIVATED Cl<sup>-</sup> CHANNELS

The two aforementioned TRPs are not the only excitatory effector ion channels involved in bradykinin-induced nociceptor firing. Many studies have shown that incapacitating TRPs by using antagonist pharmacology or gene deletion did not completely ablate the action potential discharges of nociceptors upon bradykinin exposure (Dray *et al.*, 1990; Dickenson and Dray, 1991; Zahner *et al.*, 2003; Kollarik and Undem, 2004; Rong *et al.*, 2004; Qin *et al.*, 2006; Katanosaka *et al.*, 2008; Brierley *et al.*, 2009). An explanation as to the unknown excitatory component, i.e., the role for Cl<sup>-</sup> channels has been emerging. Chemical inhibition of Ca<sup>2+</sup>-activated Cl<sup>-</sup> channels (CaCCs) was effective in reducing bradykinin-induced firing in an *ex vivo* tissue-nerve model from the guinea pig airway sensory system and in *in vitro* cultured nodose neurons (Oh and Weinreich, 2004; Lee *et al.*, 2005a). In sensory neurons, it is well known that the intracellular Cl<sup>-</sup> concentration is high and that opening of Cl<sup>-</sup> channels leads to the efflux of Cl<sup>-</sup>, resulting in depolarization, which was confirmed by the same studies. In pathologic conditions, like chronic exposure to major inflammatory pain mediators in the periphery including prostaglandins, nerve growth factor (NGF), and ATP, the intracellular Cl<sup>-</sup> concentration is known to even further be elevated owing to the facilitated activities of surface Na<sup>+</sup>-K<sup>+</sup>-2Cl<sup>-</sup> cotransporters (NKCCs), indicating that Cl<sup>-</sup> channel opening may contribute more to nociceptor firing than non-inflamed situations (Sung *et al.*, 2000; Labrakakis *et al.*, 2003; Funk *et al.*, 2008; Price *et al.*, 2009).

Delicate molecular approach was conducted by a group for determining the genetic identity of CaCC of rodent nociceptors contributing to bradykinin-induced excitation and the underlying activation mechanisms (Liu *et al.*, 2010; Jin *et al.*, 2013). The central player is anoctamin 1 (ANO1, also known as TMEM16A), a relatively newly discovered ion channel. The studies dissected how the intracellular Ca<sup>2+</sup> was affected by bradykinin stimulation and revealed two phases, and the first of which may more depend on intracellular mobilization by IP<sub>3</sub>. This indicates that bradykinin-evoked G protein signaling leads to PLC activation, resulting in productions of IP<sub>3</sub> and DAG as mentioned above (Thayer *et al.*, 1988; Burgess *et al.*, 1989; Kano *et al.*, 1994; Kozaki *et al.*, 2007), followed by the activation of IP<sub>3</sub> receptors on the endoplasmic reticulum (ER) membrane. This causes Ca<sup>2+</sup> release from the ER, completing the first Ca<sup>2+</sup> phase. This firstly elevated cytoplasmic Ca<sup>2+</sup> activates ANO1. ANO1 was shown to physically couple to IP<sub>3</sub> receptor type 1 and bradykinin B2 receptor to compose a signaling complex, enabling the efficient utilization of the locally increased levels of Ca<sup>2+</sup>. The second phase of Ca<sup>2+</sup> elevation may depend on the influx of extracellular Ca<sup>2+</sup>, resulting from activation of voltage-gated Ca<sup>2+</sup> channels by ANO1-elicited depolarization, and of TRP channels that are highly Ca<sup>2+</sup> permeable. Such ANO1-dependent bradykinin-mediated nociception was again confirmed in an *in vivo* study using tissue-specific ANO1-deficient mice (Advillin/Ano1<sup>fl/fl</sup>) that lost ANO1 expression mainly in DRG neurons (Lee *et al.*, 2014).

## K<sup>+</sup> CHANNEL INHIBITION

The decreased activity of resting K<sup>+</sup> channels may contribute to depolarization. Indeed, two studies that were mentioned

previously, exploring the outcomes of the first phase of  $\text{Ca}^{2+}$  elevation in response to bradykinin stimulation have proposed that together with CaCC activation,  $\text{K}^+$  channel inhibition is also involved in nociceptor firing during this first phase (Oh and Weinreich, 2004; Liu *et al.*, 2010). Two different  $\text{K}^+$ -permeating components were identified as contributors by the two studies respectively, as explained in the following section.

### KCNQ voltage-gated $\text{K}^+$ channels

The outward  $\text{K}^+$  current mediated by the opening of the KCNQ channel (also known as Kv7) refers to the M current as it was first found as a downstream effector of M2 muscarinic receptor signaling. A fraction of KCNQ channels open in the resting state and control the resting membrane potential and action potential rheobase (Delmas and Brown, 2005). The M current can be inhibited in the early phase of the intracellular  $\text{Ca}^{2+}$  wave caused by bradykinin exposure (Liu *et al.*, 2010). Further inhibition of the KCNQ-mediated current by a synthetic specific antagonist potentiated bradykinin-induced firing while its activation using the channel opener retigabine diminished it. Acutely pretreated retigabine also prevented nociceptive behaviors caused by intraplantar bradykinin injection in *in vivo* observations. Additionally, chelation of the early  $\text{Ca}^{2+}$  rise but not PKC or PLA2 inhibition reversed the closing of the  $\text{K}^+$  channel in *in vitro* nociceptor assays, indicating that the Gq/11-coupled-PLC-IP3- $\text{Ca}^{2+}$  cascade is required for the  $\text{K}^+$  channel contribution and that no other signaling downstream of PLC or other branches of G protein signaling seems to be involved. The genetic identity of the KCNQ subtypes responsible for the underlying molecular mechanisms involved in bradykinin-induced signaling remain to be elucidated. Very recently, KCNQ3 and KCNQ5 have been raised as major Kv7 subtypes that depolarize murine and human visceral nociceptors upon B2 receptor stimulation (Peiris *et al.*, 2017).

### $\text{Ca}^{2+}$ -activated $\text{K}^+$ channels

Another  $\text{K}^+$  component altered by bradykinin stimulation has been shown to be mediated by  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channels ( $I_{\text{KCa}}$ ). With regards to the action potential phase, these  $\text{K}^+$  currents typically compose a slow component of the afterhyperpolarization (AHP). AHP is responsible for spike frequency accommodation in repeated firing. A shortened AHP resulting from  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channel inhibition causes sustained or increased firing frequencies (Weinreich and Wonderlin, 1987; Cordoba-Rodriguez *et al.*, 1999). The contribution of the bradykinin-induced channel blockade to the alteration of nodose neuronal firing may reflect this paradigm (Oh and Weinreich, 2004).

## CONCLUSIONS

Bradykinin is among the major pain mediators during inflammation. Peripherally produced bradykinin alters the electrical functions of nociceptor sensory neurons that are the forefront initiators of the ascending signals of the sensory neural pathway for pain perception. Bradykinin generally enhances their excitability, greatly contributing to the generation and exacerbation of pain. At the cellular level, bradykinin not only acutely excites the neurons but also electrically sensitizes them. Through intracellular signaling, mostly composed of G-protein coupled ones, it has been hypothesized that

bradykinin may finally augment the depolarizing activities of some specific effector ion channels expressed in the nociceptor neurons. Currently, an array of ion channels have been shown to be affected in this paradigm. Here we overviewed six important ion channels in their role as a depolarizing effector. The opening of TRPV1, TRPA1, and ANO1 all seem to enact this role in the direct induction of neuronal firing by bradykinin. TRPV1 and TRPA1 also appear to be involved in sensitized neuronal function in a longer duration. PIEZO2 is an emerging ion channel that functions as a mechanical pain-specific sensitizing effector. KCNQ and  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channels may contribute to the initial excitation through their functional downregulation. Linker signals between bradykinin receptor activation and depolarizing effectors are currently being revealed in greater depth (summarized in Fig. 1).

The consistent expansion of information has broadened the knowledge of the molecular nature of bradykinin-induced inflammatory pain and has validated bradykinin signaling as an analgesic target. In particular, the B2 receptor inhibitor HOE 140 and capsaicin-mediated TRPV1 incapacitation seem to have promising outcomes from a multitude of clinical researches (Backonja *et al.*, 2008; Song *et al.*, 2008). Nonetheless, efforts to establish the excitatory mechanism mediated by relatively recent found effectors such as ANO1 and  $\text{K}^+$  channels are still required. Further, unknown component may be present for the nociceptive neuronal actions of bradykinin. For example, pharmacological antagonism of purinergic P2X3 ion channel has once been shown to be effective specifically at bradykinin induced mechanical hyperalgesia, which needs to be confirmed by further molecular approaches (de Oliveira Fusaro *et al.*, 2010). Among nociceptor-specific voltage-gated  $\text{Na}^+$  channels, Nav1.9 may particularly be affected under bradykinin-including pathologic condition but the mechanism remains elusive (Vaughn and Gold, 2010). Further accumulation of the knowledge will contribute to more precise understanding of the depolarization mechanisms and to development of more sophisticated painkilling strategies.

## ACKNOWLEDGMENTS

This work was supported by grants from the National Research Foundation of Korea (2017R1A2B2001817, 2017M3C7A1025600). SIC collected and analyzed the information and wrote the preliminary draft. SWH supervised the studies and wrote the manuscript. All authors read and approved the final manuscript. The authors declare that there is no conflict of interest regarding the publication of this article.

## REFERENCES

- Akopian, A. N., Ruparel, N. B., Jeske, N. A. and Hargreaves, K. M. (2007) Transient receptor potential TRPA1 channel desensitization in sensory neurons is agonist dependent and regulated by TRPV1-directed internalization. *J. Physiol.* **583**, 175-193.
- Allen, A. C., Gammon, C. M., Ousley, A. H., McCarthy, K. D. and Morell, P. (1992) Bradykinin stimulates arachidonic acid release through the sequential actions of an sn-1 diacylglycerol lipase and a monoacylglycerol lipase. *J. Neurochem.* **58**, 1130-1139.
- Amaya, F., Wang, H., Costigan, M., Allchorne, A. J., Hatcher, J. P., Egerton, J., Stean, T., Morisset, V., Grose, D., Gunthorpe, M. J., Chessell, I. P., Tate, S., Green, P. J. and Woolf, C. J. (2006) The voltage-gated sodium channel Na(v)1.9 is an effector of peripheral

- inflammatory pain hypersensitivity. *J. Neurosci.* **26**, 12852-12860.
- Andersson, D. A., Gentry, C., Moss, S. and Bevan, S. (2008) Transient receptor potential A1 is a sensory receptor for multiple products of oxidative stress. *J. Neurosci.* **28**, 2485-2494.
- Armstrong, D., Jepson, J., Keele, C. and Stewart, J. (1957) Pain-producing substance in human inflammatory exudates and plasma. *J. Physiol.* **135**, 350-370.
- Babenko, V., Graven-Nielsen, T., Svensson, P., Drewes, A. M., Jensen, T. S. and Arendt-Nielsen, L. (1999) Experimental human muscle pain induced by intramuscular injections of bradykinin, serotonin, and substance P. *Eur. J. Pain* **3**, 93-102.
- Backonja, M., Wallace, M. S., Blonsky, E. R., Cutler, B. J., Malan, P., Rauck, R. and Tobias, J. (2008) NGX-4010, a high-concentration capsaicin patch, for the treatment of postherpetic neuralgia: a randomised, double-blind study. *Lancet Neurol.* **7**, 1106-1112.
- Bae, S. W., Kim, H. S., Cha, Y. N., Park, Y. S., Jo, S. A. and Jo, I. (2003) Rapid increase in endothelial nitric oxide production by bradykinin is mediated by protein kinase A signaling pathway. *Biochem. Biophys. Res. Commun.* **306**, 981-987.
- Bandell, M., Story, G. M., Hwang, S. W., Viswanath, V., Eid, S. R., Petrus, M. J., Earley, T. J. and Patapoutian, A. (2004) Noxious cold ion channel TRPA1 is activated by pungent compounds and bradykinin. *Neuron* **41**, 849-857.
- Bang, S. and Hwang, S. W. (2009) Polymodal ligand sensitivity of TRPA1 and its modes of interactions. *J. Gen. Physiol.* **133**, 257-262.
- Barber, L. A. and Vasko, M. R. (1996) Activation of protein kinase C augments peptide release from rat sensory neurons. *J. Neurochem.* **67**, 72-80.
- Bautista, D. M., Jordt, S.-E., Nikai, T., Tsuruda, P. R., Read, A. J., Poblete, J., Yamoah, E. N., Basbaum, A. I. and Julius, D. (2006) TRPA1 mediates the inflammatory actions of environmental irritants and proalgesic agents. *Cell* **124**, 1269-1282.
- Beck, P. W. and Handwerker, H. O. (1974) Bradykinin and serotonin effects on various types of cutaneous nerve fibres. *Pflugers Arch.* **347**, 209-222.
- Bhave, G., Hu, H.-J., Glauner, K. S., Zhu, W., Wang, H., Brasier, D., Oxford, G. S. and Gereau, R. W. (2003) Protein kinase C phosphorylation sensitizes but does not activate the capsaicin receptor transient receptor potential vanilloid 1 (TRPV1). *Proc. Natl. Acad. Sci. U.S.A.* **100**, 12480-12485.
- Brierley, S., Jones, R., Xu, L., Gebhart, G. and Blackshaw, L. (2005) Activation of splanchnic and pelvic colonic afferents by bradykinin in mice. *Neurogastroenterol. Motil.* **17**, 854-862.
- Brierley, S. M., Hughes, P. A., Page, A. J., Kwan, K. Y., Martin, C. M., O'Donnell, T. A., Cooper, N. J., Harrington, A. M., Adam, B. and Liebrechts, T. (2009) The ion channel TRPA1 is required for normal mechanosensation and is modulated by algesic stimuli. *Gastroenterology* **137**, 2084-2095.e3.
- Burch, R. M. and Axelrod, J. (1987) Dissociation of bradykinin-induced prostaglandin formation from phosphatidylinositol turnover in Swiss 3T3 fibroblasts: evidence for G protein regulation of phospholipase A2. *Proc. Natl. Acad. Sci. U.S.A.* **84**, 6374-6378.
- Burgess, G. M., Mullaney, I., McNeill, M., Dunn, P. M. and Rang, H. P. (1989) Second messengers involved in the mechanism of action of bradykinin in sensory neurons in culture. *J. Neurosci.* **9**, 3314-3325.
- Camprubi-Robles, M., Planells-Cases, R. and Ferrer-Montiel, A. (2009) Differential contribution of SNARE-dependent exocytosis to inflammatory potentiation of TRPV1 in nociceptors. *FASEB J.* **23**, 3722-3733.
- Carr, M. J., Kollarik, M., Meeker, S. N. and Udem, B. J. (2003) A role for TRPV1 in bradykinin-induced excitation of vagal airway afferent nerve terminals. *J. Pharmacol. Exp. Ther.* **304**, 1275-1279.
- Cavanaugh, D. J., Lee, H., Lo, L., Shields, S. D., Zylka, M. J., Basbaum, A. I. and Anderson, D. J. (2009) Distinct subsets of unmyelinated primary sensory fibers mediate behavioral responses to noxious thermal and mechanical stimuli. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 9075-9080.
- Cesare, P., Dekker, L. V., Sardini, A., Parker, P. J. and McNaughton, P. A. (1999) Specific involvement of PKC- $\epsilon$  in sensitization of the neuronal response to painful heat. *Neuron* **23**, 617-624.
- Cesare, P. and McNaughton, P. (1996) A novel heat-activated current in nociceptive neurons and its sensitization by bradykinin. *Proc. Natl. Acad. Sci. U.S.A.* **93**, 15435-15439.
- Chahl, L. A. and Iggo, A. (1977) The effects of bradykinin and prostaglandin E1 on rat cutaneous afferent nerve activity. *Br. J. Pharmacol.* **59**, 343-347.
- Choi, S. I., Yoo, S., Lim, J. Y. and Hwang, S. W. (2014) Are sensory TRP channels biological alarms for lipid peroxidation? *Int. J. Mol. Sci.* **15**, 16430-16457.
- Chuang, H.-h., Prescott, E. D., Kong, H., Shields, S., Jordt, S.-E., Basbaum, A. I., Chao, M. V. and Julius, D. (2001) Bradykinin and nerve growth factor release the capsaicin receptor from PtdIns (4, 5) P2-mediated inhibition. *Nature* **411**, 957-962.
- Cordoba-Rodriguez, R., Moore, K. A., Kao, J. P. and Weinreich, D. (1999) Calcium regulation of a slow post-spike hyperpolarization in vagal afferent neurons. *Proc. Natl. Acad. Sci. U.S.A.* **96**, 7650-7657.
- Costa, R., Bicca, M. A., Manjavachi, M. N., Segat, G. C., Dias, F. C., Fernandes, E. S. and Calixto, J. B. (2018) Kinin receptors sensitize TRPV4 channel and induce mechanical hyperalgesia: relevance to paclitaxel-induced peripheral neuropathy in mice. *Mol. Neurobiol.* **55**, 2150-2161.
- Costello, A. and Hargreaves, K. (1989) Suppression of carrageenan-induced hyperalgesia, hyperthermia and edema by a bradykinin antagonist. *Eur. J. Pharmacol.* **171**, 259-263.
- Crandall, M., Kwash, J., Yu, W. and White, G. (2002) Activation of protein kinase C sensitizes human VR1 to capsaicin and to moderate decreases in pH at physiological temperatures in *Xenopus* oocytes. *Pain* **98**, 109-117.
- Cunha, T. M., Verri, W. A., Fukada, S. Y., Guerrero, A. T., Santodomingo-Garzon, T., Poole, S., Parada, C. A., Ferreira, S. H. and Cunha, F. Q. (2007) TNF- $\alpha$  and IL-1 $\beta$  mediate inflammatory hypernociception in mice triggered by B 1 but not B 2 kinin receptor. *Eur. J. Pharmacol.* **573**, 221-229.
- Dai, Y., Wang, S., Tominaga, M., Yamamoto, S., Fukuoka, T., Higashi, T., Kobayashi, K., Obata, K., Yamanaka, H. and Noguchi, K. (2007) Sensitization of TRPA1 by PAR2 contributes to the sensation of inflammatory pain. *J. Clin. Invest.* **117**, 1979-1987.
- De Campos, R., Henriques, M. and Calixto, J. (1998) Systemic treatment with *Mycobacterium bovis* bacillus calmette-guerin (BCG) potentiates kinin B 1 receptor agonist-induced nociception and oedema formation in the formalin test in mice. *Neuropeptides* **32**, 393-403.
- de Oliveira Fusaro, M. C. G., Pelegrini-da-Silva, A., Araldi, D., Parada, C. A. and Tambeli, C. H. (2010) P2X3 and P2X2/3 receptors mediate mechanical hyperalgesia induced by bradykinin, but not by pro-inflammatory cytokines, PGE 2 or dopamine. *Eur. J. Pharmacol.* **649**, 177-182.
- Delmas, P. and Brown, D. A. (2005) Pathways modulating neural KCNQ/M (Kv7) potassium channels. *Nat. Rev. Neurosci.* **6**, 850-862.
- Dickenson, A. and Dray, A. (1991) Selective antagonism of capsaicin by capsazepine: evidence for a spinal receptor site in capsaicin-induced antinociception. *Br. J. Pharmacol.* **104**, 1045-1049.
- Dray, A., Bettaney, J., Forster, P. and Perkins, M. (1988) Bradykinin-induced stimulation of afferent fibres is mediated through protein kinase C. *Neurosci. Lett.* **91**, 301-307.
- Dray, A., Forbes, C. and Burgess, G. (1990) Ruthenium red blocks the capsaicin-induced increase in intracellular calcium and activation of membrane currents in sensory neurones as well as the activation of peripheral nociceptors *in vitro*. *Neurosci. Lett.* **110**, 52-59.
- Dray, A., Patel, I., Perkins, M. and Rueff, A. (1992) Bradykinin-induced activation of nociceptors: receptor and mechanistic studies on the neonatal rat spinal cord-tail preparation *in vitro*. *Br. J. Pharmacol.* **107**, 1129-1134.
- Dubin, A. E., Schmidt, M., Mathur, J., Petrus, M. J., Xiao, B., Coste, B. and Patapoutian, A. (2012) Inflammatory signals enhance piezo2-mediated mechanosensitive currents. *Cell Rep.* **2**, 511-517.
- Ferreira, J., Beirith, A., Mori, M. A., Araújo, R. C., Bader, M., Pesquero, J. B. and Calixto, J. B. (2005) Reduced nerve injury-induced neuropathic pain in kinin B1 receptor knock-out mice. *J. Neurosci.* **25**, 2405-2412.

- Ferreira, J., Campos, M. M., Pesquero, J. B., Araújo, R. C., Bader, M. and Calixto, J. B. (2001) Evidence for the participation of kinins in Freund's adjuvant-induced inflammatory and nociceptive responses in kinin B 1 and B 2 receptor knockout mice. *Neuropharmacology* **41**, 1006-1012.
- Ferreira, J., Da Silva, G. L. and Calixto, J. B. (2004) Contribution of vanilloid receptors to the overt nociception induced by B2 kinin receptor activation in mice. *Br. J. Pharmacol.* **141**, 787-794.
- Ferreira, S., Lorenzetti, B., Cunha, F. and Poole, S. (1993a) Bradykinin release of TNF- $\alpha$  plays a key role in the development of inflammatory hyperalgesia. *Agents Actions* **38**, C7-C9.
- Ferreira, S., Lorenzetti, B. and Poole, S. (1993b) Bradykinin initiates cytokine-mediated inflammatory hyperalgesia. *Br. J. Pharmacol.* **110**, 1227-1231.
- Fischer, M. J., Balasuriya, D., Jeggle, P., Goetze, T. A., McNaughton, P. A., Reeh, P. W. and Edwardson, J. M. (2014) Direct evidence for functional TRPV1/TRPA1 heteromers. *Pflugers Arch.* **466**, 2229-2241.
- Fischer, M. J. and McNaughton, P. A. (2014) How anchoring proteins shape pain. *Pharmacol. Ther.* **143**, 316-322.
- Fox, A., Barnes, P., Urban, L. and Dray, A. (1993) An *in vitro* study of the properties of single vagal afferents innervating guinea-pig airways. *J. Physiol.* **469**, 21-35.
- Fox, A. J., Laloo, U. G., Belvisi, M. G., Bernareggi, M., Chung, K. F. and Barnes, P. J. (1996) Bradykinin-evoked sensitization of airway sensory nerves: A mechanism for ACE-inhibitor cough. *Nat. Med.* **2**, 814-817.
- Franco-Cereceda, A. (1989) Prostaglandins and CGRP release from cardiac sensory nerves. *Naunyn Schmiedebergs Arch. Pharmacol.* **340**, 180-184.
- Füredi, R., Bölcskei, K., Szolcsányi, J. and Pethő, G. (2010) Comparison of the peripheral mediator background of heat injury- and plantar incision-induced drop of the noxious heat threshold in the rat. *Life Sci.* **86**, 244-250.
- Funk, K., Woltecki, A., Franjic-Würtz, C., Gensch, T., Möhrlen, F. and Frings, S. (2008) Modulation of chloride homeostasis by inflammatory mediators in dorsal root ganglion neurons. *Mol. Pain* **4**, 32.
- Gabra, B. H., Benrezzak, O., Pheng, L.-H., Duta, D., Daull, P., Sirois, P., Nantel, F. and Battistini, B. (2005a) Inhibition of type 1 diabetic hyperalgesia in streptozotocin-induced Wistar versus spontaneous gene-prone BB/Worcester rats: efficacy of a selective bradykinin B1 receptor antagonist. *J. Neuropathol. Exp. Neurol.* **64**, 782-789.
- Gabra, B. H., Merino, V. F., Bader, M., Pesquero, J. B. and Sirois, P. (2005b) Absence of diabetic hyperalgesia in bradykinin B1 receptor-knockout mice. *Regul. Pept.* **127**, 245-248.
- Gabra, B. H. and Sirois, P. (2002) Role of bradykinin B 1 receptors in diabetes-induced hyperalgesia in streptozotocin-treated mice. *Eur. J. Pharmacol.* **457**, 115-124.
- Gabra, B. H. and Sirois, P. (2003a) Beneficial effect of chronic treatment with the selective bradykinin B 1 receptor antagonists, R-715 and R-954, in attenuating streptozotocin-diabetic thermal hyperalgesia in mice. *Peptides* **24**, 1131-1139.
- Gabra, B. H. and Sirois, P. (2003b) Kinin B 1 receptor antagonists inhibit diabetes-induced hyperalgesia in mice. *Neuropeptides* **37**, 36-44.
- Gammon, C. M., Allen, A. C. and Morell, P. (1989) Bradykinin stimulates phosphoinositide hydrolysis and mobilization of arachidonic acid in dorsal root ganglion neurons. *J. Neurochem.* **53**, 95-101.
- Gao, Y., Cao, E., Julius, D. and Cheng, Y. (2016) TRPV1 structures in nanodiscs reveal mechanisms of ligand and lipid action. *Nature* **534**, 347-351.
- Gibson, H. E., Edwards, J. G., Page, R. S., Van Hook, M. J. and Kauer, J. A. (2008) TRPV1 channels mediate long-term depression at synapses on hippocampal interneurons. *Neuron* **57**, 746-759.
- Gougat, J., Ferrari, B., Sarran, L., Planchenault, C., Poncelet, M., Maruani, J., Alonso, R., Cudennec, A., Croci, T. and Guagnini, F. (2004) SSR240612 [(2R)-2-(((3R)-3-(1, 3-benzodioxol-5-yl)-3-((6-methoxy-2-naphthyl) sulfonyl) amino) propanoyl) amino]-3-(4-[[2R, 6S)-2, 6-dimethylpiperidinyl] methyl] phenyl)-N-isopropyl-N-methylpropanamide hydrochloride], a new nonpeptide antagonist of the bradykinin B1 receptor: biochemical and pharmacological characterization. *J. Pharmacol. Exp. Ther.* **309**, 661-669.
- Gregus, A. M., Doolen, S., Dumlao, D. S., Buczynski, M. W., Takasusuki, T., Fitzsimmons, B. L., Hua, X.-Y., Taylor, B. K., Dennis, E. A. and Yaksh, T. L. (2012) Spinal 12-lipoxygenase-derived heptaxilin A3 contributes to inflammatory hyperalgesia via activation of TRPV1 and TRPA1 receptors. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 6721-6726.
- Griesbacher, T., Amann, R., Sametz, W., Diethart, S. and Juan, H. (1998) The nonpeptide B2 receptor antagonist FR173657: inhibition of effects of bradykinin related to its role in nociception. *Br. J. Pharmacol.* **124**, 1328-1334.
- Guo, Z.-L., Fu, L.-W., Symons, J. D. and Longhurst, J. C. (1998) Signal transduction in activation of ischemically sensitive abdominal visceral afferents: role of PKC. *Am. J. Physiol.* **275**, H1024-H1031.
- Guo, Z.-L., Symons, J. D. and Longhurst, J. C. (1999) Activation of visceral afferents by bradykinin and ischemia: independent roles of PKC and prostaglandins. *Am. J. Physiol.* **276**, H1884-H1891.
- Haake, B., Liang, Y. and Reeh, P. (1996) Bradykinin effects and receptor subtypes in rat cutaneous nociceptors, *in vitro*. *Pflugers Arch.* **431**, R15.
- Hinman, A., Chuang, H.-h., Bautista, D. M. and Julius, D. (2006) TRP channel activation by reversible covalent modification. *Proc. Natl. Acad. Sci. U.S.A.* **103**, 19564-19568.
- Hong, Y. and Abbott, F. (1994) Behavioural effects of intraplantar injection of inflammatory mediators in the rat. *Neuroscience* **63**, 827-836.
- Hwang, S. W., Cho, H., Kwak, J., Lee, S.-Y., Kang, C.-J., Jung, J., Cho, S., Min, K. H., Suh, Y.-G. and Kim, D. (2000) Direct activation of capsaicin receptors by products of lipoxygenases: endogenous capsaicin-like substances. *Proc. Natl. Acad. Sci. U.S.A.* **97**, 6155-6160.
- Ikeda, Y., Ueno, A., Naraba, H. and Oh-ishi, S. (2001a) Evidence for bradykinin mediation of carrageenin-induced inflammatory pain: a study using kininogen-deficient Brown Norway Katholiek rats. *Biochem. Pharmacol.* **61**, 911-914.
- Ikeda, Y., Ueno, A., Naraba, H. and Oh-ishi, S. (2001b) Involvement of vanilloid receptor VR1 and prostanoids in the acid-induced writhing responses of mice. *Life Sci.* **69**, 2911-2919.
- Inoue, A., Iwasa, M., Nishikura, Y., Ogawa, S., Nakasuka, A. and Nakata, Y. (2006) The long-term exposure of rat cultured dorsal root ganglion cells to bradykinin induced the release of prostaglandin E2 by the activation of cyclooxygenase-2. *Neurosci. Lett.* **401**, 242-247.
- Jackson, J. G., Usachev, Y. M. and Thayer, S. A. (2007) Bradykinin-induced nuclear factor of activated T-cells-dependent transcription in rat dorsal root ganglion neurons. *Mol. Pharmacol.* **72**, 303-310.
- Jenkins, D. W., Sellers, L. A., Feniuk, W. and Humphrey, P. P. (2003) Characterization of bradykinin-induced prostaglandin E2 release from cultured rat trigeminal ganglion neurones. *Eur. J. Pharmacol.* **469**, 29-36.
- Jin, X., Shah, S., Liu, Y., Zhang, H., Lees, M., Fu, Z., Lippiat, J. D., Beech, D. J., Sivaprasadarao, A. and Baldwin, S. A. (2013) Activation of the Cl<sup>-</sup> channel ANO1 by localized calcium signals in nociceptive sensory neurons requires coupling with the IP3 receptor. *Sci. Signal.* **6**, ra73.
- Juan, H. (1977) Mechanism of action of bradykinin-induced release of prostaglandin E. *Naunyn Schmiedebergs Arch. Pharmacol.* **300**, 77-85.
- Juan, H. and Lembeck, F. (1974) Action of peptides and other algescic agents on paravascular pain receptors of the isolated perfused rabbit ear. *Naunyn Schmiedebergs Arch. Pharmacol.* **283**, 151-164.
- Kajekar, R., Proud, D., Myers, A. C., Meeker, S. N. and Udem, B. J. (1999) Characterization of vagal afferent subtypes stimulated by bradykinin in guinea pig trachea. *J. Pharmacol. Exp. Ther.* **289**, 682-687.
- Kano, M., Kawakami, T., Hikawa, N., Hori, H., Takenaka, T. and Gotoh, H. (1994) Bradykinin-responsive cells of dorsal root ganglia in culture: cell size, firing, cytosolic calcium, and substance P. *Cell. Mol. Neurobiol.* **14**, 49-57.
- Karashima, Y., Prenen, J., Meseguer, V., Owsianik, G., Voets, T. and Nilius, B. (2008) Modulation of the transient receptor potential channel TRPA1 by phosphatidylinositol 4,5-bisphosphate manipulators. *Pflugers Arch.* **457**, 77.

- Katanosaka, K., Banik, R. K., Giron, R., Higashi, T., Tominaga, M. and Mizumura, K. (2008) Contribution of TRPV1 to the bradykinin-evoked nociceptive behavior and excitation of cutaneous sensory neurons. *Neurosci. Res.* **62**, 168-175.
- Khan, A. A., Raja, S. N., Manning, D. C., Campbell, J. N. and Meyer, R. A. (1992) The effects of bradykinin and sequence-related analogs on the response properties of cutaneous nociceptors in monkeys. *Somatosens. Mot. Res.* **9**, 97-106.
- Khasar, S. G., Green, P. G. and Levine, J. D. (1993) Comparison of intradermal and subcutaneous hyperalgesic effects of inflammatory mediators in the rat. *Neurosci. Lett.* **153**, 215-218.
- Kim, D. and Cavanaugh, E. J. (2007) Requirement of a soluble intracellular factor for activation of transient receptor potential A1 by pungent chemicals: role of inorganic polyphosphates. *J. Neurosci.* **27**, 6500-6509.
- Kim, D., Cavanaugh, E. J. and Simkin, D. (2008) Inhibition of transient receptor potential A1 channel by phosphatidylinositol-4, 5-bisphosphate. *Am. J. Physiol. Cell Physiol.* **295**, C92-C99.
- Kindgen-Milles, D., Klement, W. and Arndt, J. (1994) The nociceptive systems of skin, paravascular tissue and hand veins of humans and their sensitivity to bradykinin. *Neurosci. Lett.* **181**, 39-42.
- Kollarik, M. and Udem, B. (2004) Activation of bronchopulmonary vagal afferent nerves with bradykinin, acid and vanilloid receptor agonists in wild-type and TRPV1-/- mice. *J. Physiol.* **555**, 115-123.
- Koltzenburg, M., Kress, M. and Reeh, P. (1992) The nociceptor sensitization by bradykinin does not depend on sympathetic neurons. *Neuroscience* **46**, 465-473.
- Kozaki, Y., Kambe, F., Hayashi, Y., Ohmori, S., Seo, H., Kumazawa, T. and Mizumura, K. (2007) Molecular cloning of prostaglandin EP3 receptors from canine sensory ganglia and their facilitatory action on bradykinin-induced mobilization of intracellular calcium. *J. Neurochem.* **100**, 1636-1647.
- Kumazawa, T. and Mizumura, K. (1976) The polymodal C-fiber receptor in the muscle of the dog. *Brain Res.* **101**, 589-593.
- Kumazawa, T., Mizumura, K., Minagawa, M. and Tsujii, Y. (1991) Sensitizing effects of bradykinin on the heat responses of the visceral nociceptor. *J. Neurophysiol.* **66**, 1819-1824.
- Kwan, K. Y., Allchorne, A. J., Vollrath, M. A., Christensen, A. P., Zhang, D.-S., Woolf, C. J. and Corey, D. P. (2006) TRPA1 contributes to cold, mechanical, and chemical nociception but is not essential for hair-cell transduction. *Neuron* **50**, 277-289.
- Kwan, K. Y., Glazer, J. M., Corey, D. P., Rice, F. L. and Stucky, C. L. (2009) TRPA1 modulates mechanotransduction in cutaneous sensory neurons. *J. Neurosci.* **29**, 4808-4819.
- Labrakakis, C., Tong, C. K., Weissman, T., Torsney, C. and MacDermott, A. B. (2003) Localization and function of ATP and GABAA receptors expressed by nociceptors and other postnatal sensory neurons in rat. *J. Physiol.* **549**, 131-142.
- Lang, E., Novak, A., Reeh, P. and Handwerker, H. (1990) Chemosensitivity of fine afferents from rat skin *in vitro*. *J. Neurophysiol.* **63**, 887-901.
- Lee, M. G., MacGlashan, D. W. and Udem, B. J. (2005a) Role of chloride channels in bradykinin-induced guinea pig airway vagal C-fibre activation. *J. Physiol.* **566**, 205-212.
- Lee, S.-Y., Lee, J.-H., Kang, K. K., Hwang, S.-Y., Choi, K. D. and Oh, U. (2005b) Sensitization of vanilloid receptor involves an increase in the phosphorylated form of the channel. *Arch. Pharm. Res.* **28**, 405-412.
- Lee, B., Cho, H., Jung, J., Yang, Y. D., Yang, D. J. and Oh, U. (2014). Anoctamin 1 contributes to inflammatory and nerve-injury induced hypersensitivity. *Mol. Pain* **10**, 5.
- Lembeck, F. and Juan, H. (1974) Interaction of prostaglandins and indomethacin with algesic substances. *Naunyn Schmiedeberg's Arch. Pharmacol.* **285**, 301-313.
- Lembeck, F., Popper, H. and Juan, H. (1976) Release of prostaglandins by bradykinin as an intrinsic mechanism of its algesic effect. *Naunyn Schmiedeberg's Arch. Pharmacol.* **294**, 69-73.
- Leonard, P. A., Arunkumar, R. and Brennan, T. J. (2004) Bradykinin antagonists have no analgesic effect on incisional pain. *Anesth. Analg.* **99**, 1166-1172.
- Levy, D. and Zochodne, D. W. (2000) Increased mRNA expression of the B1 and B2 bradykinin receptors and antinociceptive effects of their antagonists in an animal model of neuropathic pain. *Pain* **86**, 265-271.
- Liang, Y. F., Haake, B. and Reeh, P. W. (2001) Sustained sensitization and recruitment of rat cutaneous nociceptors by bradykinin and a novel theory of its excitatory action. *J. Physiol.* **532**, 229-239.
- Liebmann, C., Graness, A., Ludwig, B., Adomeit, A., Boehmer, A., Boehmer, F.-D., Nürnberg, B. and Wetzker, R. (1996) Dual bradykinin B2 receptor signalling in A431 human epidermoid carcinoma cells: activation of protein kinase C is counteracted by a GS-mediated stimulation of the cyclic AMP pathway. *Biochem. J.* **313**, 109-118.
- Liu, B., Linley, J. E., Du, X., Zhang, X., Ooi, L., Zhang, H. and Gamper, N. (2010) The acute nociceptive signals induced by bradykinin in rat sensory neurons are mediated by inhibition of M-type K<sup>+</sup> channels and activation of Ca<sup>2+</sup>-activated Cl<sup>-</sup> channels. *J. Clin. Invest.* **120**, 1240-1252.
- Luiz, A. P., Schroeder, S. D., Chichorro, J. G., Calixto, J. B., Zampronio, A. R. and Rae, G. A. (2010) Kinin B1 and B2 receptors contribute to orofacial heat hyperalgesia induced by infraorbital nerve constriction injury in mice and rats. *Neuropeptides* **44**, 87-92.
- Lukacs, V., Thyagarajan, B., Varnai, P., Balla, A., Balla, T. and Rohacs, T. (2007) Dual regulation of TRPV1 by phosphoinositides. *J. Neurosci.* **27**, 7070-7080.
- Macpherson, L. J., Dubin, A. E., Evans, M. J., Marr, F., Schultz, P. G., Cravatt, B. F. and Patapoutian, A. (2007) Noxious compounds activate TRPA1 ion channels through covalent modification of cysteines. *Nature* **445**, 541-545.
- Manning, D. C., Raja, S. N., Meyer, R. A. and Campbell, J. N. (1991) Pain and hyperalgesia after intradermal injection of bradykinin in humans. *Clin. Pharmacol. Ther.* **50**, 721-729.
- Materazzi, S., Nassini, R., André, E., Campi, B., Amadesi, S., Trevisani, M., Bunnett, N. W., Patacchini, R. and Geppetti, P. (2008) Cox-dependent fatty acid metabolites cause pain through activation of the irritant receptor TRPA1. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 12045-12050.
- Mathivanan, S., Devesa, I., Changeux, J. P. and Ferrer-Montiel, A. (2016) Bradykinin induces TRPV1 exocytotic recruitment in peptidergic nociceptors. *Front. Pharmacol.* **7**, 178.
- Maubach, K. A. and Grundy, D. (1999) The role of prostaglandins in the bradykinin-induced activation of serosal afferents of the rat jejunum *in vitro*. *J. Physiol.* **515**, 277-285.
- Mayer, S., Izydorczyk, I., Reeh, P. W. and Grubb, B. D. (2007) Bradykinin-induced nociceptor sensitisation to heat depends on cox-1 and cox-2 in isolated rat skin. *Pain* **130**, 14-24.
- McGehee, D. S. and Oxford, G. S. (1991) Bradykinin modulates the electrophysiology of cultured rat sensory neurons through a pertussis toxin-insensitive G protein. *Mol. Cell. Neurosci.* **2**, 21-30.
- McGuirk, S. and Dolphin, A. (1992) G-protein mediation in nociceptive signal transduction: an investigation into the excitatory action of bradykinin in a subpopulation of cultured rat sensory neurons. *Neuroscience* **49**, 117-128.
- McGuirk, S., Vallis, Y., Pasternak, C. and Dolphin, A. (1989) Bradykinin enhances excitability in cultured rat sensory neurons by a GTP-dependent mechanisms. *Neurosci. Lett.* **99**, 85-89.
- Meotti, F. C., Figueiredo, C. P., Manjavachi, M. and Calixto, J. B. (2017) The transient receptor potential ankyrin-1 mediates mechanical hyperalgesia induced by the activation of B1 receptor in mice. *Biochem. Pharmacol.* **125**, 75-83.
- Meyer, R. A., Davis, K. D., Raja, S. N. and Campbell, J. N. (1992) Sympathectomy does not abolish bradykinin-induced cutaneous hyperalgesia in man. *Pain* **51**, 323-327.
- Mizumura, K., Koda, H. and Kumazawa, T. (1997) Evidence that protein kinase C activation is involved in the excitatory and facilitatory effects of bradykinin on canine visceral nociceptors *in vitro*. *Neurosci. Lett.* **237**, 29-32.
- Mizumura, K., Sato, J. and Kumazawa, T. (1987) Effects of prostaglandins and other putative chemical intermediaries on the activity of canine testicular polymodal receptors studied *in vitro*. *Pflügers Arch.* **408**, 565-572.
- Mizumura, K., Sugiura, T., Katanosaka, K., Banik, R. K. and Kozaki, Y. (2009) Excitation and sensitization of nociceptors by bradykinin: what do we know? *Exp. Brain Res.* **196**, 53-65.

- Moriyama, T., Higashi, T., Togashi, K., Iida, T., Segi, E., Sugimoto, Y., Tominaga, T., Narumiya, S. and Tominaga, M. (2005) Sensitization of TRPV1 by EP 1 and IP reveals peripheral nociceptive mechanism of prostaglandins. *Mol. Pain* **1**, 3.
- Nakamura, A., Fujita, M. and Shiomi, H. (1996) Involvement of endogenous nitric oxide in the mechanism of bradykinin-induced peripheral hyperalgesia. *Br. J. Pharmacol.* **117**, 407-412.
- Numazaki, M., Tominaga, T., Toyooka, H. and Tominaga, M. (2002) Direct phosphorylation of capsaicin receptor VR1 by protein kinase C $\epsilon$  and identification of two target serine residues. *J. Biol. Chem.* **277**, 13375-13378.
- Oh, E. J. and Weinreich, D. (2004) Bradykinin decreases K<sup>+</sup> and increases Cl<sup>-</sup> conductances in vagal afferent neurones of the guinea pig. *J. Physiol.* **558**, 513-526.
- Oshita, K., Inoue, A., Tang, H.-B., Nakata, Y., Kawamoto, M. and Yuge, O. (2005) CB1 cannabinoid receptor stimulation modulates transient receptor potential vanilloid receptor 1 activities in calcium influx and substance P release in cultured rat dorsal root ganglion cells. *J. Pharmacol. Sci.* **97**, 377-385.
- Pan, H.-L. and Chen, S.-R. (2004) Sensing tissue ischemia. *Circulation* **110**, 1826-1831.
- Patapoutian, A., Tate, S. and Woolf, C. J. (2009) Transient receptor potential channels: targeting pain at the source. *Nat. Rev. Drug Discov.* **8**, 55-68.
- Peiris, M., Hockley, J. R., Reed, D. E., Smith, E. S. J., Bulmer, D. C. and Blackshaw, L. A. (2017) Peripheral KV7 channels regulate visceral sensory function in mouse and human colon. *Mol. Pain* **13**, 1744806917709371.
- Perkins, M. and Kelly, D. (1993) Induction of bradykinin B1 receptors *in vivo* in a model of ultra-violet irradiation-induced thermal hyperalgesia in the rat. *Br. J. Pharmacol.* **110**, 1441-1444.
- Perkins, M. N., Campbell, E. and Dray, A. (1993) Antinociceptive activity of the bradykinin B1 and B2 receptor antagonists, des-Arg 9, [Leu 8]-BK and HOE 140, in two models of persistent hyperalgesia in the rat. *Pain* **53**, 191-197.
- Petcu, M., Dias, J., Ongali, B., Thibault, G., Neugebauer, W. and Coutrure, R. (2008) Role of kinin B1 and B2 receptors in a rat model of neuropathic pain. *Int. Immunopharmacol.* **8**, 188-196.
- Pethö, G., Derow, A. and Reeh, P. W. (2001) Bradykinin-induced nociceptor sensitization to heat is mediated by cyclooxygenase products in isolated rat skin. *Eur. J. Neurosci.* **14**, 210-218.
- Pethö, G. and Reeh, P. W. (2012) Sensory and signaling mechanisms of bradykinin, eicosanoids, platelet-activating factor, and nitric oxide in peripheral nociceptors. *Physiol. Rev.* **92**, 1699-1775.
- Petrus, M., Peier, A. M., Bandell, M., Hwang, S. W., Huynh, T., Olney, N., Jegla, T. and Patapoutian, A. (2007) A role of TRPA1 in mechanical hyperalgesia is revealed by pharmacological inhibition. *Mol. Pain* **3**, 40.
- Poole, S., Lorenzetti, B., Cunha, J., Cunha, F. and Ferreira, S. (1999) Bradykinin B1 and B2 receptors, tumour necrosis factor  $\alpha$  and inflammatory hyperalgesia. *Br. J. Pharmacol.* **126**, 649-656.
- Porreca, F., Vanderah, T. W., Guo, W., Barth, M., Dodey, P., Peyrou, V., Luccarini, J., Junien, J.-L. and Pruneau, D. (2006) Antinociceptive pharmacology of N-[[4-(4, 5-dihydro-1H-imidazol-2-yl) phenyl] methyl]-2-[2-[[4-(4-methoxy-2, 6-dimethylphenyl) sulfonyl] methylamino] ethoxy]-N-methylacetamide, fumarate (LF22-0542), a novel nonpeptidic bradykinin B1 receptor antagonist. *J. Pharmacol. Exp. Ther.* **318**, 195-205.
- Premkumar, L. S. and Ahern, G. P. (2000) Induction of vanilloid receptor channel activity by protein kinase C. *Nature* **408**, 985-990.
- Prescott, E. D. and Julius, D. (2003) A modular PIP2 binding site as a determinant of capsaicin receptor sensitivity. *Science* **300**, 1284-1288.
- Price, T. J., Cervero, F., Gold, M. S., Hammond, D. L. and Prescott, S. A. (2009) Chloride regulation in the pain pathway. *Brain Res. Rev.* **60**, 149-170.
- Qin, C., Farber, J. P., Miller, K. E. and Foreman, R. D. (2006) Responses of thoracic spinal neurons to activation and desensitization of cardiac TRPV1-containing afferents in rats. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **291**, R1700-R1707.
- Quintão, N. L., Passos, G. F., Medeiros, R., Paszcuk, A. F., Motta, F. L., Pesquero, J. B., Campos, M. M. and Calixto, J. B. (2008) Neuropathic pain-like behavior after brachial plexus avulsion in mice: the relevance of kinin B1 and B2 receptors. *J. Neurosci.* **28**, 2856-2863.
- Rang, H. and Ritchie, J. (1988) Depolarization of nonmyelinated fibers of the rat vagus nerve produced by activation of protein kinase C. *J. Neurosci.* **8**, 2606-2617.
- Reeh, P. W. and Pethö, G. (2000) Nociceptor excitation by thermal sensitization—a hypothesis. *Prog. Brain Res.* **129**, 39-50.
- Rong, W., Hillsley, K., Davis, J. B., Hicks, G., Winchester, W. J. and Grundy, D. (2004) Jejunal afferent nerve sensitivity in wild-type and TRPV1 knockout mice. *J. Physiol.* **560**, 867-881.
- Rueff, A. and Dray, A. (1993) Sensitization of peripheral afferent fibres in the *in vitro* neonatal rat spinal cord-tail by bradykinin and prostaglandins. *Neuroscience* **54**, 527-535.
- Rupniak, N. M., Boyce, S., Webb, J. K., Williams, A. R., Carlson, E. J., Hill, R. G., Borkowski, J. A. and Hess, J. F. (1997) Effects of the bradykinin B 1 receptor antagonist des-Arg 9 [Leu 8] bradykinin and genetic disruption of the B2 receptor on nociception in rats and mice. *Pain* **71**, 89-97.
- Sauer, S., Schäfer, D., Kress, M. and Reeh, P. (1998) Stimulated prostaglandin E 2 release from rat skin, *in vitro*. *Life Sci.* **62**, 2045-2055.
- Sauer, S. K., Averbeck, B. and Reeh, P. W. (2000) Denervation and NK1 receptor block modulate stimulated CGRP and PGE2 release from rat skin. *Neuroreport* **11**, 283-286.
- Schuligoi, R., Donnerer, J. and Amann, R. (1994) Bradykinin-induced sensitization of afferent neurons in the rat paw. *Neuroscience* **59**, 211-215.
- Shin, J., Cho, H., Hwang, S. W., Jung, J., Shin, C. Y., Lee, S.-Y., Kim, S. H., Lee, M. G., Choi, Y. H. and Kim, J. (2002) Bradykinin-12-lipoxygenase-VR1 signaling pathway for inflammatory hyperalgesia. *Proc. Natl. Acad. Sci. U.S.A.* **99**, 10150-10155.
- Song, I., Althoff, C., Hermann, K., Scheel, A., Knetsch, T., Burmester, G. and Backhaus, M. (2008) Contrast-enhanced ultrasound in monitoring the efficacy of a bradykinin receptor 2 antagonist in painful knee osteoarthritis compared with MRI. *Ann. Rheum. Dis.* **68**, 75-83.
- Soukhova-O'Hare, G. K., Zhang, J. W., Gozal, D. and Yu, J. (2006) Bradykinin B 2 receptors mediate pulmonary sympathetic afferents induced reflexes in rabbits. *Life Sci.* **78**, 1990-1997.
- Staruschenko, A., Jeske, N. A. and Akopian, A. N. (2010) Contribution of TRPV1-TRPA1 interaction to the single channel properties of the TRPA1 channel. *J. Biol. Chem.* **285**, 15167-15177.
- Steranka, L. R., Manning, D. C., DeHaas, C. J., Ferkany, J. W., Borosky, S. A., Connor, J. R., Vavrek, R. J., Stewart, J. M. and Snyder, S. H. (1988) Bradykinin as a pain mediator: receptors are localized to sensory neurons, and antagonists have analgesic actions. *Proc. Natl. Acad. Sci. U.S.A.* **85**, 3245-3249.
- Stevens, P. A., Pyne, S., Grady, M. and Pyne, N. J. (1994) Bradykinin-dependent activation of adenylate cyclase activity and cyclic AMP accumulation in tracheal smooth muscle occurs via protein kinase C-dependent and-independent pathways. *Biochem. J.* **297**, 233-239.
- Stucky, C., Abrahams, L. and Seybold, V. (1998) Bradykinin increases the proportion of neonatal rat dorsal root ganglion neurons that respond to capsaicin and protons. *Neuroscience* **84**, 1257-1265.
- Sugiura, T., Tominaga, M., Katsuya, H. and Mizumura, K. (2002) Bradykinin lowers the threshold temperature for heat activation of vanilloid receptor 1. *J. Neurophysiol.* **88**, 544-548.
- Sung, K.-W., Kirby, M., McDonald, M. P., Lovinger, D. M. and Delpire, E. (2000) Abnormal GABA<sub>A</sub> receptor-mediated currents in dorsal root ganglion neurons isolated from Na<sup>+</sup>-K<sup>+</sup>-2Cl<sup>-</sup> cotransporter null mice. *J. Neurosci.* **20**, 7531-7538.
- Taiwo, Y., Heller, P. and Levine, J. (1990) Characterization of distinct phospholipases mediating bradykinin and noradrenaline hyperalgesia. *Neuroscience* **39**, 523-531.
- Taiwo, Y. O. and Levine, J. D. (1988) Characterization of the arachidonic acid metabolites mediating bradykinin and noradrenaline hyperalgesia. *Brain Res.* **458**, 402-406.
- Tang, H.-B., Inoue, A., Oshita, K., Hirate, K. and Nakata, Y. (2005) Zaltoprofen inhibits bradykinin-induced responses by blocking the activation of second messenger signaling cascades in rat dorsal root ganglion cells. *Neuropharmacology* **48**, 1035-1042.

- Tang, H.-B., Inoue, A., Oshita, K. and Nakata, Y. (2004) Sensitization of vanilloid receptor 1 induced by bradykinin via the activation of second messenger signaling cascades in rat primary afferent neurons. *Eur. J. Pharmacol.* **498**, 37-43.
- Tang, H. B., Inoue, A., Iwasa, M., Hide, I. and Nakata, Y. (2006) Substance P release evoked by capsaicin or potassium from rat cultured dorsal root ganglion neurons is conversely modulated with bradykinin. *J. Neurochem.* **97**, 1412-1418.
- Thayer, S. A., Perney, T. M. and Miller, R. J. (1988) Regulation of calcium homeostasis in sensory neurons by bradykinin. *J. Neurosci.* **8**, 4089-4097.
- Vasko, M., Campbell, W. and Waite, K. (1994) Prostaglandin E2 enhances bradykinin-stimulated release of neuropeptides from rat sensory neurons in culture. *J. Neurosci.* **14**, 4987-4997.
- Vaughn, A. H. and Gold, M. S. (2010) Ionic mechanisms underlying inflammatory mediator-induced sensitization of dural afferents. *J. Neurosci.* **30**, 7878-7888.
- Vellani, V., Mapplebeck, S., Moriondo, A., Davis, J. B. and McNaughton, P. A. (2001) Protein kinase C activation potentiates gating of the vanilloid receptor VR1 by capsaicin, protons, heat and anandamide. *J. Physiol.* **534**, 813-825.
- Vellani, V., Zachrisson, O. and McNaughton, P. A. (2004) Functional bradykinin B1 receptors are expressed in nociceptive neurones and are upregulated by the neurotrophin GDNF. *J. Physiol.* **560**, 391-401.
- Voets, T., Droogmans, G., Wissenbach, U., Janssens, A., Flockerzi, V. and Nilius, B. (2004) The principle of temperature-dependent gating in cold- and heat-sensitive TRP channels. *Nature* **430**, 748-754.
- Vyklický, L., Vlachova, V., Vitaskova, Z., Dittert, I., Kabat, M. and Orkand, R. (1999) Temperature coefficient of membrane currents induced by noxious heat in sensory neurones in the rat. *J. Physiol.* **517**, 181-192.
- Walter, T., Chau, T. and Weichman, B. (1989) Effects of analgesics on bradykinin-induced writhing in mice presensitized with PGE 2. *Agents Actions* **27**, 375-377.
- Wang, M. M., Reynaud, D. and Pace-Asciak, C. R. (1999) *In vivo* stimulation of 12(S)-lipoxygenase in the rat skin by bradykinin and platelet activating factor: formation of 12(S)-HETE and hepxilins, and actions on vascular permeability. *Biochim. Biophys. Acta* **1436**, 354-362.
- Wang, S., Dai, Y., Fukuoka, T., Yamanaka, H., Kobayashi, K., Obata, K., Cui, X., Tominaga, M. and Noguchi, K. (2008) Phospholipase C and protein kinase A mediate bradykinin sensitization of TRPA1: a molecular mechanism of inflammatory pain. *Brain* **131**, 1241-1251.
- Wang, S., Joseph, J., Ro, J. Y. and Chung, M. K. (2015) Modality-specific mechanisms of protein kinase C-induced hypersensitivity of TRPV1: S800 is a polymodal sensitization site. *Pain* **156**, 931-941.
- Weinreich, D., Koschorke, G., Udem, B. and Taylor, G. (1995) Prevention of the excitatory actions of bradykinin by inhibition of PGI2 formation in nodose neurones of the guinea-pig. *J. Physiol.* **483**, 735-746.
- Weinreich, D. and Wonderlin, W. (1987) Inhibition of calcium-dependent spike after-hyperpolarization increases excitability of rabbit visceral sensory neurones. *J. Physiol.* **394**, 415-427.
- Weng, H.-J., Patel, K. N., Jeske, N. A., Bierbower, S. M., Zou, W., Tiwari, V., Zheng, Q., Tang, Z., Mo, G. C. and Wang, Y. (2015) Tmem100 is a regulator of TRPA1-TRPV1 complex and contributes to persistent pain. *Neuron* **85**, 833-846.
- Whalley, E., Clegg, S., Stewart, J. and Vavrek, R. (1987) The effect of kinin agonists and antagonists on the pain response of the human blister base. *Naunyn Schmiedeberg's Arch. Pharmacol.* **336**, 652-655.
- Wu, Z.-Z. and Pan, H.-L. (2007) Role of TRPV1 and intracellular Ca<sup>2+</sup> in excitation of cardiac sensory neurons by bradykinin. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **293**, R276-R283.
- Yamaguchi-Sase, S., Hayashi, I., Okamoto, H., Nara, Y., Matsuzaki, S., Hoka, S. and Majima, M. (2003) Amelioration of hyperalgesia by kinin receptor antagonists or kininogen deficiency in chronic constriction nerve injury in rats. *Inflamm. Res.* **52**, 164-169.
- Yanaga, F., Hirata, M. and Koga, T. (1991) Evidence for coupling of bradykinin receptors to a guanine-nucleotide binding protein to stimulate arachidonate liberation in the osteoblast-like cell line, MC3T3-E1. *Biochim. Biophys. Acta* **1094**, 139-146.
- Yu, S. and Ouyang, A. (2009) TRPA1 in bradykinin-induced mechanical hypersensitivity of vagal C fibers in guinea pig esophagus. *Am. J. Physiol. Gastrointest. Liver Physiol.* **296**, G255-G265.
- Zahner, M. R., Li, D. P., Chen, S. R. and Pan, H. L. (2003) Cardiac vanilloid receptor 1-expressing afferent nerves and their role in the cardiogenic sympathetic reflex in rats. *J. Physiol.* **551**, 515-523.
- Zhang, X., Li, L. and McNaughton, P. A. (2008) Proinflammatory mediators modulate the heat-activated ion channel TRPV1 via the scaffolding protein AKAP79/150. *Neuron* **59**, 450-461.
- Zylka, M. J., Rice, F. L. and Anderson, D. J. (2005) Topographically distinct epidermal nociceptive circuits revealed by axonal tracers targeted to Mrgprd. *Neuron* **45**, 17-25.