

## ( $p, q$ )-EXTENSION OF THE WHITTAKER FUNCTION AND ITS CERTAIN PROPERTIES

SHOWKAT AHMAD DAR AND MOHD SHADAB

ABSTRACT. In this paper, we obtain a  $(p, q)$ -extension of the Whittaker function  $M_{k, \mu}(z)$  together with its integral representations, by using the extended confluent hypergeometric function of the first kind  $\Phi_{p, q}(b; c; z)$  [recently extended by J. Choi]. Also, we give some of its main properties, namely the summation formula, a transformation formula, a Mellin transform, a differential formula and inequalities. In addition, our extension on Whittaker function finds interesting connection with the Laguerre polynomials.

### 1. Introduction and preliminaries

The simple Beta function  $B(\alpha, \beta)$  is defined by [11, p. 26, eq. (48)]

$$(1.1) \quad B(\alpha, \beta) = \begin{cases} \int_0^1 t^{\alpha-1}(1-t)^{\beta-1} dt, & (\Re(\alpha) > 0; \Re(\beta) > 0), \\ \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}, & (\Re(\alpha) < 0; \Re(\beta) < 0), \end{cases}$$

where  $\alpha, \beta \in \mathbb{C} \setminus \mathbb{Z}_0^-$ .

Gauss hypergeometric function (GHF) is defined by [12, p. 124]

$$(1.2) \quad F \left( \begin{matrix} a, b; \\ c; \end{matrix} z \right) = \sum_{\ell=0}^{\infty} \frac{(a)_{\ell}(b)_{\ell}}{(c)_{\ell}} \frac{z^{\ell}}{\ell!},$$

where  $|z| < 1$ ,  $c \in \mathbb{C}$ .

Confluent hypergeometric function (CHF) of the first kind  $\Phi(\cdot)$  is defined by [12, p. 124]

$$(1.3) \quad \Phi \left( \begin{matrix} b; \\ c; \end{matrix} z \right) = \sum_{\ell=0}^{\infty} \frac{(b)_{\ell}}{(c)_{\ell}} \frac{z^{\ell}}{\ell!}.$$

---

Received February 2, 2017; Revised August 26, 2017; Accepted November 2, 2017.

2010 *Mathematics Subject Classification.* 33C15, 33B15, 33C05, 35A22, 33C45.

*Key words and phrases.* Whittaker function, extended Beta function, Mellin transform, extended confluent hypergeometric function, Laguerre polynomials, extended Gauss hypergeometric function.

An integral representations of GHF and CHF are given by

$$(1.4) \quad F(a, b; c; z) = \frac{1}{B(b, c-b)} \int_0^1 \left[ t^{b-1} (1-t)^{c-b-1} (1-zt)^{-a} \right] dt,$$

where  $|\arg(1-z)| < \pi$ ,  $\Re(c) > \Re(b) > 0$ ,  
and

$$(1.5) \quad \Phi(b; c; z) = \frac{1}{B(b, c-b)} \int_0^1 \left[ t^{b-1} (1-t)^{c-b-1} \exp(zt) \right] dt,$$

where  $\Re(c) > \Re(b) > 0$ .

From the equations (1.4) and (1.5), using summation series of  $(1-zt)^{-a}$  and  $\exp(zt)$ , we find

$$(1.6) \quad F(a, b; c; z) = \sum_{\ell=0}^{\infty} \frac{(a)_{\ell} B(b+\ell, c-b) z^{\ell}}{B(b, c-b) \ell!}, \quad |z| < 1, \Re(c) > \Re(b) > 0,$$

and

$$(1.7) \quad \Phi(b; c; z) = \sum_{\ell=0}^{\infty} \frac{B(b+\ell, c-b) z^{\ell}}{B(b, c-b) \ell!}, \quad \Re(c) > \Re(b) > 0.$$

The definition of Whittaker function of the first kind  $M_{k,\mu}(z)$  in [11, p. 39, Eq. (23)] and [13] is given by

$$(1.8) \quad M_{k,\mu}(z) = z^{\mu+\frac{1}{2}} \exp\left(-\frac{z}{2}\right) \Phi\left(\mu-k+\frac{1}{2}; 2\mu+1; z\right),$$

where  $-\pi < \arg(z) \leq \pi$ ,  $k \in \mathbb{C}$ ,  $\Re(\mu) > -\frac{1}{2}$ ,  $\Re(\mu-k) > -\frac{1}{2}$ ,  $2\mu \in \mathbb{C} \setminus \mathbb{Z}^-$  and  $\Phi(\cdot)$  is the confluent hypergeometric function of first kind.

In 1997, Chaudhry et al. [3, p. 20, Eq. (1.7)] gave a  $p$ -extension of the Beta function  $B(x, y)$  given by

$$(1.9) \quad B(x, y; p) = \int_0^1 t^{x-1} (1-t)^{y-1} \exp\left[\frac{-p}{t(1-t)}\right] dt, \quad \Re(p) > 0$$

and they proved that this extension has connections with the Macdonald, error and Whittaker function. In 2004, Chaudhry et al. [4], given extension of GHF  $F(a, b; c; z)$  and CHF  $\Phi(b; c; z)$  based on the extended Beta function  $B(x, y; p)$ , which they were denoted by  $F_p(a, b; c; z)$  and  $\Phi_p(b; c; z)$ .

An integral representations of  $F_p(a, b; c; z)$  and  $\Phi_p(b; c; z)$  are given by [4] (1.10)

$$F_p(a, b; c; z) = \frac{1}{B(b, c-b)} \int_0^1 \left[ t^{b-1} (1-t)^{c-b-1} (1-zt)^{-a} \exp\left(-\frac{p}{t(1-t)}\right) \right] dt,$$

where  $|\arg(1-z)| < \pi$ ,  $p > 0$ ,  $p = 0$ ,  $\Re(c) > \Re(b) > 0$ ,  
and

$$(1.11) \quad \Phi_p(b; c; z) = \frac{1}{B(b, c-b)} \int_0^1 \left[ t^{b-1} (1-t)^{c-b-1} \exp\left(zt - \frac{p}{t(1-t)}\right) \right] dt,$$

where  $p > 0$ ,  $p = 0$ , and  $\Re(c) > \Re(b) > 0$ .

The extended Whittaker function  $M_{p,k,\mu}(\cdot)$  based on the extended confluent hypergeometric function  $\Phi_p(\cdot)$  is defined by [8]

$$(1.12) \quad M_{p,k,\mu}(z) = z^{\mu+\frac{1}{2}} \exp\left(-\frac{z}{2}\right) \Phi_p\left(\mu - k + \frac{1}{2}; 2\mu + 1; z\right),$$

where  $p \geq 0, -\pi < \arg(z) \leq \pi, k \in \mathbb{C}, \Re(\mu) > -\frac{1}{2}, \Re(\mu - k) > -\frac{1}{2}, 2\mu \in \mathbb{C} \setminus \mathbb{Z}^-$ . This definition clearly reduces to the simple Whittaker function, when we set  $p = 0$ .

Recently, Choi et al. [6] has given  $(p, q)$ -extension of the Beta function  $B(x, y)$  by adding one more parameter  $q$ , denoted and defined by

$$(1.13) \quad B(x, y; p, q) \equiv B_{p,q}(x, y) = \int_0^1 \left[ t^{x-1}(1-t)^{y-1} \exp\left(-\frac{p}{t} - \frac{q}{1-t}\right) \right] dt,$$

where  $\Re(p) > 0, \Re(q) > 0$  and also given extension of GHF  $F(a, b; c; z)$  and CHF  $\Phi(b; c; z)$ , based on the extended Beta function  $B_{p,q}(x, y)$ , which are defined by [6]

$$(1.14) \quad F_{p,q}(a, b; c; z) = \sum_{\ell=0}^{\infty} \frac{(a)_{\ell} B_{p,q}(b + \ell, c - b)}{B(b, c - b)} \frac{z^{\ell}}{\ell!},$$

where  $|z| < 1, p \geq 0, q \geq 0, \Re(c) > \Re(b) > 0$ , and

$$(1.15) \quad \Phi_{p,q}(b; c; z) = \sum_{\ell=0}^{\infty} \frac{B_{p,q}(b + \ell, c - b)}{B(b, c - b)} \frac{z^{\ell}}{\ell!},$$

where  $p \geq 0, q \geq 0, \Re(c) > \Re(b) > 0$ .

An integral representation of  $F_{p,q}(a, b; c; z)$  and  $\Phi_{p,q}(b; c; z)$  are given by [6]

$$(1.16) \quad F_{p,q}(a, b; c; z) = \frac{1}{B(b, c - b)} \int_0^1 \left[ t^{b-1}(1-t)^{c-b-1}(1-zt)^{-a} \exp\left(-\frac{p}{t} - \frac{q}{1-t}\right) \right] dt,$$

where  $p \geq 0, q \geq 0, \Re(c) > \Re(b) > 0, |\arg(1 - z)| < \pi$ ,

and

$$(1.17) \quad \Phi_{p,q}(b; c; z) = \frac{1}{B(b, c - b)} \int_0^1 \left[ t^{b-1}(1-t)^{c-b-1} \exp\left(zt - \frac{p}{t} - \frac{q}{1-t}\right) \right] dt,$$

where  $p \geq 0, q \geq 0, \Re(c) > \Re(b) > 0$ .

Differentiation formula of Eq. (1.15) with respect to  $z$  is given by

$$(1.18) \quad \frac{d^n}{dz^n} \{\Phi_{p,q}(b; c; z)\} = \frac{(b)_n}{(c)_n} \Phi_{p,q}(b + n; c + n; z), \quad (n \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}).$$

Transformation formula of  $\Phi_{p,q}(b; c; z)$  is defined by

$$(1.19) \quad \Phi_{p,q}(b; c; z) = \exp(z) \Phi_{q,p}(c - b; c; -z).$$

The Laguerre polynomials  $L_n^\alpha(z)$  is defined by [10, pp. 201–202]

$$(1.20) \quad L_n^\alpha(z) = \frac{(1+\alpha)_n}{n!} {}_1F_1\left(-n; 1+\alpha; z\right) = \sum_{k=0}^n \frac{(-1)^k (1+\alpha)_n z^k}{k!(n-k)!(1+\alpha)_k}.$$

In view of Eq. (1.20), the definition of simple Laguerre polynomials  $L_m(p) = L_m^0(p)$  ( $m \in \mathbb{N}_0$ ) is given by the generating function [10, p. 202]

$$(1.21) \quad \exp\left(-\frac{pt}{1-t}\right) = (1-t)^{1+\alpha} \sum_{m=0}^{\infty} t^m L_m^\alpha(p), \quad (\alpha \in \mathbb{C}, |t| < 1),$$

and for  $\alpha \geq 0$ ,  $0 \leq p < \infty$ .

Replace  $t \rightarrow 1-t$  and  $\alpha = 0$  in (1.21), which yield

$$(1.22) \quad \exp\left(-\frac{p}{t}\right) = e^{-p} \left[ t \sum_{n=0}^{\infty} L_n(p) (1-t)^n \right], \quad (t \in \mathbb{D}_1 := \{t \in \mathbb{R} : |1-t| < 1\}).$$

Also, replace  $t \rightarrow 1-t$  and  $p = q$  in (1.22), which yield

$$(1.23) \quad \exp\left(-\frac{q}{1-t}\right) = e^{-q} \left[ (1-t) \sum_{m=0}^{\infty} L_m(q) t^m \right], \quad (t \in \mathbb{D}_2 := \{t \in \mathbb{R} : |t| < 1\}),$$

comparing the equations (1.22) and (1.23), we find

$$(1.24) \quad \exp\left(-\frac{p}{t} - \frac{q}{1-t}\right) = e^{-p-q} \left[ \sum_{n,m=0}^{\infty} L_n(p) L_m(q) (1-t)^{n+1} t^{m+1} \right], \quad (t \in \mathbb{D}_1 \cap \mathbb{D}_2).$$

In this paper, we obtain a  $(p, q)$ -extension of the Whittaker function  $M_{k,\mu}(\cdot)$  together with its integral representations, which we denoted by  $M_{p,q,k,\mu}(\cdot)$ . Moreover, our work is also motivated by the following references [1, 2, 5, 9]. The plan of this paper is as follow. The  $(p, q)$ -extension of the Whittaker function in (1.12) is given in Section 2 and its integral representations are shown in Section 3. We consider some properties of this function, namely the summation formula, a transformation formula, inequalities, a Mellin transform and a differential formula are given in Sections 4–8. We introduce concluding remarks about the function are given in Section 9.

## 2. The $(p, q)$ -extended Whittaker function $M_{p,q,k,\mu}(\cdot)$

Here, we consider the following  $(p, q)$ -extension of the Whittaker function  $M_{k,\mu}(z)$  based on the extended Beta function  $B_{p,q}(x, y)$ , which we denoted by  $M_{p,q,k,\mu}(\cdot)$ , and defined by

$$(2.1) \quad M_{p,q,k,\mu}(z) = z^{\mu+\frac{1}{2}} \exp\left(-\frac{z}{2}\right) \Phi_{p,q}\left(\mu - k + \frac{1}{2}; 2\mu + 1; z\right),$$

where  $p \geq 0, q \geq 0$ ,  $-\pi < \arg(z) \leq \pi$ ,  $k \in \mathbb{C}$ ,  $\Re(\mu) > -\frac{1}{2}$ ,  $\Re(\mu - k) > -\frac{1}{2}$ ,  $2\mu \in \mathbb{C} \setminus \mathbb{Z}^-$  and  $\Phi_{p,q}(\cdot)$  is the extended confluent hypergeometric function

of the first kind defined in (1.17). This definition clearly reduces to the simple Whittaker function when  $p = 0 = q$ .

### 3. Integral representations for $M_{p,q,k,\mu}(\cdot)$

**Theorem 1.** *Each of the following integral representations of (p, q)-extended Whittaker functions hold for  $\Re(p) > 0, \Re(q) > 0, \Re(\mu) > -\frac{1}{2}, \Re(\mu \pm k) > -\frac{1}{2}$ .*

• *The integral representation of  $M_{p,q,k,\mu}(\cdot)$  can be obtained by using (1.17) in Eq. (2.1), we find*

$$\begin{aligned}
 M_{p,q,k,\mu}(z) &= \frac{z^{\mu+\frac{1}{2}} \exp\left(-\frac{z}{2}\right)}{B\left(\mu-k+\frac{1}{2}, \mu+k+\frac{1}{2}\right)} \\
 (3.1) \quad &\times \int_0^1 \left[ t^{\mu-k-\frac{1}{2}}(1-t)^{\mu+k-\frac{1}{2}} \exp\left(zt - \frac{p}{t} - \frac{q}{1-t}\right) \right] dt.
 \end{aligned}$$

• *If we set the transformation  $t \rightarrow \frac{(u-\alpha)}{(\beta-\alpha)}$ , where  $(\beta - \alpha) > 0$  [when  $t = 0, u = \alpha$  and  $t = 1, u = \beta$ ] in Eq. (3.1), we find*

$$\begin{aligned}
 M_{p,q,k,\mu}(z) &= \frac{(\beta - \alpha)^{-2\mu} z^{\mu+\frac{1}{2}} \exp\left(-\frac{z}{2}\right)}{B\left(\mu-k+\frac{1}{2}, \mu+k+\frac{1}{2}\right)} \int_{\alpha}^{\beta} \left[ (u-\alpha)^{\mu-k-\frac{1}{2}}(\beta-u)^{\mu+k-\frac{1}{2}} \right. \\
 (3.2) \quad &\times \left. \exp\left(\frac{z(u-\alpha)}{\beta-\alpha} - \frac{p(\beta-\alpha)}{u-\alpha} - \frac{q(\beta-\alpha)}{\beta-u}\right) \right] du.
 \end{aligned}$$

• *If we set  $\alpha = -1$  and  $\beta = 1$  in Eq. (3.2), we get following another integral representation*

$$\begin{aligned}
 M_{p,q,k,\mu}(z) &= \frac{(2)^{-2\mu} z^{\mu+\frac{1}{2}} \exp\left(-\frac{z}{2}\right)}{B\left(\mu-k+\frac{1}{2}, \mu+k+\frac{1}{2}\right)} \int_{-1}^1 \left[ (1+u)^{\mu-k-\frac{1}{2}}(1-u)^{\mu+k-\frac{1}{2}} \right. \\
 (3.3) \quad &\times \left. \exp\left(\frac{z(u+1)}{2} - \frac{2p}{u+1} - \frac{2q}{1-u}\right) \right] du.
 \end{aligned}$$

• *Upon setting  $t \rightarrow \frac{u}{1+u}$ , ( $u \neq -1$ ) [when  $t \rightarrow 0, u \rightarrow 0$ , and  $t \rightarrow 1, u \rightarrow \infty$ ] in Eq. (3.1), we obtain*

$$\begin{aligned}
 M_{p,q,k,\mu}(z) &= \frac{z^{\mu+\frac{1}{2}} \exp\left(-\frac{z}{2}\right)}{B\left(\mu-k+\frac{1}{2}, \mu+k+\frac{1}{2}\right)} \\
 (3.4) \quad &\times \int_0^{\infty} \left[ \frac{u^{\mu-k-\frac{1}{2}}}{(1+u)^{2\mu+1}} \exp\left(\frac{zu}{1+u} - \frac{p(1+u)}{u} - q(1+u)\right) \right] du.
 \end{aligned}$$

• *Clearly, if we consider  $p = 0 = q$  in (3.1)-(3.4), we get another following integral representations of the simple Whittaker function:*

$$\begin{aligned}
 M_{k,\mu}(z) &= \frac{z^{\mu+\frac{1}{2}} \exp\left(-\frac{z}{2}\right)}{B\left(\mu-k+\frac{1}{2}, \mu+k+\frac{1}{2}\right)} \int_0^1 \left[ t^{\mu-k-\frac{1}{2}}(1-t)^{\mu+k-\frac{1}{2}} \exp(zt) \right] dt, \\
 &= \frac{(\beta - \alpha)^{-2\mu} z^{\mu+\frac{1}{2}} \exp\left(-\frac{z}{2}\right)}{B\left(\mu-k+\frac{1}{2}, \mu+k+\frac{1}{2}\right)}
 \end{aligned}$$

$$(3.5) \quad \times \int_{\alpha}^{\beta} \left[ (u - \alpha)^{\mu - k - \frac{1}{2}} (\beta - u)^{\mu + k - \frac{1}{2}} \exp\left(\frac{z(u - \alpha)}{\beta - \alpha}\right) \right] du,$$

where  $\beta \neq \alpha$  and  $\alpha, \beta$  are two scalars.

$$(3.6) \quad \begin{aligned} M_{k, \mu}(z) &= \frac{(2)^{-2\mu} z^{\mu + \frac{1}{2}} \exp\left(-\frac{z}{2}\right)}{B\left(\mu - k + \frac{1}{2}, \mu + k + \frac{1}{2}\right)} \\ &\quad \times \int_{-1}^1 \left[ (1 + u)^{\mu - k - \frac{1}{2}} (1 - u)^{\mu + k - \frac{1}{2}} \exp\left(\frac{z(u + 1)}{2}\right) \right] du \\ &= \frac{z^{\mu + \frac{1}{2}} \exp\left(-\frac{z}{2}\right)}{B\left(\mu - k + \frac{1}{2}, \mu + k + \frac{1}{2}\right)} \\ &\quad \times \int_0^{\infty} \left[ \frac{(u)^{\mu - k - \frac{1}{2}}}{(1 + u)^{2\mu + 1}} \exp\left(\frac{zu}{1 + u}\right) \right] du, \quad u \neq -1. \end{aligned}$$

#### 4. Summation formula for $M_{p, q, k, \mu}(\cdot)$

We obtain the following summation formula of the  $(p, q)$ -extended Whittaker function  $M_{p, q, k, \mu}(\cdot)$ .

**Theorem 2.** *The following summation formula hold true:*

$$(4.1) \quad \begin{aligned} M_{p, q, k, \mu}(z) &= \frac{\exp(-p - q) z^{\mu + \frac{1}{2}} \exp\left(-\frac{z}{2}\right)}{B\left(\mu - k + \frac{1}{2}, \mu + k + \frac{1}{2}\right)} \\ &\quad \times \sum_{n, m=0}^{\infty} \left[ L_n(p) L_m(q) B\left(\mu - k + m + \frac{3}{2}, \mu + k + n + \frac{3}{2}\right) \right. \\ &\quad \left. \times \Phi\left(\mu - k + m + \frac{3}{2}; 2\mu + n + m + 3; z\right) \right], \end{aligned}$$

where  $\Re(\mu) > -\frac{1}{2}$ ,  $\Re(\mu - k + m) > -\frac{3}{2}$ ,  $\Re(\mu \pm k) > -\frac{1}{2}$ .

*Proof.* We begin with left hand side of (4.1) and involving integral representation (3.1), which yield

$$(4.2) \quad \begin{aligned} M_{p, q, k, \mu}(z) &= \frac{z^{\mu + \frac{1}{2}} \exp\left(-\frac{z}{2}\right)}{B\left(\mu - k + \frac{1}{2}, \mu + k + \frac{1}{2}\right)} \\ &\quad \times \int_0^1 \left[ t^{\mu - k - \frac{1}{2}} (1 - t)^{\mu + k - \frac{1}{2}} dt \exp(zt) \exp\left(-\frac{p}{t} - \frac{q}{1 - t}\right) \right] dt, \end{aligned}$$

we find (4.3) upon using the result (1.24) in Eq. (4.2)

$$(4.3) \quad \begin{aligned} M_{p, q, k, \mu}(z) &= \frac{\exp(-p - q) z^{\mu + \frac{1}{2}} \exp\left(-\frac{z}{2}\right)}{B\left(\mu - k + \frac{1}{2}, \mu + k + \frac{1}{2}\right)} \sum_{n, m=0}^{\infty} \left[ L_n(p) L_m(q) \right. \\ &\quad \left. \times \int_0^1 t^{\mu - k + m + \frac{3}{2} - 1} (1 - t)^{\mu + k + n + \frac{3}{2} - 1} \exp(zt) \right] dt. \end{aligned}$$

Finally, applying the confluent hypergeometric function (1.5) in the above Eq. (4.3), we get right hand side of (4.1) which is our required result.  $\square$

**5. Transformation formula for  $M_{p,q,k,\mu}(\cdot)$**

**Theorem 3.** *The following transformation formula hold true:*

$$(5.1) \quad M_{p,q,k,\mu}(-z) = (-1)^{\mu+\frac{1}{2}} M_{p,q,-k,\mu}(z),$$

where  $p \geq 0, q \geq 0, \Re(\mu) > -\frac{1}{2}, 2\mu \neq -1, -2, -3, \dots$

*Proof.* We take left hand side of (5.1), and applying the definition of Whittaker function given in (2.1). Then we find

$$(5.2) \quad M_{p,q,k,\mu}(-z) = (-1)^{\mu+\frac{1}{2}} (z)^{\mu+\frac{1}{2}} \exp\left(\frac{z}{2}\right) \Phi_{p,q}\left(\mu - k + \frac{1}{2}; 2\mu + 1; -z\right).$$

Using transformation formula of the extended confluent hypergeometric function (1.19) in the above Eq. (5.2), we obtain

$$(5.3) \quad M_{p,q,k,\mu}(-z) = (-1)^{\mu+\frac{1}{2}} (z)^{\mu+\frac{1}{2}} \exp\left(-\frac{z}{2}\right) \Phi_{q,p}\left(\mu + k + \frac{1}{2}; 2\mu + 1; z\right).$$

Lastly, applying (2.1) in the above Eq. (5.3), we get right hand side of (5.1).  $\square$

**6. Inequalities for  $M_{p,q,k,\mu}(\cdot)$**

**Theorem 4.** *The following inequalities hold true.*

$$(6.1) \quad M_{p,q,k,\mu}(z) \leq \exp(-2p - 2q) M_{k,\mu}(z),$$

where  $p > 0, q > 0, \Re(\mu) > -\frac{1}{2}, \Re(\mu \pm k) > -\frac{1}{2}$ .

*Proof.* We start with left hand side of (6.1) and involving the integral representation (3.4), we get

$$(6.2) \quad M_{p,q,k,\mu}(z) = \frac{z^{\mu+\frac{1}{2}} \exp\left(-\frac{z}{2}\right)}{B\left(\mu - k + \frac{1}{2}, \mu + k + \frac{1}{2}\right)} \times \int_0^\infty \left[ \frac{(u)^{\mu-k-\frac{1}{2}}}{(1+u)^{2\mu+1}} \exp\left(\frac{zu}{1+u}\right) \exp\{-p(1+u^{-1})\} \times \exp\{-q(1+u)\} \right] du.$$

For  $p > 0, q > 0$ , we have

$$(6.3) \quad p(1+u^{-1}) \geq 2p; u \in (0, 1],$$

$$(6.4) \quad \Rightarrow \exp\{-p(1+u^{-1})\} \leq \exp(-2p),$$

also

$$(6.5) \quad q(1+u) \geq 2q; u \in [1, \infty),$$

$$(6.6) \quad \Rightarrow \exp\{-q(1+u)\} \leq \exp(-2q).$$

Using inequalities (6.4) and (6.6) in the above Eq. (6.2), which yield

$$(6.7) \quad M_{p,q,k,\mu}(z) \leq \frac{\exp(-2p-2q) z^{\mu+\frac{1}{2}} \exp\left(-\frac{z}{2}\right)}{B\left(\mu-k+\frac{1}{2}, \mu+k+\frac{1}{2}\right)} \times \int_0^\infty \left[ \frac{u^{\mu-k-\frac{1}{2}}}{(1+u)^{2\mu+1}} \exp\left(\frac{zu}{1+u}\right) \right] du.$$

Finally, using (3.6) in the above Eq. (6.7), we get right hand side of (6.1), which is our required result.

Here, we obtain an another interesting result by using the following inequality:

$$(6.8) \quad \exp(x) \geq 1 + \frac{x^n}{n!}, \quad n > 0, \quad x > 0,$$

in the above Eq. (3.1), we find

$$(6.9) \quad M_{p,q,k,\mu}(z) \geq \frac{z^{\mu+\frac{1}{2}} \exp\left(-\frac{z}{2}\right)}{B\left(\mu-k+\frac{1}{2}, \mu+k+\frac{1}{2}\right)} \times \left[ \int_0^1 t^{\mu-k-\frac{1}{2}} (1-t)^{\mu+k-\frac{1}{2}} \exp\left(-\frac{p}{t} - \frac{q}{1-t}\right) dt + \frac{z^n}{n!} \int_0^1 t^{\mu-k+n-\frac{1}{2}} (1-t)^{\mu+k-\frac{1}{2}} \exp\left(-\frac{p}{t} - \frac{q}{1-t}\right) dt \right].$$

Using the extended Beta function (1.13) in the above Eq. (6.9), we get

$$(6.10) \quad M_{p,q,k,\mu}(z) \geq \frac{z^{\mu+\frac{1}{2}} \exp\left(-\frac{z}{2}\right)}{B\left(\mu-k+\frac{1}{2}, \mu+k+\frac{1}{2}\right)} \left[ B\left(\mu-k+\frac{1}{2}, \mu+k+\frac{1}{2}; p, q\right) + \frac{z^n}{n!} B\left(\mu-k+n+\frac{1}{2}, \mu+k+\frac{1}{2}; p, q\right) \right]. \quad \square$$

## 7. Mellin transforms for $M_{p,q,k,\mu}(\cdot)$

We obtain the Mellin transform of  $M_{p,q,k,\mu}(\cdot)$  by multiply the both sides of Eq. (2.1) with  $p^{r-1}$  and  $q^{s-1}$ , then integrate with respect to  $p$  and  $q$  over  $(0, \infty)$ , we get the following result.

**Theorem 5.** *The following Mellin transformation hold true*

$$(7.1) \quad \int_0^\infty \int_0^\infty p^{r-1} q^{s-1} M_{p,q,k,\mu}(z) dp dq = \frac{\Gamma(r)\Gamma(s)B\left(\mu-k+r+s+\frac{1}{2}, \mu+k+r+s+\frac{1}{2}\right) z^{-r-s}}{B\left(\mu-k+\frac{1}{2}, \mu+k+\frac{1}{2}\right)} M_{p,q,k,\mu+r+s}(z),$$

where  $\Re(r) > 0$ ,  $\Re(s) > 0$ ,  $\Re(\mu \pm k + r + s) > -\frac{1}{2}$ ,  $\Re(\mu \pm k) > -\frac{1}{2}$ .



*Proof.* We take left hand side of (7.1) upon using (2.1), we get

$$\begin{aligned}
(7.2) \quad & \int_0^\infty \int_0^\infty p^{r-1} q^{s-1} M_{p,q,k,\mu}(z) dpdq \\
&= \int_0^\infty \int_0^\infty \left[ p^{r-1} q^{s-1} z^{\mu+\frac{1}{2}} \exp\left(-\frac{z}{2}\right) \Phi_{p,q}\left(\mu-k+\frac{1}{2}; 2\mu+1; z\right) \right] dpdq.
\end{aligned}$$

Applying (1.17) in Eq. (7.2) and change the order of integration, we find

$$\begin{aligned}
(7.3) \quad & \int_0^\infty \int_0^\infty p^{r-1} q^{s-1} M_{p,q,k,\mu}(z) dpdq \\
&= \frac{z^{\mu+\frac{1}{2}} \exp\left(-\frac{z}{2}\right)}{B\left(\mu-k+\frac{1}{2}, \mu+k+\frac{1}{2}\right)} \int_0^1 \left[ t^{\mu-k-\frac{1}{2}} (1-t)^{\mu+k-\frac{1}{2}} dt \exp(zt) \right. \\
&\quad \left. \times \left\{ \int_0^\infty p^{r-1} \exp\left(-\frac{p}{t}\right) dp \int_0^\infty q^{s-1} \exp\left(-\frac{q}{1-t}\right) dq \right\} \right] dt.
\end{aligned}$$

If we set  $\frac{p}{t} \rightarrow z$  [when  $p = 0, z = 0; p \rightarrow \infty, z \rightarrow \infty$ ] and  $\frac{q}{1-t} \rightarrow u$  [when  $q = 0, u = 0; q \rightarrow \infty, u \rightarrow \infty$ ] in the above Eq. (7.3), we find

$$\begin{aligned}
(7.4) \quad & \int_0^\infty \int_0^\infty p^{r-1} q^{s-1} M_{p,q,k,\mu}(z) dpdq \\
&= \frac{\Gamma(r)\Gamma(s)z^{\mu+\frac{1}{2}} \exp\left(-\frac{z}{2}\right)}{B\left(\mu-k+\frac{1}{2}, \mu+k+\frac{1}{2}\right)} \int_0^1 \left[ t^{\mu-k+r-\frac{1}{2}} (1-t)^{\mu+k+s-\frac{1}{2}} \exp(zt) \right] dt.
\end{aligned}$$

Upon using confluent hypergeometric function (1.5) in Eq. (7.4), we obtain

$$\begin{aligned}
(7.5) \quad & \int_0^\infty \int_0^\infty p^{r-1} q^{s-1} M_{p,q,k,\mu}(z) dpdq \\
&= \frac{z^{\mu+r+s+\frac{1}{2}} \exp\left(-\frac{z}{2}\right) \Gamma(r)\Gamma(s) z^{-r-s}}{B\left(\mu-k+\frac{1}{2}, \mu+k+\frac{1}{2}\right)} \Phi\left(\mu-k+r+\frac{1}{2}; 2\mu+r+s+1; z\right) \\
&\quad \times B\left(\mu-k+r+\frac{1}{2}, \mu+k+s+\frac{1}{2}\right).
\end{aligned}$$

Applying the classical Whittaker function (1.8) in Eq. (7.5), which gives

$$\begin{aligned}
(7.6) \quad & \int_0^\infty \int_0^\infty p^{r-1} q^{s-1} M_{p,q,k,\mu}(z) dpdq \\
&= \frac{\Gamma(r)\Gamma(s)B\left(\mu-k+r+\frac{1}{2}, \mu+k+s+\frac{1}{2}\right) z^{-r-s}}{B\left(\mu-k+\frac{1}{2}, \mu+k+\frac{1}{2}\right)} M_{k,\mu+r+s}(z).
\end{aligned}$$

If we take  $r = 1 = s$  in (7.6), we get the following another interesting result

$$(7.7) \quad \int_0^\infty \int_0^\infty M_{p,q,k,\mu}(z) dpdq = \frac{B\left(\mu-k+\frac{3}{2}, \mu+k+\frac{3}{2}\right) z^{-2}}{B\left(\mu-k+\frac{1}{2}, \mu+k+\frac{1}{2}\right)} M_{k,\mu+2}(z). \quad \square$$

**Theorem 6.** *If  $p \geq 0, q \geq 0, (2\alpha - \beta) > 0$  and  $\Re(\mu \pm \alpha) > -\frac{1}{2}$ , then*

$$(7.8) \quad \int_0^\infty \exp(-\alpha x) x^{\alpha-1} M_{p,q,k,\mu}(\beta x) dx$$

$$= \frac{\Gamma(\alpha + \mu + \frac{1}{2})}{(\alpha + \frac{\beta}{2})^{\alpha + \mu + \frac{1}{2}}} F_{p,q} \left( \mu + k + \frac{1}{2}, \mu - k + \frac{1}{2}; 2\mu + 1; \frac{2\alpha}{2\alpha + \beta} \right),$$

where  $|\arg(1 - 2\beta/2\alpha + \beta)| < \pi$ .

*Proof.* We begin with left hand side of (7.8) and using the definition of extended Whittaker function (2.1), which yields

$$(7.9) \quad \int_0^\infty \exp(-\alpha x) x^{\alpha-1} M_{p,q,k,\mu}(\beta x) dx \\ = \int_0^\infty \left[ \exp(-\alpha x) x^{\alpha-1} (\beta x)^{\mu+\frac{1}{2}} \exp\left(-\frac{\beta x}{2}\right) \Phi_{p,q}\left(\mu - k + \frac{1}{2}; 2\mu + 1; \beta x\right) \right] dx.$$

Applying the extended confluent hypergeometric function (1.17) in the above Eq. (7.9) and change the order of integration (with uniform convergence), which gives

$$(7.10) \quad M_{p,q,k,\mu}(z) = \frac{\beta^{\mu+\frac{1}{2}}}{B(\mu - k + \frac{1}{2}, \mu + k + \frac{1}{2})} \int_0^1 \left[ t^{\mu-k-\frac{1}{2}} (1-t)^{\mu+k-\frac{1}{2}} \right. \\ \left. \times \exp\left(-\frac{p}{t} - \frac{q}{1-t}\right) \int_0^\infty \exp\left(-\left(\alpha + \frac{\beta}{2} - \beta t\right)x\right) x^{\alpha+\mu-\frac{1}{2}} dx \right] dt.$$

Applying the Euler's integral formula [7, Eq. (5)],

$$(7.11) \quad \int_0^\infty e^{-st} t^{z-1} dt = \frac{\Gamma(z)}{s^z}, \quad \Re(z) > 0, \quad \Re(s) > 0$$

in the above Eq. (7.10), we find

$$(7.12) \quad M_{p,q,k,\mu}(z) = \frac{\beta^{\mu+\frac{1}{2}} \Gamma(\mu + \alpha + \frac{1}{2})}{(\alpha + \frac{\beta}{2})^{\mu+\alpha+\frac{1}{2}} B(\mu - k + \frac{1}{2}, \mu + k + \frac{1}{2})} \\ \times \int_0^1 \left[ t^{\mu-k-\frac{1}{2}} (1-t)^{\mu+k-\frac{1}{2}} \left(1 - \frac{2\beta t}{2\alpha + \beta}\right)^{-(\mu+\alpha+\frac{1}{2})} \right. \\ \left. \times \exp\left(-\frac{p}{t} - \frac{q}{1-t}\right) \right] dt,$$

where  $|\arg(1 - 2\beta/2\alpha + \beta)| < \pi$ .

Finally, applying the integral representation (1.16) in Eq. (7.12), we get right hand side of (7.8).  $\square$

**Corollary.** If we take  $\beta = 2\alpha$  in the above Eq. (7.8), we find

$$(7.13) \quad \int_0^\infty \exp(-\alpha x) x^{\alpha-1} M_{p,q,k,\mu}(2\alpha x) dx = \frac{\Gamma(\mu + \alpha + \frac{1}{2}) B(\mu - k + \frac{1}{2}, k - \alpha; p, q)}{B(\mu - k + \frac{1}{2}, \mu + k + \frac{1}{2}) (2\alpha)^\alpha},$$

where  $\alpha > 0$ ;  $q, p > 0$ , and  $\Re(\alpha + \mu) > -\frac{1}{2}$ .

### 8. Differentiation formula for $M_{p,q,k,\mu}(\cdot)$

Differentiation formula of (p, q)-extended Whittaker function  $M_{p,q,k,\mu}(\cdot)$  upon using the following formula:

$$(8.1) \quad (\lambda)_{1+K} = (\lambda)_1(\lambda + 1)_K.$$

**Theorem 7.** *The following differential formula hold true:*

$$(8.2) \quad \begin{aligned} & \frac{d^n}{dz^n} \left\{ z^{-\mu-\frac{1}{2}} \exp\left(\frac{z}{2}\right) M_{p,q,k,\mu}(z) \right\} \\ &= \frac{(\mu - k + \frac{1}{2})_n}{(2\mu + 1)_n} \exp\left(\frac{z}{2}\right) z^{-\mu-\frac{1}{2}-\frac{n}{2}} M_{p,q,k-\frac{n}{2},\mu+\frac{n}{2}}(z), \end{aligned}$$

where  $n = 0, 1, 2, 3, \dots$

*Proof.* Using (2.1) in the above Eq. (8.2), which yield

$$(8.3) \quad z^{-\mu-\frac{1}{2}} \exp\left(\frac{z}{2}\right) M_{p,q,k,\mu}(z) = \Phi_{p,q}\left(\mu - k + \frac{1}{2}; 2\mu + 1; z\right).$$

From (8.3) we call  $n^{th}$  derivative on both sides and using result (8.1), which yield

$$(8.4) \quad \begin{aligned} & \frac{d^n}{dz^n} \left\{ z^{-\mu-\frac{1}{2}} \exp\left(\frac{z}{2}\right) M_{p,q,k,\mu}(z) \right\} \\ &= \frac{(\mu - k + \frac{1}{2})_n}{(2\mu + 1)_n} \Phi_{p,q}\left(\mu - k + \frac{1}{2} + n; 2\mu + 1 + n; z\right). \end{aligned}$$

Finally, applying (2.1) in the above Eq. (8.4), we get right hand side of (8.2).  $\square$

### 9. Conclusion remark

The Whittaker function has various applications in the field of Mathematical physics and engineering sciences. Here in this paper, we have obtained (p, q)-extension of the Whittaker function together with its integral representation. We consider some properties for this function, namely the summation formula, a transformation formula, a Mellin transform, a differential formula and inequalities. Therefore, these evaluated results may be useful in the field of Engineering sciences, Mathematical physics, and so on. Also, this extended Whittaker function have some relations with Laguerre polynomials, Macdonald function, error function, Modified Bessel function and G-function.

**Acknowledgement.** We (authors) are expressing sincere thanks to the referees for give their valuable suggestions to us, which let to a better presentation of the paper.

### References

- [1] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions*. Dover publications. Newyork, 1972.
- [2] G. E. Andrews, R. Askey, and R. Roy, *Special Functions*, Encyclopedia of Mathematics and its Applications, **71**, Cambridge University Press, Cambridge, 1999.
- [3] M. A. Chaudhry, A. Qadir, M. Rafique, and S. M. Zubair, *Extension of Euler's beta function*, J. Comput. Appl. Math. **78** (1997), no. 1, 19–32.
- [4] M. A. Chaudhry, A. Qadir, H. M. Srivastava, and R. B. Paris, *Extended hypergeometric and confluent hypergeometric functions*, Appl. Math. Comput. **159** (2004), no. 2, 589–602.
- [5] J. Choi, M. Ghayasuddin, and N. Khan, *Generalized extended Whittaker function and its properties*, Appl. Math. Sci. **9** (2015), no. 131, 6529–6541.
- [6] J. Choi, A. K. Rathie, and R. K. Parmar, *Extension of extended Beta, hypergeometric function and confluent hypergeometric function*, Honam Math. J. **36** (2014), no. 2, 357–385.
- [7] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi, *Higher Transcendental Functions*, McGraw-Hill, New York, Toronto and London, 1953.
- [8] D. K. Nagar, R. A. M. Vásquez, and A. K. Gupta, *Properties of the extended Whittaker function*, Progress in Applied Mathematics **6** (2013), no. 2, 70–80.
- [9] E. Özergin, M. A. Özarslan, and A. Altn, *Extension of gamma, beta and hypergeometric functions*, J. Comput. Appl. Math. **235** (2011), no. 16, 4601–4610.
- [10] E. D. Rainville, *Special Functions*. Macmillan Company, New York, 1960; Reprinted by Chelsea Publishing Company, Bronx, New York, 1971.
- [11] H. M. Srivastava and H. L. Monocha, *A Treatise on Generating Functions*. Halsted Press (Ellis Horwood Limited, Chichester, U.K.) John Wiley and Sons, New york, Chichester, Brisbane and Toronto, 1984.
- [12] L. J. Slater, *Generalized Hypergeometric Functions*, Cambridge University Press, Cambridge, 1966.
- [13] E. T. Whittaker and G. N. Watson, *A Course of Modern Analysis*, reprint of the fourth (1927) edition, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1996.

SHOWKAT AHMAD DAR  
 DEPARTMENT OF APPLIED SCIENCES AND HUMANITIES  
 FACULTY OF ENGINEERING AND TECHNOLOGY  
 JAMIA MILLIA ISLAMIA (CENTRAL UNIVERSITY)  
 NEW DELHI, 110025, INDIA  
 Email address: [showkatjmi34@gmail.com](mailto:showkatjmi34@gmail.com)

MOHD SHADAB  
 DEPARTMENT OF APPLIED SCIENCES AND HUMANITIES  
 FACULTY OF ENGINEERING AND TECHNOLOGY  
 JAMIA MILLIA ISLAMIA (CENTRAL UNIVERSITY)  
 NEW DELHI, 110025, INDIA  
 Email address: [shadabmohd786@gmail.com](mailto:shadabmohd786@gmail.com)