

Invited Mini Review

The role of necroptosis in the treatment of diseases

Young Sik Cho*

Department of Pharmacy, Keimyung University, Daegu 42601, Korea

Necroptosis is an emerging form of programmed cell death occurring via active and well-regulated necrosis, distinct from apoptosis morphologically, and biochemically. Necroptosis is mainly unmasked when apoptosis is compromised in response to tumor necrosis factor alpha. Unlike apoptotic cells, which are cleared by macrophages or neighboring cells, necrotic cells release danger signals, triggering inflammation, and exacerbating tissue damage. Evidence increasingly suggests that programmed necrosis is not only associated with pathophysiology of disease, but also induces innate immune response to viral infection. Therefore, necroptotic cell death plays both physiological and pathological roles. Physiologically, necroptosis induce an innate immune response as well as premature assembly of viral particles in cells infected with virus that abrogates host apoptotic machinery. On the other hand, necroptosis per se is detrimental, causing various diseases such as sepsis, neurodegenerative diseases and ischemic reperfusion injury. This review discusses the signaling pathways leading to necroptosis, associated necroptotic proteins with target-specific inhibitors and diseases involved. Several studies currently focus on protective approaches to inhibiting necroptotic cell death. In cancer biology, however, anticancer drug resistance severely hampers the efficacy of chemotherapy based on apoptosis. Pharmacological switch of cell death finds therapeutic application in drug-resistant cancers. Therefore, the possible clinical role of necroptosis in cancer control will be discussed in brief. [BMB Reports 2018; 51(5): 219-224]

INTRODUCTION

Cell survival is constantly in dynamic equilibrium with cell death for tissue homeostasis. Aside from cell survival, cell death occurs in various modes such as programmed or random mechanisms upon exposure to various stresses.

*Corresponding author. Tel: +82-53-580-6466; Fax: +82-53-580-6645; E-mail: yscho123@kmu.ac.kr

<https://doi.org/10.5483/BMBRep.2018.51.5.074>

Received 20 March 2018

Keywords: Apoptosis, Cancer, Necroptosis, Necroptosis modulator, Regulator

Programmed cell death or apoptosis is activated via intrinsic or extrinsic pathways in response to death stimuli. Both pathways involve a series of signaling molecules with activated caspases and is termed caspase-dependent cell death. Meanwhile, necroptosis is classified as type III programmed cell death with apoptotic and necrotic features. It is also known as programmed necrosis or caspase-independent cell death according to morphological or molecular features. Initially, necroptosis was considered as undesirable passive cell death. However, it is currently considered as specialized cell death since it is orchestrated under a caspase-compromised condition. Similar to apoptosis, necroptosis is executed via distinctive signaling mechanism comprising a cascade of specified proteins, resulting in regulated necrotic cell death. Despite unknown mechanisms and pathological significance compared with apoptosis, the discovery of pharmacological inhibitors targeting necroptosis has been extensively pursued. Here, we introduce the concept of necroptotic cell death induced under various pathophysiological conditions and delineate the pathological mechanisms unmasking necroptosis in a well-orchestrated fashion. In this review, necroptosis-associated diseases and the underlying pathogenesis will be discussed.

SIGNALING PATHWAYS LEADING TO NECROPTOSIS INDUCTION AND BIOCHEMICAL CHANGES

Typically, necroptosis occurs in cells exposed to extrinsic stimuli such as TNF α , FASL, and TRAIL in combination with compromised caspase-8 (C-8). C-8 inhibition is induced by genetic defects, viral proteins, and treatment with a pan-caspase inhibitor zVAD (1-3). Necroptosis-regulating proteins have been identified and biologically validated comprehensively. Genome-wide siRNA screening yielded a few candidate genes associated with necroptosis (4). In addition to RIP1 as a necroptosis regulator, RIP3, mixed lineage kinase domain-like protein (MLKL) and PGAM5 were identified (5-8). The signaling pathway leading to necroptosis is illustrated in Fig. 1. TNF α ligation to its cognate receptor triggers caspase-dependent cell death as a default cell death mode. Caspase-compromised conditions drive cell death via necroptosis with accompanying RIP1 activation (9). Upon stimulation, RIP1 interacts with RIP3 leading to the formation of a necrosome complex via RIP homotypic interaction motif (RHIM) within cells (10). MLKL pseudokinase was identified as




Fig. 1. Necroptosis regulators and signaling pathways in necroptosis activation. Upon TNF receptor ligation, a signal transduction mediated via RIP1, caspase activation, and tBid cleavage occurs, resulting in apoptotic cell death. Active caspase-8 inhibits necroptosis via cleavage of RIP1 and RIP3. Inhibition of caspase leads to the formation of a physical complex of RIP1 with RIP3 to trigger downstream signaling events including MLKL and PGAM5 recruitment, and transduction of cytosolic death signals to mitochondria, resulting in necroptosis.

a substrate of RIP3 under necroptotic condition. Phosphorylated RIP3 induces MLKL phosphorylation at serine or threonine sites. The RIP3-MLKL complex translocates to mitochondria-associated membrane mediated via phosphoglycerate mutase 5 (PGAM5), which is a necrosome-associated protein regulating dynamic-related protein (Drp1) in turn (8). Finally, Drp1 induces mitochondrial fragmentation, a crucial event for necroptosis (11).

Biochemical changes during the early stages of necroptosis include ATP depletion, ROS generation, calcium overload, and loss of mitochondrial permeability transition (12). At the cellular level, necroptosis is characterized by loss of plasma membrane integrity as well as organelle swelling, finally leading to cellular collapse (10). Notably, necroptotic cells release various damage-associated molecular patterns (DAMPs) such as HMGB1, cytokines and histones into extracellular media. Particularly, a disulfide form of HMGB1 by oxidation contributes to the inflammatory response (13). HMGB1 binds to a cognate receptor on endothelial cells and macrophages to transduce cellular responses such as release of proinflammatory cytokines and chemotactic cell migration (14).

NECROPTOSIS: TARGET PROTEINS AND INHIBITORS

As described above, a series of necroptosis-associated proteins were identified and further validated as targets of necroptosis. Subsequently, a few small molecules were successfully

Table 1. Target proteins associated with necroptosis and their specific inhibitors

Necroptosis Target proteins	Inhibitors	Mode of action/function
RIP-1	Nec-1 Furo[2,3-d]pyrimidine derivatives GSK'963	Allosteric inhibitor SAR reported >10000-fold selective for RIP1
RIP-3	GSK'840, GSK'843, GSK'872	Caspase activity activated
MLKL	Necrosulfonamide TC13172 GW806742X	Covalent binding Covalent binding (nanomolar potency) Binds to pseudokinase domain (ATP-mimetic)
Others	Ponatinib & pazopanib Sorafenib TPCK Hydroxyanisole Diphenyleneiodonium (DPI)	Interfere with signaling proteins upstream of MLKL Target unknown UCH-L1 activator ROS scavenger ROS suppression

discovered, specifically targeting necroptosis proteins. Target-specific small molecules that can modulate necroptosis are listed in Table 1.

RIP-1 inhibitors

Necrostatin-1 (Nec-1), 5-(1H-indol-3-ylmethyl)-2-thiohydantoin 1, was first discovered by screening of necroptosis inhibitors and later identified as an allosteric inhibitor of RIP1 via stabilization of an inactive conformation of the kinase domain (KD) (15). It has been widely validated in various necroptosis-associated animal models (16-19). Accordingly, Nec-1 plays a crucial role in prevention or alleviation of necroptotic damage caused by various stimuli via targeting RIP1. Another potent Nec-1 derivative Nec-1s (7-Cl-O-Nec-1) showed greater specificity to RIP1 compared with other kinases. Other inhibitors have been developed as RIP1 inhibitors, including furo[2,3-d]pyrimidines and GSK'963 (20, 21).

RIP3 inhibitors

Recent studies proposed that necroptosis is mediated in a RIP1- or RIP3-dependent manner (22, 23). Unlike RIP1, RIP3 is essential for necroptosis, but not apoptosis, and targeting RIP3 appears to be more specific in controlling necroptosis. Interestingly, silencing of RIP3 substantially protected cells and tissues against necroptosis (24). Small molecules such as GSK'840, GSK'843 and GSK'872 have been reported to suppress RIP3-dependent necroptosis (25). Mechanistically, these molecules form a complex with RIP3 to restore caspase

activity. Consequently, RIP3 inhibitors potentially protect cells from diverse stimuli than RIP1 inhibitors.

MLKL inhibitors

Necrosulfonamide ((E)-N-(4-(N-(3-methoxy-pyrazin-2-yl) sulfamoyl) phenyl)-3-(5-nitrothiophene-2-yl) acrylamide, NAS) protects cells against TNF-induced necroptosis via covalent modification of a cysteine in MLKL (7). A compound TC13172 is an MLKL inhibitor with single nanomolar potency. It induced covalent binding at Cys-86 of MLKL (26). In addition, GW806742X targets the pseudo-kinase domain of MLKL with nanomolar inhibitory activity, protecting against necroptosis, although it shows off-target effects against other kinases (26).

Additional agents have been developed as inhibitors of necroptosis acting directly via targets that have yet to be designated. Ponatinib and pazopanib (27) are two anti-cancer agents identified from a representative panel of United States Food and Drug Administration (FDA)-approved drugs through phenotypic screening. Both drugs abrogate phosphorylation of MLKL during TNF α -induced necroptosis, suggesting interference with the signaling proteins upstream of MLKL. However, they do not rescue cells from apoptosis. Interestingly, ponatinib inhibits both RIP1 and RIP3 while pazopanib acts against preferential inhibitor of RIP1. Further, sorafenib is a multi-targeting kinase inhibitor with potent inhibitory activity against B-RAK, and is widely used for the treatment of acute leukemia. It blocks signaling target proteins upstream of MLKL such as RIP1 and RIP3, although its real target remains elusive (28). Therefore, sorafenib can be harnessed for fine-tuning of necroptosis-inducing agents.

TPCK

N-tosyl-L-phenylalanine-chloromethyl ketone (TPCK), a serine protease inhibitor, protects cells from TNF-mediated necroptosis. Using a TPCK probe, HtrA2/Omi has been identified as a target and it acts as an activator of ubiquitin C-terminal hydrolase (UCH-L1) resulting in necroptosis (29).

ROS scavengers

Reactive oxygen species (ROS) play a role in diverse signaling pathways, due to high reactivity with biomolecules such as proteins, DNA and lipids. Upon exposure of L929 cells to TNF α , necroptotic signaling generates ROS via mitochondrial transport chain but not cytosolic enzyme (30), which was strongly supported by reports suggesting that butylated hydroxyanisole blocks ROS accumulation and cell death (31). An NADPH oxidase inhibitor diphenyleneiodonium (DPI) protects renal tubular epithelial cells against necroptosis by blocking ROS generation (32).

PHYSIOLOGICAL AND PATHOLOGICAL SIGNIFICANCE OF NECROPTOSIS

Necroptosis was originally considered as undesirable cell

death upon exposure to stimuli, inducing tissue damage. Furthermore, it is the alternative cell death induced under conditions of defective apoptosis. However, growing evidence suggests that necroptosis is mediated via an orchestrated and specialized pathway (33-35). Well-regulated cell death occurs in a range of biological phenomena such as development, immunology and differentiation. Further, extrinsic apoptosis and necroptosis contribute to host defense mechanism against microbial infection. Viruses such as adenoviruses, poxviruses, and herpes viruses evade host apoptosis machinery, and perpetuate if host cells only execute caspase-dependent default cell death (36-38). For instance, vaccinia viruses encode a caspase 8 inhibitor to block apoptotic cell death upon infection. Under this caspase-compromised condition, cells are committed to alternative necroptosis (6). The resulting necroptosis is vital to provoke innate immune response by killing virus-infected cells and releasing danger signals from host cells into external milieu. Furthermore, necroptosis in T cells regulates antigen-activated T-cell proliferation and survival. Caspase-8 negatively regulates necroptosis, promoting survival of activated T cells under physiological conditions. In mice lacking caspase-8, T cells fail to show immune response when infected with murine hepatitis virus (39).

Pathologically, evidence suggests that necroptotic cell death leads to various diseases. Sepsis is mainly caused by Gram-negative bacteria, which release endotoxin that elicits systemic inflammation via release of TNF α and IL-1 inflammatory cytokines. Necroptosis occurs in ischemia-reperfusion (I/R) injury and neurodegenerative diseases. During restoration of blood flow into tissues, tissue damage occurs with severe neutrophil infiltration and cytokine production. Furthermore, necroptosis is involved in traumatic brain and spinal cord injuries (19, 40). Excitotoxicity, Huntington's disease and retinal degeneration are closely associated with necroptosis. Exposure to a specific RIP inhibitor Nec-1 effectively protects cells from necroptosis. Nec-1 protects hippocampal HT-22 cells against glutamate-induced oxytosis (41). Inhibition of RIP1 kinase or RIP3 silencing significantly rescues necroptotic cell death (42). Furthermore, Nec-1 reduces or delays necroptotic damage in transgenic mice expressing mutant Huntingtin protein, astrocytes from amyotrophic lateral sclerosis, and retinal pigment epithelium (42-44). Interestingly, microbial infection induces necroptosis in host cells. Microbial proliferation occurs by circumventing default programmed cell death in the host. Necroptosis is induced by viral infections such as vaccinia virus (VV). VV expresses an inhibitor of caspase-1 and -8, diverting the host response toward necroptosis in a RIP3-dependent pathway (45). Further, Sendai virus induces necroptosis in neuroblastoma cells (17). Macrophage infected with *S. typhimurium* induces necroptotic cell death in a RIP1- and RIP3-dependent manner.

PERSPECTIVES OF NECROPTOSIS AND CONCLUSIONS

Necroptosis has been recognized as an alternative to apoptosis when cells are exposed to various stimuli under specific conditions.

As mentioned above, it induces a variety of pathological conditions including septic shock, acute pancreatitis and neuronal degeneration. Apoptotic death of damaged cells is removed by phagocytosis in macrophages or neighboring cells. Conversely, because loss of membrane integrity occurs in cells undergoing necroptosis, intracellular substances including heat-shock proteins and HMGB1 are released into extracellular media to provoke inflammation and immune responses.

In response to infectious viruses or intracellular bacterial pathogens escaping apoptotic cell death, host cells actively switch to necroptosis, causing premature assembly of virus particles or bacteria progeny and release of critical components triggering an immune response (46-48).

Similarly, necroptosis was demonstrated under conditions of chronic sterile inflammation such as alcoholic-induced liver injury and atherosclerosis (18, 49). Under these pathological conditions, necroptotic cells release DAMPS to trigger sterile inflammation, via unknown mechanism.

A potent and specific regulation of necroptosis is therefore, needed. Indeed, pharmacological blockade of necroptosis is of primary concern in the treatment of various diseases. Further studies will be extensively undertaken to identify the target molecules mediating the signal transduction leading to necroptosis and facilitate the discovery of mechanism-based inhibitors.

On the other hand, necroptosis induction can be actively harnessed via molecular switch or unmasking. Generally, cancer cells grow in an uncontrolled manner and further acquire mechanisms to evade cell death intrinsically or extrinsically. Chemotherapy or radiotherapy is mainly based on apoptosis through caspase activation. However, many cancers have developed strategies to disarm apoptotic machinery, including dysregulated apoptosis, activation of pro-survival signaling pathways and upregulation of drug transporters. Altered apoptosis contributes to drug resistance of cancer during chemotherapy. To overcome drug-resistant cancers, alternative cell death mechanisms such as necroptosis or autophagy can be considered. In previous reports, cancer cells that are refractory to apoptotic agents were shown to succumb to necroptosis-inducing agents (50). It is conceivable that resistance to apoptosis can be overcome by necroptosis, because necroptotic pathway is distinct from apoptotic mechanisms. Furthermore, several reports suggest that necroptosis contributes to suppression of tumorigenesis. Particularly, mutations in the *CYLD* gene aggressively facilitate carcinoma via upregulation of angiogenic factors (51). Further, *RIP3* gene polymorphisms occur in non-Hodgkin's lymphoma

and levels of a spliced variant *RIP3-γ* are relatively high in colon and lung cancers (52, 53).

Practically, a few inducers of necroptosis have been reported to trigger necroptotic cell death in malignant cancers although their targets remain to be identified. Caspase activation is a prerequisite for apoptosis induction, and apoptotic cell death fails if caspases are blocked or compromised by unknown mechanisms. Under those conditions, cells activate necroptosis in response to cell death stimuli. In fact, a natural product shikonin induces necroptotic cell death in MCF-7 breast cancer cells that express Bcl-2 or Bcl-xL, which acquire multidrug resistance. Obatoclax, an antagonist of Bcl-2 family members, promotes necroptosis based on autophagy in acute lymphoblastic leukemia resistant to glucocorticoids. In addition, combined treatment of pancaspase inhibitors with 5-fluorouracil drives necroptosis in colorectal cancer (54). Furthermore, necroptotic cells trigger adaptive immunity in dendritic cells (DCs), which in turn activate CD8⁺ T cells for antitumor immunity (55).

However, necroptosis-based cancer therapy still remains elusive. It has been demonstrated that necroptosis in tumor microenvironment contributes to inflammation and cancer metastasis (16). Furthermore, the low expression of key modulators of necroptosis in specific cancers fails to induce necroptosis (56-58), resulting in cancer evasion.

In conclusion, suppression or enhancement of necroptosis is therapeutically effective in specific diseases. A comprehensive insight into the underlying mechanisms is needed to facilitate the diagnosis, biomarkers, and drug development in necroptosis-associated diseases. Most studies have focused on the identification of specific inhibitors targeting necroptosis, and their underlying mechanisms of regulation. As a result, a few small molecules have been discovered from the chemical library, and optimized for further clinical use.

CONFLICTS OF INTEREST

The authors have no conflicting interests.

REFERENCES

1. Jouan-Lanhuet S, Arshad MI, Piquet-Pellorce C et al (2012) TRAIL induces necroptosis involving RIPK1/RIPK3-dependent PARP-1 activation. *Cell Death Differ* 19, 2003-2014
2. Moriwaki K, Bertin J, Gough PJ, Orlowski GM and Chan FK (2015) Differential roles of RIPK1 and RIPK3 in TNF-induced necroptosis and chemotherapeutic agent-induced cell death. *Cell Death Dis* 6, e1636
3. Miao B and Degterev A (2009) Methods to analyze cellular necroptosis. *Methods Mol Biol* 559, 79-93
4. Hitomi J, Christofferson DE, Ng A et al (2008) Identification of a molecular signaling network that regulates a cellular necrotic cell death pathway. *Cell* 135, 1311-1323

5. Degtrev A, Hitomi J, Gernscheid M et al (2008) Identification of RIP1 kinase as a specific cellular target of necrostatins. *Nat Chem Biol* 4, 313-321
6. Cho YS, Challa S, Moquin D et al (2009) Phosphorylation-driven assembly of the RIP1-RIP3 complex regulates programmed necrosis and virus-induced inflammation. *Cell* 137, 1112-1123
7. Sun L, Wang H, Wang Z et al (2012) Mixed lineage kinase domain-like protein mediates necrosis signaling downstream of RIP3 kinase. *Cell* 148, 213-227
8. Wang Z, Jiang H, Chen S, Du F and Wang X (2012) The mitochondrial phosphatase PGAM5 functions at the convergence point of multiple necrotic death pathways. *Cell* 148, 228-243
9. Feoktistova M and Leverkus M (2015) Programmed necrosis and necroptosis signalling. *FEBS J* 282, 19-31
10. Belizario J, Vieira-Cordeiro L and Enns S (2015) Necroptotic Cell Death Signaling and Execution Pathway: Lessons from Knockout Mice. *Mediators Inflamm* 2015, 128076
11. Frank S, Gaume B, Bergmann-Leitner ES et al (2001) The role of dynamin-related protein 1, a mediator of mitochondrial fission, in apoptosis. *Dev Cell* 1, 515-525
12. Vanlangenakker N, Vanden Berghe T, Krysko DV, Festjens N and Vandenabeele P (2008) Molecular mechanisms and pathophysiology of necrotic cell death. *Curr Mol Med* 8, 207-220
13. Janko C, Filipovic M, Munoz LE et al (2014) Redox modulation of HMGB1-related signaling. *Antioxid Redox Signal* 20, 1075-1085
14. Stern D, Yan SD, Yan SF and Schmidt AM (2002) Receptor for advanced glycation endproducts: a multiligand receptor magnifying cell stress in diverse pathologic settings. *Adv Drug Deliv Rev* 54, 1615-1625
15. Degtrev A, Maki JL and Yuan J (2013) Activity and specificity of necrostatin-1, small-molecule inhibitor of RIP1 kinase. *Cell Death Differ* 20, 366
16. Liu ZY, Wu B, Guo YS et al (2015) Necrostatin-1 reduces intestinal inflammation and colitis-associated tumorigenesis in mice. *Am J Cancer Res* 5, 3174-3185
17. Nomura M, Ueno A, Saga K, Fukuzawa M and Kaneda Y (2014) Accumulation of cytosolic calcium induces necroptotic cell death in human neuroblastoma. *Cancer Res* 74, 1056-1066
18. Roychowdhury S, McMullen MR, Pisano SG, Liu X and Nagy LE (2013) Absence of receptor interacting protein kinase 3 prevents ethanol-induced liver injury. *Hepatology* 57, 1773-1783
19. Wang Y, Wang H, Tao Y, Zhang S, Wang J and Feng X (2014) Necroptosis inhibitor necrostatin-1 promotes cell protection and physiological function in traumatic spinal cord injury. *Neuroscience* 266, 91-101
20. Harris PA, Bandyopadhyay D, Berger SB et al (2013) Discovery of small molecule RIP1 kinase inhibitors for the treatment of pathologies associated with necroptosis. *ACS Med Chem Lett* 4, 1238-1243
21. Berger SB, Harris P, Nagilla R et al (2015) Characterization of GSK'963: a structurally distinct, potent and selective inhibitor of RIP1 kinase. *Cell Death Discov* 1, 15009
22. Vandenabeele P, Declercq W, Van Herreweghe F and Vanden Berghe T (2010) The role of the kinases RIP1 and RIP3 in TNF-induced necrosis. *Sci Signal* 3, re4
23. Park SY, Shim JH and Cho YS (2014) Distinctive roles of receptor-interacting protein kinases 1 and 3 in caspase-independent cell death of L929. *Cell Biochem Funct* 32, 62-69
24. Fettweis G, Di Valentin E, L'Homme L et al (2017) RIP3 antagonizes a TSC2-mediated pro-survival pathway in glioblastoma cell death. *Biochim Biophys Acta* 1864, 113-124
25. Mandal P, Berger SB, Pillay S et al (2014) RIP3 induces apoptosis independent of pronecrotic kinase activity. *Mol Cell* 56, 481-495
26. Yan B, Liu L, Huang S et al (2017) Discovery of a new class of highly potent necroptosis inhibitors targeting the mixed lineage kinase domain-like protein. *Chem Commun (Camb)* 53, 3637-3640
27. Fauster A, Rebsamen M, Huber KV et al (2015) A cellular screen identifies ponatinib and pazopanib as inhibitors of necroptosis. *Cell Death Dis* 6, e1767
28. Feldmann F, Schenk B, Martens S, Vandenabeele P and Fulda S (2017) Sorafenib inhibits therapeutic induction of necroptosis in acute leukemia cells. *Oncotarget* 8, 68208-68220
29. Sosna J, Voigt S, Mathieu S et al (2013) The proteases HtrA2/Omi and UCH-L1 regulate TNF-induced necroptosis. *Cell Commun Signal* 11, 76
30. Ardestani S, Deskins DL and Young PP (2013) Membrane TNF-alpha-activated programmed necrosis is mediated by ceramide-induced reactive oxygen species. *J Mol Signal* 8, 12
31. Lin Y, Choksi S, Shen HM et al (2004) Tumor necrosis factor-induced nonapoptotic cell death requires receptor-interacting protein-mediated cellular reactive oxygen species accumulation. *J Biol Chem* 279, 10822-10828
32. Dong W, Li Z, Chen Y et al (2017) NADPH oxidase inhibitor, diphenyleneiodonium prevents necroptosis in HK-2 cells. *Biomed Rep* 7, 226-230
33. Fulda S (2016) Regulation of necroptosis signaling and cell death by reactive oxygen species. *Biol Chem* 397, 657-660
34. Galluzzi L and Kroemer G (2008) Necroptosis: a specialized pathway of programmed necrosis. *Cell* 135, 1161-1163
35. Liu C, Zhang K, Shen H, Yao X, Sun Q and Chen G (2018) Necroptosis: A novel manner of cell death, associated with stroke (Review). *Int J Mol Med* 41, 624-630
36. Benedict CA, Norris PS, Prigozy TI et al (2001) Three adenovirus E3 proteins cooperate to evade apoptosis by tumor necrosis factor-related apoptosis-inducing ligand receptor-1 and -2. *J Biol Chem* 276, 3270-3278
37. Nichols DB, De Martini W and Cottrell J (2017) Poxviruses Utilize Multiple Strategies to Inhibit Apoptosis. *Viruses* 9
38. Jerome KR, Fox R, Chen Z, Sears AE, Lee H and Corey L (1999) Herpes simplex virus inhibits apoptosis through the action of two genes, Us5 and Us3. *J Virol* 73, 8950-8957

39. Lu JV, Weist BM, van Raam BJ et al (2011) Complementary roles of Fas-associated death domain (FADD) and receptor interacting protein kinase-3 (RIPK3) in T-cell homeostasis and antiviral immunity. *Proc Natl Acad Sci U S A* 108, 15312-15317
40. Wang YQ, Wang L, Zhang MY et al (2012) Necrostatin-1 suppresses autophagy and apoptosis in mice traumatic brain injury model. *Neurochem Res* 37, 1849-1858
41. Xu X, Chua CC, Kong J et al (2007) Necrostatin-1 protects against glutamate-induced glutathione depletion and caspase-independent cell death in HT-22 cells. *J Neurochem* 103, 2004-2014
42. Zhu S, Zhang Y, Bai G and Li H (2011) Necrostatin-1 ameliorates symptoms in R6/2 transgenic mouse model of Huntington's disease. *Cell Death Dis* 2, e115
43. Re DB, Le Verche V, Yu C et al (2014) Necroptosis drives motor neuron death in models of both sporadic and familial ALS. *Neuron* 81, 1001-1008
44. Murakami Y, Matsumoto H, Roh M et al (2014) Programmed necrosis, not apoptosis, is a key mediator of cell loss and DAMP-mediated inflammation in dsRNA-induced retinal degeneration. *Cell Death Differ* 21, 270-277
45. Veyer DL, Carrara G, Maluquer de Motes C and Smith GL (2017) Vaccinia virus evasion of regulated cell death. *Immunol Lett* 186, 68-80
46. Robinson N, McComb S, Mulligan R, Dudani R, Krishnan L and Sad S (2012) Type I interferon induces necroptosis in macrophages during infection with *Salmonella enterica* serovar Typhimurium. *Nat Immunol* 13, 954-962
47. Sridharan H and Upton JW (2014) Programmed necrosis in microbial pathogenesis. *Trends Microbiol* 22, 199-207
48. Kaminsky V and Zhivotovsky B (2010) To kill or be killed: how viruses interact with the cell death machinery. *J Intern Med* 267, 473-482
49. Lin J, Li H, Yang M et al (2013) A role of RIP3-mediated macrophage necrosis in atherosclerosis development. *Cell Rep* 3, 200-210
50. Han W, Li L, Qiu S et al (2007) Shikonin circumvents cancer drug resistance by induction of a necroptotic death. *Mol Cancer Ther* 6, 1641-1649
51. Alameda JP, Moreno-Maldonado R, Navarro M et al (2010) An inactivating CYLD mutation promotes skin tumor progression by conferring enhanced proliferative, survival and angiogenic properties to epidermal cancer cells. *Oncogene* 29, 6522-6532
52. Yang Y, Hu W, Feng S, Ma J and Wu M (2005) RIP3 beta and RIP3 gamma, two novel splice variants of receptor-interacting protein 3 (RIP3), downregulate RIP3-induced apoptosis. *Biochem Biophys Res Commun* 332, 181-187
53. Wu W, Liu P and Li J (2012) Necroptosis: an emerging form of programmed cell death. *Crit Rev Oncol Hematol* 82, 249-258
54. Oliver Metzger M, Fuchs D, Tagscherer KE, Grone HJ, Schirmacher P and Roth W (2016) Inhibition of caspases primes colon cancer cells for 5-fluorouracil-induced TNF-alpha-dependent necroptosis driven by RIP1 kinase and NF-kappaB. *Oncogene* 35, 3399-3409
55. Najafov A, Chen H and Yuan J (2017) Necroptosis and Cancer. *Trends Cancer* 3, 294-301
56. Wu W, Zhu H, Fu Y et al (2014) Clinical significance of down-regulated cylindromatosis gene in chronic lymphocytic leukemia. *Leuk Lymphoma* 55, 588-594
57. Feng X, Song Q, Yu A, Tang H, Peng Z and Wang X (2015) Receptor-interacting protein kinase 3 is a predictor of survival and plays a tumor suppressive role in colorectal cancer. *Neoplasma* 62, 592-601
58. Ruan J, Mei L, Zhu Q, Shi G and Wang H (2015) Mixed lineage kinase domain-like protein is a prognostic biomarker for cervical squamous cell cancer. *Int J Clin Exp Pathol* 8, 15035-15038