

FIXED POINT THEOREMS IN FUZZY METRIC SPACES FOR MAPPINGS WITH SOME CONTRACTIVE TYPE CONDITIONS

BIJOY PATIR, NILAKSHI GOSWAMI, AND LAKSHMI NARAYAN MISHRA*

ABSTRACT. In this paper, we derive some fixed point theorems in fuzzy metric spaces for self mappings satisfying different contractive type conditions. Some of these theorems generalize some results of Wairojjana et al. (*Fixed Point Theory and Applications* (2015) 2015:69). Several examples in support of the theorems are also presented here.

1. Introduction and preliminaries

The concept of fuzzy metric spaces was introduced by Kramosil and Michalek [23] in 1975. The results of Banach and Edelstein [3], [14] plays an effective role in several fixed point theorems. The study of fixed point theory in fuzzy metric spaces was started with the work of Grabiec in 1988 [18], by extending the existing results to fuzzy metric spaces. Further, a new notion of complete fuzzy metric space was introduced by George and Veeramani [16], [17] by defining a Hausdorff and first countable topology for some specific fuzzy metric space. In this paper, we follow this definition of completeness.

The concept of altering distance was earlier introduced by M.S Khan et al. [22] in 1984, with a view to enhance the metric fixed point theory.

Received February 21, 2018. Revised June 5, 2018. Accepted June 10, 2018.

2010 Mathematics Subject Classification: 54H25, 47H10.

Key words and phrases: fuzzy metric space, altering distance, t-norm, fixed point.

* Corresponding author.

© The Kangwon-Kyungki Mathematical Society, 2018.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

Recently many outstanding research workers have developed several interesting results on fixed point theory in relation with different operators (refer to [1], [2], [4]- [10], [12], [15], [19], [20], [21], [24], [26]- [33], [35]- [38]) as well as fuzzy fixed point theory (refer to [11], [13], [25], [34], [39], [40], [41],etc) with many applications in different branches of applied science. In this paper, our aim is to derive some fixed point theorems in fuzzy metric spaces using mappings with different contractive type conditions defined by altering distance function. Some well known results are also generalized here with suitable examples.

DEFINITION 1.1. [37] A mapping $T : [0, 1] \times [0, 1] \longrightarrow [0, 1]$ is called a triangular norm or t-norm if

- (i) $T(a, 1) = a$ for all $a \in [0, 1]$,
- (ii) $T(a, b) = T(b, a)$ for all $a, b \in [0, 1]$,
- (iii) $a \geq b, c \geq d \Rightarrow T(a, c) \geq T(b, d), \forall a, b, c, d \in [0, 1]$,
- (iv) $T(a, T(b, c)) = T(T(a, b), c), \forall a, b, c \in [0, 1]$.

$T_m(a, b) = \min(a, b)$, $T_p(a, b) = a.b$, $T_L(a, b) = \max(a + b - 1, 0)$ are some commonly used t-norms.

DEFINITION 1.2. [16] For an arbitrary set X , let T be a continuous t-norm and M be a fuzzy set on $X^2 \times (0, \infty)$. The 3-tuple (X, M, T) is called a fuzzy metric space if :

- (i) $M(a, b, s) > 0$, for all $a, b \in X, s > 0$,
- (ii) $M(a, b, s) = 1$, for all $s > 0 \Leftrightarrow a = b$,
- (iii) $M(a, b, s) = M(b, a, s)$, for all $a, b \in X, s > 0$,
- (iv) $T(M(a, b, s), M(b, c, p)) \leq M(a, c, s + p)$,
for all $a, b, c \in X, s, p > 0$,
- (v) $M(a, b, \cdot) : (0, \infty) \longrightarrow [0, 1]$ is continuous for all $a, b \in X$.

EXAMPLE 1.3. For $X = \mathbb{R}$, taking the usual metric $d(x, y) = |x - y|$ and the fuzzy t-norm $M(x, y, t) = \frac{t}{t+d(x,y)}, t \in (0, \infty)$, we have, (X, M, T) is a fuzzy metric space with respect to the t-norm $T_p(x, y) = x.y$, $x, y \in X$

LEMMA 1.4. [18] *With respect to t , $M(x, y, t)$ is a non-decreasing function for all x, y in X ,*

DEFINITION 1.5. [12] [16] [18] For a fuzzy metric space (X, M, T) ,

- (i) A sequence $\{a_p\}$ in X is said to be a Cauchy sequence if for any $\epsilon \in (0, 1), t > 0$, there exists a positive integer k satisfying $M(a_p, a_q, t) >$

- $1 - \epsilon, \forall p, q \geq k$
- (ii) A sequence $\{a_p\}$ in X converges to a if for any $\epsilon \in (0, 1), t > 0$, there exists a positive integer k satisfying $M(a_p, a, t) > 1 - \epsilon, \forall p \geq k$.
 - (iii) X is said to be complete if and only if every Cauchy sequence converges in X .
 - (iv) X is called compact if and only if every sequence in X has a convergent subsequence.

DEFINITION 1.6. [38] Let $\phi : [0, \infty[\rightarrow [0, \infty[$ be a mapping. If

- (i) ϕ is strictly decreasing and left continuous,
 - (ii) $\phi(\lambda) = 0$ if and only if $\lambda = 1$
- i.e, $\lim_{\phi \rightarrow 1^-} \phi(1) = 0$.

then, the function ϕ is called an altering distance function.

In [38], Shen et.al, proved the following fixed point theorem in complete fuzzy metric spaces.

THEOREM 1.7. [38] For a complete fuzzy metric space (X, M, T) , let f be a self map on X . Let ϕ be the altering distance function and k be a function from $(0, \infty)$ into $(0, 1)$. If for any $t > 0$, f satisfies the following condition:

$$(1) \quad \phi(M(f(x), f(y), t)) \leq k(t) \cdot \phi(M(x, y, t))$$

$x, y \in X$ and $x \neq y$, then there exists a unique fixed point for f in X .

The following theorem by Wairojjana et al. comes as a generalization of the Theorem 1.7

THEOREM 1.8. [41] For a complete fuzzy metric space (X, M, T) , let f be a self map on X . If there exist functions $k_1, k_2 : (0, \infty) \rightarrow [0, 1)$, $k_3 : (0, \infty) \rightarrow (0, 1)$, $\sum_{i=1}^3 k_i(t) < 1$, and an altering distance function ϕ such that the following condition is satisfied:

$$(2) \quad \begin{aligned} \phi(M(f(x), f(y), t)) &\leq k_1(t) \cdot \phi(M(x, f(x), t)) + k_2(t) \cdot \phi(M(y, f(y), t)) \\ &\quad + k_3(t) \cdot \phi(M(x, y, t)) \end{aligned}$$

$x, y \in X, x \neq y$, and $t > 0$, then there exists a unique fixed point for f .

Taking a minimum condition, Wairojjana et al., also proved the following result.

THEOREM 1.9. *For a complete fuzzy metric space (X, M, T) , let T be a triangular norm and f be a self map on X . If there exist $k_1, k_2 : (0, \infty) \rightarrow (0, 1)$, and an altering distance function ϕ such that the following condition holds:*

$$(3) \quad \begin{aligned} & \phi(M(f(x), f(y), t)) \\ & \leq k_1(t) \cdot \min\{\phi(M(x, y, t)), \phi(M(x, f(x), t)), \phi(M(x, f(y), 2t)), \\ & \phi(M(y, f(y), t))\} + k_2(t)\phi(M(f(x), y, 2t)) \end{aligned}$$

$x, y \in X$, $x \neq y$ and $t > 0$, then there exists a unique point for f .

2. Results and discussion

Considering the function $k : (0, \infty) \rightarrow (0, 1]$, in Theorem 1.8, we take the case when $k(t) = 1$. Using a condition as in [38], we proved the following:

THEOREM 2.1. *Let (X, M, T) be a complete fuzzy metric space and ϕ be the altering distance function. Let f be a self-map on X satisfying*

$$(4) \quad \phi(M(f(x), f(y), t)) \leq \phi(M(x, y, t)), \forall x, y \in X, t > 0.$$

For $x_0 \in X$, let the sequence $\{x_n\}$ in X is constructed by: $x_{n+1} = f(x_n), n = 0, 1, 2, \dots$

Then $x_n \rightarrow p$ in X , p being the unique fixed point of f if and only if for any two sub sequences $\{x_{p_n}\}$ and $\{x_{q_n}\}$ with $x_{p_n} \neq x_{q_n} \forall n$, we have that,

$$(5) \quad \begin{aligned} Q_n &= \frac{\phi(M(f(x_{p_n}), f(x_{q_n}), t))}{\phi(M(x_{p_n}, x_{q_n}, t))} \rightarrow 1 \\ &\text{only if } \phi(M(x_{p_n}, x_{q_n}, t)) \rightarrow 0 \text{ as } n \rightarrow \infty \end{aligned}$$

Proof. First we assume that the sequence $\{x_n\}$ converges to the unique fixed point p of f . Let $\{x_{p_n}\}$ and $\{x_{q_n}\}$ be any two sub sequences of $\{x_n\}$.

Then

$$\begin{aligned} \lim_{n \rightarrow \infty} \phi(M(x_{p_n}, x_{q_n}, t)) &= \phi(M(p, p, t)) = \phi(1) = 0 \\ \text{and } Q_n &= \frac{\phi(M(f(x_{p_n}), f(x_{q_n}), t))}{\phi(M(x_{p_n}, x_{q_n}, t))} \\ &\rightarrow \frac{\phi(M(f(p), f(p), t))}{\phi(M(p, p, t))} = \frac{\phi(M(p, p, t))}{\phi(M(p, p, t))} \quad (\text{since } f(p) = p) \\ &= 1 \text{ as } n \rightarrow \infty \end{aligned}$$

So, the condition (5) is satisfied.

Conversely, for $x_0 \in X$, $f(x_n) = x_{n+1}$, $n = 0, 1, 2, \dots$

Let $\phi_n = \phi(M(x_n, x_{n+1}, t))$, $n = 0, 1, 2, \dots$. Then,

$$\begin{aligned} \phi_{n+1} &= \phi(M(f(x_n), f(x_{n+1}), t)) \\ &\leq \phi(M(x_n, x_{n+1}, t)) \quad (\text{by(4)}) \\ &= \phi_n \end{aligned}$$

Thus $\{\phi_n\}$ is a decreasing sequence of non negative real numbers and so, is convergent to some real, say $\epsilon \geq 0$.

Assume that $\epsilon > 0$. Taking $p_n = n$ and $q_n = n + 1$,

$$\begin{aligned} &\phi(M(x_{p_n}, x_{q_n}, t)) \\ &= \phi(M(x_n, x_{n+1}, t)) = \phi_n \rightarrow \epsilon > 0. \\ \text{But } Q_n &= \frac{\phi(M(f(x_n), f(x_{n+1}), t))}{\phi(M(x_n, x_{n+1}, t))} \\ &= \frac{\phi(M(x_{n+1}, x_{n+2}, t))}{\phi(M(x_n, x_{n+1}, t))} \rightarrow 1 \text{ as } n \rightarrow \infty, \end{aligned}$$

violating the given condition (5).

Hence $\epsilon = 0$, i.e., $\phi_n \rightarrow 0$

$$(6) \quad \text{i.e., } M(x_n, x_{n+1}, t) \rightarrow 1 \text{ as } n \rightarrow \infty.$$

Now, to show that the sequence $\{x_n\}$ is Cauchy :

If not, $\exists 0 < \epsilon < 1$ and two increasing sequences of integers $\{a_n\}$ and $\{b_n\}$ such that for every $n \in \mathbb{N} \cup \{0\}$, we have $a_n > b_n > n$

$$\begin{aligned} M(x_{a_n}, x_{b_n}, t) &\leq 1 - \epsilon \text{ and} \\ M(x_{a_{n-1}}, x_{b_n}, t) &> 1 - \epsilon, \quad t > 0. \end{aligned}$$

Let $S_n(t) = M(x_{a_n}, x_{b_n}, t)$ for each $n \in \mathbb{N} \cup \{0\}$.

$$\begin{aligned} \text{Then } 1 - \epsilon \geq S_n(t) &\geq T(M(x_{a_n}, x_{a_{n-1}}, t/2), M(x_{a_{n-1}}, x_{b_n}, t/2)) \\ &\geq T(M(x_{a_n}, x_{a_{n-1}}, t/2), 1 - \epsilon) \\ &\longrightarrow T(1, 1 - \epsilon) \quad (\text{by(6)}) \\ &= 1 - \epsilon \text{ as } n \longrightarrow \infty \end{aligned}$$

Thus, $S_n(t) \longrightarrow 1 - \epsilon$ as $n \longrightarrow \infty$

Therefore, $\phi(S_n(t)) = \phi(M(x_{a_n}, x_{b_n}, t)) \longrightarrow \phi(1 - \epsilon)$ as $n \longrightarrow \infty$

$$\text{But } Q_n = \frac{\phi(M(x_{a_{n+1}}, x_{b_{n+1}}, t))}{\phi(M(x_{a_n}, x_{b_n}, t))} \longrightarrow 1 \text{ as } n \longrightarrow \infty,$$

which again violates the condition (5).

Therefore, $\{x_n\}$ is a Cauchy sequence in X . Since (X, M, T) is complete, so, $\exists p \in X$ such that $x_n \longrightarrow p$.

$$\begin{aligned} \text{Now, } 0 \leq \phi(M(f(x_n), f(p), t)) &= \phi(M(x_{n+1}, f(p), t)) \\ &\leq \phi(M(x_n, p, t)) \end{aligned}$$

Taking $n \longrightarrow \infty$,

$$\begin{aligned} \phi(M(p, f(p), t)) &\leq \phi(M(p, p, t)) = \phi(1) = 0 \\ &\Rightarrow \phi(M(p, f(p), t)) = 0 \\ &\Rightarrow M(p, f(p), t) = 1 \\ &\Rightarrow f(p) = p, \end{aligned}$$

showing that p is a fixed point for f .

To show the uniqueness :

If p_1 is another fixed point for f ,

$$\phi(M(p, p_1, t)) = \phi(M(f(p), f(p_1), t)) < \phi(M(p, p_1, t))$$

a contradiction. So, p is the unique fixed point for f . □

EXAMPLE 2.2. Let $X = [0, 1]$ and $d(x, y) = |x - y| \forall x, y \in X$. Then (X, d) is a complete metric space. Let M be a fuzzy set on $X^2 \times [0, \infty[$ given by $M(x, y, t) = \frac{t}{t + d(x, y)}$, if $t > 0$ and $M(x, y, 0) = 0$.

Then (X, M, T) is a complete fuzzy metric space with respect to the t-norm : $T(a, b) = \min\{a, b\}$, $a, b \in [0, 1]$. We take $f : X \longrightarrow X$ given

by $f(x) = \frac{x}{7} \forall x \in X$ and $\phi(\lambda) = 1 - \lambda$.

$$\begin{aligned} \text{Now, } M(f(x), f(y), t) &= \frac{t}{t + \frac{1}{7}|x - y|} \\ &= M(x, y, 7t), \quad t > 0 \\ &\geq M(x, y, t) \\ &\text{(as } M(x, y, \cdot) \text{ is non-decreasing)} \end{aligned}$$

So, $\phi(M(f(x), f(y), t)) \leq \phi(M(x, y, t))$

For $x_0 \in [0, 1]$, we consider the sequence $\{\frac{x_0}{7^n}\}$ which is obtained by: $x_{n+1} = f(x_n), n = 0, 1, 2, 3, \dots$. Now, for any two sub sequences of $\{\frac{x_0}{7^n}\}$, the condition (5) is obviously satisfied. Hence f has a unique fixed point which is clearly 0, here.

Omitting the completeness condition in Theorem 2.1, we get

THEOREM 2.3. *Let (X, M, T) be a fuzzy metric space and ϕ be the altering distance function. Let f be a self map on X satisfying*

- (i) $\phi(M(f(x), f(y), t)) \leq k_1(x, y, t)\phi(M(x, f(x), t)) + k_2(x, y, t)\phi(M(y, f(y), t)) + k_3(x, y, t)\phi(M(x, y, t))$, where k_1, k_2, k_3 are functions from $X^2 \times (0, \infty)$ to $(0, 1)$ such that $0 < k_1 + k_2 + k_3 < 1, k_i = k_i(x, y, t)$.
- (ii) f is continuous at some point $p \in X$
- (iii) \exists a point $z \in X$ such that the sequence of iterates $\{f^n(z)\}$ has a subsequence $\{f^{n_k}(z)\}$ converging to p .

Then p is the unique fixed point for f .

Proof. Let $x_0 = z \in X$. we define

$$x_{n+1} = f(x_n), \quad n = 0, 1, 2, \dots$$

If $f^n(x_0) = f^{n+1}(x_0)$ for some n , then there exists $y \in X$ such that $y = f^k(x_0)$ for all $k \geq n$. [In that case, y is the unique fixed point.]

So, we assume that $f^n(x_0) \neq f^{n+1}(x_0)$, for any n .

By (iii), $\lim_{k \rightarrow \infty} f^{n_k}(z) = p$, i.e, $\lim_{k \rightarrow \infty} x_{n_k} = p$.

Since, f is continuous at p , (by(ii)), so,

$$\begin{aligned} f(p) &= \lim_{k \rightarrow \infty} f^{n_k+1}(z) = \lim_{k \rightarrow \infty} f^{n_k+1}(x_0) \\ &= \lim_{k \rightarrow \infty} x_{n_k+1} \end{aligned}$$

By (i),

$$\begin{aligned}
\phi(M(f(x_{n-1}), f(x_n), t)) &\leq k_1\phi(M(x_{n-1}, f(x_{n-1}), t)) \\
&\quad + k_2\phi(M(x_n, f(x_n), t)) + k_3\phi(M(x_{n-1}, x_n, t)) \\
\Rightarrow \phi(M(x_n, x_{n+1}, t)) &\leq k_1\phi(M(x_{n-1}, x_n, t)) \\
&\quad + k_2\phi(M(x_n, x_{n+1}, t)) + k_3\phi(M(x_{n-1}, x_n, t)) \\
\Rightarrow \phi(M(x_n, x_{n+1}, t)) &\leq \left(\frac{k_1 + k_3}{1 - k_2}\right)\phi(M(x_{n-1}, x_n, t)) \\
(7) \quad \Rightarrow \phi(M(x_n, x_{n+1}, t)) &< \phi(M(x_{n-1}, x_n, t))
\end{aligned}$$

So, $\phi(M(f(x_{n_k+1}), f(f(p)), t)) < \phi(M(x_{n_k+1}, f(p), t))$

Since, ϕ is left continuous and M is continuous,

$$\begin{aligned}
\phi(M(\lim_{k \rightarrow \infty} x_{n_k+2}, f^2(p), t)) &\leq \phi(M(\lim_{k \rightarrow \infty} x_{n_k+1}, f(p), t)) \\
&= \phi(M(f(p), f(p), t)) \\
&= 0 \\
(8) \quad \Rightarrow \lim_{k \rightarrow \infty} x_{n_k+2} &= f^2(p)
\end{aligned}$$

Again,

$$\begin{aligned}
\phi(M(p, f(p), t)) &= \lim_{k \rightarrow \infty} \phi(M(x_{n_k}, x_{n_k+1}, t)) \\
&= \lim_{k \rightarrow \infty} \phi(M(x_{n_k+1}, x_{n_k+2}, t)) \\
(9) \quad &= \phi(M(f(p), f^2(p), t)) \text{ (by(8))}
\end{aligned}$$

If $f(p) \neq p$, then from (7),

$$\phi(M(f(p), f^2(p), t)) < \phi(M(p, f(p), t)),$$

a contradiction to (9). Therefore, $f(p) = p$, showing that p is a fixed point for f .

Uniqueness follows easily from (7). \square

EXAMPLE 2.4. Let $X = [0, 1]$ and $d(x, y) = |e^x - e^y|$ for every $x, y \in X$. Then (X, d) is an incomplete metric space. Let M be a fuzzy set on $X^2 \times [0, \infty[$ given by $M(x, y, t) = \frac{t}{t + d(x, y)}$, $t > 0$ and $M(x, y, 0) = 0$.

Let, $T(x, y) = xy$. Then (X, M, T) is an incomplete fuzzy metric space. We take $\phi(\lambda) = 1 - \lambda$, $\lambda \in [0, 1]$ and $f : X \rightarrow X$ is defined by $f(x) = \frac{2+x}{3}$.

Clearly, f is continuous at the point $1 \in X$.

Let $z = x_0 \in X$ and we consider the sequence of iterates $\{f^n(x_0)\}$ having a subsequence $\{f^{n_k}(x_0)\}$, where $n_k = 2k - 1, k \in N$

Now, $\lim_{k \rightarrow \infty} f^{n_k}(x_0) = 1$

We take,
$$k_1(x, y, t) = \begin{cases} \frac{1 + \frac{t}{e^{\frac{2+x}{3}} + e^x}}{1 + \frac{t}{e^{\frac{2+x}{3}} - e^{\frac{2+y}{3}}}}, & \text{if } x > y \\ \frac{1 + \frac{t}{e^{\frac{2+x}{3}} + e^x}}{1 + \frac{t}{e^{\frac{2+y}{3}} - e^{\frac{2+x}{3}}}}, & \text{if } y > x \end{cases}$$

and $k_2(x, y, t) = k_3(x, y, t) = \frac{1 - k_1(x, y, t)}{3}$

Again,
$$\phi(M(f(x), f(y), t)) = \begin{cases} \frac{e^{\frac{2+x}{3}} - e^{\frac{2+y}{3}}}{t + (e^{\frac{2+x}{3}} - e^{\frac{2+y}{3}})}, & \text{if } x > y \\ \frac{e^{\frac{2+y}{3}} - e^{\frac{2+x}{3}}}{t + (e^{\frac{2+y}{3}} - e^{\frac{2+x}{3}})}, & \text{if } y > x \end{cases}$$

$$\phi(M(x, f(x), t)) = \frac{e^{\frac{2+x}{3}} - e^x}{t + e^{\frac{2+x}{3}} - e^x}$$

Similarly,

$$\phi(M(y, f(y), t)) = \frac{e^{\frac{2+y}{3}} - e^y}{t + e^{\frac{2+y}{3}} - e^y}$$

and
$$\phi(M(x, y, t)) = \begin{cases} \frac{e^x - e^y}{t + e^x - e^y}, & \text{for } x > y \\ \frac{e^y - e^x}{t + e^y - e^x}, & \text{for } y > x \end{cases}$$

Now,
$$\begin{aligned} & k_1(x, y, t)\phi(M(x, f(x), t)) + k_2(x, y, t)\phi(M(y, f(y), t)) \\ & + k_3(x, y, t)\phi(M(x, y, t)) \\ & \geq \phi(M(f(x), f(y), t)) \end{aligned}$$

So, all the conditions of Theorem 2.3 are satisfied and f has a unique fixed point. Clearly, 1 is the unique fixed point for f here.

REMARK 2.5. Introducing another function $k_4 : X^2 \times (0, \infty) \rightarrow (0, 1)$, the condition (i) in the Theorem 2.3 can be replaced by:

$$\begin{aligned} \phi(M(f(x), f(y), t)) &\leq k_1\phi(M(x, f(x), t)) + k_2\phi(M(y, f(y), t)) \\ &+ k_3\phi(M(x, y, t)) + k_4 \frac{\phi(M(y, f(y), t)) (1 + \phi(M(x, f(x), t)))}{1 + \phi(M(x, y, t))} \\ &\text{with } 0 \leq k_1 + k_2 + k_3 + k_4 < 1 \end{aligned}$$

In this case,

$$\begin{aligned} \phi(M(f(x_{n-1}), f(x_n), t)) &\leq k_1\phi(M(x_{n-1}, f(x_{n-1}), t)) \\ &+ k_2\phi(M(x_n, f(x_n), t)) + k_3\phi(M(x_{n-1}, x_n, t)) \\ &\quad + k_4 \frac{\phi(M(x_n, f(x_n), t)) (1 + \phi(M(x_{n-1}, f(x_{n-1}), t)))}{1 + \phi(M(x_{n-1}, x_n, t))} \\ &= k_1\phi(M(x_{n-1}, x_n, t)) + k_2\phi(M(x_n, x_{n+1}, t)) \\ &\quad + k_3\phi(M(x_{n-1}, x_n, t)) + k_4 \frac{\phi(M(x_n, x_{n+1}, t)) (1 + \phi(M(x_{n-1}, x_n, t)))}{1 + \phi(M(x_{n-1}, x_n, t))} \\ &= (k_1 + k_3)\phi(M(x_{n-1}, x_n, t)) + (k_2 + k_4)\phi(M(x_n, x_{n+1}, t)) \\ &\Rightarrow \phi(M(x_n, x_{n+1}, t)) \leq \frac{k_1 + k_3}{1 - (k_2 + k_4)} \phi(M(x_n, x_{n+1}, t)) \\ &\quad < \phi(M(x_n, x_{n+1}, t)) \end{aligned}$$

The other part is same as in Theorem 2.3.

Similar example as 2.4 can be constructed for this also.

In the following theorem, we use the condition $\lim_{t \rightarrow \infty} M(x, y, t) = 1$.

THEOREM 2.6. Let (X, M, T) be a complete fuzzy metric space and ϕ be the altering distance function. Let f be a self map on X satisfying the following: for every positive integer n and $t > 0$,

$$(10) \quad \phi(M(f^n(x), f^n(y), k_n t)) \leq \phi(M(x, y, t)),$$

for all $x, y \in X$, $k_n > 0$ being independent of x and y . If $k_n \rightarrow 0$ as $n \rightarrow \infty$ then f has a unique fixed point in X .

Proof. Let $x_0 \in X$. We construct a sequence $\{x_n\}$ in X by $x_1 = f(x_0)$, $x_2 = f(x_1) = f^2(x_0)$, \dots , $x_n = f(x_{n-1}) = f^n(x_0)$, \dots , $n \in \mathbb{N}$
From (10),

$$\phi(M(f^n(x), f^n(y), k_n t)) \leq \phi(M(x, y, t))$$

Since, ϕ is strictly decreasing, so,

$$(11) \quad M(f^n(x), f^n(y), k_n t) \geq M(x, y, t)$$

Again, for $p \in \mathbb{N}$,

$$\begin{aligned} 1 &\geq M(x_n, x_{n+p}, t) \\ &\geq T(T(\dots T(M(x_n, x_{n+1}, t/p), M(x_{n+1}, x_{n+2}, t/p)\dots), M(x_{n+p-1}, x_{n+p}, t/p)) \\ &\geq T(T(\dots T(M(x, x_1, \frac{t}{pk_n}), M(x, x_1, \frac{t}{pk_{n+1}}), \dots), M(x, x_1, \frac{t}{pk_{n+p-1}}))) \text{ (by (11))} \\ &\quad \longrightarrow T(T(\dots T(1, 1), \dots), 1) \text{ as } n \longrightarrow \infty \\ &\quad \text{(Since } k_n \longrightarrow 0 \text{ as } n \longrightarrow \infty \text{ and } \lim_{t \longrightarrow \infty} M(x, y, t) = 1) \\ &= 1 \end{aligned}$$

Thus, $\lim_{n \rightarrow \infty} M(x_n, x_{n+p}, t) = 1 \forall t > 0$

So, $\{x_n\}$ is a Cauchy sequence in X .
As X is complete, $\exists p \in X$ such that $x_n \rightarrow p$ as $n \rightarrow \infty$.
Now,

$$\begin{aligned} 0 &\leq \phi(M(x_{n+1}, f(p), t)) = \phi(M(f(x_n), f(p), t)) \\ &\leq \phi(M(x_n, p, t/k_1)) \text{ (by (10)) } \quad \forall t > 0 \end{aligned}$$

Taking $n \rightarrow \infty$, we have,

$$\begin{aligned} \phi(M(p, f(p), t)) &\leq \phi(M(p, p, t/k_1)) = 0 \\ \Rightarrow \phi(M(p, f(p), t)) &= 0 \quad \forall t > 0 \\ \Rightarrow p &= f(p), \end{aligned}$$

showing that p is a fixed point for f .

To show uniqueness:

Let $q \in X$ be such that $f(q) = q$. Now,

$$\begin{aligned} 0 &\leq \phi(M(p, q, t)) = \phi(M(f^n(p), f^n(q), t)) \\ &\leq \phi(M(p, q, t/k_n)) \longrightarrow 0 \text{ as } n \longrightarrow \infty \quad \forall t > 0 \\ \Rightarrow \phi(M(p, q, t)) &= 0 \quad \forall t > 0 \\ \Rightarrow p &= q. \end{aligned}$$

Thus, f has the unique fixed point p in X . □

EXAMPLE 2.7. We take the same complete fuzzy metric space (X, M, T) as in Example 2.2. Let f be a self map on X defined by : $f(x) =$

$\frac{3+x}{4}$, $x \in X$ and $\phi(\lambda) = 1 - \lambda$, $\lambda \in [0, 1]$. For $n \in \mathbb{N}$, let $k_n = \frac{1}{n}$. Then $k_n \rightarrow 0$ as $n \rightarrow \infty$.

For $x \neq y$,

$$\begin{aligned} \phi(M(f^n(x), f^n(y), k_n t)) &= \frac{\left|\frac{x-y}{4^n}\right|}{\frac{t}{n} + \left|\frac{x-y}{4^n}\right|} \\ &= \frac{|x-y|}{\frac{4^n}{n}t + |x-y|} < \frac{|x-y|}{t + |x-y|} = \phi(M(x, y, t)) \end{aligned}$$

For $x = y$, $\phi(M(f^n(x), f^n(y), k_n t)) = \phi(M(x, y, t))$.

Thus, the condition of Theorem 2.6 is satisfied and so, f has a unique fixed point.

Clearly, 1 is the unique fixed point for f in X .

COROLLARY 2.8. *Let (X, M, T) be a complete metric space, ϕ be the altering distance function and f be a self map on X satisfying*

$$\phi(M(f(x), f(y), kt)) \leq \phi(M(x, y, t)) \quad \forall x, y \in X, t > 0$$

and $0 < k < 1$.

Then, f has a unique fixed point in X .

Now, instead of taking the constants $k_n (n \in \mathbb{N})$ in Theorem 2.6, we take $k(t)$ as a function of t and prove the following:

THEOREM 2.9. *Let (X, M, T) be a complete fuzzy metric space, ϕ be the altering distance function and f be a self map on X . Let $k : (0, \infty) \rightarrow (0, \infty)$ be a bijective function such that for each $t \in (0, \infty)$, $\{k^n(t)\}$ is a decreasing sequence with $\lim_{n \rightarrow \infty} k^n(t) = 0$. If f satisfies:*

$$(12) \quad \phi(M(f^n(x), f^n(y), k^n(t))) \leq \phi(M(x, y, t))$$

for all $x, y \in X$ and $t > 0$, $n \in \mathbb{N}$, then f has a unique fixed point in X .

Proof. For $x_0 \in X$, we construct the sequence $\{x_n\}$ in X as in Theorem 2.6.

Now,

$$\begin{aligned} 0 &\leq \phi(M(x_n, x_{n+m}, t)) \\ &\leq \phi(M(x, x_m, k^{-n}t)) \quad (\text{Replacing } t \text{ by } k^{-n}t \text{ in (12)}) \\ &\quad \text{for each } m \in \mathbb{N}. \end{aligned}$$

Again, $\lim_{n \rightarrow \infty} \phi(M(x, x_m, k^{-n}(t))) = 0$ (Since ϕ is left continuous) $\forall t > 0$

$$\begin{aligned} \text{So, } \phi(M(x_n, x_{n+m}, t)) &\longrightarrow 0 \text{ as } n \longrightarrow \infty \\ \Rightarrow M(x_n, x_{n+m}, t) &\longrightarrow 1 \text{ as } n \longrightarrow \infty \quad \forall t > 0 \end{aligned}$$

Thus $\{x_n\}$ is a Cauchy sequence in X and so, converges to some $p \in X$.

$$\begin{aligned} \text{Now, } 0 \leq \phi(M(x_{n+1}, f(p), t)) &= \phi(M(f(x_n), f(p), t)) \\ &\leq \phi(M(x_n, p, k^{-1}(t))) \quad \forall t > 0 \end{aligned}$$

Taking $n \longrightarrow \infty$,

$$\begin{aligned} \phi(M(p, f(p), t)) &\leq \phi(M(p, p, k^{-1}(t))) = 0 \quad \forall t > 0 \\ \Rightarrow M(p, f(p), t) &= 1 \quad \forall t > 0 \\ \Rightarrow f(p) &= p. \end{aligned}$$

showing that p is a fixed point for f .

Uniqueness can be shown in a similar manner. □

EXAMPLE 2.10. For the same fuzzy metric space (X, M, T) and the same altering distance function as in Example 2.2, let $k : (0, \infty) \longrightarrow (0, \infty)$ be defined by: $k(t) = \frac{t}{2}$, $t > 0$. Then $k(t)$ is bijective. Also, for each t , $\{k^n(t)\}$ is a decreasing sequence and $\lim_{n \rightarrow \infty} k^n(t) = \lim_{n \rightarrow \infty} \frac{t}{2^n} = 0$.

We take $f : X \longrightarrow X$ as $f(x) = \frac{2+x}{3}$.

So, $f^n(x) = \frac{2}{3} + \frac{2}{3^2} + \dots + \frac{2}{3^n} + \frac{x}{3^n}$, and

$$\begin{aligned} \phi(M(f^n(x), f^n(y), k^n(t))) &= \frac{|x - y|}{\frac{3^n}{2^n}t + |x - y|} \\ &< \frac{|x - y|}{t + |x - y|} \\ &= \phi(M(x, y, t)) \quad \forall x, y \in X, t \in (0, \infty), n \in \mathbb{N} \end{aligned}$$

Thus, all the conditions of Theorem 2.9 are satisfied and so, f has a unique fixed point which is clearly 1 here.

The following theorem can be taken as a generalization of Theorem 1.9 by Wairojjana et al. [41] :

THEOREM 2.11. *Let (X, M, T) be a complete fuzzy metric space, f be a self map on X , and ϕ be the altering distance function. If there*

exist $k_1, k_2 : (0, \infty) \rightarrow (0, 1)$ such that $k_1(t) + k_2(t) < 1$, and

$$(13) \quad \begin{aligned} & \phi(M(f(x), f(y), t)) \\ & \leq k_1(t) \cdot \min\{\phi(M(x, y, t)), \phi(M(x, f(x), t)), \phi(M(x, f(y), 2t)), \\ & \quad \phi(M(y, f(y), t))\} + k_2(t) \cdot \max\{\phi(M(x, y, t)), \phi(M(x, f(x), t)), \\ & \quad \phi(M(f(x), y, 2t)), \phi(M(y, f(y), t))\}, \quad x, y \in X, \quad x \neq y, \quad t > 0, \end{aligned}$$

then f has a unique fixed point.

Proof. For $x_0 \in X$, we consider the sequence $\{x_n\}$ in X , where $x_{n+1} = f(x_n)$, $n = 0, 1, 2, \dots$

If there exists $n_0 \in \mathbb{N} \cup \{0\}$ such that $x_{n_0+1} = x_{n_0}$, then x_{n_0} will be a fixed point for f . So, we assume that $x_n \neq x_{n+1}$ for any n . Then $0 < M(x_n, x_{n+1}, t) < 1$, $t > 0$, $n \in \mathbb{N} \cup \{0\}$.

Substituting $x = x_{n-1}$, $y = x_n$ in (13), we have,

$$(14) \quad \begin{aligned} & \phi(M(x_n, x_{n+1}, t)) \\ & \leq k_1(t) \cdot \min\{\phi(M(x_{n-1}, x_n, t)), \phi(M(x_{n-1}, x_n, t)), \phi(M(x_{n-1}, x_{n+1}, 2t)), \\ & \quad \phi(M(x_n, x_{n+1}, t))\} + k_2(t) \cdot \max\{\phi(M(x_{n-1}, x_n, t)), \phi(M(x_{n-1}, x_n, t)), \\ & \quad \phi(M(x_n, x_n, 2t)), \phi(M(x_n, x_{n+1}, t))\} \\ & = k_1(t) \cdot \min\{\phi(M(x_{n-1}, x_n, t)), \phi(M(x_{n-1}, x_{n+1}, 2t)), \phi(M(x_n, x_{n+1}, t))\} \\ & \quad + k_2(t) \cdot \max\{\phi(M(x_{n-1}, x_n, t)), \phi(M(x_n, x_{n+1}, t))\} \\ & \leq k_1(t) \cdot \min\{\phi(M(x_{n-1}, x_n, t)), \phi(T(M(x_{n-1}, x_n, t)), (M(x_n, x_{n+1}, t))), \\ & \quad \phi(M(x_n, x_{n+1}, t))\} + k_2(t) \cdot \max\{\phi(M(x_{n-1}, x_n, t)), \\ & \quad \phi(M(x_n, x_{n+1}, t))\} \end{aligned}$$

Again,

$$\phi(a) \leq \phi(T(a, b)) \text{ and } \phi(b) \leq \phi(T(a, b))$$

$$(\text{Since, } a = T(a, 1) \geq T(a, b) \text{ and } b = T(b, 1) \geq T(a, b))$$

$$\text{So, } \min\{\phi(a), \phi(b), \phi(T(a, b))\} = \min\{\phi(a), \phi(b)\} \quad \forall a, b \in [0, 1].$$

Hence, (14) $\Rightarrow \phi(M(x_n, x_{n+1}, t))$

$$\begin{aligned} & \leq k_1(t) \cdot \min\{\phi(M(x_{n-1}, x_n, t)), \phi(M(x_n, x_{n+1}, t))\} \\ & \quad + k_2(t) \cdot \max\{\phi(M(x_{n-1}, x_n, t)), \phi(M(x_n, x_{n+1}, t))\} \end{aligned}$$

If $\min\{\phi(M(x_{n-1}, x_n, t)), \phi(M(x_n, x_{n+1}, t))\} = \phi(M(x_{n-1}, x_n, t))$, then,

$$\begin{aligned} \phi(M(x_n, x_{n+1}, t)) &\leq k_1(t)\phi(M(x_{n-1}, x_n, t)) + k_2(t)\phi(M(x_n, x_{n+1}, t)) \\ \Rightarrow \phi(M(x_n, x_{n+1}, t)) &\leq \frac{k_1(t)}{1 - k_2(t)}\phi(M(x_{n-1}, x_n, t)) \\ &< \phi(M(x_{n-1}, x_n, t)), \quad \text{a contradiction.} \end{aligned}$$

So, $\min\{\phi(M(x_{n-1}, x_n, t)), \phi(M(x_n, x_{n+1}, t))\} = \phi(M(x_n, x_{n+1}, t))$.

$$\begin{aligned} \text{Then, } \phi(M(x_n, x_{n+1}, t)) &\leq \frac{k_2(t)}{1 - k_1(t)}\phi(M(x_{n-1}, x_n, t)) \\ &< \phi(M(x_{n-1}, x_n, t)) \end{aligned}$$

Thus, $\phi(M(x_n, x_{n+1}, t)) < \phi(M(x_{n-1}, x_n, t))$.

Since ϕ is strictly decreasing, so, it follows that $\{M(x_n, x_{n+1}, t)\}_n$ is strictly increasing for every $t > 0$.

Now, proceeding as in Theorem 2.1 together with the given condition(13), it can be easily shown that f has a unique fixed point in X . \square

EXAMPLE 2.12. Let $X = \{P, Q, R, S\} \subseteq \mathbb{R}^2$, where P, Q and R are the points $(1, 1), (2, 1)$ and $(2, 5)$ respectively and S is an arbitrary point on the unit circle with centre at R . Let $f : X \rightarrow X$ be defined by: $f(P) = f(Q) = f(S) = P, f(R) = Q$.

Let $M(x, y, t) = \frac{t}{t+d(x,y)}, t > 0$, where $d(x, y)$ is the Euclidean distance in \mathbb{R}^2 . The altering distance ϕ is given by $\phi(\lambda) = 1 - \lambda, \lambda \in [0, 1]$. With $T(a, b) = \min\{a, b\}$, (X, M, T) becomes a complete fuzzy metric space. Let $k_1, k_2 : (0, \infty) \rightarrow (0, 1)$ be defined by:

$$k_2(t) = \begin{cases} \frac{1}{1+\frac{t}{7}}, & t \in (0, 1] \\ \frac{t}{t+\frac{1}{7}}, & t \in (1, \infty) \end{cases}$$

and $k_1(t) = (1 - k_2(t))/2$

Then, $k_1(t) + k_2(t) < 1$.

Now, it can be shown that every pair of points $(P, Q), (P, S), (Q, S), (P, R), (Q, R)$ and (R, S) satisfies the condition (13).

Hence by Theorem 2.11 the function has a unique fixed point.

Next, we obtain the following generalization of Theorem 1.9 :

THEOREM 2.13. Let (X, M, T) be a complete fuzzy metric space, f be a self map on X , and ϕ be the altering distance function. If there

exist functions $k_1, k_2, k_3, k_4 : (0, \infty) \longrightarrow (0, 1)$ with $k_1(t) + k_2(t) + k_3(t) + k_4(t) < 1$, such that

$$\begin{aligned}
 (15) \quad & \phi(M(f(x), f(y), t)) \\
 & \leq k_1(t) \cdot \min\{\phi(M(x, f(x), t)), \phi(M(x, y, t))\} \\
 & \quad + k_2(t) \cdot \min\{\phi(M(y, f(y), t)), \phi(M(x, f(x), t))\} \\
 & \quad + k_3(t) \cdot \min\{\phi(M(x, y, t)), \phi(M(x, f(y), 2t)), \phi(M(y, f(y), t))\} \\
 & \quad + k_4(t)\phi(M(f(x), y, 2t)), \quad x, y \in X, \quad x \neq y, \quad t > 0
 \end{aligned}$$

then f has a unique fixed point.

Proof. For $x_0 \in X$, we take the sequence $\{x_n\}$ with $x_{n+1} = f(x_n)$ as earlier. Taking $x_n \neq x_{n+1}$, for any n i.e., $0 < M(x_n, x_{n+1}, t) < 1, n \in \mathbb{N} \cup \{0\}, t > 0$, we have,

$$\begin{aligned}
 & \phi(M(x_n, x_{n+1}, t)) \\
 \leq & k_1(t) \cdot \min\{\phi(M(x_{n-1}, x_n, t)), \phi(M(x_{n-1}, x_n, t))\} \\
 & + k_2(t) \cdot \min\{\phi(M(x_n, x_{n+1}, t)), \phi(M(x_{n-1}, x_n, t))\} \\
 & + k_3(t) \cdot \min\{\phi(M(x_{n-1}, x_n, t)), \phi(M(x_{n-1}, x_{n+1}, 2t)), \phi(M(x_n, x_{n+1}, t))\} \\
 & + k_4(t)\phi(M(x_n, x_n, t)) \\
 \leq & k_1(t) \cdot \phi(M(x_{n-1}, x_n, t)) + k_2(t) \cdot \min\{\phi(M(x_n, x_{n+1}, t)), \phi(M(x_{n-1}, x_n, t))\} \\
 & + k_3(t) \cdot \min\{\phi(M(x_{n-1}, x_n, t)), \phi(T(M(x_{n-1}, x_n, t), M(x_n, x_{n+1}, t))), \\
 & \quad \phi(M(x_n, x_{n+1}, t))\}
 \end{aligned}$$

Again,

$$\min\{\phi(a), \phi(b), \phi(T(a, b))\} = \min\{\phi(a), \phi(b)\} \quad \forall a, b \in [0, 1]$$

Therefore,

$$\begin{aligned}
 \phi(M(x_n, x_{n+1}, t)) \leq & k_1(t) \cdot \phi(M(x_{n-1}, x_n, t)) + k_2(t) \cdot \min\{\phi(M(x_n, x_{n+1}, t)), \\
 & \phi(M(x_{n-1}, x_n, t))\} + k_3(t) \cdot \min\{\phi(M(x_n, x_{n+1}, t)), \phi(M(x_{n-1}, x_n, t))\}
 \end{aligned}$$

$$\begin{aligned}
&\text{If } \min\{\phi(M(x_n, x_{n+1}, t)), \phi(M(x_{n-1}, x_n, t))\} = \phi(M(x_n, x_{n+1}, t)), \\
&\text{then } \phi(M(x_n, x_{n+1}, t)) \leq k_1(t) \cdot \phi(M(x_{n-1}, x_n, t)) + k_2(t) \cdot \phi(M(x_n, x_{n+1}, t)) \\
&\quad + k_3(t) \cdot \phi(M(x_n, x_{n+1}, t)) \\
&\Rightarrow \phi(M(x_n, x_{n+1}, t)) \leq \frac{k_1(t)}{1 - k_2(t) - k_3(t)} \phi(M(x_{n-1}, x_n, t)) \\
&\quad < \phi(M(x_{n-1}, x_n, t))
\end{aligned}$$

$$\begin{aligned}
&\text{If } \min\{\phi(M(x_n, x_{n+1}, t)), \phi(M(x_{n-1}, x_n, t))\} = \phi(M(x_{n-1}, x_n, t)), \\
&\text{then } \phi(M(x_n, x_{n+1}, t)) \leq (k_1(t) + k_2(t) + k_3(t)) \cdot \phi(M(x_{n-1}, x_n, t)) \\
&\quad < \phi(M(x_{n-1}, x_n, t))
\end{aligned}$$

Thus, in both cases, $\phi(M(x_n, x_{n+1}, t)) \leq \phi(M(x_{n-1}, x_n, t))$.

So, proceeding as in the previous theorem, f has a unique fixed point. \square

EXAMPLE 2.14. We consider the same fuzzy metric space (X, M, T) together with the self map f and the altering distance function ϕ as Example 2.12. Here we take:

$$k_4(t) = \begin{cases} \frac{7}{t+7}, & t \in (0, 1] \\ \frac{t}{t+\frac{1}{7}}, & t \in (1, \infty) \end{cases}$$

and $k_1(t) = k_2(t) = k_3(t) = \frac{1-k_4(t)}{3}$

Then every pair of points satisfies the condition (15). Hence by Theorem 2.13, f has a unique fixed point.

3. Conclusion

In this paper, we have obtained different fixed point theorems for fuzzy metric spaces with several contractive type mappings. In 2016, Lael and Heidarpour [24] introduced a new class of generalized nonexpansive mappings and obtained some fixed point theorems in Banach spaces. Here we can pose the following problem:

Can we obtain similar type of results as in Theorem 2.1, 2.3, etc., in case of nonexpansive mappings in fuzzy metric spaces ?

Similar problems can also be raised for other types of mappings on fuzzy metric spaces.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally to the writing of this paper. All authors read and approved the final manuscript.

References

- [1] M. Abbas, T. Nazir, B.E. Rhoades, *Fixed points of multivalued mapping satisfying ciric type contractive conditions in G -metric spaces*, Hacettepe Journal of Mathematics and Statistic, Volume **42** (1) (2013), 21–29.
- [2] A. Aghajani, M. Abbas, J. R. Roshan, *Common fixed point of generalized weak contractive mapping in partially ordered G_b -metric spaces*, Filomat **28**:6 (2014), 1087–1101.
- [3] S. Banach, *Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales*, Fund. Math. **3** (1922), 133–181.
- [4] R. Chen, X. Wang, *Fixed point of nonlinear contractions in modular spaces*, Journal of Inequalities and Applications (2013), 2013:399.
- [5] D. Das and N. Goswami, *Fixed points of different contractive type mappings on tensor product spaces*, IJIRSET **3** (2009), 14512–14519.
- [6] D. Das and N. Goswami, *Fixed points of mappings satisfying a weakly contractive type condition*, Journal of Mathematical Research with Applications **36** (1) (2016), 70–78.
- [7] D. Das and N. Goswami, *Some fixed point theorems on the sum and product of operators in tensor product spaces*, International Journal of Pure and Applied Mathematics **109** (3) (2016), 651–663.
- [8] D. Das, N. Goswami, V.N. Mishra, *Some Results on Fixed Point Theorems in Banach Algebras*, Int. J. Anal. Appl. **13** (1) (2017), 32–40.
- [9] N.R. Das and M.L. Saha, *On fixed points in fuzzy metric spaces*, Annals of Fuzzy Mathematics and Informatics **7** (2) (Feb 2014), 313–318.
- [10] Deepmala, *A Study on Fixed Point Theorems for Nonlinear Contractions and its Applications*, Ph.D. Thesis (2014), Pt. Ravishankar Shukla University, Raipur 492 010, Chhatisgarh, India.
- [11] D. Dey, M. Saha, *An extension of Banach fixed point theorem in fuzzy metric space*, Bol. Soc. Paran. Mat. **32** (1) (2014), 299–304.
- [12] M. Dinarvand, *Some fixed point results for admissible Geraghty contraction type mappings in fuzzy metric spaces*, Iranian journal of fuzzy systems **14** (3) (2017), 161–177.
- [13] T. Došenović, D. Rakić, M. Brdar, *Fixed point theorem in fuzzy metric space using altering distance*, Filomat **28**:7 (2014), 1517–1524.
- [14] M. Edelstein, *On fixed and periodic points under contractive mappings*, J. Lond. Math. Soc. **37** (1962), 74–79.

- [15] L. Gajic, M. Stojakovic, *Sehgal-Thomas type fixed point theorems in generalized metric spaces*, Filomat **31**:11 (2017), 3347–3356.
- [16] A. George, P. Veeramani, *On some results in fuzzy metric spaces*, Fuzzy Set. Syst. **64** (1994), 395–399.
- [17] A. George, P. Veeramani, *On some results of analysis for fuzzy metric spaces*, Fuzzy set. Syst. **90** (1997), 365–368.
- [18] M. Grabiec, *Fixed points in fuzzy metric spaces*, Fuzzy Set. Syst. **27** (1988), 385–389.
- [19] J. Harjani, B. López and K. Sadarangani, *Fixed point theorems for cyclic ϕ -contractions in ordered metric space*, Fixed Point Theory **14** (2) (2013), 359–368.
- [20] N. Hussain, V. Parvaneh, B. Samet and C. Vetro, *Some fixed point theorems for generalized contractive mappings in complete metric spaces*, Fixed Point Theory and Application(2015), 2015:185.
- [21] E. Karapinar, *Edelstein type fixed point theorems*, Fixed Point Theory and Application(2012), 2012:107.
- [22] M. S. Khan, M. Swaleh, S. Sessa, *Fixed point theorems by altering distances between the points*, Bull. Aust. Math. Soc. **30** (1984), 1–9.
- [23] I. Kramosil and J. Michalek, *Fuzzy metrics and statistical metric spaces*, Kybernetika(Prague) **11** (5)(1975), 336–344.
- [24] F. Lael and Z. Heidarpour, *Fixed point theorems for a class of generalized non-expansive mappings*, Fixed Point Theory and Applications (2016), 2016:82.
- [25] X. Li, M. Guo, Y. Su, *On the intuitionistic fuzzy metric spaces and the intuitionistic fuzzy normed spaces*, J. Nonlinear Sci. Appl. www.tjnsa.com.
- [26] V.N. Mishra, *Some Problems on Approximations of Functions in Banach Spaces*, Ph.D. Thesis (2007), Indian Institute of Technology, Roorkee 247 667, Uttarakhand, India.
- [27] L.N. Mishra, *On existence and behavior of solutions to some nonlinear integral equations with Applications*, Ph.D. Thesis (2017), National Institute of Technology, Silchar 788 010, Assam, India.
- [28] V.N. Mishra, L.N. Mishra, *Trigonometric Approximation of Signals (Functions) in L_p ($p \geq 1$)-norm*, International Journal of Contemporary Mathematical Sciences **7** (19) (2012), 909–918.
- [29] L.N. Mishra, K. Jyoti, A. Rani, Vandana, *Fixed point theorems with digital contractions image processing*, Nonlinear Sci. Lett. A **9** (2) (2018), 104–115.
- [30] L.N. Mishra, H.M. Srivastava, M. Sen, *On existence results for some nonlinear functional-integral equations in Banach algebra with applications*, Int. J. Anal. Appl. **11** (1) (2016), 1–10.
- [31] L.N. Mishra, S.K. Tiwari, V.N. Mishra, *Fixed point theorems for generalized weakly S -contractive mappings in partial metric spaces*, Journal of Applied Analysis and Computation **5** (4) (2015), 600–612. doi:10.11948/2015047.
- [32] L.N. Mishra, S.K. Tiwari, V.N. Mishra, I.A. Khan, *Unique Fixed Point Theorems for Generalized Contractive Mappings in Partial Metric Spaces*, Journal of Function Spaces, Volume 2015 (2015), Article ID 960827, 8 pages.

- [33] J. R. Morales, E. M. Rojas, *Some generalizations of Jungck's fixed point theorem*, IJMMS (2012), Vol:2012, Article ID 213876.
- [34] R. M. Patel, R. Bhardwaj, *Some common fixed point theorems for compatible mapping in fuzzy metric spaces for Integral type mapping*, Mathematical Theory and modeling **3** (6) (2013).
- [35] H.K. Pathak and Deepmala, *Common fixed point theorems for PD-operator pairs under Relaxed conditions with applications*, Journal of Computational and Applied Mathematics, **239** (2013), 103-113.
- [36] A. Sadiku, *Schauder bases and locally complemented subspaces of Banach spaces*, Master's thesis, University of Agder, (2014).
- [37] B. Schweizer, A. Sklar, *Statistical metric spaces*, Pacific J. Math. **10** (1960), 314–334.
- [38] Y. Shen, D. Qiu, W. Chen, *Fixed point theorems in fuzzy metric spaces*, Appl. Math. Lett. **25** (2012), 138–141.
- [39] S. Shukla and M. Abbas, *Fixed point results in fuzzy metric-like spaces*, Iranian Journal of Fuzzy Systems **11** (5) (2014), 81–92.
- [40] H. Vosoughi, S. J. Hosseini Ghoncheh, *A remark on fuzzy contractions*, ICM (2012), 11-14 March, Faculty.uae.ac.ae.
- [41] N. Wairojjana, T. Došenović, D. Rakić, D. Gopal and P. Kumam, *An altering distance function in fuzzy metric space theorems*, Fixed Point Theory and Applications (2015), 2015:69.

Bijoy Patir

Department of Mathematics
 Gauhati University
 Guwahati-781014, Assam, India
E-mail: bpatir07@gmail.com

Nilakshi Goswami

Department of Mathematics
 Gauhati University
 Guwahati-781014, Assam, India
E-mail: nila.g2003@yahoo.co.in

Lakshmi Narayan Mishra

Department of Mathematics
 School of Advanced Sciences
 Vellore Institute of Technology (VIT) University
 Vellore 632 014, Tamil Nadu, India
 L. 1627 Awadh Puri Colony Beniganj, Phase-III
 Opposite-Industrial Training Institute (I.T.I.)
 Ayodhya Main Road, Faizabad 224 001, U.P., India
E-mail: lakshminarayanmishra04@gmail.com, l_n_mishra@yahoo.co.in