

ARTICLE

Development of a Rapid Method for the Screening of Conjugated Linoleic Acid (CLA)-Producing Strains of *Bifidobacterium breve*

Sun-Hae Choi[†], Kyoung-Min Lee[†], Kwan-Hu Kim, and Geun-Bae Kim*

Department of Animal Science and Technology, Chung-Ang University, Anseong 17546, Korea



OPEN ACCESS

Received July 30, 2018
Revised August 7, 2018
Accepted August 8, 2018

*Corresponding author : Geun-Bae Kim
Department of Animal Science and
Technology, Chung-Ang University,
Anseong 17546, Korea
Tel: +82-31-670-3027
Fax: +82-31-676-5986
E-mail: kimgeun@cau.ac.kr

[†] These authors contributed equally to this work.

Abstract This study was performed to isolate some strains of *Bifidobacterium breve* from fecal materials of neonates and to screen them for the biotransformation activity of converting linoleic acid into conjugated linoleic acid (CLA). Fecal samples were collected from twenty healthy neonates between 14 and 100 days old, and four hundred colonies were randomly selected from a *Bifidobacterium* selective transoligosaccharide medium. A duplex polymerase chain reaction technique was developed for the rapid and accurate molecular characterization of the *B. breve* strains that have been reported to show the species-specific characteristic of CLA production. They are identified by 16S ribosomal DNA, fructose-6-phosphate phosphoketolase encoding genes (*xfp*), and rapid pulsed field gel electrophoresis. Thirty-six isolates were identified as *B. breve*, and just two of the 12 neonates were harboring *B. breve* strains. Each isolate showed different CLA-producing ability in the spectrophotometric assay. All of the positive strains from the primary spectrophotometric assay were confirmed for their CLA-producing activities using gas-chromatographic analysis, and their conversion rates were different, depending on the strain isolated in this study. Some strains of *B. breve* were successfully isolated and characterized based on the CLA-producing activity, and further studies are necessary to characterize the enzyme and the gene responsible for the enzyme activity.

Keywords probiotics, *Bifidobacterium breve*, duplex PCR, conjugated linoleic acid, rapid screening

Introduction

The intestinal flora of an individual is composed of 100 trillion viable bacteria, representing 100 or more different bacterial species. These organisms live together in symbiotic or antagonistic relationships, growing on the food components ingested and biocomponents secreted into the alimentary tract by the host (Mitsuoka, 2000). Bifidobacteria are normal components of the intestinal flora throughout the life cycle. They are ubiquitous, endosymbiotic inhabitants of the gastrointestinal tract, vagina, and mouth. Bifidobacteria are the dominant species with *Eubacterium*, *Clostridium*, and

Bacteroides in the colon flora, existing from 10^8 to 10^{11} per 1 g of colon contents (Mitsuoka, 1978).

Members of the genus *Bifidobacterium* are believed to exert various positive health benefits on their host. These beneficial effects include anticarcinogenic activity (Biffi et al., 1997; Rafter et al., 2007; Reddy and Rivenson, 1993; Rowland et al., 1998), modulation of immune response (Yasui and Ohwaki, 1991), improvement of the digestive system (Saavedra et al., 1994), and lowering incidences of necrotizing enterocolitis in preterm neonates (Patole et al., 2016). According to a recent report, there are some *Bifidobacterium* strains that can convert linoleic acid into bioactive conjugated linoleic acids (CLAs) (Coakley et al., 2003; O'Shea et al., 2012).

The term "CLA" refers to a group of positional and geometric isomers of essential fatty-acid linoleic acid (LA, *cis*9, *cis*12- $C_{18:2}$) with conjugated double bonds, among which, the *cis*-9-,*trans*-11-ontadecadienoic acid (c9, t11 isomer) is approximately 80% of all possible CLA isomers (Pariza et al., 2001).

The conversion of linoleic acid to CLA takes place naturally in the rumen as part of the biohydrogenation process by the action of rumen bacteria such as *Butyrivibrio fibrisolvens* (Grinari and Bauman, 1999). Therefore, CLA is present as a natural component of ruminant meat as well as dairy products. There is increasing evidence of the potential health-promoting properties of CLA isomers, including antiatherosclerosis (Lee et al., 1994), anticarcinogenesis (Tian et al., 2011), enhancement of immunological function (Hayek et al., 1999), and reduction of body fat (Chaplin et al., 2015; Park et al., 1997; West et al., 1998).

Many probiotic bacteria, especially some bifidobacteria, have been reported to exhibit bioconversion activity from LA (linoleic acid) into CLA; however, this activity could be a strain-specific characteristic. From the screening of CLA production with the major species of bifidobacteria, it has been demonstrated that the most-efficient CLA producers belong to the species of *B. breve* (Barrett et al., 2007; Gorissen et al., 2010; Raimondi et al., 2016). In this study, a species-specific duplex polymerase chain reaction (PCR) technique for the selection of many strains belonging to the species of *B. breve* was developed, and this technique was successfully combined with a spectrophotometric assay for the rapid screening of probiotic candidates harboring CLA-producing activity.

Materials and Methods

***Bifidobacterium* isolation from fecal materials of neonates**

First, traditional colonies from neonatal feces were propagated under anaerobic conditions in transoligosaccharide (TOS) agar (Yakult Honsha, Japan) at 37°C for 48 h (Thitaram et al., 2005).

Bifidobacterium isolates were activated by successive subculturing into a Man-Rogosa-Sharpe (MRS) (Difco, USA) broth supplemented with 0.05% L-cysteine·HCl and stored in 10% skim milk supplemented with glycerol in a -80°C deep freezer. As control strains, *B. breve* BB5 (Korea Yakult R & D Center) and *B. breve* LMC520 (Choi et al., 2008) were used for the CLA production activity.

Identification of the isolates

Duplex PCR

Colonies grown on TOS agar were selected, and their genomic DNA was extracted with Lyse-N-Go (Pierce, USA) reagent according to the manufacturer's protocol. The primers used in the duplex PCR were a genus-specific primer based on *xfp* genes, which encode fructose-6-phosphate phosphoketolase, and species-specific primers based on 16S rDNA from *B. breve* (Table 1).

Table 1. Primers used in this study

Primers	Description	Sequence (5' to 3')	Product size (bp)	References
GP-107	Forward; 16S rDNA for <i>B. breve</i>	CCGGATGCTCCATCACAC	250	Matsuki et al. (2002)
GP-108	Reverse; 16S rDNA for <i>B. breve</i>	ACAAAGTGCCTTGCTCCCT		Matsuki et al. (2002)
GP-109	Forward 1; xfp for <i>Bifidobacterium</i>	TGGCAGTCCAACAAGCTC	923	This study
GP-111	Reverse ; xfp for <i>Bifidobacterium</i>	TAGGAGCTCCAGATGCCGTG		This study
GP-110	Forward 2; xfp for <i>Bifidobacterium</i>	CATCGACGGCAAGAAGACCG	582	This study
GP-111	Reverse ; xfp for <i>Bifidobacterium</i>	TAGGAGCTCCAGATGCCGTG		This study
GP-116	Forward; 27F, universal for bacteria	AGAGTTTGTATCCTGGCTCAG	1,450	
GP-120	Reverse; 1492R, universal for bacteria	GGTTACCTTGTTACGACTT		
GP-145	Forward; BSH gene for <i>B. breve</i>	ATGTGCACTGGTGTCCGTTTC	954	This study
GP-146	Reverse; BSH gene for <i>B. breve</i>	TCATCGGGCGACGCTGCTG		This study

16S rDNA partial sequencing

Colonies grown on TOS agar were selected, and the genomic DNA was extracted with Lyse-N-Go reagent, as recommended. Then, DNA was added in a mixture of 25 μ L HotSTARTaq Master mix (Qiagen, USA), and a 1 μ L forward and reverse primer set (GP116 & GP120) was used. The PCR product was purified using a PCR purification kit (Qiagen, USA) and identified by a Basic Local Alignment Search Tool (BLAST) search.

F6PPK encoding gene sequencing

Primers used in the duplex PCR were genus-specific primers (GP109 & GP111) based on the *xfp* gene. The PCR product was ligated into the pGEM-T easy vector (Promega, USA). The ligation product was transformed *E. coli* JM109 (Takara, Japan) and grown in Luria-Bertani (LB) agar supplemented with 50 μ g/mL of ampicillin and X-gal (LAX agar) plate at 37°C overnight. White colonies selected were inoculated in LB broth supplemented with 50 μ g/mL of ampicillin and grown under 150 rpm at 37°C for 8 h. Plasmids from recombinant colonies were prepared by a miniprep kit (Qiagen, USA) and sequenced with a SP6 and T7 primer pair (Macrogen, Korea). The sequence was identified by BLAST search.

Rapid pulsed field gel electrophoresis (PFGE)

According to the method of Briczinski and Roberts (2006), 2 mL of the overnight cultured cell was centrifuged at 14,000 rpm for 10 min and resuspended with 600 μ L of 100 mM Tris EDTA buffer (pH 7.6). 160 μ L of sample was added to 40 μ L of lysozyme (100 mg/mL) and 10 μ L of proteinase K (20 mg/mL) and mixed with an equal volume of 1.6% InCert agarose (Bio-Rad Laboratories, USA) prepared in 0.1% SDS (International Biotechnologies, USA). Blended samples were dispensed into disposable plug molds. The plugs were incubated in 0.5 M EDTA, 1% sarkosyl buffer (pH 9.0) supplemented by 2 mL of lysozyme (4 mg/mL) and mutanolysin (200 unit/mL) at 55°C for 90 min. After lysis, samples were cultured in fresh sarkosyl buffer supplemented with proteinase K (0.5 mg/mL) at 55°C for 1 h and washed in preheated distilled water at 50°C for 15 min. The plugs were washed three times with 10 mM EDTA buffer (pH 7.6) at 50°C for 15 min under 75 rpm in a shaking water bath. After lysis, the plugs were sliced at comb size and then transferred to 200 μ L of a fresh restriction digest mixture containing 50 units of *Xba* I and incubated at 37°C for 2 h.

Electrophoresis was performed on 1% InCert agarose gel using a 0.5 \times TBE buffer (Bio-Rad Laboratories, USA). A lambda

ladder (Bio-Rad Laboratories, USA) was included as a molecular size marker. An electrophoresis machine was operated using a CHEF DR-III (Bio-Rad Laboratories, USA). Operation times were increased linearly at 6 V/cm and 14°C. After electrophoresis, the gel was stained with ethidium bromide (0.4 mg/L; Promega, USA) for 1 h and then destained for 2 h with distilled water. The band pattern was confirmed using a UV transilluminator.

Screening of CLA-producing lactic-acid bacteria

Spectrophotometric assay

According to the method of Barrett et al. (2007), strains were incubated anaerobically in an mMRS broth containing linoleic acid (0.5 mg/mL) and 2% (wt/vol) Tween 80 at 37°C for 24 h. Following incubation, 1 mL of culture was centrifuged at 20,800×g for 1 min, the pellet was discarded, and the supernatant (0.5 mL) was mixed with 0.5 mL of hexane by vortexing for 2 min. 0.2 mL of the fatty acids extracted by vortexing the solution was mixed with 0.8 mL of methanol. The presence of CLA in the culture supernatant was assayed using a spectrophotometer (smart plus SP-1900PC, Youngwoo, Korea) at 233 nm.

Gas chromatography (GC)

According to the method of Jung et al. (2006), strains were incubated anaerobically in a mMRS broth containing linoleic acid (0.5 mg/mL) and 2% (wt/vol) Tween 80 at 37°C for 24 h. 0.5 mL of the sample was mixed with 20 µL of stock (8 mL of hexane, 100 mg of internal standard, C_{17:0}) and 0.5 mL of hexane. After tilting for 10 min, the hexane layer was extracted by centrifuge. The hexane layer was dried under nitrogen gas at 60°C for 30 min.

Fatty acid was methylated in 1% HCl at 60°C for 30 min. Fatty-acid methyl esters were produced by adding 2 mL of saturated NaCl and 1 mL of hexane. The solvent was dried under nitrogen gas and concentrated with 100 µL of hexane. Methylated fatty acids were analyzed with a gas chromatographer (Varian STAR 3400, USA) equipped with an SP-2560 column and a flame ionization detector.

Results and Discussion

Selection of bifidobacteria

Twelve neonatal feces were plated on TOS agar and incubated under anaerobic conditions at 37°C for 48 h. Four hundred typical milky-white colonies were selected from agar plate and were subcultured in MRS broth. Isolated strains were stored in 10% skim-milk-supplemented glycerol in a -80°C deep freezer.

Identification of the genus *Bifidobacterium*

Duplex PCR

Conventional biochemical tests have some limitations in identifying specific species and discriminating a large number of isolates. Thus, in this study, duplex PCR that can simultaneously accomplish the rapid identification and the correct differentiation of *B. breve* was developed.

The PCR assay was conducted with genus-specific primers (Table 1; GP109 & GP111 and GP111 & GP110) for the genus *Bifidobacterium*, producing products of 923 bp and 582 bp, respectively. As a result of PCR with species-specific primers (GP107 & GP108), it was confirmed that 250 bp PCR product was obtained from only *B. breve* strains (Fig. 1).




Fig. 1. Agarose gel electrophoresis of PCR products. (A) establishment of duplex PCR condition M: 1 kb ladder; lane 1: *Bifidobacterium*-specific PCR product (923 kb) & *B. breve*-specific PCR product (250 bp); lane 2: *Bifidobacterium*-specific PCR product (582 bp) & *B. breve*-specific PCR product (250 bp); lane 3: *B. breve*-specific PCR product (250 bp); lane 4: *Bifidobacterium*-specific PCR product (923 kb); lane 5: *Bifidobacterium*-specific PCR product (582 bp). (B) Identification of *Bifidobacterium breve*. M: 1 kb ladder; lane 1–10: *Bifidobacterium* strains isolated from infant feces. Dual bands of PCR products (lanes 1, 2, 4, 6, 7, 8, 9) represent *B. breve* strains and the others (lanes 3, 5, and 10) are belonging to the other species of *Bifidobacterium*. PCR, polymerase chain reaction.

To increase the specificity of the PCR, a duplex PCR was conducted with multiplex primer sets, and two pairs of genus-specific primer (GP109 & GP111, GP145 & GP146) and a pair of species-specific primer (GP107 & GP108) were used. After the evaluation process, primer pairs (GP109 & GP111 and GP107 & GP108) producing products of 923 bp and 250 bp were selected because this set was better than the other sets producing products of 582 bp and 250 bp in terms of the band separation (Fig. 1A).

Through the duplex PCR assay developed in this study, we were able to confirm 36 strains *B. breve* from total 400 strains of *Bifidobacterium* isolated from new born infants. Some of the typical PCR band pattern for *B. breve* strains, which produce the PCR products of 923 bp and 250 bp are presented in Fig. 1B.

It is generally accepted that the predominant microorganisms in the colon of breast-fed babies are bifidobacteria, while the colons of bottle-fed babies contain various other bacteria. Mitsou et al. (2007) reported that *Bifidobacterium longum* and *Bifidobacterium breve* were the most frequently detected *Bifidobacterium* species in breast-fed infants.

In this study, *B. breve* was isolated from two infants among 12 infants. Only one strain among 40 strains from one person was identified as a *B. breve*, but 31 strains among 40 strains from another person were identified as *B. breve*, proving significant differences depending on the host.

16S rDNA partial sequencing

As a result of sequencing the 16S rDNA of *B. breve* by duplex PCR, all strains had high homology with *B. breve*, indicating that all strains were *B. breve*. If the sequence of the 16S rRNA gene has more than 97% species sequence identity, they are considered to be the same species (Stackebrandt and Goebel, 1994). *Bifidobacterium breve* ATCC 15700^T and commercial strain BB5 have 98% sequence identity, and *B. breve* ATCC 15700^T and IB84 strain have 99% sequence identity. As a result of comparing the nucleotide sequence by clustal W to confirm the difference between the strains of isolated *B. breve*, all strains have the same 16S rDNA except BB5. Although it is difficult to identify the strain because of the high homology of the DNA sequence between genus *Bifidobacterium*, studies, such as those on multiple PCR (Kwon et al., 2005; Ventura et al., 2004) and DGGE (Satokari et al., 2001), have been conducted to identify *Bifidobacterium*, and it was found to be a convenient and accurate method for the identification of strain level.

F6PPK encoding gene sequencing

PCRs were conducted with genus-specific primers, and the PCR products were cloned into a pGEM-T easy vector. As a result of comparing the sequence by clustal W, the IB sample corresponded with *B. breve* ATCC 15700^T, and the BB5 sample and *B. breve* ATCC 15700^T showed difference in six nucleotides.

Rosselló-Mora and Amman (2001) reported that the 16S rRNA gene sequence cannot distinguish clearly between difference species belonging to the same genus of bacteria. Berthoud et al. (2005) identified *Bifidobacterium* by the *xfp* gene and reported that the *xfp* gene sequencing is more accurate than the 16S rRNA gene sequencing, except for *B. thermophilum*, *B. thermacidophilum*, and *B. boum*. In this study, it was also confirmed that in addition to 16S rRNA gene sequencing, *xfp* gene sequencing could be used as a tool of identification for the isolated *B. breve* strains (data not shown).

Rapid PFGE

For the genotyping and genetic fingerprinting of the bacterial isolates, PFGE was applied using *Xba* I as restriction enzymes. As shown in Fig. 2, PFGE analysis indicated that there was high correlation between new isolated strains (lane 2 to lane 6) and different band pattern was observed from KCTC strains (lane 8 to lane 11). Briczinski and Roberts (2006) reported that there is no difference between a PFGE protocol consuming 5–7 days and a rapid protocol that takes 24 h. In this study, PFGE was carried out using a rapid protocol, and it was confirmed that the band of the sample isolated from the same infants also showed the same patterns (Fig. 2).

Screening of CLA producing lactic-acid bacteria

Spectrophotometric assay

The standard gas liquid chromatography-based screening process is laborious and time-consuming and can be a limiting factor when a large number of strains are being tested. In this study, a simple and straightforward spectrophotometric method was used for screening a large number of culture supernatants for the CLA production. Through the spectrophotometric assay, the present study confirmed that *B. breve* isolated from neonatal feces produces CLA, but three species of *B. breve* received




Fig. 2. PFGE pattern of genomic DNA of *Bifidobacterium* strains after digestion by *Xba* I. Lane M, lambda ladder; lane 1, BB5; lane 2, IB26; lane 3, IB51; lane 4, IB55; lane 5, IB 62; lane 6, LMC520; lane 7, *B. breve* 2003; lane 8, *B. breve* KCTC 3220; lane 9, *B. breve* KCTC 3419; lane 10, *B. breve* KCTC 5081; lane 11, *B. breve* KCTC 3441; lane 12, *B. longum* BBL. PFGE, rapid pulsed field gel electrophoresis.

from Korean Collection of Type Cultures (*B. breve* KCTC 3220, *B. breve* KCTC 3419, *B. breve* KCTC 5018) and *B. breve* 2003 do not produce CLA (Fig. 3).

As a result of the spectrophotometric assay, CLA-producing ability was different depending on the strains tested, and it was observed that some of those isolates had higher CLA production activity than that of LMC520, which is one of the best CLA-producing strains (Choi et al., 2008).

Pariza and Yang (1999) reported that CLA detection is possible through the spectrophotometric assay, and Barrett et al. (2007) isolated and identified CLA-producing strains. In this study, the prescreening process was successful through the spectrophotometric assay, and the fatty acid analysis was carried out by GC for a more accurate quantitative analysis (Fig. 3).

Gas chromatography (GC)

Strains that had a high optical density value in the spectrophotometer assay were analyzed by GC to find excellent strains for the CLA production. First, the peak pattern of fatty-acid methyl ester (FAME) was analyzed using linoleic acid and conjugated methyl ester (Sigma, USA).

It was observed that linoleic acid was converted to CLA by *B. breve* isolated from neonatal feces through chromatograms (Fig. 4) with c9, t11-form as a major form of isomers of conjugated CLA. As shown in Fig. 5, conversion rates from LA into CLA was different depending on the strains. Among the isolated *B. breve* strains, IB52 and IB84 strains showed more than 65% conversion rate, which is comparable with LMC 520 strain (Choi et al., 2008). Many studies reported that free linoleic acid inhibits the growth of CLA-producing strains (Jiang et al., 1998; Kim et al., 2000; Rainio et al., 2001), and Tween 80 neutralizes the antibacterial activity and increases the solubility of the fatty acid in the liquid medium (Jiang et al., 1998; Rainio et al., 2001).

The study of the effect of the content of Tween 80 confirmed that the conversion rate of the medium containing 5% Tween 80 was higher than that of 2% Tween 80 (data not shown). The conversion rates of the isolates IB52, IB84, and LMC 520 were almost the same as that of LMC 520, and the conversion rate of IB84 was slightly higher than that of LMC 520.

Coakley et al. (2003) reported that *B. breve* and *B. dentium* have the highest productivity, comparing the conversion rate of CLA by *Bifidobacterium* isolated from neonatal feces, and Oh et al. (2003) confirmed that *B. breve* and *B. pseudocatenulatum*




Fig. 3. Screening of *B. breve* isolates for the production of CLA using spectrophotometric assay. Negative control: *B. longum* ATCC 15700^T; Positive control: LMC520; Commercial strain: BB5; Isolated strains from infant feces: IB7, IB13, IB29, IB30, IB31, IB32, IB52, IB54, IB55, IB75, IB80, IB84. CLA, conjugated linoleic acid.




Fig. 4. Chromatogram of conjugated linoleic acid (CLA) analysis with gas chromatograph. Heptadecanoic acid ($C_{17:0}$) was used as an internal standard. After 24 hr incubation, most of linoleic acid was converted into c9, t11-CLA and minor portion of t10, c12-CLA was also detected.




Fig. 5. Conversion rates of conjugated linoleic acids (*cis*-9, *trans*-11 isomer) from 24 h growth culture of *Bifidobacterium breve* strains in MRS broth. The linoleic acid was added as a 50 mg/mL stock solution containing 2% (v/v) Tween 80 and was previously filter sterilized through a 0.45 μ m filter. MRS, Man-Rogosa-Sharpe.

produce CLA. In this study, only *B. breve* had CLA-producing ability among 400 *Bifidobacterium* strains from neonatal feces.

Conclusion

Among the 400 strains identified out of 12 fecal isolates, 36 strains of *B. breve* were selected through duplex PCR. The screening study showed that IB52, IB80, and IB84 had CLA-producing ability converted from linoleic acid. In this study, using a duplex PCR for the selection of *B. breve* strains was a useful and precise method of selecting CLA-producing strains from human feces. Selected strains are expected to be widely used in materials of fermented milk products and health food. There are many studies that reported the efficacy of CLA-producing strains, including *B. breve*, through clinical trials. Further studies are necessary to clarify CLA-producing ability from linoleic acid by *B. breve* in the body.

Acknowledgement

This research was supported by the Chung-Ang University Research Scholarship Grants in 2012.

References

- Barrett E, Ross RP, Fitzgerald GF, Stanton C. 2007. Rapid screening method for analyzing the conjugated linoleic acid production capabilities of bacterial cultures. *Appl Environ Microbiol* 73:2333-2337.
- Berthoud H, Chavagnat F, Haueter M, Casey MG. 2005. Comparison of partial gene sequences encoding a phosphoketolase for the identification of bifidobacteria. *LWT-Food Sci Technol* 38:101-105.
- Biffi A, Coradini D, Larsen R, Riva L, Fronzo GD. 1997. Antiproliferative effect of fermented milk on the growth of a human cancer cell line. *Nutr Cancer* 28:93-99.
- Briczinski EP, Roberts RF. 2006. A rapid pulsed-field gel electrophoresis method for analysis of bifidobacteria. *J Dairy Sci* 89:2424-2427.
- Chaplin A, Parra P, Serra F, Palou A. 2015. Conjugated linoleic acid supplementation under a high-fat diet modulates stomach protein expression and intestinal microbiota in adult mice. *PLoS ONE* 10:e0125091.
- Choi NJ, Park HG, Kim YJ, Kim IH, Kang HS, Yoon CS, Yoon HG, Park SI, Lee JW, Chung SH. 2008. Utilization of monolinolein as a substrate for conjugated linoleic acid production by *Bifidobacterium breve* LMC 520 of human neonatal origin. *J Agric Food Chem* 56:10908-10912.
- Coakley M, Ross RP, Nordgren M, Fitzgerald G, Devery D, Stanton C. 2003. Conjugated linoleic acid biosynthesis by human-derived *Bifidobacterium* species. *J Appl Microbiol* 94:138-145.
- Gorissen L, Raes K, Weckx S, Dannenberger D, Leroy F, De Vuyst L, De Smet S. 2010. Production of conjugated linoleic acid and conjugated linolenic acid isomers by *Bifidobacterium* species. *Appl Microbiol Biotechnol* 87:2257-2266.
- Griinari JM, Bauman DE. 1999. Biosynthesis of conjugated linoleic acid and its incorporation into meat and milk in ruminants. In *Advances in conjugated linoleic acid research*. Yurawez MP, Mossoba MM, Kramer JKG, Pariza MW, Nelson GJ (ed). AOCS Press, Champaign, IL. pp 180-200.
- Hayek MG, Han SN, Wu D, Watkins BA, Meydani M, Dorsey JL, Smith DE, Meydani SN. 1999. Dietary conjugated linoleic acid influences the immune response of young and old C57BL/6NCrIBR mice. *J Nutr* 129:32-38.
- Jiang J, Björck L, Fonden R. 1998. Production of conjugated linoleic acid by dairy starter cultures. *J App Microbiol* 81:95-102.
- Jung MY, Kim GB, Jang ES, Jung YK, Park SY, Lee BH. 2006. Improved extraction method with hexane for gas chromatographic analysis of conjugated linoleic acids. *J Dairy Sci* 89:90-94.
- Kim YJ, Liu RH, Bond DR, Russel JB. 2000. Effect of linoleic acid concentration on conjugated linoleic acid production by *Butyrivibrio fibrisolvens* A38. *Appl Environ Microbiol* 66:5226-5230.
- Kwon HS, Yang EH, Lee SH, Yeon SW, Kang BH, Kim TY. 2005. Rapid identification of potentially probiotic *Bifidobacterium* species by multiplex PCR using species-specific primers based on the region extending from 16S rRNA through 23S rRNA. *FEMS Microbiol Lett* 250:55-62.
- Lee KN, Kritchevsky D, Pariza MW. 1994. Conjugated linoleic acid and atherosclerosis in rabbits. *Atherosclerosis* 108:19-25.
- Matsuki T, Watanabe K, Tanaka R. 2002. Genus- and species-specific PCR primers for the detection and identification of bifidobacteria. *Curr Issues Intest Microbiol* 4:61-69.
- Mitsuoka T. 1978. *Intestinal bacteria and health: An introductory narrative*. Iwanami Shoten Publishers, Tokyo, Japan.
- Mitsuoka T. 2000. Significance of dietary modulation of intestine flora and intestinal environment. *Biosci Microflora* 19:15-25.
- Oh D, Hong G, Lee Y, Min S, Sin H, Kim S. 2003. Production of conjugated linoleic acid by isolated *Bifidobacterium* strains. *J Microbiol Biotechnol* 19:907-912.

- O'Shea EF, Cotter PD, Stanton C, Ross RP, Hill C. 2012. Production of bioactive substances by intestinal bacteria as a basis of explaining probiotic mechanisms: Bacteriocins and conjugated linoleic acid. *Int J Food Microbiol* 152:189-205.
- Pariza MW, Yang XY. 1999. Method of producing conjugated fatty acids. US Patent 5,856,149.
- Park Y, Albright KJ, Liu W, Storkson JM, Cook ME, Pariza MW. 1997. Effect of conjugated linoleic acid on body composition in mice. *Lipids* 32:853-858.
- Pariza MW, Park Y, Cook ME. 2001. The biologically active isomers of conjugated linoleic acid. *Prog Lipid Res* 40:283-298.
- Patole SK, Rao SC, Keil AD, Nathan EA, Doherty DA, Slimmer KN. 2016. Benefits of *Bifidobacterium breve* M-16V supplementation in preterm neonates - A retrospective cohort study. *PLoS ONE* 11:e0150775.
- Rafter J, Bennett M, Caderni G, Clune Y, Hughes R, Karlsson PC, Klinder A, O'Riordan M, O'Sullivan GC, Pool-Zobel B, Rechkemmer G, Roller M, Rowland I, Salvadori M, Thijs H, Loo JV, Watzl B, Collins JK. 2007. Dietary synbiotics reduce cancer risk factors in polypectomized and colon cancer patients. *Am J Clin Nutr* 85:488-496.
- Raimondi S, Amaretti A, Leonardi A, Quartieri A, Gozzoli C, Rossi M. 2016. Conjugated linoleic acid production by bifidobacteria: Screening, kinetic, and composition. *Biomed Res Int* 2016:8654317.
- Rainio A, Vahvaselkä M, Suomalainen T, Laakso S. 2001. Reduction of linoleic acid inhibition in production of conjugated linoleic acid by *Propionibacterium freudenreichii* spp. *shermanii*. *Can J Microbiol* 47:735-740.
- Reddy BS, Rivenson A. 1993. Inhibitory effect of *Bifidobacterium longum* on colon, mammary, and liver carcinogenesis induced by 2-amino-3-methylimidazo [4,5f] quinoline, a food mutagen. *Cancer Res* 53:3914-3918.
- Rosselló-Mora R, Amman R. 2001. The species concept for prokaryotes. *FEMS Microbiol Rev* 25:39-67.
- Rowland IR, Rumney CJ, Coutts JT, Lievens LC. 1998. Effect of *Bifidobacterium longum* and inulin on gut bacterial metabolism and carcinogen-induced aberrant crypt foci in rats. *Carcinogenesis* 19:281-285.
- Saavedra JM, Bauman NA, Perma JA, Yolken RH, Saavedra JM, Bauman NA, Oung I. 1994. Feeding of *Bifidobacterium bifidum* and *Streptococcus thermophilus* to infants in hospital for prevention of diarrhea and shedding of rotavirus. *Lancet* 344:1046-1049.
- Satokari RM, Vaughan EE, Akkermans ADL, Saarela M, De Vos WM. 2001. Bifidobacterial diversity in human feces detected by genus-specific PCR and denaturing gradient gel electrophoresis. *Appl Environ Microbiol* 67:504-513.
- Stackebrandt E, Goebel BM. 1994. A place for DNA-DNA reassociation and 16S rRNA sequence analysis in the present species definition in bacteriology. *Int J Syst Bacteriol* 44:846-849.
- Thitaram SN, Siragusa GR, Hinton A Jr. 2005. Bifidobacterium-selective isolation and enumeration from chicken caeca by a modified oligosaccharide antibiotic-selective agar medium. *Lett Appl Microbiol* 41:355-360.
- Tian M, Kliewer KL, Asp ML, Stout MB, Belury MA. 2011. c9,t11-Conjugated linoleic acid-rich oil fails to attenuate wasting in colon-26 tumor-induced late-stage cancer cachexia in male CD2F1 mice. *Mol Nutr Food Res* 55:268-277.
- Ventura M, van Sinderen D, Fitzgerald GF, Zink R. 2004. Insights into the taxonomy, genetics and physiology of bifidobacteria. *Antonie van Leeuwenhoek* 86:205-223.
- West DB, Delany JP, Camet PM, Bolhm F, Truett AA, Scimeca JA. 1998. Effects of conjugated linoleic acid on body fat and energy metabolism in the mouse. *Am J Physiol* 275:667-672.
- Yasui H, Ohwaki M. 1991. Enhancement of immune response in Peyer's patch cells cultured with *Bifidobacterium breve*. *J Dairy Sci* 74:1187-1195.