

大韓醫療氣功學會

대한의료기공학회지

J. OF KOREAN ACADEMY OF MEDICAL GI-GONG

Vol.18 No.1 2018.12.31.

A Study on the Relationship between the Brain and the Pelvis

정재훈*, 필감매*, 안훈모*

* : 대한의료기공학회

ABSTRACT

A Study on the Relationship between the Brain and the Pelvis

Jae Hun Jung*, Chien Mei Pi*, Hun Mo Ahn*

* : The Member of the Korean Academy of Medical Gi-Gong

Objective : The purpose of this study is to examine the relationship between the brain and the pelvis.

Methods : The relationship between the pelvis and the brain was searched in PubMed, and these searching studies were reviewed.

· Received : 10 Nov. 2018 · Revised : 16 Nov. 2018 · Accepted : 25 Nov. 2018

Correspondence to : 안훈모(Hun Mo Ahn)

경기도 김포시 월곶면 애기봉로 7번길 20-12번지 월곶한의원

Tel. 031-987-8471 Fax. 031-987-8472 E-mail : ahnpig@gmail.com

Conclusions :

1. Urinary disorder is influenced by brain.
2. Brain is influenced by luteinizing hormone.
3. Pelvic floor muscles are influenced by brain.
4. Urological Chronic Pelvic Pain Syndrome(UCPPS) is influenced by brain.
5. Brain is influenced by the low intensity laser acupuncture stimulating thirteen ghost acupoints(includes CV1)

* **Keywords** : brain, pelvis, review article, 골반, 뇌, 종설 논문

I . INTRODUCTION

The safe performance of balance- and mobility-related activities during daily life, such as standing while performing manual tasks, rising from a chair and walking, requires adequate balance control mechanisms[1]. The disorders in balance control can be a consequence of pathologies, such as neurological disease, stroke, diabetes disease or a specific vestibular deficit, or can be due to age-related processes, such as a decline in muscle strength[2, 3], sensory functioning[4], or in generating appropriate sensorimotor responses[5]. Balance and mobility disorders can have serious consequences regarding physical functioning (e.g. reduced ability to perform activities of daily living) as well as psycho-social functioning (e.g. activity avoidance, social isolation, fear of falls) and may even lead to fall-related injuries. Advanced age alone carries an increased risk of perioperative morbidity and mortality[6, 7] Impaired cognition has also been correlated with adverse postoperative health outcomes, including an increased incidence of complications, delirium, a longer length of hospital

stay, and higher 6 month mortality[8]. Thus, more emphasis is being directed at cognition testing in the standard panel of preoperative risk assessment testing[6, 7]. ER Trowbridge et al.[9] say that screening for potential cognitive impairment in an ambulatory urogynecology population is feasible and useful in clinical practices.

For example, while walking, individuals with Huntington's disease(HD) demonstrated greater pelvic movement than controls in two conditions, peak anterior excursion and total anterior-posterior excursion. This indicates that during walking, individuals with HD tended to move farther into anterior pelvic tilt than the controls, while moving into an equivalent amount of posterior pelvic tilt. This finding is consistent with findings in elderly adults and individuals with traumatic brain injury[10, 11]. So pelvic condition is important for the quality of life.

Through the local reflex and central nervous system, acupuncture induces endocrine and autonomic responses to meet therapeutic effects[12, 13]. In East Asia, acupuncture has been used for a long time to treat many kinds of disorders especially including the neuropsychiatric illnesses[14, 15]. For decades, acupuncture was becoming increasingly popular in western countries and has aroused interests of scientists[16, 17]. Many studies[18-20] adopted fMRI or positron emission tomography (PET) to investigate the effects on human brain functions caused by acupuncture at the particular acupoint.

We researched about the relation of the brain and the pelvis by one principle of Korean medicine, 'shang bing xia qu, xia bing shang qu'(上病下取, 下病上取).

II. METHOD OF RESEARCH

We searched the keywords, 'brain' and 'pelvic' in PubMed. We selected the papers from the

results (2505 papers at 2018.9.28) which contained 'brain' and 'pelvic' at once. After reading their abstracts, we selected suitable some of them which were suitable with our objective. And of their reference, we selected suitable papers in same way. By reviewing them, we have come up with some conclusion.

III. REVIEW

1. Research about Relation of the Brain and the Pelvic Floor.

1) Urinary Disorder.

Camryn Harvie et al.[21] likelihood estimation map yielded six clusters of activation in the pons, cerebellum, insula, anterior cingulate cortex, thalamus, and the inferior frontal gyrus. Regions of the brain involved in executive control (frontal cortex), interoception (anterior cingulate cortex, insula), motor control (cerebellum, thalamus), and brainstem (pons) are involved in micturition. This analysis provides insight into the supraspinal control of voiding in healthy adults and a framework to understand dysfunctional voiding. Underactive bladder is a common voiding dysfunction in which the detrusor muscle is unable to generate an effective contraction sufficient to empty the bladder[22]. As many as 9–48% of men and 12 – 45% of women undergoing evaluation for lower urinary tract symptoms are diagnosed with underactive bladder[23, 24]. Leading theories suggested that the failure of the bladder to generate a sufficient bladder contraction during the voiding phase is likely due to an underlying neurological mechanism[25]. Evidence from animal experiments[26] suggests that during the voiding phase regions in the caudal brainstem coordinate the contraction of the detrusor muscle with the relaxation of the urethral sphincter but the initiation of micturition is determined in

regions of the forebrain. In humans, several neuroimaging studies have been performed to determine central mechanisms of micturition.

Using activation likelihood estimation method (ALE), Arya et al.[27] performed a meta-analysis of the regions of the brain activated during the storage phase of micturition. Afferent signals from the PAG are relayed via the thalamus to the insula, and likely the ACC which ultimately allows storage of urine in the bladder through increased sympathetic discharge, relaxation of the detrusor muscle relaxation of the bladder, and urethral sphincter contraction[28]. Using ALE meta-analysis, Arya et al.[29] reported on the regions of the brain that are activated during the filling or the storage phase of micturition.

C. Harvie et al.[30] provided the first voxel based meta-analysis to identify brain regions that are activated during the voiding phase of micturition. These regions include the brainstem, inferior frontal gyrus, insula, anterior cingulate cortex, thalamus, and cerebellum. Their finding that the anterior cingulate cortex (ACC) and the insula are activated in the voiding phase of micturition is clinically important. The ACC and the insula are also activated during bladder filling both in healthy controls as well as in subjects with overactive bladder, a condition of urinary urgency with or without incontinence[31, 32].

The insula, an area of the cerebral cortex that lies deep to the lateral sulcus receives afferent homeostatic information from the viscera, such as heart rate, respiration, digestive processes and micturition[33]. Also called the sensory cortex of the autonomic nervous system, activation of the insula during the storage phase represents afferent signaling from the bladder to the CNS. The ACC, located on the medial surface of the frontal lobe and surrounding the anterior genu of the corpus callosum, is called the motor cortex of the autonomic nervous system because it is responsible for

emotional, behavioral, and motor responses to visceral stimuli[34]. Alternatively, activation of the ACC and the insula during voiding may represent tight coupling between afferent and efferent centers in the brain[30]. Activation of the PFC has been reported both in subjects who could void successfully as well as those who were unable to void in the scanner despite trying vigorously to do so[35–37]. Some findings suggested that the decision to void is likely generated in the PFC. Prefrontal cortex is part of the attention network and is also concerned with the highest level of brain function including executive decision making[38]. On the other hand, stimulation of the cell group situated more ventrally and laterally in the dorsal pontine tegmentum results in contraction of the pelvic floor including the urethral sphincter through the Onuf's nucleus in the sacral spinal cord and has been called the L-region or the continence center[26]. As part of the attention and awareness networks, thalamic activation during storage determines which afferent signals from the bladder should reach the cerebral cortex[39]. Animal studies have demonstrated extensive connections of the cerebellum to other parts of the central nervous system including those involved in micturition and lesions of the cerebellum in humans are associated with both storage and voiding symptoms[40].

BD Clarkson et al.[41] evaluated older women with Urgency urinary incontinence(UUI) receiving biofeedback-assisted pelvic floor muscle therapy (PFMT), which focuses on identifying the correct muscles via biofeedback, urge suppression strategies, and behavioral techniques to avoid panic reactions to urgency and compared them to a similar continent group (“controls”) who did not receive treatment using three different approaches. Women who responded well to therapy had different baseline brain activation patterns associated with urgency[42] and these patterns tended to normalize among responders, but not among non-responders. Their results suggested that there

may be at least two mechanisms contributing to UUI: one in which a breakdown in central control plays the greater role and is thus responsive to a behaviorally targeted therapy like PFMT and another in which central control is still largely intact and for which a brain/behavior-targeted approach is less useful.

2) Luteinizing Hormone

Luteinizing hormone receptor(LHR) expression was also detected in nonpregnant human uterus (glandular and luminal epithelial cells, stromal cells, myometrial cells and vascular smooth muscle cells), syncytiotrophoblast of midterm but not term human placenta, fetal membranes and decidua[43], human and rat brain[44, 45], human trophoblast and syncytiotrophoblast from early pregnancies[46], porcine fallopian tubes and umbilical cord[47, 48], ovarian, decidual, endometrial and luteal macrophages[49, 50], and ovarian cancer cell cultures and tumor tissues[51]. Since the brain was reported to express functional LHR[44, 45], there is also a possibility that high LH levels in aging individuals may in some way participate in the development and progression of neurodegenerative process of several degenerative neurological diseases, including Alzheimer's disease.

Strong LHR expression in microglial cells in the brain cortex is of particular interest. Microglial cells are resident macrophages in the central nervous system, and LHR expression has been described in other types of human resident macrophages (ovary, decidua, endometrium and corpora lutea)[49, 50] Pathological activation of microglia has been reported in a wide range of conditions such as Alzheimer's disease, cerebral ischemia, prion diseases, multiple sclerosis, AIDS dementia, and other degenerative neurological diseases[52, 53]. Some of these degenerative diseases are associated with advanced age and high levels of circulating LH. High LH levels might pass the

presumptive LH barrier of brain vessels expressing LHR, or vascular LHR expression might diminish with age, as in abnormal term placentae. Yet, elevated maternal mid-trimester chorionic gonadotropin is associated with fetal cerebral blood flow redistribution and adverse perinatal outcome[54]. In addition, estrogens, which are known to cause a diminution of high LH levels in postmenopausal women, have been reported to be effective in the prevention and treatment of Alzheimer's disease[55–57]. Since CG increases secretion of a variety of cytokines by monocytes, and induces their inflammatory reaction and phagocytic activity[58–61], high LH levels in aging individuals may also activate resident macrophages in the central nervous system and contribute to the development of Alzheimer's disease and other inflammation-mediated neurodegenerative diseases. Outside of the area occupied by microglial cells in the grey matter of the cerebral cortex, weaker LHR expression was also detected in some nerve cells. A Bukovsky et al.[62] said that such nerve cells could be directly influenced by LH/CG proteohormones (with unknown consequences), while the nerve cells among microglial branching processes are protected from such an effect.

3) Pelvic Floor Muscles Disorder

Pelvic floor muscles play a critical role in reproductive and excretory processes[63]. A link between stress urinary incontinence(SUI) and nerve injuries affecting the urethral viscerosomatic reflex has been reported[64]. Consequently, a number of therapies to manage SUI, ranging from Kegel exercises to electrical stimulation of sacral and pudendal nerves, have been proposed as potential therapies to improve the function of pelvic floor musculature and its innervation[65]

According to Williams G et al.[10] and Lee LW et al. [11], individuals with Huntington's disease(HD) demonstrated greater pelvic movement than controls in two

conditions, peak anterior excursion and total anterior–posterior excursion while walking. This indicates that during walking, individuals with HD tended to move farther into anterior pelvic tilt than controls, while moving into an equivalent amount of posterior pelvic tilt. This finding is consistent with findings in elderly adults and individuals with traumatic brain injury.

4) Urological Chronic Pelvic Pain Syndrome (UCPPS)

UCPPS represents a group of pain syndromes relating to male and female pelvic pain that are poorly understood and for which treatment is largely unsatisfactory[66–68]. Since 2007, the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) [69] has combined Chronic Prostatitis/Chronic Pelvic Pain Syndrome (CP/CPPS) and Interstitial Cystitis/Painful Bladder Syndrome (IC/PBS) into a single classification of UCPPS. The complex and largely unknown etiology, as well as the varied symptom presentation in UCPPS, presents tremendous difficulties in both diagnosis and classification[70–72]. This complexity inherent in UCPPS suggests that some potential combination of peripheral and central nervous system biomarkers may be the best approach to diagnosis and may lead to a better understanding of the condition itself. For example, a functional magnetic resonance imaging (fMRI) brain study has recently shown functional differences between CP/ CPPS patients and healthy controls (HCs) that seem unique to this specific pathophysiology[73]. Other non-invasive neuroimaging studies have also suggested structural and functional differences may exist between various pain conditions and HCs, including Irritable Bowel Syndrome (IBS)[74, 75], chronic back pain [76–78], chronic pancreatitis[79], and chronic complex regional pain syndrome (CRPS)[80]. The Multidisciplinary Approach to the Study of Chronic Pelvic Pain (MAPP) Research Network (The MAPP Research Network

Study Group.) was developed to better understand the underlying pathophysiology of UCPPS and inform clinical management of patients[66, 81]. One important goal of the MAPP Network is to identify potential biological markers for UCPPS[82].

DE Harper et al. [83] interested in determining whether brain metabolite levels in several pain processing regions differ in UCPPS patients and controls. Additionally, they assessed whether regional metabolite levels related to functional connectivity of these regions. The results showed lower GABA and higher choline levels in the ACC of UCPPS patients, compared to age- and gender-matched control participants.

In patients, higher choline levels were related negative mood and higher ACC-to- limbic system connectivity. In addition, posterior insula Glx levels were related to higher pIC-to-ACC connectivity in patients compared to controls, in a region where significantly different metabolite levels were detected across groups. Their results add to the growing body of literature indicating central abnormalities in chronic pelvic pain and are the first to showing possible deficits in inhibitory neurotransmission in UCPPS.

The evidence for centralized components of chronic pelvic pain has been provided by brain neuroimaging studies, many from the MAPP Research Network, showing significant structural and functional anomalies in patients including white and gray matter differences, altered resting state functional connectivity[84–88] and heightened pain-evoked brain activity in and functional connectivity between pain-processing brain regions.

Furthermore, quantitative sensory testing has demonstrated widespread hyperalgesia, allodynia, and impaired endogenous analgesia in individuals with chronic pelvic pain, including UCPPS[89–97], which also strongly suggested involvement of the central nervous system.

Altered levels of brain metabolites have been observed in chronic pain patients with

endometriosis[89] and in other non-pelvic pain conditions[76, 98-114].

However, mounting evidence from neuroimaging suggests that the pain in many UCPPS patients may be related to alterations in the function and morphology of the central nervous system (CNS)[82, 84-89, 115-117]. Similar abnormalities of the CNS occur in other chronic overlapping pain conditions that are frequently comorbid with UCPPS[118-121], including fibromyalgia (FM), irritable bowel syndrome, chronic fatigue syndrome, and temporomandibular disorders.

The evidence for centralized components of chronic pelvic pain has been provided by brain neuroimaging studies, many from the MAPP Research Network, showing significant structural and functional anomalies in patients including white and gray matter differences[82, 115, 116, 122, 123] altered resting state functional connectivity[84-88], and heightened pain-evoked brain activity in and functional connectivity between pain-processing brain regions[85, 124]. Furthermore, quantitative sensory testing has demonstrated widespread hyperalgesia, allodynia, and impaired endogenous analgesia in individuals with chronic pelvic pain, including UCPPS[89-97] which also strongly suggest involvement of the central nervous system. UCPPS patients presented with significantly higher choline and lower GABA in the ACC, compared to healthy controls.

2. Acupuncture about brain

In recent years, the low intensity laser acupuncture (LA) emerged as a new modality of acupuncture and became increasingly popular due to its painless and noninvasive advantages[125, 126] and its operations in a precise and controlled manner when performing acupuncture[125] since the stimulating depth and strength of LA are independent of the acupuncturist's experience. Some studies had demonstrated low intensity LA at particular acupoint could cause certain effect on brain activities[125,

127–129]. For instance, in Siedentopf's study[125], laser acupuncture at an acupoint in foot (a target for visual disease treatment) caused the activation of the occipital cortex. Besides, Biella et al.[128], Hui et al.[129], Napadow et al.[127] found activation of the hypothalamic–limbic system when stimulated the analgesic acupoints by using LA. Thus, the LA is appropriate to be introduced for stimulating thirteen ghost acupoints(TGA). Jian Lv et al.'s[130] understanding the effect of acupuncture on brain functions is mainly based on employing fMRI which allows us to investigate the activation of the brain by recording the blood oxy–genation level–dependent (BOLD) signals obtained from the time series collected when subjects are performing certain cognitive tasks[131] or at rest[132]. In their study, they employed rs–fMRI and evaluated the ReHo and ALFF values to investigate the alterations of brain spontaneous activation in healthy participants after laser acupuncture at TGA.

IV. CONCLUSION

Treating the lower part of the body to heal the upper and treating the upper to heal the lower ('上病下取, 下病上取', 'shang bing xia qu, xia bing shang qu' – literally, 'upper disease, below treat, lower disease, upper treat') has been a widely applied treatment principle in Chinese medicine since ancient times. We explore and analyse the features of this treatment principle based on published clinical case studies and current research evidence.

We conclude that

1. Urinary disorder is influenced by brain.
2. Brain is influenced by luteinizing hormone.

3. Pelvic floor muscles is influenced by brain.
4. Urological Chronic Pelvic Pain Syndrome(UCPPS) is influenced by brain.
5. Brain is influenced by the low intensity laser acupuncture stimulating thirteen ghost acupoints(includes CV1).

The brain is mostly undiscovered part in the body. In Korean medicine, there are many principles about the brain. So we can discover them about the brain.

V. REFERENCES

- 1) Zijlstra A, Mancini M, Chiari L, Zijlstra W. *Biofeedback for training balance and mobility tasks in older populations: a systematic review*. Journal of neuroengineering and rehabilitation. 2010;7(1):58.
- 2) Horlings CG, Van Engelen BG, Allum JH, Bloem BR. *A weak balance: the contribution of muscle weakness to postural instability and falls*. Nature Reviews Neurology. 2008;4(9):504.
- 3) Moreland JD, Richardson JA, Goldsmith CH, Clase CM. *Muscle weakness and falls in older adults: a systematic review and meta-analysis*. Journal of the American Geriatrics Society. 2004;52(7):1121-9.
- 4) Shaffer SW, Harrison AL. *Aging of the somatosensory system: a translational perspective*. Physical therapy. 2007;87(2):193-207.
- 5) Bugnariu N, Fung J. *Aging and selective sensorimotor strategies in the regulation of upright balance*. Journal of neuroengineering and rehabilitation. 2007;4(1):19.
- 6) Hughes S, Leary A, Zweizig S, Cain J. *Surgery in elderly people: preoperative, operative and postoperative care to assist healing*. Best Practice & Research Clinical Obstetrics & Gynaecology. 2013;27(5):753-65.
- 7) Cook DJ, Rooke GA. *Priorities in perioperative geriatrics*. Anesthesia & Analgesia. 2003;96(6):1823-36.
- 8) Robinson TN, Wu DS, Pointer LF, Dunn CL, Moss M. *Preoperative cognitive dysfunction is*

- related to adverse postoperative outcomes in the elderly.* Journal of the American College of Surgeons. 2012;215(1):12-7.
- 9) Trowbridge ER, Kim D, Barletta K, Fitz V, Larkin S, Hullfish KL. *Prevalence of positive screening test for cognitive impairment among elderly urogynecologic patients.* American journal of obstetrics and gynecology. 2016;215(5):663. e1-. e6.
- 10) Williams G, Morris ME, Schache A, McCrory PR. *Incidence of gait abnormalities after traumatic brain injury.* Archives of physical medicine and rehabilitation. 2009;90(4):587-93.
- 11) Lee LW, Zavarei K, Evans J, Lelas JJ, Riley PO, Kerrigan DC. *Reduced hip extension in the elderly: dynamic or postural?* Archives of physical medicine and rehabilitation. 2005;86(9):1851-4.
- 12) Kim H, Park H-J, Han S-M, Hahn D-H, Lee H-J, Kim K-S, et al. The effects of acupuncture stimulation at PC6 (Neiguan) on chronic mild stress-induced biochemical and behavioral responses. *Neuroscience letters.* 2009;460(1):56-60.
- 13) Samuels N, Gropp C, Singer SR, Oberbaum M. *Acupuncture for psychiatric illness: a literature review.* Behavioral Medicine. 2008;34(2):55-64.
- 14) Courbasson CM, de Sorkin AA, Dullerud B, Van Wyk L. Acupuncture treatment for women with concurrent substance use and anxiety/depression: an effective alternative therapy? *Family & community health.* 2007;30(2):112-20.
- 15) Gallagher S, Allen J, Hitt S, Schnyer R, Manber R. *Six-month depression relapse rates among women treated with acupuncture.* Complementary therapies in medicine. 2001;9(4):216-8.
- 16) Pilkington K. *Acupuncture therapy for psychiatric illness.* International review of neurobiology. 111: Elsevier; 2013. p. 197-216.
- 17) Zeng B-Y, Salvage S, Jenner P. *Effect and mechanism of acupuncture on Alzheimer's disease.* International review of neurobiology. 111: Elsevier; 2013. p. 181-95.
- 18) Napadow V, Makris N, Liu J, Kettner NW, Kwong KK, Hui KK. *Effects of electroacupuncture versus manual acupuncture on the human brain as measured by fMRI.* Human brain mapping. 2005;24(3):193-205.
- 19) Napadow V, Dhond R, Purdon P, Kettner N, Makris N, Kwong K, et al., editors. *Correlating acupuncture fMRI in the human brainstem with heart rate variability.* 2005

IEEE Engineering in Medicine and Biology 27th Annual Conference; 2006: IEEE.

- 20) Dhond RP, Yeh C, Park K, Kettner N, Napadow V. Acupuncture modulates resting state connectivity in default and sensorimotor brain networks. *Pain*. 2008;136(3):407-18.
- 21) Harvie C, Weissbart SJ, Kadam-Halani P, Rao H, Arya LA. Brain Activation during the Voiding phase of Micturition in Healthy Adults: A Meta-analysis of Neuroimaging Studies. *Clinical Anatomy*. 2018.
- 22) Osman NI, Chapple CR. Contemporary concepts in the aetiopathogenesis of detrusor underactivity. *Nature Reviews Urology*. 2014;11(11):639.
- 23) Abarbanel J, Marcus E-L. Impaired detrusor contractility in community-dwelling elderly presenting with lower urinary tract symptoms. *Urology*. 2007;69(3):436-40.
- 24) Resnick NM, Brandeis GH, Baumann MM, DuBeau CE, Yalla SV. *Misdiagnosis of urinary incontinence in nursing home women: prevalence and a proposed solution*. *Neurourology and Urodynamics: Official Journal of the International Continence Society*. 1996;15(6):599-618.
- 25) Osman NI, Chapple CR. Fowler's syndrome—a cause of unexplained urinary retention in young women? *Nature Reviews Urology*. 2014;11(2):87.
- 26) Holstege G, Griffiths D, De Wall H, Dalm E. Anatomical and physiological observations on suprapinal control of bladder and urethral sphincter muscles in the cat. *Journal of Comparative Neurology*. 1986;250(4):449-61.
- 27) Arya NG, Weissbart SJ, Xu S, Rao H. Brain activation in response to bladder filling in healthy adults: An activation likelihood estimation meta-analysis of neuroimaging studies. *Neurourology and urodynamics*. 2017;36(4):960-5.
- 28) Griffiths D, Fowler C. *The micturition switch and its forebrain influences*. *Acta physiologica*. 2013;207(1):93-109.
- 29) Arya NG, Weissbart SJ. Central control of micturition in women: Brain-bladder pathways in continence and urgency urinary incontinence. *Clinical anatomy (New York, NY)*. 2017;30(3):373-84.
- 30) Harvie C, Weissbart SJ, Kadam-Halani P, Rao H, Arya LA. Brain activation during the voiding phase of micturition in healthy adults: A meta-analysis of neuroimaging studies. *Clinical anatomy (New York, NY)*. 2018.
- 31) Griffiths D, Tadic SD, Schaefer W, Resnick NM. *Cerebral control of the bladder in*

- normal and urge-incontinent women*. *NeuroImage*. 2007;37(1):1–7.
- 32) Komesu YM, Ketai LH, Mayer AR, Teshiba TM, Rogers RG. *Functional MRI of the Brain in Women with Overactive Bladder: Brain Activation During Urinary Urgency*. *Female pelvic medicine & reconstructive surgery*. 2011;17(1):50–4.
- 33) Craig AD. How do you feel? Interoception: the sense of the physiological condition of the body. *Nature reviews Neuroscience*. 2002;3(8):655–66.
- 34) Bush G, Luu P, Posner MI. *Cognitive and emotional influences in anterior cingulate cortex*. *Trends in cognitive sciences*. 2000;4(6):215–22.
- 35) Blok BF, Willemsen AT, Holstege G. *A PET study on brain control of micturition in humans*. *Brain : a journal of neurology*. 1997;120 (Pt 1):111–21.
- 36) Blok BF, Sturms LM, Holstege G. *Brain activation during micturition in women*. *Brain : a journal of neurology*. 1998;121 (Pt 11):2033–42.
- 37) Michels L, Blok BF, Gregorini F, Kurz M, Schurch B, Kessler TM, et al. *Supraspinal Control of Urine Storage and Micturition in Men—An fMRI Study*. *Cerebral cortex (New York, NY : 1991)*. 2015;25(10):3369–80.
- 38) Rossi AF, Pessoa L, Desimone R, Ungerleider LG. *The prefrontal cortex and the executive control of attention*. *Experimental brain research*. 2009;192(3):489–97.
- 39) Arya NG, Weissbart SJ, Xu S, Rao H. Brain activation in response to bladder filling in healthy adults: An activation likelihood estimation meta-analysis of neuroimaging studies. *Neurourol Urodyn*. 2017;36(4):960–5.
- 40) Sakakibara R, Uchida Y, Uchiyama T, Yamanishi T, Hattori T. Reduced cerebellar vermis activation during urinary storage and micturition in multiple system atrophy: 99mTc-labelled ECD SPECT study. *European journal of neurology*. 2004;11(10):705–8.
- 41) Clarkson BD, Karim HT, Griffiths DJ, Resnick NM. *Functional connectivity of the brain in older women with urgency urinary incontinence*. *Neurourol Urodyn*. 2018;37(8):2763–75.
- 42) Griffiths D, Clarkson B, Tadic SD, Resnick NM. Brain Mechanisms Underlying Urge Incontinence and its Response to Pelvic Floor Muscle Training. *J Urol*. 2015;194(3):708–15.
- 43) Reshef E, Lei ZM, Rao CV, Pridham DD, Chegini N, Luborsky JL. *The presence of gonadotropin receptors in nonpregnant human uterus, human placenta, fetal membranes, and decidua*. *The Journal of clinical endocrinology and metabolism*. 1990;70(2):421–30.

- 44) Lei ZM, Rao CV, Kornyei JL, Licht P, Hiatt ES. Novel expression of human chorionic gonadotropin/luteinizing hormone receptor gene in brain. *Endocrinology*. 1993;132(5):2262-70.
- 45) AA AL-H, Lei ZM, Rao CV. Neurons from fetal rat brains contain functional luteinizing hormone/chorionic gonadotropin receptors. *Biology of reproduction*. 1997;56(5):1071-6.
- 46) Tao YX, Lei ZM, Hofmann GE, Rao CV. Human intermediate trophoblasts express chorionic gonadotropin/luteinizing hormone receptor gene. *Biology of reproduction*. 1995;53(4):899-904.
- 47) Gawronska B, Pauku T, Huhtaniemi I, Wasowicz G, Ziecik AJ. *Oestrogen-dependent expression of LH/hCG receptors in pig Fallopian tube and their role in relaxation of the oviduct*. *Journal of reproduction and fertility*. 1999;115(2):293-301.
- 48) Wasowicz G, Derecka K, Stepien A, Pelliniemi L, Doboszynska T, Gawronska B, et al. *Evidence for the presence of luteinizing hormone-chorionic gonadotrophin receptors in the pig umbilical cord*. *Journal of reproduction and fertility*. 1999;117(1):1-9.
- 49) Bukovsky A, Chen TT, Wimalasena J, Caudle MR. Cellular localization of luteinizing hormone receptor immunoreactivity in the ovaries of immature, gonadotropin-primed and normal cycling rats. *Biology of reproduction*. 1993;48(6):1367-82.
- 50) Zhang YM, Rao Ch V, Lei ZM. *Macrophages in human reproductive tissues contain luteinizing hormone/chorionic gonadotropin receptors*. *American journal of reproductive immunology (New York, NY : 1989)*. 2003;49(2):93-100.
- 51) Gawronska B, Leuschner C, Enright FM, Hansel W. Effects of a lytic peptide conjugated to beta HCG on ovarian cancer: studies in vitro and in vivo. *Gynecologic oncology*. 2002;85(1):45-52.
- 52) Liu B, Hong JS. Role of microglia in inflammation-mediated neurodegenerative diseases: mechanisms and strategies for therapeutic intervention. *The Journal of pharmacology and experimental therapeutics*. 2003;304(1):1-7.
- 53) Nakamura Y. *Regulating factors for microglial activation*. *Biological & pharmaceutical bulletin*. 2002;25(8):945-53.
- 54) Hershkovitz R, Erez O, Sheiner E, Landau D, Mankuta D, Mazor M. *Elevated maternal mid-trimester chorionic gonadotropin > or =4 MoM is associated with fetal cerebral blood flow redistribution*. *Acta obstetricia et gynecologica Scandinavica*. 2003;82(1):22-7.

- 55) Fillit H. *Estrogens in the pathogenesis and treatment of Alzheimer's disease in postmenopausal women*. Annals of the New York Academy of Sciences. 1994;743:233-8; discussion 8-9.
- 56) Green PS, Simpkins JW. Estrogens and estrogen-like non-feminizing compounds. Their role in the prevention and treatment of Alzheimer's disease. Annals of the New York Academy of Sciences. 2000;924:93-8.
- 57) Monk D, Brodaty H. *Use of estrogens for the prevention and treatment of Alzheimer's disease*. Dementia and geriatric cognitive disorders. 2000;11(1):1-10.
- 58) Schäfer A, Pauli G, Friedmann W, Dudenhausen JW. Human choriongonadotropin (hCG) and placental lactogen (hPL) inhibit interleukin-2 (IL-2) and increase interleukin-1 β (IL-1 β), -6 (IL-6) and tumor necrosis factor (TNF- α) expression in monocyte cell cultures. Journal of Perinatal Medicine-Official Journal of the WAPM. 1992;20(3):233-40.
- 59) Reinisch N, Sitte B, Kähler C, Wiedermann C. Human chorionic gonadotrophin: a chemoattractant for human blood monocytes, neutrophils and lymphocytes. Journal of endocrinology. 1994;142(1):167-70.
- 60) Shirshv S, Lyalina O, Zamorina S. *Role of potassium ions in monocyte-regulating effects of chorionic gonadotropin*. Bulletin of experimental biology and medicine. 2000;130(5):1099.
- 61) Kosaka K, Fujiwara H, Tatsumi K, Yoshioka S, Sato Y, Egawa H, et al. Human chorionic gonadotropin (HCG) activates monocytes to produce interleukin-8 via a different pathway from luteinizing hormone/HCG receptor system. The Journal of Clinical Endocrinology & Metabolism. 2002;87(11):5199-208.
- 62) Bukovsky A, Indrapichate K, Fujiwara H, Cekanova M, Ayala ME, Dominguez R, et al. Multiple luteinizing hormone receptor (LHR) protein variants, interspecies reactivity of anti-LHR mAb clone 3B5, subcellular localization of LHR in human placenta, pelvic floor and brain, and possible role for LHR in the development of abnormal pregnancy, pelvic floor disorders and Alzheimer's disease. Reproductive Biology and Endocrinology. 2003;1(1):46.
- 63) Castelán F, López-García K, Moreno-Pérez S, Zempoalteca R, Corona-Quintanilla DL, Romero-Ortega MI, et al. *Multiparity affects conduction properties of pelvic floor nerves in rabbits*. Brain and behavior. 2018;8(10):e01105.

- 64) de Aguiar Cavalcanti G, Manzano GM, Nunes KF, Giuliano LMP, de Menezes TA, Bruschini H. *Electrophysiological evaluation of the pudendal nerve and urethral innervation in female stress urinary incontinence*. International urogynecology journal. 2013;24(5):801-7.
- 65) Elser DM. Stress urinary incontinence and overactive bladder syndrome: current options and new targets for management. Postgraduate medicine. 2012;124(3):42-9.
- 66) Clemens JQ, Mullins C, Kusek JW, Kirkali Z, Mayer EA, Rodríguez LV, et al. *The MAPP research network: a novel study of urologic chronic pelvic pain syndromes*. BMC urology. 2014;14(1):57.
- 67) Allsop SA, Erstad DJ, Brook K, Bhai SF, Cohen JM, Dimitrakoff JD. *The DABBECC phenotyping system: towards a mechanistic understanding of CP/CPPS*. Nature reviews Urology. 2011;8(2):107.
- 68) Hanno P, Andersson KE, Birder L, Elneil S, Kanai A, Pontari M. *Chronic pelvic pain syndrome/bladder pain syndrome: Taking stock, looking ahead: ICI-RS 2011*. Neurourology and urodynamics. 2012;31(3):375-83.
- 69) Nicke JC. A new approach to understanding and managing chronic prostatitis and interstitial cystitis. Reviews in urology. 2010;12(1):67.
- 70) Shoskes D, Nickel J, Rackley R, Pontari M. Clinical phenotyping in chronic prostatitis/chronic pelvic pain syndrome and interstitial cystitis: a management strategy for urologic chronic pelvic pain syndromes. Prostate cancer and prostatic diseases. 2009;12(2):177.
- 71) Lai HH, North CS, Andriole GL, Sayuk GS, Hong BA. *Polysymptomatic, polysyndromic presentation of patients with urological chronic pelvic pain syndrome*. The Journal of urology. 2012;187(6):2106-12.
- 72) Dimitrakov J, Guthrie D. *Genetics and phenotyping of urological chronic pelvic pain syndrome*. The Journal of urology. 2009;181(4):1550-7.
- 73) Farmer MA, Chanda ML, Parks EL, Baliki MN, Apkarian AV, Schaeffer AJ. *Brain functional and anatomical changes in chronic prostatitis/chronic pelvic pain syndrome*. The Journal of urology. 2011;186(1):117-24.
- 74) Chen JY-W, Blankstein U, Diamant NE, Davis KD. White matter abnormalities in irritable bowel syndrome and relation to individual factors. Brain research. 2011;1392:121-31.
- 75) Ellingson BM, Mayer E, Harris RJ, Ashe-McNally C, Naliboff BD, Labus JS, et al.

Diffusion tensor imaging detects microstructural reorganization in the brain associated with chronic irritable bowel syndrome. *PAIN®*. 2013;154(9):1528–41.

- 76) Grachev ID, Fredrickson BE, Apkarian AV. Abnormal brain chemistry in chronic back pain: an in vivo proton magnetic resonance spectroscopy study. *Pain*. 2000;89(1):7–18.
- 77) Apkarian AV, Sosa Y, Sonty S, Levy RM, Harden RN, Parrish TB, et al. *Chronic back pain is associated with decreased prefrontal and thalamic gray matter density*. *Journal of neuroscience*. 2004;24(46):10410–5.
- 78) Baliki MN, Chialvo DR, Geha PY, Levy RM, Harden RN, Parrish TB, et al. Chronic pain and the emotional brain: specific brain activity associated with spontaneous fluctuations of intensity of chronic back pain. *Journal of Neuroscience*. 2006;26(47):12165–73.
- 79) Frøkjær JB, Olesen SS, Gram M, Yavarian Y, Bouwense SA, Wilder-Smith OH, et al. *Altered brain microstructure assessed by diffusion tensor imaging in patients with chronic pancreatitis*. *Gut*. 2011;60(11):1554–62.
- 80) Geha PY, Baliki MN, Harden RN, Bauer WR, Parrish TB, Apkarian AV. The brain in chronic CRPS pain: abnormal gray–white matter interactions in emotional and autonomic regions. *Neuron*. 2008;60(4):570–81.
- 81) Landis JR, Williams DA, Lucia MS, Clauw DJ, Naliboff BD, Robinson NA, et al. *The MAPP research network: design, patient characterization and operations*. *BMC urology*. 2014;14(1):58.
- 82) Woodworth D, Mayer E, Leu K, Ashe-McNalley C, Naliboff BD, Labus JS, et al. Unique Microstructural Changes in the Brain Associated with Urological Chronic Pelvic Pain Syndrome (UCPPS) Revealed by Diffusion Tensor MRI, Super-Resolution Track Density Imaging, and Statistical Parameter Mapping: A MAPP Network Neuroimaging Study. *PLoS one*. 2015;10(10):e0140250.
- 83) Harper DE, Ichesco E, Schrepf A, Halvorson M, Puiu T, Clauw DJ, et al. Relationships between brain metabolite levels, functional connectivity, and negative mood in urologic chronic pelvic pain syndrome patients compared to controls: A MAPP research network study. *NeuroImage Clinical*. 2018;17:570–8.
- 84) Kilpatrick LA, Kutch JJ, Tillisch K, Naliboff BD, Labus JS, Jiang Z, et al. Alterations in resting state oscillations and connectivity in sensory and motor networks in women with interstitial cystitis/painful bladder syndrome. *J Urol*. 2014;192(3):947–55.

- 85) Kleinhans NM, Yang CC, Strachan ED, Buchwald DS, Maravilla KR. Alterations in Connectivity on Functional Magnetic Resonance Imaging with Provocation of Lower Urinary Tract Symptoms: A MAPP Research Network Feasibility Study of Urological Chronic Pelvic Pain Syndromes. *J Urol*. 2016;195(3):639-45.
- 86) Kutch JJ, Labus JS, Harris RE, Martucci KT, Farmer MA, Fenske S, et al. Resting-state functional connectivity predicts longitudinal pain symptom change in urologic chronic pelvic pain syndrome: a MAPP network study. *Pain*. 2017;158(6):1069-82.
- 87) Kutch JJ, Yani MS, Asavasopon S, Kirages DJ, Rana M, Cosand L, et al. Altered resting state neuromotor connectivity in men with chronic prostatitis/chronic pelvic pain syndrome: A MAPP: Research Network Neuroimaging Study. *NeuroImage Clinical*. 2015;8:493-502.
- 88) Martucci KT, Shirer WR, Bagarinao E, Johnson KA, Farmer MA, Labus JS, et al. The posterior medial cortex in urologic chronic pelvic pain syndrome: detachment from default mode network—a resting-state study from the MAPP Research Network. *Pain*. 2015;156(9):1755-64.
- 89) As-Sanie S, Kim J, Schmidt-Wilcke T, Sundgren PC, Clauw DJ, Napadow V, et al. *Functional Connectivity is Associated With Altered Brain Chemistry in Women With Endometriosis-Associated Chronic Pelvic Pain*. *The journal of pain : official journal of the American Pain Society*. 2016;17(1):1-13.
- 90) Clauw DJ, Schmidt M, Radulovic D, Singer A, Katz P, Bresette J. *The relationship between fibromyalgia and interstitial cystitis*. *Journal of psychiatric research*. 1997;31(1):125-31.
- 91) Grinberg K, Granot M, Lowenstein L, Abramov L, Weissman-Fogel I. A common pronociceptive pain modulation profile typifying subgroups of chronic pelvic pain syndromes is interrelated with enhanced clinical pain. *Pain*. 2017;158(6):1021-9.
- 92) Iacovides S, Avidon I, Baker FC. Women with dysmenorrhoea are hypersensitive to experimentally induced forearm ischaemia during painful menstruation and during the pain-free follicular phase. *European journal of pain (London, England)*. 2015;19(6):797-804.
- 93) Ness TJ, Powell-Boone T, Cannon R, Lloyd LK, Fillingim RB. *Psychophysical evidence of hypersensitivity in subjects with interstitial cystitis*. *The Journal of urology*.

2005;173(6):1983–7.

- 94) Ness TJ, Lloyd LK, Fillingim RB. An endogenous pain control system is altered in subjects with interstitial cystitis. *The Journal of urology*. 2014;191(2):364–70.
- 95) Powell–Boone T, Ness TJ, Cannon R, Lloyd LK, Weigent DA, Fillingim RB. *Menstrual cycle affects bladder pain sensation in subjects with interstitial cystitis*. *The Journal of urology*. 2005;174(5):1832–6.
- 96) Slater H, Paananen M, Smith AJ, O'Sullivan P, Briggs AM, Hickey M, et al. Heightened cold pain and pressure pain sensitivity in young female adults with moderate-to-severe menstrual pain. *Pain*. 2015;156(12):2468–78.
- 97) Stratton P, Khachikyan I, Sinaii N, Ortiz R, Shah J. *Association of chronic pelvic pain and endometriosis with signs of sensitization and myofascial pain*. *Obstetrics and gynecology*. 2015;125(3):719–28.
- 98) Aguila MR, Rebbeck T, Leaver AM, Lagopoulos J, Brennan PC, Hubscher M, et al. *The Association Between Clinical Characteristics of Migraine and Brain GABA Levels: An Exploratory Study*. *The journal of pain : official journal of the American Pain Society*. 2016;17(10):1058–67.
- 99) Cao B, Stanley JA, Selvaraj S, Mwangi B, Passos IC, Zunta-Soares GB, et al. Evidence of altered membrane phospholipid metabolism in the anterior cingulate cortex and striatum of patients with bipolar disorder I: A multi-voxel (1)H MRS study. *Journal of psychiatric research*. 2016;81:48–55.
- 100) Chang L, Munsaka SM, Kraft–Terry S, Ernst T. *Magnetic resonance spectroscopy to assess neuroinflammation and neuropathic pain*. *Journal of neuroimmune pharmacology : the official journal of the Society on Neuroimmune Pharmacology*. 2013;8(3):576–93.
- 101) Fayed N, Garcia–Campayo J, Magallon R, Andres–Bergareche H, Luciano JV, Andres E, et al. Localized 1H-NMR spectroscopy in patients with fibromyalgia: a controlled study of changes in cerebral glutamate/glutamine, inositol, choline, and N-acetylaspartate. *Arthritis research & therapy*. 2010;12(4):R134.
- 102) Feraco P, Bacci A, Pedrabissi F, Passamonti L, Zampogna G, Pedrabissi F, et al. *Metabolic abnormalities in pain-processing regions of patients with fibromyalgia: a 3T MR spectroscopy study*. *AJNR American journal of neuroradiology*. 2011;32(9):1585–90.
- 103) Foerster BR, Petrou M, Edden RA, Sundgren PC, Schmidt–Wilcke T, Lowe SE, et al.

- Reduced insular gamma-aminobutyric acid in fibromyalgia.* Arthritis and rheumatism. 2012;64(2):579-83.
- 104) Harris RE, Clauw DJ. Imaging central neurochemical alterations in chronic pain with proton magnetic resonance spectroscopy. *Neuroscience letters.* 2012;520(2):192-6.
- 105) Harris RE, Napadow V, Huggins JP, Pauer L, Kim J, Hampson J, et al. *Pregabalin rectifies aberrant brain chemistry, connectivity, and functional response in chronic pain patients.* Anesthesiology. 2013;119(6):1453-64.
- 106) Harris RE, Sundgren PC, Craig AD, Kirshenbaum E, Sen A, Napadow V, et al. *Elevated insular glutamate in fibromyalgia is associated with experimental pain.* Arthritis and rheumatism. 2009;60(10):3146-52.
- 107) Harris RE, Sundgren PC, Pang Y, Hsu M, Petrou M, Kim SH, et al. Dynamic levels of glutamate within the insula are associated with improvements in multiple pain domains in fibromyalgia. *Arthritis and rheumatism.* 2008;58(3):903-7.
- 108) Ito T, Tanaka-Mizuno S, Iwashita N, Tooyama I, Shiino A, Miura K, et al. Proton magnetic resonance spectroscopy assessment of metabolite status of the anterior cingulate cortex in chronic pain patients and healthy controls. *Journal of pain research.* 2017;10:287-93.
- 109) Petrou M, Harris RE, Foerster BR, McLean SA, Sen A, Clauw DJ, et al. Proton MR spectroscopy in the evaluation of cerebral metabolism in patients with fibromyalgia: comparison with healthy controls and correlation with symptom severity. *AJNR American journal of neuroradiology.* 2008;29(5):913-8.
- 110) Petrou M, Pop-Busui R, Foerster BR, Edden RA, Callaghan BC, Harte SE, et al. *Altered excitation-inhibition balance in the brain of patients with diabetic neuropathy.* Academic radiology. 2012;19(5):607-12.
- 111) Reckziegel D, Raschke F, Cottam WJ, Auer DP. Cingulate GABA levels inversely correlate with the intensity of ongoing chronic knee osteoarthritis pain. *Molecular pain.* 2016;12.
- 112) Valdes M, Collado A, Bargallo N, Vazquez M, Rami L, Gomez E, et al. Increased glutamate/glutamine compounds in the brains of patients with fibromyalgia: a magnetic resonance spectroscopy study. *Arthritis and rheumatism.* 2010;62(6):1829-36.
- 113) Widerstrom-Noga E, Pattany PM, Cruz-Almeida Y, Felix ER, Perez S, Cardenas DD, et al. Metabolite concentrations in the anterior cingulate cortex predict high neuropathic pain

- impact after spinal cord injury. *Pain*. 2013;154(2):204–12.
- 114) Zhao X, Xu M, Jorgenson K, Kong J. Neurochemical changes in patients with chronic low back pain detected by proton magnetic resonance spectroscopy: A systematic review. *NeuroImage Clinical*. 2017;13:33–8.
- 115) Bagarinao E, Johnson KA, Martucci KT, Ichesco E, Farmer MA, Labus J, et al. *Preliminary structural MRI based brain classification of chronic pelvic pain: A MAPP network study*. *Pain*. 2014;155(12):2502–9.
- 116) Farmer MA, Huang L, Martucci K, Yang CC, Maravilla KR, Harris RE, et al. *Brain White Matter Abnormalities in Female Interstitial Cystitis/Bladder Pain Syndrome: A MAPP Network Neuroimaging Study*. *The Journal of urology*. 2015;194(1):118–26.
- 117) Mayer EA, Gupta A, Kilpatrick LA, Hong JY. *Imaging brain mechanisms in chronic visceral pain*. *Pain*. 2015;156 Suppl 1:S50–63.
- 118) Kuner R, Flor H. *Structural plasticity and reorganisation in chronic pain*. *Nature reviews Neuroscience*. 2016;18(1):20–30.
- 119) Maixner W, Fillingim RB, Williams DA, Smith SB, Slade GD. *Overlapping Chronic Pain Conditions: Implications for Diagnosis and Classification*. *The journal of pain : official journal of the American Pain Society*. 2016;17(9 Suppl):T93–t107.
- 120) Schmidt-Wilcke T. *Neuroimaging of chronic pain*. *Best practice & research Clinical rheumatology*. 2015;29(1):29–41.
- 121) Walitt B, Ceko M, Gracely JL, Gracely RH. *Neuroimaging of Central Sensitivity Syndromes: Key Insights from the Scientific Literature*. *Current rheumatology reviews*. 2016;12(1):55–87.
- 122) Huang L, Kutch JJ, Ellingson BM, Martucci KT, Harris RE, Clauw DJ, et al. Brain white matter changes associated with urological chronic pelvic pain syndrome: multisite neuroimaging from a MAPP case-control study. *Pain*. 2016;157(12):2782–91.
- 123) Kairys AE, Schmidt-Wilcke T, Puiu T, Ichesco E, Labus JS, Martucci K, et al. Increased brain gray matter in the primary somatosensory cortex is associated with increased pain and mood disturbance in patients with interstitial cystitis/painful bladder syndrome. *The Journal of urology*. 2015;193(1):131–7.
- 124) Deutsch G, Deshpande H, Frolich MA, Lai HH, Ness TJ. Bladder Distension Increases Blood Flow in Pain Related Brain Structures in Subjects with Interstitial Cystitis. *The*

- Journal of urology. 2016;196(3):902-10.
- 125) Siedentopf CM, Golaszewski SM, Mottaghy FM, Ruff CC, Felber S, Schlager A. Functional magnetic resonance imaging detects activation of the visual association cortex during laser acupuncture of the foot in humans. *Neuroscience letters*. 2002;327(1):53-6.
- 126) Siedentopf CM, Koppelstaetter F, Haala IA, Haid V, Rhomberg P, Ischebeck A, et al. *Laser acupuncture induced specific cerebral cortical and subcortical activations in humans*. *Lasers in medical science*. 2005;20(2):68-73.
- 127) Napadow V, Makris N, Liu J, Kettner NW, Kwong KK, Hui KK. *Effects of electroacupuncture versus manual acupuncture on the human brain as measured by fMRI*. *Human brain mapping*. 2005;24(3):193-205.
- 128) Biella G, Sotgiu ML, Pellegata G, Paulesu E, Castiglioni I, Fazio F. *Acupuncture produces central activations in pain regions*. *NeuroImage*. 2001;14(1 Pt 1):60-6.
- 129) Hui KK, Liu J, Makris N, Gollub RL, Chen AJ, Moore CI, et al. Acupuncture modulates the limbic system and subcortical gray structures of the human brain: evidence from fMRI studies in normal subjects. *Human brain mapping*. 2000;9(1):13-25.
- 130) Lv J, Shi C, Deng Y, Lou W, Hu J, Shi L, et al. The brain effects of laser acupuncture at thirteen ghost acupoints in healthy individuals: A resting-state functional MRI investigation. *Computerized medical imaging and graphics : the official journal of the Computerized Medical Imaging Society*. 2016;54:48-54.
- 131) Zhang S, Li CS. A neural measure of behavioral engagement: task-residual low-frequency blood oxygenation level-dependent activity in the precuneus. *NeuroImage*. 2010;49(2):1911-8.
- 132) Rosazza C, Minati L. *Resting-state brain networks: literature review and clinical applications*. *Neurological sciences : official journal of the Italian Neurological Society and of the Italian Society of Clinical Neurophysiology*. 2011;32(5):773-85.