

Infection Status with *Metagonimus* spp. Metacercariae in Fishes from Seomjin-gang and Tamjin-gang in Republic of Korea

Woon-Mok Sohn^{1,*}, Byoung-Kuk Na¹, Shin-Hyeong Cho², Jung-Won Ju², Cheon-Hyeon Kim³, Ki-Bok Yoon⁴

¹Department of Parasitology and Tropical Medicine, and Institute of Health Sciences, Gyeongsang National University College of Medicine, Jinju 52727, Korea; ²Division of Vectors and Parasitic Diseases, Centers for Disease Control and Prevention, Osong 28159, Korea; ³Division of Microorganism, Jeollabuk-do Institute of Health and Environment, Imsil 55928, Korea; ⁴Division of Microbiology, Jeollanam-do Institute of Health and Environment, Muan 58568, Korea

Abstract: To grasp the infection status of *Metagonimus* spp. metacercariae (MsMc), the freshwater fishes were surveyed from Seomjin-gang (river) and Tamjin-gang in the Republic of Korea. Total 1,604 fishes from 7 local sites of Seomjin-gang and 1,649 fishes from 2 sites of Tamjin-gang were examined for 6 years (2012-2017) by the artificial digestion method. MsMc were detected in fishes from 7 sites, i.e., Osucheon in Imsil-gun (36.3% fish in 6 spp.), Seomjin-gang in Sunchang-gun (49.8% in 18 spp.), Songdaecheon in Namwon-si (64.5% in 8 spp.), Seomjin-gang in Gokseong-gun (72.4% in 14 spp.) and in Gurye-gun (78.8% in 17 spp.), Hoengcheon (75.9% in 11 spp.) and Namsancheon (58.9% in 7 spp.) in Hadong-gun. Their average densities were 4.2, 86.8, 39.9, 43.1, 246.5, 173.6 and 67.5 per fish infected respectively. Prevalence with MsMc in rasborinid fish from Seomjin-gang was 89.2% and their intensity was 73.6 per fish infected. Endemicities of MsMc in rasborinid fish from Seomjin-gang were significantly higher in fishes from the lower reaches (prevalence: 98.7%; density: 137) rather than in fishes from the middle (93.5%; 38) and upper (72.4%; 13) reaches. MsMc were also detected in 56.1% and 66.4% fishes from 2 surveyed areas, i.e., the middle reaches in Jangheung-gun and the lower reaches in Gangjin-gun, of Tamjin-gang, and their densities were 147 and 121 per fish infected. In susceptible fishes from Tamjin-gang, the prevalence was 84.1% and density was 227 per fish infected. By the present study, it was confirmed that MsMc is highly prevalent in the fishes from Seomjin-gang and Tamjin-gang in Korea.

Key words: *Metagonimus* spp., metacercaria, Seomjin-gang, Tamjin-gang

INTRODUCTION

Members in the genus *Metagonimus* Katsurada, 1912 (Digenea: Heterophyidae) are comprised more than 7 nominal species, i.e., *M. yokogawai* Katsurada, 1912, *M. takahashii* Suzuki, 1930, *M. minutus* Katsuta, 1932, *M. katsuradai* Izumi, 1935, *M. otsurui* Saito and Shimizu, 1968, *M. miyatai* Saito et al., 1997 and *M. hakubensis* Shimazu, 1999. Among 7 *Metagonimus* species, 3 ones, i.e., *M. yokogawai*, *M. takahashii*, and *M. miyatai*, are known to distribute in the Republic of Korea (Korea) [1,2]. Human infection by these species of fluke, metagonimiasis, is an important endemic disease together with clonorchiasis in

Korea [1-3]. Infections by *M. yokogawai* are chiefly prevalent in the riverside areas of eastern and southern coast of Korean peninsula [3-7]. Human cases by *M. takahashii* were first reported in inhabitants of Eumseong-gun (gun = county), Chungcheongnam-do (do = Province), along the upper reaches of the Namhan-gang [6]. Endemic areas of *M. miyatai* were confirmed among peoples residing around lakes and along the rivers and/or streams in inland of Korea [9-11]. These *Metagonimus* species give rise to severe gastrointestinal troubles and chronic diarrhea in heavily infected cases [1,2,12].

As the infection sources of metagonimiasis, lots of fish species have been reported in Korea [13]. The sweet smelt (*Plecoglossus altivelis*), the sea rundace (*Tribolodon hakonensis*) and the Japanese seabass (*Lateolabrax japonicus*), are known to be the second intermediate hosts of *M. yokogawai* [13-16]. The crucian carp (*Carassius auratus*), common carp (*Cyprinus carpio*), sea rundace, and Japanese seabass are reported as the second intermediate hosts of *M. takahashii* [13,16,17]. As the second

•Received 12 May 2018, revised 3 July 2018, accepted 29 July 2018.

*Corresponding author (wmsohn@gnu.ac.kr)

© 2018, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

intermediate hosts of *M. miyatai*, many species of fish, including the sea rundace, pale chub, (*Zacco platypus*) and dark chub (*Z. temminckii*), are listed in Korea [13,16,18].

On the other hand, many Korean workers have been investigated the infection status with zoonotic trematodes (ZT), i.e., *C. sinensis*, *Metagonimus* spp. including *M. yokogawai*, *Centrocestus armatus* and *Echinostoma* spp., metacercariae in fishes from various endemic areas to estimate the endemicities of these trematode infections [1,13]. In case of *Metagonimus* spp., most of studies were performed on the infection status of *M. yokogawai* in sweet smelts from the specific regions [19-25]. Recently, Cho et al. [26] surveyed on the infection status of ZT metacercariae including *Metagonimus* spp. in freshwater fish from Gangwon-do, Korea. Sohn et al. [27] investigated the infection status of digenetic trematode metacercariae in freshwater fish from the water systems of Hantan-gang and Imjin-gang located in relatively northern regions of Korea. Sohn et al. [28] and Yoon et al. [29] reported the infection status with *C. sinensis* metacercariae in fishes from Seomjin-gang and Tamjin-gang. However, the infection status with MsMc in fish from Seomjin-gang and Tamjin-gang has not been widely and systematically examined yet, although these 2 riverside areas has been known as the endemic areas of heterophyid flukes including *M. yokogawai* [3,4,30-32]. Therefore, we performed the present study to know the infection status with MsMc in fishes from 2 rivers, Seomjin-gang and Tamjin-gang, located in the southern parts of Korea.

MATERIALS AND METHODS

Collection sites of fish

We collected total 1,604 freshwater fishes in 7 local sites of Seomjin-gang, i.e., Osucheon ("cheon" means stream) (Latitude: 35.528473; Longitude: 127.328177) in Imsil-gun, Seomjin-gang (35.409674; 127.219528) in Sunchang-gun, Songdaecheon (35.352539; 127.189819) in Namwon-si, Jeollabuk-do, Seomjin-gang (35.212588; 127.371886) in Gokseong-gun, Seomjin-gang (35.200735; 127.491839) in Gurye-gun, Jeollanam-do, Hoengcheon (35.107203; 127.807894) and Namsancheon (35.095020; 127.798093) in Hadong-gun, Gyeongsangnam-do, for 5 years (2012-2016) (Fig. in [28]). We also collected total 1,132 fishes (22 species) in the middle reaches of Tamjin-gang in Jangheung-gun (Latitude: 34.425719; Longitude: 126.543227) for 4 years (2014-2017) and total 517 fishes (17 species) in the lower reaches of Tamjin-gang in Gangjin-gun (Latitude: 34.380531; Longitude: 126.485115),

Jeollanam-do, Korea in 2104 and 2017.

Fishes examined in the upper reaches of Seomjin-gang

Total 236 freshwater fish (11 species) from Osucheon in Imsil-gun, Jeollabuk-do were examined in 2012 and 2013. Fish species (No. of fish) examined were *Squalidus japonicus coreanus* (20), *Microphysogobio jeoni* (6), *Pseudogobio esocinus* (5), *Carassius auratus* (3), and *Hemibarbus longirostris* (1) including 6 ones with MsMc in Table 1. Total 310 freshwater fish (29 species) from Seomjin-gang in Sunchang-gun, Jeollabuk-do were examined in 2014 and 2015. Fish species (No. of fish) examined were *Coreoperca herzi* (19), *Acheilognathus koreanis* (12), *Odontobutis platycephala* (9), *Coreoleuciscus splendidus* (4), *Acheilognathus yamatsutae* (3), *Abbottina springeri* (3), *Cobitis tetralin-*

Table 1. Infection status of *Metagonimus* spp. metacercariae in freshwater fish from the upper reaches of Seomjin-gang (River)

Locality and fish sp.	No. of fish examined	No. (%) of fish infected	No. of MsMc detected	
			Range	Average
Osucheon in Imsil-gun				
<i>Zacco platypus</i>	80	48 (60.0)	1-32	4.2
<i>Hemibarbus labeo</i>	56	1 (1.8)	-	1.0
<i>Squalidus chankaensis</i>	35	8 (22.9)	1-10	3.3
<i>Opsariichthys uncirostris</i>	21	13 (61.9)	1-25	5.1
<i>Acanthorhodeus gracilis</i>	6	1 (16.7)	-	3.0
<i>Rhodeus ocellatus</i>	3	2 (66.7)	2-4	3.0
Subtotal	201	73 (36.3)	1-32	4.2
Seomjin-gang in Sunchang-gun				
<i>Pungtungia herzi</i>	56	13 (23.2)	1-9	1.9
<i>Zacco platypus</i>	51	45 (88.2)	1-200	25.8
<i>Pseudogobio esocinus</i>	32	13 (40.6)	1-60	6.9
<i>Zacco koreanus</i>	15	11 (73.3)	1-36	7.9
<i>Opsariichthys uncirostris</i>	14	14 (100)	1-45	9.9
<i>Carassius auratus</i>	14	1 (7.1)	-	5.0
<i>Sarcocheilichthys variegatus</i>	12	4 (33.3)	1-6	2.8
<i>Squalidus japonicus coreanus</i>	11	3 (27.3)	-	1.0
<i>Acheilognathus rhombeus</i>	10	3 (30.0)	1-4	2.3
<i>Siniperca scherzeri</i>	10	1 (10.0)	-	2.0
<i>Squalidus gracilis majimae</i>	6	1 (16.7)	-	1.0
<i>Acanthorhodeus gracilis</i>	6	5 (83.3)	7-40	16.2
<i>Gnathopogon strigatus</i>	5	4 (80.0)	5-14	7.5
<i>Hemibarbus longirostris</i>	5	4 (80.0)	2-14	8.0
<i>Hemibarbus labeo</i>	4	1 (25.0)	-	2.0
<i>Plecoglossus altivelis</i>	3	3 (100)	282-6,750	3,146
<i>Cyprinus carpio</i>	2	1 (50.0)	-	1.0
<i>Hemiculter leucisculus</i>	1	1 (100)	-	1.0
Subtotal	257	128 (49.8)	1-6,750	86.8
Total	458	201 (43.9)	1-6,750	56.8

eata (2), *Acheilognathus lanceolatus* (1), *Acanthorhodeus macrop-terus* (1), *Cobitis lutheri* (1), and *Misgurnus anguillicaudatus* (1) including 18 ones with MsMc in Table 1.

Fishes examined in the middle reaches of Seomjingang

Total 196 freshwater fish (14 species) from Songdaechon in Namwon-si, Jeollabuk-do were examined in 2012 and 2013. Fish species (No. of fish) examined were *C. herzi* (6), *Pseudorasbora parva* (2), *A. springeri* (2), *P. esocinus* (1), *Iksookimia longicorpus* (1), and *Lepomis macrochirus* (1) including 8 ones with MsMc in Table 2. Total 289 freshwater fish (15 species) from Seomjin-gang

Table 2. Infection status of *Metagonimus* spp. metacercariae in freshwater fish from the middle reaches of Seomjin-gang (River)

Locality and fish sp.	No. of fish examined	No. (%) of fish infected	No. of MsMc detected	
			Range	Average
Songdaechon in Namwon-si				
<i>Zacco koreanus</i>	57	56 (98.3)	2-334	60.5
<i>Pungtungia herzi</i>	47	10 (21.3)	1-4	2.0
<i>Zacco platypus</i>	46	45 (97.8)	1-175	28.4
<i>Rhynchocypris oxycephalus</i>	21	1 (4.8)	-	1.0
<i>Microphysogobio koeensis</i>	5	1 (20.0)	-	4.0
<i>Carassius auratus</i>	3	1 (33.3)	-	1.0
<i>Gnathopogon strigatus</i>	2	2 (100)	1-8	4.5
<i>Squalidus gracilis majimae</i>	2	2 (100)	-	1.0
Subtotal	183	118 (64.5)	1-334	39.9
Seomjin-gang in Gokseong-gun				
<i>Zacco koreanus</i>	52	41 (78.9)	1-124	18.0
<i>Zacco platypus</i>	43	43 (100)	1-130	23.4
<i>Coreoleuciscus splendidus</i>	37	15 (40.5)	1-6	1.9
<i>Sarcocheilichthys variegatus</i>	31	21 (67.7)	1-15	3.8
<i>Squalidus japonicus coreanus</i>	27	22 (81.5)	1-10	2.7
<i>Pungtungia herzi</i>	22	13 (59.1)	1-5	2.3
<i>Hemibarbus longirostris</i>	20	19 (95.5)	1-33	12.0
<i>Microphysogobio koeensis</i>	16	8 (50.0)	1-4	2.4
<i>Pseudogobio esocinus</i>	16	13 (81.3)	1-65	12.9
<i>Coreoperca herzi</i>	11	2 (18.2)	2-5	3.5
<i>Acheilognathus majusculus</i>	5	4 (80.0)	3-9	5.0
<i>Plecoglossus altivelis</i>	2	2 (100)	1,370-4,380	2,875
<i>Hemibarbus labeo</i>	2	2 (100)	2-9	5.5
<i>Opsariichthys amurensis</i>	2	2 (100)	11-762	386.5
Subtotal	286	207 (72.4)	1-4,380	43.1
Total	469	325 (69.3)	1-4,380	41.9

in Gokseong-gun, Jeollanam-do were examined in 2015 and 2016. Fish species (No. of fish) examined were *Ladislavia taczanowskii* (1) including 14 ones with MsMc in Table 2.

Fishes examined in the lower reaches of Seomjin-gang

Total 183 freshwater fish (21 species) from Seomjin-gang in Gurye-gun, Jeollanam-do were examined in 2014. Fish species (No. of fish) examined were *C. auratus* (5), *O. platycephala* (4), *A. lanceolatus* (3), and *S. scherzeri* (1) including 17 ones with MsMc in Table 3. Total 156 freshwater fish (13 species) from Hoengcheon in Hadong-gun, Gyeongsangnam-do were examined in 2014 and 2016. Fish species (No. of fish) examined were *C. auratus* (1) and *O. platycephala* (1) including 11 ones with MsMc in Table 3. Total 234 freshwater fish (12 species) from Namsancheon in Hadong-gun, Gyeongsangnam-do were examined in 2015 and 2016. Fish species (No. of fish) examined were *C. auratus* (16), *A. koreensis* (7), *A. gracilis* (2), *C. splendidus* (1), and *O. platycephala* (1) including 7 ones with MsMc in Table 3.

Fishes examined in Tamjin-gang

In the middle reaches in Jangheung-gun, total 1,132 fishes in 22 species were examined for 4 years (2014-2017). Fish species (No. of fish) examined were *Micropterus salmoides* (8), *A. koreensis* (7), and *Cyprinus carpio* (2) including 19 ones with MsMc in Table 5. In the lower reaches in Gangjin-gun, a total of 517 fishes in 17 species were examined in 2014 and 2017. Fish species (No. of fish) examined were *A. lanceolatus* (22), *C. herzi* (3), and *Mugil cephalus* (1) including 14 ones with MsMc in Table 5.

Examination methods

All collected fishes with ice were transferred to the laboratory of the Department of Parasitology and Tropical Medicine, Gyeongsang National University College of Medicine, Jinju, Korea. After the identification of fish species, they were individually ground with a mortar or grinder. Each ground fish meat was mixed with artificial gastric juice and the mixture was incubated at 36°C for 2 hr. The digested material was filtered with 1 × 1 mm of mesh, and washed with 0.85% saline until the supernatant is clear. The sediment was carefully examined under a stereomicroscope. The metacercariae of *Metagonimus* spp. (MsMc) were separately collected by the general feature [13], and they were counted to get hold of infection rates (%) and densities (No. of MsMc per fish infected) by fish species.

Table 3. Infection status of *Metagonimus* spp. metacercariae in freshwater fish from the lower reaches of Seomjin-gang (River)

Locality and fish sp.	No. of fish examined	No. (%) of fish infected	No. of MsMc detected	
			Range	Average
Seomjin-gang in Gurye-gun				
<i>Zacco platypus</i>	24	24 (100)	6-520	126.3
<i>Pungtungia herzi</i>	21	14 (66.7)	1-8	2.6
<i>Squalidus japonicus coreanus</i>	15	12 (80.0)	1-10	3.7
<i>Zacco koreanus</i>	14	11 (78.6)	3-290	64.0
<i>Sarcocheilichthys nigripinnis</i>	14	13 (92.9)	1-52	13.6
<i>Opsariichthys uncirostris</i>	13	13 (100)	52-495	143.8
<i>Acheilognathus rhombeus</i>	11	10 (90.9)	17-150	54.0
<i>Coreoperca herzi</i>	10	4 (40.0)	1-3	1.8
<i>Hemibarbus labeo</i>	10	3 (30.0)	-	1.0
<i>Squalidus gracilis majimae</i>	8	5 (62.5)	1-2	1.4
<i>Pseudogobio esocinus</i>	7	7 (100)	6-62	28.1
<i>Microphysogobio koeensis</i>	6	6 (100)	2-22	6.2
<i>Acanthorhodeus gracilis</i>	5	4 (80.0)	10-58	37.5
<i>Abbottina rivularis</i>	5	1 (20.0)	-	2.0
<i>Plecoglossus altivelis</i>	3	3 (100)	6,280-10,750	8,727
<i>Pseudorasbora parva</i>	3	3 (100)	4-31	15.7
<i>Hemibarbus longirostris</i>	1	1 (100)	-	3.0
Subtotal	170	134 (78.8)	1-10,750	246.5
Hoengcheon in Hadong-gun				
<i>Zacco koreanus</i>	35	35 (100)	1-5,860	369.2
<i>Zacco platypus</i>	27	27 (100)	2-1,510	115.3
<i>Acheilognathus koreensis</i>	20	1 (5.0)	-	1.0
<i>Pungtungia herzi</i>	16	7 (43.8)	1-3	2.1
<i>Coreoperca herzi</i>	11	5 (45.5)	1-2	1.4
<i>Zacco temminckii</i>	10	10 (100)	6-480	156.5
<i>Hemibarbus longirostris</i>	8	8 (100)	2-145	63.0
<i>Pseudogobio esocinus</i>	7	6 (85.7)	4-16	8.5
<i>Abbottina springeri</i>	7	4 (57.1)	3-358	97.8
<i>Coreoleuciscus splendidus</i>	4	3 (75.0)	-	1.0
<i>Squalidus gracilis majimae</i>	1	1 (100)	-	1.0
Subtotal	146	107 (73.3)	1-5,860	173.6
Namsancheon in Hadong-gun				
<i>Zacco koreanus</i>	57	57 (100)	1-2,860	104.3
<i>Pseudogobio esocinus</i>	46	4 (8.7)	1-2	1.3
<i>Zacco platypus</i>	41	41 (100)	3-101	27.4
<i>Pungtungia herzi</i>	35	7 (20.0)	1-7	2.3
<i>Hemibarbus longirostris</i>	14	1 (7.1)	-	1.0
<i>Zacco temminckii</i>	11	11 (100)	2-580	104.5
<i>Squalidus gracilis majimae</i>	3	1 (33.3)	-	1.0
Subtotal	207	122 (58.9)	1-2,860	67.5
Total	523	363 (69.4)	1-10,750	165

RESULTS

Infection status with MsMc in the upper reaches of Seomjin-gang

The metacercariae of *Metagonimus* spp. (MsMc) were detected in 201 (43.9%) out of 458 fishes in 20 species from the upper reaches of Seomjin-gang, i.e., Osucheon in Imsil-gun and Seomjin-gang in Sunchang-gun, Jeollabuk-do, and their average density was 56.8 per fish infected. The infection status by the fish species and surveyed areas was detailedly shown in Table 1.

Infection status with MsMc in the middle reaches of Seomjin-gang

MsMc were detected in 325 (69.3%) out of 469 fishes in 18 species from the middle reaches of Seomjin-gang, i.e., Songdaecheon in Namwon-si, Jeollabuk-do and Seomjin-gang in Gokseong-gun, Jeollanam-do, and their average density was 41.9 per fish infected. The infection status by the fish species and surveyed areas was detailedly revealed in Table 2.

Infection status with MsMc in the lower reaches of Seomjin-gang

MsMc were detected in 363 (69.4%) out of 523 fishes in 21 species from the lower reaches of Seomjin-gang, i.e., Seomjin-gang in Gurye-gun, Jeollanam-do, Hoengcheon and Namsancheon in Hadong-gun, Gyeongsangnam-do, and their average

Table 4. Infection status of *Metagonimus* spp. metacercariae in rasborinid fish from Seomjin-gang (River)

Locality and fish sp.	No. of fish examined	No. (%) of fish infected	No. of MsMc detected	
			Range	Average
Upper reaches				
<i>Zacco platypus</i>	131	93 (71.0)	1-200	14.6
<i>Zacco koreanus</i>	15	11 (73.3)	1-36	7.9
<i>Opsariichthys uncirostris</i>	35	27 (77.1)	1-45	7.6
Subtotal	181	131 (72.4)	1-200	12.6
Middle reaches				
<i>Zacco platypus</i>	89	88 (98.9)	1-175	26.0
<i>Zacco koreanus</i>	109	97 (89.0)	1-334	42.5
<i>Opsariichthys amurensis</i>	2	2 (100)	11-762	386.5
Subtotal	200	187 (93.5)	1-762	38.4
Lower reaches				
<i>Zacco platypus</i>	92	92 (100)	2-1,510	79.0
<i>Zacco koreanus</i>	106	103 (97.2)	1-5,860	190.0
<i>Zacco temminckii</i>	21	21 (100)	2-580	129.2
<i>Opsariichthys amurensis</i>	13	13 (100)	52-495	143.8
Subtotal	232	229 (98.7)	1-5,860	137.2
Total	613	547 (89.2)	1-5,860	73.6

density was 165 per fish infected. The infection status by the fish species and surveyed areas was detailedly shown in Table 3.

Infection status with MsMc in rasborinid fish from Seomjin-gang

MsMc were detected in 547 (89.2%) out of total 613 rasborinid fishes from Seomjin-gang, and their average density was

Table 5. Infection status of *Metagonimus* spp. metacercariae in fishes from Tamjin-gang (River) in Jeollanam-do, Korea

Locality and fish sp.	No. of fish examined	No. (%) of fish infected	No. of MsMc detected	
			Range	Average
Middle reaches in Jangheung-gun				
<i>Zacco temminckii</i>	152	143 (94.1)	1-2,460	65.2
<i>Pungtungia herzi</i>	152	51 (33.6)	1-20	2.4
<i>Zacco platypus</i>	138	110 (79.7)	1-140	16.3
<i>Carassius auratus</i>	116	63 (54.3)	1-924	126.6
<i>Pseudogobio esocinus</i>	86	70 (81.4)	1-86	12.0
<i>Coreoperca kawamebari</i>	84	31 (36.9)	1-61	13.9
<i>Hemibarbus longirostris</i>	77	49 (63.6)	1-45	7.3
<i>Sarcocheilichthys variegatus</i>	54	17 (31.5)	1-7	2.0
<i>Plecoglossus altivelis</i>	52	51 (98.1)	1-5,320	1,339
<i>Acheilognathus lanceolatus</i>	52	1 (1.9)	-	2.0
<i>Odontobutis platycephala</i>	48	7 (14.6)	1-8	3.1
<i>Sarcocheilichthys nigripinnis</i>	29	6 (13.3)	1-32	8.3
<i>Acheilognathus yamatsutae</i>	23	2 (8.7)	4-5	4.5
<i>Coreoperca herzi</i>	16	3 (18.8)	-	1.0
<i>Acanthorhodeus gracilis</i>	11	10 (90.9)	1-55	31.9
<i>Acheilognathus rhombeus</i>	10	6 (60.0)	1-1,400	417.7
<i>Siniperca scherzeri</i>	6	1 (16.7)	-	1.0
<i>Hemiculter eigenmanni</i>	6	2 (33.3)	-	4.0
<i>Squalidus gracilis majimae</i>	3	3 (100)	1-5	2.7
Subtotal	1,115	626 (56.1)	1-5,320	147
Lower reaches in Gangjin-gun				
<i>Zacco platypus</i>	75	70 (93.3)	1-58	8.1
<i>Pungtungia herzi</i>	70	23 (32.9)	1-8	2.4
<i>Carassius auratus</i>	62	54 (87.1)	1-780	38.1
<i>Pseudogobio esocinus</i>	51	41 (80.4)	1-20	5.4
<i>Plecoglossus altivelis</i>	40	40 (100)	26-4,280	841
<i>Sarcocheilichthys nigripinis</i>	33	14 (42.4)	1-17	3.2
<i>Acanthorhodeus macropterus</i>	28	12 (42.9)	1-10	3.2
<i>Zacco temminckii</i>	26	25 (96.2)	1-875	110.3
<i>Odontobutis platycephala</i>	26	5 (19.2)	1-6	2.8
<i>Coreoperca kawamebari</i>	25	9 (36.0)	1-5	1.9
<i>Hemibarbus longirostris</i>	24	19 (79.2)	1-15	4.9
<i>Sarcocheilichthys variegatus</i>	20	3 (15.0)	2-4	3.0
<i>Acheilognathus rhombeus</i>	9	9 (100)	2-67	25.8
<i>Lateolabrax japonicus</i>	2	2 (100)	1-4	2.5
Subtotal	491	326 (66.4)	1-4,280	121
Total	1,606	952 (59.3)	1-5,320	138

73.6 per fish infected. The infection status by the rasborinid fish species and surveyed reaches of river was detailedly revealed in Table 4.

Infection status with MsMc in fishes from Tamjin-gang

MsMc were detected in 626 (56.1%) out of 1,115 fishes in positive fish species from the middle reaches in Jangheung-gun and their average density was 147 per fish infected. MsMc were found in 326 (66.4%) out of 491 fishes in positive fish species from the lower reaches in Gangjin-gun and their average density was 121 per fish infected. The infection status by the fish species and surveyed areas was detailedly revealed in Table 5.

Infection status with MsMc in the susceptible fish species from Tamjin-gang

MsMc were detected in 556 (84.1%) out of total 661 susceptible fishes from Tamjin-gang, and their average density was 227 per fish infected. The infection status by the fish species and surveyed areas was detailedly revealed in Table 6.

DISCUSSION

By the present study, it was confirmed that MsMc are more or less prevalent in fishes from Seomjin-gang and Tamjin-gang. The positive rates with MsMc were very similar, 55.4% and 59.3%, in fishes from 2 rivers, but average metacercarial densities were 96 and 138 per fish infected, higher in fishes from Tamjin-gang. We couldn't compare the endemicity of

Table 6. Infection status of *Metagonimus* spp. metacercariae in susceptible fishes from Tamjin-gang (River) in Jeollanam-do, Korea

Locality and fish sp.	No. of fish examined	No. (%) of fish infected	No. of MsMc detected	
			Range	Average
Tamjin-gang in Jangheung-gun				
<i>Zacco platypus</i>	138	110 (79.7)	1-140	16.3
<i>Zacco temminckii</i>	152	143 (94.1)	1-2,460	65.2
<i>Carassius auratus</i>	116	63 (54.3)	1-924	126.6
<i>Plecoglossus altivelis</i>	52	51 (98.1)	1-5,320	1,339
Subtotal	458	367 (80.1)	1-5,320	238
Tamjin-gang in Gangjin-gun				
<i>Zacco platypus</i>	75	70 (93.3)	1-58	8.1
<i>Zacco temminckii</i>	26	25 (96.2)	1-875	110.3
<i>Carassius auratus</i>	62	54 (87.1)	1-780	38.1
<i>Plecoglossus altivelis</i>	40	40 (100)	26-4,280	841
Subtotal	203	189 (93.1)	1-4,280	205
Total	661	556 (84.1)	1-5,320	227

Table 7. Comparison of the infection status^a with *Metagonimus* spp. metacercariae in fishes from Seomjin-gang and Tamjin-gang

Items	Infection status of MsMc in fishes from		
	Seomjin-gang	Tamjin-gang	Total
No. (%) of fish examined	1,604 (49.3)	1,649 (50.7)	3,253 (100)
Overall positive rate (%)	889/1,604 (55.4)	952/1,649 (57.7)	1,841/3,253 (56.6)
Total metacercarial density	95.5	138.1	117.5
No. (%) ^b of rasborinids examined	613 (38.2)	391 (23.7)	1,004 (30.9)
No. (%) of rasborinids infected	547 (89.2)	348 (89.0)	895 (89.1)
MsMc density in rasborinids	73.6	41.5	61.1

^aPositive rate: No. of fish infected/No. of fish examined×100; metacercarial density: mean No. of MsMc per fish infected.

^bNo. of rasborinid fish/Total No. of fish examined×100.

MsMc in the most susceptible fish species, sweet smelt (*P. altivelis*), so the number of fish examined (8 from Seomjin-gang and 92 from Tamjin-gang) was too much different in 2 rivers. However, their prevalences were 100% and 98.9% and densities were 5,171 and 1,120 per fish infected in sweet smelts from Seomjin-gang and Tamjin-gang each. In another MsMc susceptible fish group, rasborinid fish such as *Zacco* spp. and *O. uncirostris amurensis*, the prevalences were very similar, 89.2% and 89.0%, in 2 rivers, but average metacercarial densities were 74 and 42 per fish infected, more or less higher in Seomjin-gang (Table 7). On the other hand, in the rasborinid fish from Seomjin-gang, the prevalences were 72.4% (the upper reaches), 93.5% (the middle reaches), and 98.7% (the lower reaches), and metacercarial densities were 12.6, 38.4, and 137.2 per fish infected respectively. These findings suggested that the endemicity with MsMc is more higher in fish from the lower reaches than in fish from the upper and middle reaches of Seomjin-gang. In the susceptible fish species, i.e., *Zacco* spp., *C. auratus* and *P. altivelis*, from Tamjin-gang, the prevalence was 84.1% (80.1% in the middle reaches and 93.1% in the lower reaches) and MsMc density was 227 (238 in the middle reaches and 205 in the lower reaches) per fish infected. Therefore, the endemicity with MsMc is similar in fish from 2 surveyed sites of Tamjin-gang.

With regard to the infection status of MsMc in sweet smelts, *P. altivelis*, from Seomjin-gang and Tamjin-gang, at least 4 studies were available [4,21,22,25]. In 1977, Chai et al. [4] reported 100% prevalence and 15,688 MsMc per fish in 20 sweet smelts from a water reservoir nearby Tamjin-gang in Jangheung-gun. Seo et al. [22] detected av. 14,887 MsMc in all 16 sweet smelts from Tamjin-gang in Gangjin-gun. In 1985, Song et al. [21] reported 92.3% prevalence and 636 MsMc per fish in 12 sweet smelts from Tamjin-gang in Gangjin-gun. Cho et al. [25] detected av. 1,037 and 1,511 MsMc per fish in 16 sweet smelts from

Tamjin-gang in Gangjin-gun and in 20 ones from Seomjin-gang in Gurye-gun, Jeollanam-do. Song et al. [21] also reported 2,724 and 2,412 MsMc per fish in each 5 sweet smelts from Seomjin-gang in Gokseong-gun and Gurye-gun, Jeollanam-do. In this study, all 8 sweet smelts from Seomjin-gang were infected with av. 5,171 MsMc, and 91 (98.9%) sweet smelts from Tamjin-gang were infected with 1,120 MsMc per fish infected. From the above findings of present and previous studies, we can suppose that the sweet smelts naturally produced in 2 rivers are highly infected with MsMc and the riverside areas are to be the highly endemic regions of metagonimiasis.

The rasborinid fish, i.e., *Z. platypus*, *Z. koreanus*, *Z. temminckii* and *O. uncirostris amurensis*, are known as the susceptible fish hosts of MsMc in both regions without and/or with sweet smelts in Korea. These fish species are also known as the second intermediate hosts of *M. miyatai* [9,18]. One of them, *Z. platypus*, is the most dominant species in the water systems of Korea. And then the rasborinid fish, especially *Zacco* spp., are highly recommended as the index fish of MsMc to evaluate the endemicity of metagonimiasis. In the present study, total 1004 (30.9%) rasborinid fish (613 from Seomjin-gang: 38.2% and 391 from Tamjin-gang: 23.7%) were examined, and 895 (89.1%) ones (547 in Seomjin-gang: 89.2% and 348 in Tamjin-gang: 89.0%) were to be infected with MsMc. Their density was 61.1 per fish infected (73.6 in Seomjin-gang and 41.5 in Tamjin-gang) (Table 7). In the water systems of Gangwon-do, the prevalences with MsMc were 23.5-100% (85.2% in average) in rasborinid fish and their densities were 3.4-108 (29.7 in average) in 10 surveyed areas [26]. Sohn et al. [27] reported 73.9% and 72.1% MsMc prevalences in the rasborinid fish from the water systems of Hantan-gang and Imjin-gang. They also reported av. 43.1 and 18.6 MsMc densities in the rasborinid fish from 2 surveyed areas [27]. From the aforementioned findings, we can suppose that the endemicities with MsMc in

fishes from Seomjin-gang and Tamjin-gang are more higher than those from the water systems of Gangwon-do, Hantan-gang and Imjin-gang.

Among fishes from Seomjin-gang, some rasborinid fishes, i.e., *Z. koreanus* and *Z. temminckii* from Hoengcheon and Nam-sancheon in Hadong-gun, *O. uncirostris amurensis* from Seomjin-gang in Gurye-gun and *Z. platypus* from Seomjin-gang in Gurye-gun and Hoengcheon in Hadong-gun, were revealed 100% prevalences and more than 100 MsMc densities. In fishes from Tamjin-gang, crasian carp, *C. auratus*, from Jangheung-gun and dark chub, *Z. temminckii*, from Gangjin-gun were highly and heavily infected with MsMc. Among fishes from the water systems of Gangwon-do, 90.0% sea rundace, *T. hakonensis*, from Namdaechon in Yangyang-gun and 97.4% dark chub, *Z. temminckii*, from Joyang-gang in Jeongseon-gun were infected with 449 and 130 MsMc per fish infected [26]. Sohn et al. [27] reported 92.3% and 100% prevalences and 132 and 102 MsMc densities in 26 *P. esocinus* and 22 *Z. platypus* from Hantan-gang in Cheorwon-gun, Gangwon-do. These findings on the high infection status with MsMc will be helpful to perform a study on the experimental metagonimiasis to be needed massive MsMc.

More than 7 valid species have been reported in the genus *Metagonimus* flukes in the literatures. All of them mainly distributed in Asian countries such as Japan, Korea, China and Taiwan [2]. At least 3 human infecting species, i.e., *M. yokogawai*, *M. takahashii* and *M. miyatai*, are known to exist commonly in Japan and Korea [1,2]. They are morphologically differentiated in adult stage by the locations of uterus and vitellaria, and the size of eggs, but not in metacercarial stages in the fish intermediate hosts. The cercariae of *Metagonimus* spp. naturally shed from freshwater snails, *Semisulcospira coreana* and *S. libertina*, which are known as the first intermediate hosts in Korea, and they penetrate into the second intermediate hosts. As the second intermediate hosts of *Metagonimus* spp., numerous species of fish have been reported in Korea. However, some fish species are known as hosts of specific *Metagonimus* species, i.e., *P. altivelis*, *T. hakonensis* and *L. japonicus* for *M. yokogawai* [13-16]; *C. auratus*, *C. carpio*, *T. hakonensis* and *L. japonicus* for *M. takahashii* [13,16,17]; *Z. platypus* and *Z. temminckii* for *M. miyatai* [13,18]. Does each species of *Metagonimus* spp. cercariae have the host-specificity? How does each species of cercariae encyst in the favorable fish hosts only in even highly endemic environment like Seomjin-gang and Tamjin-gang? Studies on the host-specificity of each *Metagonimus* sp. in fish hosts should be

clarified in the near future through the morphological analysis of adult worms recovered from experimental animals, which are infected with MsMc from some susceptible species of fish, i.e., sweet smelt, sea rundace, crasian carp, pale chub and dark chub from a same endemic area.

Conclusively, by the present study, it was confirmed that MsMc are more prevalent in fishes from Seomjin-gang and Tamjin-gang than from any other rivers in Korea. Peoples residing in riverside areas of the 2 rivers should pay attention to the raw consumption of the susceptible fish hosts like chubs, *Zacco* spp., and crasian carp, *C. auratus* as well as sweet smelt, *P. altivelis*.

ACKNOWLEDGMENTS

This study was supported by an anti-communicable diseases control program, 2014E5400200 (Investigation of fish-borne parasites and acquisition of their biological resources in the southern regions of Korea) of National Institute of Health (NIH), Korea Centers for Disease Control and Prevention (KCDCP). We thank Jung-A Kim and Hee-Joo Kim (Department of Parasitology and Tropical Medicine, Gyeongsang National University College of Medicine, Jinju, Korea), for their help in the examination of fish.

CONFLICT OF INTEREST

The authors have no conflicts of interest concerning the work reported in this paper.

REFERENCES

1. Chai JY, Lee SH. Food-borne intestinal trematode infections in the Republic of Korea. *Parasitol Int* 2002; 51: 129-154.
2. Chai JY. *Metagonimus*. In Xiao L, Ryan U, Feng Y eds, *Biology of Foodborne Parasites*. Food Microbiology Series. Boca Raton, USA. CRC Press. pp 427-443.
3. Seo BS, Lee SH, Cho SY, Chai JY, Hong ST, Han IS, Sohn JS, Cho BH, Ahn SR, Lee SK, Chung SC, Kang KS, Shim HS, Hwang IS. An epidemiologic study on clonorchiasis and metagonimiasis in riverside areas in Korea. *Korean J Parasitol* 1981; 19: 137-150.
4. Chai JY, Cho SY, Seo BS. Study on *Metagonimus yokogawai* (Katsurada, 1912) in Korea: IV. An epidemiological investigation along Tamjin River basin, South Cholla Do, Korea. *Korean J Parasitol* 1977; 15: 115-120.
5. Soh CT, Ahn YK. Epidemiological study on *Metagonimus yokogawai* infection along Boseong River in Jeonra Nam Do, Korea. *Korean J Parasitol* 1978; 16: 1-13 (in Korean).

6. Ahn YK, Chung PR, Lee KT, Soh CT. Epidemiological survey on *Metagonimus yokogawai* infection in the eastern coast area of Kangwon province, Korea. *Korean J Parasitol* 1987; 25: 59-68 (in Korean).
7. Chai JY, Han ET, Park YK, Guk SM, Kim JL, Lee SH. High endemicity of *Metagonimus yokogawai* infection among residents of Samchok-shi, Kangwon-do. *Korean J Parasitol* 2000; 38: 33-36.
8. Chai JY, Huh S, Yu JR, Kook J, Jung KC, Park EC, Sohn WM, Hong ST, Lee SH. An epidemiological study of metagonimiasis along the upper reaches of the Namhan River. *Korean J Parasitol* 1993; 31: 99-108.
9. Kim CH, Kim NM, Lee CH, Park JS. Studies on the *Metagonimus* fluke in the Daechong reservoir and the upper stream of Geum river, Korea. *Korean J Parasitol* 1987; 25: 69-82 (in Korean).
10. Park MS, Kim SW, Yang YS, Park CH, Lee WT, Kim CU, Lee EM, Lee SU, Huh S. Intestinal parasite infections in the inhabitants along the Hantan River, Chorwon-gun. *Korean J Parasitol* 1993; 31: 375-378.
11. Ahn YK, Ryang YS. Epidemiological studies on *Metagonimus* infection along the Hongcheon river, Kangwon Province. *Korean J Parasitol* 1988; 26: 207-213 (in Korean).
12. Kino H, Suzuki T, Oishi H, Suzuki S, Yamagiwa S, Ishiguro M. Geographical distribution of *Metagonimus yokogawai* and *M. miyatai* in Shizuoka Prefecture, Japan, and their site preferences in the sweetfish, *Plecoglossus altivelis*, and hamsters. *Parasitol Int* 2006; 55: 201-206.
13. Sohn WM. Fish-borne zoonotic trematode metacercariae in the Republic of Korea. *Korean J Parasitol* 2009; 47 (suppl): 103-113.
14. Chun SK. A study on *Metagonimus yokogawai* from *Plecoglossus altivelis* in the Miryang River. *Bull Pusan Fish Coll* 1960a; 3: 24-32 (in Korean).
15. Ahn YK. *Lateolabrax japonicus*, a role of the second intermediate host of *Metagonimus yokogawai*. *New Med J* 1983; 26: 135-139.
16. Chai JY, Sohn WM, Kim MH, Hong ST, Lee SH. 1991. Three morphological types of the genus *Metagonimus* encysted in the dace, *Tribolodon taczanowskii*, caught from the Sumjin River. *Korean J Parasitol* 1991; 29: 217-225.
17. Kim DG, Kim TS, Cho SH, Song HJ, Sohn WM. Heterophyid metacercarial infections in brackish water fishes from Jinju-man (Bay), Kyongsangnam-do, Korea. *Korean J Parasitol* 2006; 44: 7-13.
18. Saito S, Chai JY, Kim KH, Lee SH, Rim HJ. *Metagonimus miyatai* sp. nov. (Digenea: Heterophyidae), a new intestinal trematode transmitted by freshwater fishes in Japan and Korea. *Korean J Parasitol* 1997; 35: 223-232.
19. Hwang JI, Choi DW. Metacercarial density of *Metagonimus yokogawai* in *Plecoglossus altivelis* in Kyungpook Province, Korea. *Korean J Parasitol* 1977; 15: 30-35.
20. Suh JW, Choi DW. Demonstration of *Metagonimus yokogawai* metacercariae from *Plecoglossus altivelis* in river Ahnseong, Kyungpook Province, Korea. *Korean J Parasitol* 1979; 17: 45-50.
21. Song CY. Studies on the Yokogawa's fluke *Metagonimus yokogawai* (Katsurada, 1912) in Korea. I. Geographical distribution of sweetfish and their infection status with *Metagonimus* metacercariae in Gangwon do. *Chung-Ang J Med* 1981; 6: 121-126 (in Korean).
22. Seo BS, Hong ST, Chai JY, Lee SH. Studies on *Metagonimus yokogawai* (Katsurada, 1912) in Korea. VI. The geographical distribution of metacercarial infection in sweetfish along the East and South coast. *Korean J Parasitol* 1982; 20: 28-32 (in Korean).
23. Song CY, Lee SH, Jeon SR. Studies on the intestinal fluke, *Metagonimus yokogawai* Katsurada, 1912 in Korea IV. Geographical distribution of sweetfish and infection status with *Metagonimus* metacercaria in south-eastern area of Korea. *Korean J Parasitol* 1985; 23: 123-138 (in Korean).
24. Sohn WM, Hong ST, Chai JY, Lee SH. Infection status of sweetfish from Kwangjung-stream and Namdae-stream in Yangyang-gun, Kangwon-do with the metacercariae of *Metagonimus yokogawai*. *Korean J Parasitol* 1990; 28: 253-255 (in Korean).
25. Cho SH, Kim TS, Na BK, Sohn WM. Prevalence of *Metagonimus* metacercariae in sweetfish, *Plecoglossus altivelis*, from eastern and southern coastal areas in Korea. *Korean J Parasitol* 2011; 49: 161-165.
26. Cho SH, Lee WJ, Kim TS, Seok WS, Lee T, Jeong K, Na BK, Sohn WM. Prevalence of zoonotic trematode metacercariae in freshwater fish from Gangwon-do, Korea. *Korean J Parasitol* 2014; 52: 399-412.
27. Sohn WM, Na BK, Cho SH, Lee SW, Choi SB, Seok WS. Trematode metacercariae in freshwater fish from water systems of Hantangang and Imjingang in Republic of Korea. *Korean J Parasitol* 2015; 53: 289-298.
28. Sohn WM, Na BK, Cho SH, Park MY, Kim CH, Hwang MA, No KW, Yoon KB, Lim HC. Prevalence of *Clonorchis sinensis* metacercariae in fish from water systems of Seomjin-gang (River). *Korean J Parasitol* 2017; 55: 305-312.
29. Yoon KB, Lim HC, Jeon DY, Park S, Cho SH, Ju JW, Shin SS, Na BK, Sohn WM. Infection status with *Clonorchis sinensis* metacercariae in fish from Tamjin-gang (River) in Jeollanam-do, Republic of Korea. *Korean J Parasitol* 2018; 56: 183-188.
30. Chai JY, Park JH, Han ET, Shin EH, Kim JL, Guk SM, Hong KS, Lee SH, Rim HJ. Prevalence of *Heterophyes nocens* and *Pygidiopsis summa* infections among residents of the western and southern coastal islands of the Republic of Korea. *Am J Trop Med Hyg* 2004; 71: 617-622.
31. Guk SM, Shin EH, Kim JL, Sohn WM, Hong KS, Yoon CH, Lee SH, Rim HJ, Chai JY. A survey of *Heterophyes nocens* and *Pygidiopsis summa* metacercariae in mullets and gobies along the coastal areas of the Republic of Korea. *Korean J Parasitol* 2007; 45: 205-211.
32. Park JH, Kim JL, Shin EH, Guk SM, Park YK, Chai JY. A new endemic focus of *Heterophyes nocens* and other heterophyid infections in a coastal area of Gangjin-gun, Jeollanam-do. *Korean J Parasitol* 2007; 45: 33-38.