

RESEARCH ARTICLE

Penicillifer diparietisporus: a New Record from Field Soil in Korea


Kallol Das¹, Chang-Gi Back², Seung-Yeol Lee¹, Hee-Young Jung^{1*}¹School of Applied Biosciences, College of Agriculture and Life Sciences, Kyungpook National University, Daegu 41566, Korea²Horticultural and Herbal Crop Environment Division, National Institute of Horticultural and Herbal Science, Wanju 55365, Korea

*Corresponding author: heeyoung@knu.ac.kr

ABSTRACT

A fungus was isolated from field soil collected from Daegu, Korea. The colony of the isolated fungus showed short, branched, and light to dark yellow pigments with hyaline, yellowish red to orange brown aerial mycelia. In addition, the fungus produced solitary to aggregated perithecia, ovoid to pyriform, short neck, and asci as well as biserially arranged ascospores. Phylogenetic analysis using the internal transcribed spacer region and translation elongation factor 1- α sequences and morphological characteristics identified the isolated fungus as *Penicillifer diparietisporus*, which belongs to the family Nectriaceae. To our knowledge, this is the first report of *Penicillifer diparietisporus* in Korea.

Keywords: Nectriaceae, *Penicillifer diparietisporus*, Soil fungi



OPEN ACCESS

pISSN : 0253-651X
eISSN : 2383-5249Kor. J. Mycol. 2018 September, 46(3): 227-233
<https://doi.org/10.4489/KJM.20180029>**Received:** August 11, 2018**Revised:** August 18, 2018**Accepted:** August 18, 2018

© 2018 THE KOREAN SOCIETY OF MYCOLOGY.



This is an Open Access article distributed under the terms of the Creative Commons Attribution-Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Approximately 2,700 fungal species from 240 genera are currently recognized in eight families in the order Hypocreales [1, 2]. The species produce light- to bright-colored, ostiolate, perithecial ascomata containing unitunicate asci with hyaline ascospores. The asexual morphs found in nature are, most frequently, moniliaceous and phialidic [3-7]. The fungal species are globally found in various environments; they are of great importance in agriculture and medicine and extensively exploited for industrial and commercial applications [5]. Moreover, several species have been reported as important opportunistic human pathogens [8-10], whereas others produce mycotoxins that are of medical concern [5].

The purpose of this study was to screen for unreported fungal species in field soil in Korea. This study provides information for further studies on the use of such organisms in agriculture and medicine as well as for industrial applications. On the basis of morphological and molecular characteristics, the strain KNU16-010, identified as *Penicillifer diparietisporus*, was reported for the first time in Korea.

MATERIALS AND METHODS

Collection of soil samples

In 2016, samples were collected from field soil during the screening of fungal species in Daegu, Korea (35°54'11.3"N, 128°36'56.1"E). A sterile trowel and spatula were used to collect the soil samples from a depth of 15~30 cm. The collected soil samples were placed in polythene zipper bags and transferred to the laboratory. Then, the soil samples were prepared for serial dilutions, and 1 g of soil was diluted with 10 mL of sterile distilled water. The serial dilutions were performed until a concentration of 10^{-3} was achieved, and then, 100 μ L of each sample was spread on potato dextrose agar (PDA; Difco, Detroit, MI, USA) plates and incubated for 2~3 days at 25°C. Single germinating fungal colonies were transferred to fresh PDA plates and incubated under the same conditions. The strain KNU16-10 was selected for further morphological and molecular phylogenetic analyses.

Morphological characterization

To study the cultural and morphological characteristics, PDA, oatmeal agar (OA; Difco), and malt extract agar (MEA; Difco) were used; the incubation period was 10 days at 25°C. Then, cultural characteristics such as colony color, fungal growth, and texture were observed and recorded, and the morphological characteristics were observed under a light microscope (BX-50; Olympus, Tokyo, Japan).

Genomic DNA extraction, PCR amplification, and sequencing

Genomic DNA was extracted from 7-day-old colonies of the strain KNU16-010 grown on PDA by using the HiGene Genomic DNA Prep kit (BIOFACT, Daejeon, Korea), according to the manufacturer's instructions. Partial gene sequences were amplified by the internal transcribed spacer (ITS) region, ITS1/ITS4, and translation elongation factor (TEF) 1- α gene, EF1-728F/EF2, as described previously [11-13]. Then, the amplified PCR products were purified with EXOSAP-IT (Thermo Fisher Scientific, Waltham, MA, USA) and sequenced by Solgent (Daejeon, Korea). The similarities of the sequences were analyzed using BLAST of NCBI. The sequences obtained from KNU16-010 were deposited in NCBI GenBank (accession numbers LC387549 and LC387601 for the ITS region and TEF gene, respectively).

Phylogenetic analysis

The consensus sequences were compared with other sequences in the NCBI database by using BLAST to determine the percentage of shared sequence identity with other sequences of fungal species. The alignments were performed using MEGA 7.0 [14] with 1,000 bootstrap replicates, and the evolutionary distance matrices were generated based on Kimura's neighbor-joining algorithm model [15].

RESULTS AND DISCUSSION

Morphology of the strain KNU16-010

To observe the morphological characteristics, the strain KNU16-010 was cultured on PDA, OA, and MEA at 25°C for 10 days, and the diameter of the colonies was 28.7~29.7 mm, 32.0~32.7 mm, and 31.9~32.5 mm, respectively (Fig. 1). The morphology of the strain KNU16-010 was compared with previous descriptions of *Penicillifer diparietisporus* [16] and *Pseudonectria diparietispora* [17] (Table 1). *Pseudonectria diparietispora* produced reddish brown colonies on PDA [17]. Moreover, *Neocosmospora arxii* (basionym: *Pseudonectria diparietispora*) colonies are composed of hyaline or sometimes yellow-pigmented, branched, septate, smooth-walled, hyphae that are 2.5~5 µm wide, and they produce semi-immersed to immersed perithecia, scattered or aggregated in small groups, yellowish orange to yellowish brown, broadly ovoid to somewhat pyriform, 220~340 × 160~270 µm in diameter, almost glabrous, conical shaped with short neck [18]. However, our strain formed orange brown to yellowish red colonies in all the media, and PDA was used to observe the mycological characteristics (Fig. 1). The fungal colonies showed wavy mycelia at the edges and rough wart-like structures. They also formed septate, irregular, and short hyphae, and the average width of the hyphae was 2.42 µm. They produced submerged perithecia, solitary to aggregated, yellowish orange to yellowish brown, ovoid to pyriform, smooth, conical shaped with short neck (Fig. 2A, 2C). Asci clavate, 70.9~81.1 × 16.3~19.7 µm, thin wall, and 8-spored (Fig. 2D, 2E). Ascospores were oval, septate, guttulate, 2-celled, distichous, double wall, biseriolate, tubular, hyaline and pale green, smooth and blunt at both ends with a diameter (n = 30) of 17.9~30.7 × 6.7~10.9 µm and no conidia (Fig. 2F, 2G). The ascospores were distichous, elliptical, continuous, hyaline, with thick, double wall, smooth 18~24 × 12~14 µm and no




Fig. 1. Cultural characteristics of KNU16-010. A, B, Front and reverse sides of the colony on potato dextrose agar; C, D, Front and reverse sides of the oatmeal agar; E, F, Front and reverse sides of the malt extract agar.




Fig. 2. Morphological characteristics of KNU16-010. A, B, Mature perithecia on potato dextrose agar; C, Perithecium; D, E, Asci; F, G, Ascospores (scale bars: A, B = 500 μ m, C = 50 μ m, D-G = 10 μ m).

Table 1. Morphological characteristics of the strain KNU16-010 with reference to *Penicillifer diparietisporus*

Characteristics		KNU16-010 ^a	<i>Penicillifer diparietisporus</i> ^b	<i>Pseudonectria diparietispora</i> ^c
Colony	Color	Yellowish to orange and dark brown in reverse on PDA. Light yellow in center, white edge and brown color in reverse on MEA.	Reddish white edge, changing to greenish yellow on OA and yellowish brown to brown, light brownish orange to dark brown on PDA. Dark brown color in reverse both media.	Reddish-brown color on PDA and colonies composed of tangled masses of conidial chains.
	Size (Diam.)	PDA: 28.7~29.7 mm; OA: 32.0~32.7 mm and MEA: 31.9~32.5 mm in 10 days at 25°C.	PDA: 22mm; OA:30 mm in 10 days at 25°C; MEA: N/A	N/A
	Shape	Edge margin irregular, wavy and inadequate aerial mycelium.	Edges submerged into irregular margin and scant of aerial mycelium.	Lightly floccose or plane and appressed and thin.
Asci	Size (Diam.) and Shape	Clavate, 70.9~81.1 \times 16.3~19.7 μ m, thin wall and 8-spored.	Clavate, 60~85 \times 12~25 μ m, unitunicate and 8-spored.	Broadly clavate to spindle shaped with very thin wall, short stalked, 60~80 \times 19~22 μ m and 8-spored.
Ascospores	Size (Diam.)	17.9~30.7 \times 6.7~10.9 μ m.	21~25 \times 12~15 μ m.	18~24 \times 12~14 μ m.
	Shape and Color	Oval, septate, distichous, double wall, tubular, hyaline to pale green, smooth and blunt in both ends.	Biseriate, thick walled, broadly ellipsoidal, surrounded by a thin walled sheath collapsing at maturity and sometimes giving the ascospores roughened appearance.	Distichous, elliptical, continuous, hyaline, with thick, double wall, smooth.
Perithecia		Submerged, solitary to aggregated, yellowish orange to yellowish brown, ovoid to pyriform, smooth, conical shaped and short neck.	Solitary to densely aggregated, superficial, globose, ovoid to pyriform, 270~300 \times 240~270 μ m diameter, brown, red-orange to orange.	Superficial, transparent wall, globose-conoid with very short neck, smooth, yellowish-red, 400~500 \times 300~400 μ m.

PDA, potato dextrose agar; MEA, malt extract agar; OA, oatmeal agar; Diam., diameter; N/A, not available in previous references.

^aFungal strain studied in this paper.

^bSources of the descriptions [16].

^cSources of the descriptions [17].

conidia [17]. Thus, these morphological characteristics of the strain KNU16-010 were very similar to those of *Penicillifer diparietisporus*.

Notes: Currently, the genus *Viridispota* is composed of four species, *Viridispota diparietispora* (= *Penicillifer furcatus*), *Viridispota alata* (= *Penicillifer bipapillatus*), *Viridispota fragariae* (= *Penicillifer fragariae*), and *Viridispota penicilliferi* (= *Penicillifer macrosporus*), each with its own *Penicillifer* asexual morphs [7, 16, 19, 20]. Lombard et al. [21] reported that the epithet *Pseudonectria diparietispora* (1957) pre-dates that of *Penicillifer furcatus* (1991), and the basionyms are provided below.

Penicillifer diparietisporus (J.H. Miller, Giddens & A.A. Foster) Rossman, L. Lombard & Crous, comb. nov.

Basionym: *Pseudonectria diparietispora* J.H. Miller, Giddens & A.A. Foster, *Mycologia* 49:793. 1957.

≡ *Neocosmospora diparietispora* (J.H. Miller, Giddens & A.A. Foster) Rossman, Samuels & Lowen, *Mycologia* 85:699. 1993.

≡ *Viridispota diparietispora* (J.H. Miller, Giddens & A.A. Foster) Samuels & Rossman, *Stud. Mycol.* 42:167. 1999.

= *Neocosmospora arxii* Udagawa, Horie & P. Cannon, *Sydowia* 41:353. 1989.

= *Neocosmospora endophytica* Polishook, Bills & Rossman, *Mycologia* 83:798. 1991.

= *Penicillifer furcatus* Polishook, Bills & Rossman, *Mycologia* 83:798. 1991.

Molecular phylogeny of the strain KNU16-010

Genetic sequences of the ITS regions and TEF1- α were analyzed to determine the evolutionary relationships (Fig. 3) with the strains obtained from GenBank (Table 2). After analyzing the nucleotide sequences, 523 bp and 438 bp sequences were obtained from the ITS regions and TEF1- α , respectively. The BLAST results showed 100% similarity with *Penicillifer diparietisporus* (CBS 376.59) and 99% with *Penicillifer diparietisporus* (KM231861) in the ITS regions and TEF1- α , respectively. The combined BLAST results suggested that the strain KNU16-010 showed 100% similarity with the strain *Viridispota diparietispora* (= *Penicillifer diparietisporus*) (CBS 102797) and *Penicillifer diparietisporus* (CBS 376.59, NR154310). The phylogenetic tree was constructed on the basis of the ITS regions and TEF1- α to the strain *Penicillifer diparietisporus* [21]. The strain KNU16-010 was closely clustered with *Penicillifer diparietisporus* (CBS 376.59), with a 100% bootstrap value (Fig. 3). The phylogenetic results support the fact that the strain KNU16-010 is *Penicillifer diparietisporus*.

Members of the family Nectriaceae are pleomorphic fungi that display both asexual and sexual morphs during their life cycles [22]. In this study, we used molecular analysis, which provides a considerable challenge to conventional fungal systematics. Thus, more genomic studies of members of the family Nectriaceae are urgently required. To our knowledge, this is the first report of *Penicillifer diparietisporus* isolated from field soil in Korea.




Fig. 3. Neighbor-joining phylogenetic tree based on the combined internal transcribed spacer sequences and translation elongation factor 1- α . *Pseudonectria buxi* CBS 324.56 was used as an outgroup. The strain isolated in this study is indicated in bold, and the bootstrap values are based on 1,000 replications. Bar, 0.02 substitutions per nucleotide position.

Table 2. GenBank accession numbers used in this study for the phylogenetic analyses

Species	Strains	GenBank accession numbers	
		ITS	TEF
<i>Penicillifer bipapillatus</i>	CBS 420.88	KM231740	KM231860
<i>Penicillifer diparietisporus</i>	CBS 376.59	NR 154310	KM231861
<i>Penicillifer macrosporus</i>	CBS 423.88	KM231739	KM231859
<i>Penicillifer pulcher</i>	CBS 560.67	KM231742	KM231862
<i>Penicillifer diparietisporus</i>	KNU16-010	LC387549	LC387601
<i>Pseudonectria buxi</i>	CBS 324.56	KM231778	KM231909

ITS, internal transcribed spacer; TEF, translation elongation factor.

ACKNOWLEDGEMENTS

This research was supported by a grant from the National Institute of Biological Resources (NIBR), funded by the Ministry of Environment (MOE) of the Republic of Korea for the project on survey and discovery of indigenous fungal species.

REFERENCES

1. Kirk PM, Cannon PF, Minter DW, Stalpers JA, Ainsworth GC, Bibsy GR. Ainsworth & Bisby's dictionary of the fungi. 10th ed. Wallingford: CAB International; 2008.
2. Crous PW, Shivas RG, Quaedvlieg W, van der Bank M, Zhang Y, Summerell BA, Guarro J, Wingfield MJ, Wood AR, Alfenas AC, et al. Fungal planet description sheets: 214-280. *Persoonia* 2014;32:184-306.

3. Rogerson CT. The hypocrealean fungi (Ascomycetes, Hypocreales). *Mycologia* 1970;62:865-910.
4. Samuels GJ, Seifert KA. Taxonomic implications of variation among hypocrealean anamorphs. In: Sugiyama J, editor. *Pleomorphic fungi: the diversity and its taxonomic implications*. Amsterdam: Elsevier Science; 1987. p. 29-56.
5. Rossman AY. Morphological and molecular perspectives on systematics of the Hypocreales. *Mycologia* 1996;88:1-19.
6. Rossman AY. Towards monophyletic genera in the holomorphic Hypocreales. *Stud Mycol* 2000;45:27-34.
7. Rossman AY, Samuels GJ, Rogerson CT, Lowen R. Genera of Bionectriaceae, Hypocreaceae and Nectriaceae (Hypocreales, Ascomycetes). *Stud Mycol* 1999;42:1-248.
8. Chang DC, Grant GB, O'Donnell K, Wannemuehler KA, Noble-Wang J, Rao CY, Jacobson LM, Crowell CS, Sneed RS, Lewis FM et al. Multistate outbreak of *Fusarium keratitis* associated with use of a contact lens solution. *JAMA* 2006;296:953-63.
9. de Hoog GS, Guarro J, Gene J, Figueras MJ. *Atlas of clinical fungi* (CD-ROM). 3rd ed. Utrecht: CBS-KNAW Fungal Biodiversity Centre; 2011.
10. Guarro J. Fusariosis, a complex infection caused by a high diversity of fungal species refractory to treatment. *Eur J Clin Microbiol Infect Dis* 2013;32:1491-500.
11. White TJ, Bruns T, Lee S, Taylor J. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ, editors. *PCR protocols: a guide to methods and applications*. San Diego: Academic Press; 1990. p. 315-22.
12. Carbone I, Kohn LM. A method for designing primer sets for speciation studies in filamentous ascomycetes. *Mycologia* 1999;91:553-6.
13. O'Donnell K, Kistler HC, Cigelnik E, Ploetz RC. Multiple evolutionary origins of the fungus causing panama disease of banana: concordant evidence from nuclear and mitochondrial gene genealogies. *Proc Natl Acad Sci USA* 1998;95:2044-9.
14. Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol* 2016;33:1870-4.
15. Kimura M. A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* 1980;16:111-20.
16. Polishook JD, Bills GF, Rossman AY. A new species of *Neocosmospora* with a *Penicillifer* anamorph. *Mycologia* 1991;83:797-804.
17. Miller JH, Giddens JE, Foster AA. A survey of the fungi of forest and cultivated soils of Georgia. *Mycologia* 1957;49:779-808.
18. Udagawa SI, Horie Y, Cannon PF. Two new species of *Neocosmospora* from Japan with a key to the currently accepted species. *Sydowia* 1989;41:349-59.
19. Samuels GJ. *Nectria* and *Penicillifer*. *Mycologia* 1989;81:347-55.
20. Watanabe T. Three new *Nectria* species from Japan. *Trans Mycol Soc Jpn* 1990;31:227-36.
21. Lombard L, van der Merwe NA, Groenewald JZ, Crous PW. Generic concepts in Nectriaceae. *Stud Mycol* 2015;80:189-245.
22. Cannon PF, Kirk PM. The philosophy and practicalities of amalgamating anamorph and teleomorph concepts. *Stud Mycol* 2000;45:19-25.