

## A FIXED POINT APPROACH TO THE STABILITY OF 3-LIE HOMOMORPHISMS AND 3-LIE DERIVATIONS

SIRILUK PAOKANTA<sup>a</sup> AND JUNG RYE LEE<sup>b,\*</sup>

ABSTRACT. Using the fixed point method, we prove the Hyers-Ulam stability of 3-Lie homomorphisms and 3-Lie derivations in 3-Lie algebras for Cauchy-Jensen functional equation.

### 1. INTRODUCTION AND PRELIMINARIES

The stability problem of functional equations originated from a question of Ulam [15] concerning the stability of group homomorphisms. Hyers [8] gave a first affirmative partial answer to the question of Ulam for Banach spaces. Hyers' Theorem was generalized by Aoki [1] for additive mappings and by Rassias [14] for linear mappings by considering an unbounded Cauchy difference. A generalization of the Rassias theorem was obtained by Găvruta [7] by replacing the unbounded Cauchy difference by a general control function in the spirit of Rassias' approach.

A Lie algebra is a Banach algebra endowed with the Lie product

$$[x, y] := \frac{(xy - yx)}{2}.$$

Similarly, a 3-Lie algebra is a Banach algebra endowed with the product

$$[[x, y], z] := \frac{[x, y]z - z[x, y]}{2}.$$

Let  $A$  and  $B$  be two 3-Lie algebras. A  $\mathbb{C}$ -linear mapping  $H : A \rightarrow B$  is called a 3-Lie homomorphism if

$$H([[x, y], z]) = [[H(x), H(y)], H(z)]$$

---

Received by the editors July 19, 2019. Accepted November 04, 2019

2010 *Mathematics Subject Classification*. Primary 39B52, 47H10, 39B82, 22D25, 17A40.

*Key words and phrases*. Jensen functional equation, fixed point method, 3-Lie algebra, 3-Lie homomorphisms, 3-Lie derivation, Hyers-Ulam stability.

\*Corresponding author.

for all  $x, y, z \in A$ . A  $\mathbb{C}$ -linear mapping  $D : A \rightarrow A$  is called a 3-Lie derivation if

$$D\left([[x, y], z]\right) = [[D(x), y], z] + [[x, D(y)], z] + [[x, y], D(z)]$$

for all  $x, y, z \in A$  (see [17]).

We recall a fundamental result in fixed point theory.

**Theorem 1.1** ([2, 5]). *Let  $(X, d)$  be a complete generalized metric space and let  $J : X \rightarrow X$  be a strictly contractive mapping with Lipschitz constant  $\alpha < 1$ . Then for each given element  $x \in X$ , either*

$$d(J^n x, J^{n+1} x) = \infty$$

for all nonnegative integers  $n$  or there exists a positive integer  $n_0$  such that

- (1)  $d(J^n x, J^{n+1} x) < \infty$ ,  $\forall n \geq n_0$ ;
- (2) the sequence  $\{J^n x\}$  converges to a fixed point  $y^*$  of  $J$ ;
- (3)  $y^*$  is the unique fixed point of  $J$  in the set  $Y = \{y \in X \mid d(J^{n_0} x, y) < \infty\}$ ;
- (4)  $d(y, y^*) \leq \frac{1}{1-\alpha} d(y, Jy)$  for all  $y \in Y$ .

In 1996, Isac and Rassias [9] were the first to provide applications of stability theory of functional equations for the proof of new fixed point theorems with applications. By using fixed point methods, the stability problems of several functional equations have been extensively investigated by a number of authors (see [3, 4, 6, 12, 13]).

Throughout this paper, we suppose that  $A$  and  $B$  are two 3-Lie algebras. For convenience, we use the following abbreviation for a given mapping  $f : A \rightarrow B$

$$D_\mu f(x, y, z) := f\left(\frac{\mu x + \mu y}{2} + \mu z\right) + f\left(\frac{\mu x + \mu z}{2} + \mu y\right) + f\left(\frac{\mu y + \mu z}{2} + \mu x\right) \\ - 2\mu f(x) - 2\mu f(y) - 2\mu f(z)$$

for all  $\mu \in \mathbb{T}^1 := \{\lambda \in \mathbb{C} : |\lambda| = 1\}$  and all  $x, y, z \in A$ .

Throughout this paper, assume that  $A$  is a 3-Lie algebra with norm  $\|\cdot\|$  and that  $B$  is a 3-Lie algebra with norm  $\|\cdot\|$ .

## 2. STABILITY OF 3-LIE HOMOMORPHISMS IN 3-LIE ALGEBRAS

We need the following lemmas which have been given in for proving the main results.

**Lemma 2.1** ([16]). *Let  $X$  be a uniquely 2-divisible abelian group and  $Y$  be linear space. A mapping  $f : X \rightarrow Y$  satisfies*

$$(2.1) \quad f\left(\frac{x+y}{2} + z\right) + f\left(\frac{x+z}{2} + y\right) + f\left(\frac{y+z}{2} + x\right) = 2[f(x) + f(y) + f(z)]$$

for all  $x, y, z \in X$  if and only if  $f : X \rightarrow Y$  is additive.

**Lemma 2.2** ([10]). *Let  $X$  and  $Y$  be linear spaces and let  $f : X \rightarrow Y$  be a mapping such that*

$$(2.2) \quad D_\mu f(x, y, z) = 0$$

for all  $\mu \in \mathbb{T}^1$  and all  $x, y, z \in A$ . Then the mapping  $f : X \rightarrow Y$  is  $\mathbb{C}$ -linear.

Using the fixed point method, we investigate the Hyers-Ulam stability of 3-Lie homomorphisms in 3-Lie algebras associated to the functional equation (2.1).

**Theorem 2.3.** *Let  $\varphi : A^3 \rightarrow [0, \infty)$  be a function such that there exists an  $L < 1$  with*

$$(2.3) \quad \varphi(x, y, z) \leq 2L\varphi\left(\frac{x}{2}, \frac{y}{2}, \frac{z}{2}\right)$$

for all  $x, y, z \in A$ . Suppose that  $f : A \rightarrow B$  is a mapping satisfying

$$(2.4) \quad \|D_\mu f(x, y, z)\| \leq \varphi(x, y, z),$$

$$(2.5) \quad \|f([[x, y], z]) - [[f(x), f(y)], f(z)]\| \leq \varphi(x, y, z)$$

for all  $\mu \in \mathbb{T}^1$  and all  $x, y, z \in A$ . Then there exists a unique 3-Lie homomorphism  $H : A \rightarrow B$  such that

$$(2.6) \quad \|f(x) - H(x)\| \leq \frac{1}{6(1-L)}\varphi(x, x, x)$$

for all  $x \in A$ .

*Proof.* Letting  $\mu = 1$  and  $x = y = z$  in (2.4), we get

$$(2.7) \quad \|3f(2x) - 6f(x)\| \leq \varphi(x, x, x)$$

for all  $x \in A$ .

Consider the set

$$S := \{g : A \rightarrow B, \quad g(0) = 0\}$$

and introduce the generalized metric on  $S$ :

$$d(g, h) = \inf \{ \mu \in \mathbb{R}_+ : \|g(x) - h(x)\| \leq \mu\varphi(x, x, x), \quad \forall x \in A \},$$

where, as usual,  $\inf \phi = +\infty$ . It is easy to show that  $(S, d)$  is complete (see [11]).

Now we consider the linear mapping  $J : S \rightarrow S$  such that

$$Jg(x) := \frac{1}{2}g(2x)$$

for all  $x \in A$ .

It follows from (2.7) that

$$\left\| g(x) - \frac{1}{2}g(2x) \right\| \leq \frac{1}{6}\varphi(x, x, x)$$

for all  $x \in A$ .

Let  $g, h \in S$  be given such that  $d(g, h) = \varepsilon$ . Then

$$\|g(x) - h(x)\| \leq \varepsilon\varphi(x, x, x)$$

for all  $x \in A$ . Since

$$\left\| \frac{1}{2}g(2x) - \frac{1}{2}h(2x) \right\| \leq \frac{1}{2}\varepsilon\varphi(2x, 2x, 2x) \leq L\varepsilon\varphi(x, x, x)$$

for all  $x \in A$ ,  $d(Jg, Jh) \leq L\varepsilon$ . This means that

$$d(Jg, Jh) \leq Ld(g, h)$$

for all  $g, h \in S$ .

It follows from (2.7) that  $d(f, Jf) \leq \frac{1}{6}$ .

By Theorem 1.1, there exists a mappings  $H : A \rightarrow A$  satisfying the following:

(1)  $H$  is a fixed point of  $J$ , i.e.,

$$(2.8) \quad H(x) = \frac{1}{2}H(2x)$$

for all  $x \in A$ . The mapping  $H$  is a unique fixed point of  $J$ . This implies that  $H$  is a unique mapping satisfying (2.8) such that there exists a  $\mu \in (0, \infty)$  satisfying

$$\|f(x) - H(x)\| \leq \mu\varphi(x, x, x)$$

for all  $x \in A$ ;

(2)  $d(J^l f, H) \rightarrow 0$  as  $l \rightarrow \infty$ . This implies the equality

$$\lim_{l \rightarrow \infty} \frac{1}{2^l} f(2^l x) = H(x)$$

for all  $x \in A$ ;

(3)  $d(f, H) \leq \frac{1}{1-L}d(f, Jf)$ , which implies

$$\|f(x) - H(x)\| \leq \frac{1}{6(1-L)}\varphi(x, x, x)$$

for all  $x \in A$ . Thus we get the inequality (2.6).

It follows from (2.3) that

$$\begin{aligned} \|D_\mu H(x, y, z)\| &= \lim_{n \rightarrow \infty} \frac{1}{2^n} \|D_\mu f(2^n x, 2^n y, 2^n z)\| \\ &\leq \lim_{n \rightarrow \infty} \frac{1}{2^n} \varphi(2^n x, 2^n y, 2^n z) = 0 \end{aligned}$$

for all  $x, y, z \in A$  and all  $\mu \in \mathbb{T}^1$ . So  $D_\mu H(x, y, z) = 0$  for all  $\mu \in \mathbb{T}^1$  and all  $x, y, z \in A$ . By Lemma 2.2, the mapping  $H : A \rightarrow B$  is  $\mathbb{C}$ -linear.

It follows from (2.5) that

$$\begin{aligned} &\|H([x, y], z) - [[H(x), H(y)], H(z)]\| \\ &= \lim_{n \rightarrow \infty} \frac{1}{8^n} \|f([2^n x, 2^n y], 2^n z) - [[f(2^n x), f(2^n y)], f(2^n z)]\| \\ &\leq \lim_{n \rightarrow \infty} \frac{1}{8^n} \varphi(2^n x, 2^n y, 2^n z) \leq \lim_{n \rightarrow \infty} \frac{1}{2^n} \varphi(2^n x, 2^n y, 2^n z) = 0 \end{aligned}$$

for all  $x, y, z \in A$ . Thus

$$H([x, y], z) = [[H(x), H(y)], H(z)]$$

for all  $x, y, z \in A$ .

Therefore, the mapping  $H : A \rightarrow B$  is a 3-Lie homomorphism. □

**Corollary 2.4.** *Let  $\theta, r$  be positive real numbers with  $r < 1$ . Suppose that  $f : A \rightarrow B$  is a mapping such that*

$$(2.9) \quad \|D_\mu f(x, y, z)\| \leq \theta(\|x\|^r + \|y\|^r + \|z\|^r),$$

$$(2.10) \quad \|f([x, y], z) - [[f(y), f(z)], f(x)]\| \leq \theta(\|x\|^r + \|y\|^r + \|z\|^r)$$

for all  $\mu \in \mathbb{T}^1$  and all  $x, y, z \in A$ . Then there exists a unique 3-Lie homomorphism  $H : A \rightarrow B$  such that

$$(2.11) \quad \|f(x) - H(x)\| \leq \frac{\theta}{2 - 2^r} \|x\|^r$$

for all  $x \in A$ .

*Proof.* The proof follows from Theorem 2.3 by taking  $L = 2^{r-1}$  and  $\varphi(x, y, z) = \theta(\|x\|^r + \|y\|^r + \|z\|^r)$  for all  $x, y, z \in A$ . □

**Theorem 2.5.** *Let  $\varphi : A^3 \rightarrow [0, \infty)$  be a function such that there exists an  $L < 1$  with*

$$(2.12) \quad \varphi\left(\frac{x}{2}, \frac{y}{2}, \frac{z}{2}\right) \leq \frac{L}{2} \varphi(x, y, z)$$

for all  $x, y, z \in A$ . Suppose that  $f : A \rightarrow B$  is a mapping such that

$$\|D_\mu f(x, y, z)\|_B \leq \varphi(x, y, z),$$

$$\|f([[x, y], z]) - [[f(x), f(y)], f(z)]\| \leq \varphi(x, y, z)$$

for all  $\mu \in \mathbb{T}^1$  and all  $x, y, z \in A$ . Then there exists a unique 3-Lie homomorphism  $H : A \rightarrow B$  such that

$$(2.13) \quad \|f(x) - H(x)\| \leq \frac{L}{6(1-L)}\varphi(x, x, x)$$

for all  $x \in A$ .

*Proof.* Let  $(S, d)$  be the generalized metric space defined in the proof of Theorem 2.3.

Now we consider the linear mapping  $J : S \rightarrow S$  such that

$$Jg(x) := 2g\left(\frac{x}{2}\right)$$

for all  $x \in A$ .

It follows from (2.7) that

$$\left\|f(x) - 2f\left(\frac{x}{2}\right)\right\| \leq \frac{1}{3}\varphi\left(\frac{x}{2}, \frac{x}{2}, \frac{x}{2}\right) \leq \frac{L}{6}\varphi(x, x, x)$$

for all  $x \in A$ .

The rest of the proof is similar to the proof of Theorem 2.3.  $\square$

**Corollary 2.6.** *Let  $\theta r$  and  $q_3$  be non-negative real numbers with  $r > 1$ . Suppose that  $f : A \rightarrow B$  is a mapping satisfying (2.9) and (2.10). Then there exists a unique 3-Lie homomorphism  $H : A \rightarrow B$  such that*

$$(2.14) \quad \|f(x) - H(x)\| \leq \frac{\theta}{2^r - 2}\|x\|^r$$

for all  $x \in A$ .

*Proof.* The proof follows from Theorem 2.5 by taking  $L = 2^{1-r}$  and  $\varphi(x, y, z) = \theta(\|x\|^r + \|y\|^r + \|z\|^r)$  for all  $x, y, z \in A$ .  $\square$

### 3. STABILITY OF 3-LIE DERIVATIONS ON 3-LIE ALGEBRAS

Using the fixed point method, we investigate the Hyers-Ulam stability of 3-Lie derivations in 3-Lie algebras associated to the functional equation (2.1).

**Theorem 3.1.** *Let  $\varphi : A^3 \rightarrow [0, \infty)$  be a function satisfying (2.3). Suppose that  $f : A \rightarrow A$  is a mapping satisfying*

$$\|D_\mu f(x, y, z)\| \leq \varphi(x, y, z),$$

$$(3.1) \quad \|f([[x, y], z]) - [[f(x), y], z] - [[x, f(y)], z] - [[x, y], f(z)]\| \leq \varphi(x, y, z)$$

for all  $\mu \in \mathbb{T}^1$  and all  $x, y, z \in A$ . Then there exists a unique 3-Lie derivation  $D : A \rightarrow A$  satisfying (2.6).

*Proof.* By the proof of Theorem 2.3, there exists a unique  $\mathbb{C}$ -linear mapping  $D : A \rightarrow A$  satisfying (2.6) and

$$D(x) := \lim_{n \rightarrow \infty} \frac{1}{2^n} f(2^n x)$$

for all  $x \in A$ . It follows from (3.1) that

$$\begin{aligned} & \|D([[x, y], z]) - [[D(x), y], z] - [[x, D(y)], z] - [[x, y], D(z)]\| \\ &= \lim_{n \rightarrow \infty} \frac{1}{8^n} \|f([[2^n x, 2^n y], 2^n z]) - [[f(2^n x), 2^n y], 2^n z] - [[2^n x, f(2^n y)], 2^n z] - [[2^n x, 2^n y], f(2^n z)]\| \\ &\leq \lim_{n \rightarrow \infty} \frac{1}{8^n} \varphi(2^n x, 2^n y, 2^n z) = 0 \end{aligned}$$

for all  $x, y, z \in A$ . So

$$D([[x, y], z]) = [[D(x), y], z] + [[x, D(y)], z] + [[x, y], D(z)]$$

for all  $x, y, z \in A$ . Therefore, the mapping  $D : A \rightarrow A$  is a 3-Lie derivation.  $\square$

**Corollary 3.2.** *Let  $\theta, r$  be positive real numbers with  $r < 1$ . Suppose that  $f : A \rightarrow A$  is a mapping satisfying (2.9) and (2.10). Then there exists a unique 3-Lie derivation  $D : A \rightarrow A$  satisfying (2.11).*

**Theorem 3.3.** *Let  $\psi : A^3 \rightarrow [0, \infty)$  be a function satisfying (2.12). Suppose that  $f : A \rightarrow A$  is a mapping satisfying*

$$\|D_\mu f(x, y, z)\| \leq \psi(x, y, z),$$

$$\|f([[x, y], z]) - [[f(x), y], z] - [[x, f(y)], z] - [[x, y], f(z)]\| \leq \psi(x, y, z)$$

for all  $\mu \in \mathbb{T}^1$  and all  $x, y, z \in A$ . Then there exists a unique 3-Lie derivation  $D : A \rightarrow A$  satisfying (2.13).

*Proof.* By the proof of Theorem 2.5, there exists a unique  $\mathbb{C}$ -linear mapping  $D : A \rightarrow A$  satisfying (2.13) and

$$D(x) := \lim_{n \rightarrow \infty} 2^n f\left(\frac{x}{2^n}\right)$$

for all  $x \in A$ .

The rest of proof is similar to the proof Theorem 3.1. □

**Corollary 3.4.** *Let  $\theta, r$  be non-negative real numbers with  $r > 1$ . Suppose that  $f : A \rightarrow B$  is a mapping satisfying (2.9) and (2.10). Then there exists a unique 3-Lie derivation  $D : A \rightarrow A$  satisfying (2.14).*

#### ACKNOWLEDGMENTS

This work was supported by Daejin University.

#### REFERENCES

1. T. Aoki: On the stability of the linear transformation in Banach spaces. *J. Math. Soc. Japan* **2** (1950), 64-66.
2. L. Cădariu & V. Radu: Fixed points and the stability of Jensen's functional equation. *J. Inequal. Pure Appl. Math.* **4**, no. 1, Art. ID 4 (2003).
3. ———: On the stability of the Cauchy functional equation: a fixed point approach. *Grazer Math. Ber.* **346** (2004), 43-52.
4. ———: Fixed point methods for the generalized stability of functional equations in a single variable. *Fixed Point Theory Appl.* **2008**, Art. ID 749392 (2008).
5. J. Diaz & B. Margolis: A fixed point theorem of the alternative for contractions on a generalized complete metric space. *Bull. Am. Math. Soc.* **74** (1968), 305-309.
6. I. EL-Fassi: Generalized hyperstability of a Drygas functional equation on a restricted domain using Brzdek's fixed point theorem. *J. Fixed Point Theory Appl.* **19** (2017), 2529-2540.
7. P. Găvruta: A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings. *J. Math. Anal. Appl.* **184** (1994), 431-436.
8. D.H. Hyers: On the stability of the linear functional equation, *Proc. Nat. Acad. Sci. U.S.A.* **27** (1941), 222-224.
9. G. Isac & Th.M. Rassias: Stability of  $\psi$ -additive mappings: Applications to nonlinear analysis. *Internat. J. Math. Math. Sci.* **19** (1996), 219-228.
10. V. Keshavarz, S. Jahedi, S. Aslani, J. Lee & C. Park: On the stability of 3-Lie homomorphisms and 3-Lie derivations. *J. Comput. Anal. Appl.* (in press).
11. D. Miheţ & V. Radu: On the stability of the additive Cauchy functional equation in random normed spaces. *J. Math. Anal. Appl.* **343** (2008), 567-572.
12. C. Park: Fixed point method for set-valued functional equations. *J. Fixed Point Theory Appl.* **19** (2017), 2297-2308.



13. V. Radu: The fixed point alternative and the stability of functional equations. *Fixed Point Theory* **4** (2003), 91–96.
14. Th.M. Rassias: On the stability of the linear mapping in Banach spaces. *Proc. Am. Math. Soc.* **72** (1978), 297-300.
15. S.M. Ulam: *A Collection of the Mathematical Problems*. Interscience Publ. New York, 1960.
16. A. Najati & A. Ranjbari: Stability of homomorphisms for a 3D CauchyJensen type functional equation on  $C^*$ -ternary algebras. *J. Math. Anal. Appl.* **341** (2008), 62-79.
17. H. Yuan & L. Chen: Lie  $n$  superderivations and generalized Lie  $n$  superderivations of superalgebras. *Open Math.* **16** (2018), 196-209.

<sup>a</sup>DEPARTMENT OF MATHEMATICS, RESEARCH INSTITUTE FOR NATURAL SCIENCES, HANYANG UNIVERSITY, SEOUL 04763, KOREA  
*Email address:* siriluk22@hanyang.ac.kr

<sup>b</sup>DEPARTMENT OF MATHEMATICS, DAEJIN UNIVERSITY, KYEONGGI 11159, KOREA  
*Email address:* jrlee@daejin.ac.kr