

RESEARCH

Open Access



Ecophysiology of seed dormancy and germination in four *Lonicera* (Caprifoliaceae) species native to Korea

HyungBin Park^{1†}, ChungHo Ko^{1†}, SeungYoun Lee^{1,2†}, SangYong Kim¹, JongCheol Yang¹ and KiCheol Lee^{1*}

Abstract

Background: To exploit the ornamental and medicinal purposes of *Lonicera harae* Makino, *L. subsessilis* Rehder, *L. praeflorens* Batalin, and *L. insularis* Nakai, native to Korea, it is necessary to understand their seed ecology for propagation. In this study, we investigated the seed dormancy type and germination characteristics of seeds of the four Korean native *Lonicera* species.

Results: The seeds of the four *Lonicera* species imbibed water readily, suggesting that the species do not have physical dormancy. Furthermore, the seeds exhibited underdeveloped embryos with only about 15–25% of the length of the seeds at dispersal. The embryos grew to the critical length with approximately 50–80% of the length of the seeds' development before radicle protrusion. Further, 94.4% and 61.1% of freshly matured seeds of *L. insularis* and *L. harae* germinated within 4 weeks after sowing at 15 °C and 20 °C, respectively. Contrarily, *L. praeflorens* and *L. subsessilis* seeds did not germinate within 4 weeks under all temperature treatments. At 15 °C, *L. praeflorens* seeds started to germinate from 5 weeks and the final germination rate was 51.1% at 13 weeks. At 15 °C, *L. subsessilis* seeds started to germinate from 5 weeks after sowing and the final germination rate was 85.6% at 17 weeks after sowing. Embryo growth and germination of *L. praeflorens* and *L. subsessilis* occurred at a relatively high temperature (≥ 15 °C).

Conclusions: Overall, *L. insularis* seeds have only morphological dormancy. The seeds of *L. harae* have approximately 60% and 40% of morphological dormancy and morphophysiological dormancy, respectively. Contrarily, *L. praeflorens* and *L. subsessilis* exhibited non-deep simple-type morphophysiological dormancy that requires relatively high temperature (≥ 15 °C) for embryo growth and dormancy breaking. The optimum temperature for the germination of seeds of *L. insularis*, *L. harae*, *L. praeflorens*, and *L. subsessilis* was 15 °C, 20 °C, 15 °C, and 20 °C, respectively. There was interspecific variation in seed dormancy and germination patterns in the four *Lonicera* species. The difference in these characteristics within the four *Lonicera* species could be useful for understanding the seed ecophysiological mechanisms of *Lonicera* species.

Keywords: Ecological adaption, *Lonicera* species, Morphological dormancy, Morphophysiological dormancy, Non-deep simple morphophysiological dormancy

* Correspondence: yloim@korea.kr

[†]Hyung Bin Park, Chung Ho Ko and Seung Youn Lee contributed equally to this work.

¹Division of Plant Resources, Korea National Arboretum, Yangpyeong 12519, South Korea

Full list of author information is available at the end of the article



Background

The genus *Lonicera* of the family Caprifoliaceae includes more than 200 species that are mostly arching shrubs or small trees cultivated as ornamental crops (Theis et al. 2008). *Lonicera* is mostly distributed in the Northern Hemisphere such as North America, Europe, and Asia (Naugžemys et al. 2007). Not only the plants, known as “honeysuckles,” are useful resources for ornamental and medicinal purposes, but also their extracts can be used as a herbal medicine for inflammation in China (Theis et al. 2008; Ryuk et al. 2012; Yuan et al. 2012; Kim et al. 2016). There are 17 *Lonicera* species that are native to Korea (Lee 2003). Especially, *L. insularis* Nakai and *L. subsessilis* Rehder are Korean endemic plants that are distributed along the shore of Ulleungdo and from Pyeongannam-do to Jeollanam-do in Korea, respectively (Jeong et al. 2014; KNA 2019). To exploit these plant resources, investigating germination characteristics and seed dormancy type is needed.

Dormant seeds are viable seeds that do not germinate for a period of time even under environmental conditions that are favorable for germination, and thus, they can avoid unfavorable environmental conditions (Finch-Savage and Leubner-Metzger 2006). Seed dormancy is divided into the following five categories: (1) physical dormancy (PY), (2) physiological dormancy (PD), (3) morphological dormancy (MD), (4) morphophysiological dormancy (MPD), and (5) combinational dormancy (PY+PD) (Baskin and Baskin 1998). Among them, seeds with MD and MPD have underdeveloped embryos during seed dispersal (Nikolaeva 1977; Baskin and Baskin 1998).

Seeds with MD have underdeveloped embryo and these seeds are not physiologically dormant (Baskin and Baskin 1998). Embryo of the seeds with MD must grow to a critical length before radicle protrusion. However, the seeds with MD do not require pretreatments for dormancy breaking; they only need time to grow full-sized embryo before radicle protrusion (Baskin and Baskin 2004a). Morphophysiological dormancy is a combination of MD and PD. Thus, MPD require time not only to grow embryo to a critical size, but also to break PD by cold and/or warm stratification. Morphophysiological dormancy in seeds is classified into eight types according to requirements of environmental conditions for seed dormancy breaking and embryo growth and germination responses to gibberellic acid (Baskin and Baskin 2004a).

From an ecological perspective, seed dormancy plays an important role in how germination timing is controlled in a natural habitat and how plants adapt to their natural environments (Geneve 2003). Thus, seed dormancy and germination requirements can provide insights into how plants determine where they can grow (Santiago et al. 2012). Furthermore, because the type of

seed dormancy varies within the genus and species, classification of seed dormancy type among species in the same genus provides insights into how each species adapts to different natural habitats. In a previous study, the seeds of four *Lonicera* species native to North America were investigated to classify seed dormancy type (Hidayati et al. 2000b). The seeds of these species had MPD or MD or a combination of MD and MPD. *Lonicera caerulea* var. *emphylocalyx* has only MD (Phartyal et al. 2009). On the contrary, *L. fragrantissima* has deep simple MPD and *L. japonica* has non-deep simple MPD (Baskin and Baskin 1998). Half of the seeds of *L. maackii* and *L. morrowii* have MD and the other half has non-deep simple MPD (Hidayati et al. 2000b; Santiago et al. 2012). However, seed dormancy and germination characteristics of Korean native *Lonicera* species have not been studied. The aim of the present study was to investigate seed dormancy type and germination characteristics of four *Lonicera* species native to Korea. These results will be useful to understand seed ecology and to identify optimum temperature treatments for propagating these *Lonicera* species.

Materials and methods

Seed source

To investigate the seed dormancy type and germination characteristics of the four *Lonicera* species, the fruits of these species were collected from 2016 to 2017. The seeds of *L. insularis* were collected in June 2016 from plants growing in Ulleungdo, Korea. The seeds of *L. subsessilis* Rehder were collected in September 2016 from plants growing in the Korea National Arboretum, Pocheon, Korea. The seeds of *L. harae* Makino and *L. praeflorens* Batalin were collected in May 2017 from plants growing in the Korea National Arboretum, Pocheon, Korea. The seeds were removed from the pulp and dried at ambient room temperature (approximately 25 °C) for 1–2 weeks, packed and sealed plastic containers, and stored at 4 °C until further analyses.

Water imbibition test

To investigate the PY of the seeds, a water imbibition test was conducted on March 4, 2018. Three replicates of 20 seeds each were used. The dry matter of the seeds was measured, and then the seeds were placed in 9-cm-diameter Petri dishes (Cell Culture Dish; SPL Life Sciences Co., Ltd., Gyeonggi-do, Korea) with two layers of filter papers (Whatman No.2; GE Healthcare, Buckinghamshire, UK) moistened with distilled water. The seeds were incubated at room temperature (approximately 25 °C). The fresh weight of seeds was measured after 3, 6, 9, 12, 24, and 48 h of incubation. The water uptake by seeds was calculated using the water uptake formula (Baskin et al. 2004b).

Water absorption (%) = $[(W_2 - W_1)/W_1] \times 100$

where, W_2 is the mass of the seeds after imbibition for a given interval and W_1 is the initial seed mass.

Seed morphology

To investigate MD, the seed morphology measurement was carried out on April 5, 2018 to June 12, 2018. The seeds of *L. insularis* and *L. praeflorens* were incubated at 15 °C, whereas the seeds of *L. harae* and *L. subsessilis* were incubated at 20 °C. The seeds were halved using a razor blade (stainless blade; Dorco, Seoul, KR), and then the length of the embryo of the seeds at dispersal and just before germination was measured using a USB microscope (AM 3111 Dino-Lite Premier; AnMo Electronics Co., Taiwan). Thereafter, the embryo/seed ratio (E:S ratio) was calculated and compared between seed dispersal and just before germination.

Temperature treatments

To investigate the seed dormancy type and optimum temperatures for germination, temperature experiments were carried out from May 25, 2017, to December 8, 2017. Three replicates with 30 seeds were used. The seeds were placed on two sheets of filter papers moistened with distilled water in 9-cm-diameter Petri dishes. All Petri dishes were sealed with Parafilm (PM-996; Bemis Company Inc., USA) to reduce water loss during incubation. Temperature and light-controlled multi-room chambers (WIM-R L4; Daihan scientific Co. Ltd., Wonju, Korea) were used in this experiment. The growth chambers were set at constant temperatures of 5 °C, 15 °C, 20 °C and 25 °C and 12-h light/dark photoperiod using cool white fluorescent lamps, producing a photon flux density of approximately 15–20 $\mu\text{mol}^{-2} \text{s}^{-1}$.

For *L. subsessilis*, seasonal temperature cycling treatment known as move-along experiment was performed from May 25, 2017 to February 10, 2018 (Baskin and Baskin 2003). In the move-along experiment, one set (move A treatment) of seeds was placed at 5 °C. After 12 weeks, they were moved from 5 °C to 15 °C (4 weeks) → 20 °C (4 weeks) → 25 °C (12 weeks). The second set (move B treatment) of seeds was placed initially at 25 °C. After 12 weeks, they were moved from 25 °C to 20 °C (4 weeks) → 15 °C (4 weeks) → 5 °C (12 weeks). The germinated seeds were counted every week and removed from the Petri dishes. Sterilized water was frequently supplied to the Petri dishes to maintain moist condition. Rotten seeds were removed and excluded from the calculation of germination rate.

Statistical analysis

Statistical Analysis System (SAS) version 9.4 (SAS Institute Inc., Cary, NC, USA) was used for statistical analyses of the data. Differences between the mean final

germination rate of seed under four temperature treatments were assessed using Tukey's honestly significant difference tests. Differences between the E:S ratio of seeds at seed dispersal and just before germination were assessed using paired *t* tests. The results with *p* values < 0.05 were considered statistically significant.

Results

Water imbibition test

The mass of the seeds of *L. insularis*, *L. harae*, *L. praeflorens*, and *L. subsessilis* was increased by approximately 50%, 90%, 35%, and 40% after 3 h of water imbibition and more than 60%, 170%, 50%, and 70% after 48 h of water imbibition, respectively (Fig. 1).

Seed morphology

The seeds of the four *Lonicera* species has undeveloped embryo at dispersal (Table 1). The E:S ratio of the seeds of *L. insularis*, *L. praeflorens*, *L. harae*, and *L. subsessilis* at dispersal was 0.26 ± 0.02 , 0.13 ± 0.07 , 0.14 ± 0.01 , and 0.16 ± 0.01 mm, respectively. The E:S ratio of the seeds of the four *Lonicera* species just before germination was 0.82 ± 0.04 , 0.81 ± 0.02 , 0.46 ± 0.13 , and 0.74 ± 0.01 mm, respectively (Fig. 4).

Temperature treatments

The seeds of the four *Lonicera* species presented different germination characteristics under the four temperature treatments (Fig. 2). The seeds of the four *Lonicera* species did not germinate at 5 °C. At 15 °C, the freshly matured seeds of *L. insularis* and *L. harae* germinated to 94.4% and 61.1% at 4 weeks after sowing, respectively. The final germination rate of the seeds of *L. harae* was 81.1% at 23 weeks after sowing at 15 °C (Fig. 2). On the contrary, the seeds of *L. praeflorens* and *L. subsessilis* did not germinate at the four temperature regimes within 4 weeks after sowing at all temperature regimes. At 15 °C, the seeds of *L. praeflorens* and *L. subsessilis* started to germinate from 5 weeks after sowing and the final germination rate of the seeds was 51.1% and 80.0% at 13 and 17 weeks after sowing, respectively (Fig. 2). At 20 °C, the seeds of *L. insularis* and *L. harae* germinated to 88.9% and 61.1% at 4 weeks after sowing, respectively, and the final germination rate of *L. harae* was 97.8% at 16 weeks after sowing (Fig. 2). At 20 °C, the seeds of *L. praeflorens* and *L. subsessilis* started to germinate from 8 and 9 weeks after sowing, respectively, and the final germination rate of the seeds was 40.0% and 85.6% at 19 and 14 weeks after sowing, respectively (Fig. 2). At 25 °C, the final germination rate of *L. insularis*, *L. harae*, and *L. subsessilis* was 73.3%, 67.8%, and 76.7% at 9, 27, and 27 weeks after sowing, respectively. Contrarily, the seeds of *L. praeflorens* did not germinate at 25 °C during incubation (Fig. 2).




Fig. 1 Water uptake by intact seeds of *Lonicera insularis* Nakai, *Lonicera praeflorens* Batalin, *Lonicera harae* Makino, and *Lonicera subsessilis* Rehder with increase in mass. Seeds were incubated at ambient room temperature (approximately 25 °C) on filter papers moistened with distilled water for 48 h. The vertical error bars represent SE ($n = 3$)

Move-along experiments

In the move-along experiment of *L. subsessilis*, the seeds of move A treatment (start of winter) did not germinate until 24 weeks after sowing. Germination was initiated from 5 weeks after moving the seeds from 20 °C to 25 °C, and the final germination rate of the seeds was 74.4% at 29 weeks after sowing (Fig. 3). The seeds of move B treatment (start of summer) started to germinate from 3 weeks after moving the seeds from 25 °C to 20 °C, and the final germination rate of the seeds was 83.3% at 16 weeks after sowing. The final germination rate of the seeds of move B treatment was higher than that of the seeds of move A treatment (Fig. 3).

Discussion









In general, the germination of seeds with PY is prevented by water impermeability of the seeds or fruit coats due to the presence of water-impermeable palisade layers of cells (Baskin and Baskin 2004b). If the mass of seeds increases to $\leq 20\%$ in the water imbibition test, the seeds are considered impermeable to water (Baskin and Baskin 2003). This dormancy can be broken by mechanical or chemical scarification such as dry heating, dipping in boiling water, and treating with sulfuric acid (Baskin et al. 2004; Finch-Savage and Leubner-Metzger 2006; De Souza et al. 2012). In the present study, the seeds of the four *Lonicera* species imbibed water readily, suggesting that the species have no PY.

Several previous studies have reported that the seeds of *Lonicera* species have undeveloped embryo at dispersal (Martin 1946; Hidayati et al. 2000a; Hidayati et al. 2000b). In this study, we found that the embryo of the four *Lonicera* species was small and undeveloped, and they grew to the critical length before radicle protrusion. In general, underdeveloped embryo requires time to grow to the critical length. This indicates that the seeds of the species have MD or MPD (Baskin and Baskin 1998; Baskin and Baskin 2004b).

According to Baskin and Baskin (2004b), seeds with MD germinate within about 30 days. Thus, the seeds of *L. insularis* have only MD. On the contrary, approximately 60% and 40% of the seeds of *L. harae* exhibited MD and MPD, respectively. In the case of *L. praeflorens*, the germination of seeds was initiated at a relatively high temperature (15 °C) from 5 weeks after sowing. At a high temperature (25 °C), the seeds of *L. praeflorens* did not germinate during incubation. At 15 °C, the seeds of *L. subsessilis* started to germinate from 5 weeks after sowing and the final germination rate was 80.0% at 17 weeks after sowing. At 20 °C, the seeds started to germinate from 9 weeks after sowing and the final germination rate was 85.6% at 14 weeks after sowing. The seeds of *L. praeflorens* and *L. subsessilis* did not germinate within 4 weeks after sowing (Fig. 2). Thus, the seeds of these two species have MPD.


Morphophysiological dormancy is subdivided into eight types (Nikolaeva 1977; Baskin and Baskin 1998;

Table 1 Internal seed morphology of four *Lonicera* species native to Korea. Undeveloped embryos at dispersal and fully developed embryos just before germination are shown. *pr* pericarp, *sc* seed coat, *en* endosperm, *em* embryo. Scale bars represent 1 mm

| Species | Seed morphology | |
|-------------------------------------|---|---|
| | At dispersal | Just before germination |
| <i>Lonicera insularis</i> Nakai |  |  |
| <i>Lonicera praeflorens</i> Batalin |  |  |
| <i>Lonicera harae</i> Makino |  |  |
| <i>Lonicera subsessilis</i> Rehder |  |  |

Walck et al. 2002; Baskin and Baskin 2004b). It is broadly divided into the following two categories: (1) simple-type MPD that requires relatively high temperature ($\geq 15^\circ\text{C}$) for embryo growth and (2) complex-type MPD that requires only low temperature ($0\text{--}10^\circ\text{C}$) for embryo growth (Baskin and Baskin 1998; Hidayati et al. 2000b). The simple and complex MPD can be sub-divided based on the level of PD: non-deep, intermediate, and deep. Non-deep simple MPD requires

warm or cold stratification to break dormancy. Intermediate and deep simple MPD require warm stratification followed by cold stratification to break PD (Nikolaeva 1977; Baskin and Baskin 1998; Hidayati et al. 2000b). In a previous study, *L. fragrantissima*, *L. japonica*, *L. maackii*, and *L. morrowii* have been reported to have MPD. *L. fragrantissima* had deep simple MPD and required warm plus cold stratification to break PD. *L. japonica*, *L. maackii*, and *L. morrowii* had non-deep simple



MPD that required stratification (warm or cold) to break PD, in which about 50% of the seeds of *L. maackii* and *L. morrowii* had only MD (Hidayati et al. 2000b). Approximately 50% of the seeds of *L. praeflorens* require only relatively high temperature (15 °C) to grow embryo and germinate without cold stratification, whereas approximately 50% did not germinate during the experiment. Thus, approximately 50% of the seeds of *L. praeflorens* is estimated to have non-deep simple-type MPD. Because approximately 50% of the seeds of *L. praeflorens* did not germinate in this study, further studies are needed to accurately classify dormancy type in ungerminated seeds. Germination and embryo growth of *L. subsessilis* seeds occurred only at relatively high temperature (≥ 15 °C) without cold stratification. In the move-along experiment, the seeds of *L. subsessilis* germinated from 3 weeks after moving them from 25 to 20 °C in move B treatment (start of summer), whereas the seeds in move A treatment (start of winter) germinated from 5 weeks after moving them from 20 to 25 °C. The result of move B treatment was similar to that at 25 °C (Fig. 3). The seeds of move A treatment did not germinate until 12 weeks after sowing at 5 °C, and the seeds started to germinate after exposure to relatively high temperature. Thus, 85.6% of *L. subsessilis* seeds is also considered to have non-deep simple MPD. The seeds of the four *Lonicera* species showed a variation in time to

germination. Various states of dormancy within the same seed population result in a slow germination rate which is an ecologically advantageous strategy for unpredictable environmental conditions (Doussi and Thanos 2002).

In a previous study, the dispersal pattern of four *Lonicera* seeds corresponded with their dormancy and germination patterns (Hidayati et al. 2000b). The seeds of *L. insularis* are dispersed from late July to August (late summer) in the natural environment. In the present study, they were exposed to temperatures that are considered optimum (15–20 °C) for germination. Because the dispersed seeds have only MD, they will germinate within 30 days after dispersal; however, if the seeds are dispersed under high temperature conditions (≥ 25 °C), their germination will be delayed. The seeds of *L. harae* are dispersed from June. When the seeds are exposed to relatively high temperature (20 °C), which was considered optimum for germination in this study, approximately 60% of the seeds will germinate within 30 days after dispersal as they have approximately 60% MD and the remaining will germinate within 16 weeks. The seeds of *L. praeflorens* begin to disperse from June, and then they are exposed to warm stratification. Because the optimum temperature to germinate is 15 °C, than 20 °C, germination of the dispersed seeds can be delayed. The seeds of *L. subsessilis* are dispersed from late September




Fig. 3 Cumulative germination rate of *Lonicera subsessilis* Rehder seeds in the move-along experiment. Move A started at 5°C (5°C (12 weeks) → 15°C (4 weeks) → 20°C (4 weeks) → 25°C (12 weeks)). Move B started at 25°C (25°C (12 weeks) → 20°C (4 weeks) → 15°C (4 weeks) → 5°C (12 weeks)). The vertical bars represent SE ($n = 3$)

to early October (in autumn). They might be exposed to relatively high temperatures before germination. Thus, they will germinate via exposure to relatively high temperature ($\geq 15^\circ\text{C}$) in the early autumn. If they are exposed to low temperatures, germination will be delayed until they are exposed to a relatively high temperature ($\geq 15^\circ\text{C}$), as evidenced in seeds of the move A treatment (start of winter) in the move-along experiment.

In the same genus and species, seed dormancy can vary according to the native habitat (Hidayati et al. 2000a). There was stasis of ecophysiological traits such as seed dormancy and germination and ecological divergence in seed dormancy and germination characteristics

of the disjunct species of the genera *Viburnum* and *Siphisia* (Adams et al. 2005; Walck et al. 2012). The disjunct species *Osmorhiza* and *Erythronium* have different seed dormancy. *Osmorhiza chilensis*, *Osmorhiza occidentalis*, and *Erythronium grandiflorum* have deep complex MPD, whereas *Osmorhiza longistylis*, *Osmorhiza claytonii*, *Erythronium albidum*, and *Erythronium americanum* have non-deep complex MPD (Baskin and Baskin 1984; Baskin and Baskin 1991; Baskin et al. 1995). In the present study, the four *Lonicera* species exhibited common morphological traits such as undeveloped embryo. However, there was ecological divergence in seed dormancy and germination characteristics.




Fig. 4 Embryo/seed ratio (E:S ratio) in the seeds of *Lonicera insularis* Nakai, *Lonicera praeflorens* Batalin, *Lonicera harae* Makino, and *Lonicera subsessilis* Rehder at seed dispersal and at just before germination. The seeds of *L. insularis*, *L. praeflorens*, *L. harae*, and *L. subsessilis* were incubated at 15 °C, 15 °C, 20 °C, and 25 °C, respectively. The vertical error bars represent SE ($n = 3$). Each E:S ratio at dispersal and just before germination was compared using paired t test. *** $p < 0.001$

Conclusion

The results of this study revealed that the seeds of *L. insularis* have only MD and the seeds of *L. harae* have approximately 60% MD and 40% MPD. On the contrary, the seeds of *L. praeflorens* and *L. subsessilis* have non-deep simple-type MPD that requires only relatively high temperature (≥ 15 °C) for embryo growth and dormancy breaking. The optimum temperature for the germination of seeds of *L. insularis*, *L. harae*, *L. praeflorens*, and *L. subsessilis* was 15 °C, 20 °C, 15 °C, and 20 °C, respectively (Fig. 4). There was interspecific variation in seed dormancy and germination in the four *Lonicera* species. Our results will be useful for understanding seed eco-physiological mechanisms in a habitat and propagating *Lonicera* species.

Abbreviations

E:S ratio: Embryo/seed ratio; MD: Morphological dormancy; MPD: Morphophysiological dormancy; PD: Physiological dormancy; PY: Physical dormancy

Acknowledgements

We thank Dohyun Kim and Kyusung Choi for the assistance in the field sample collections and observations. We also thank the anonymous reviewers for providing helpful comments on this paper.

Authors' contributions

PHB, GCH, and LSY analyzed the data and wrote the manuscript. LKC designed the study and analyzed the data. YJC and KSY designed the study and coordinated the overall research. All authors read and approved the final manuscript.

Funding

This study was funded by the project "Useful Plant Collection and Development of Mass Propagation Protocol for the Establishment of Foundation of Convergence Platform using Forest Plants (KNA1-2-25, 16-3)" funded by the Korea National Arboretum.

Availability of data and materials

The datasets used and/or analyzed in the present study are available from Author 1 (phb1274@korea.kr) on reasonable request.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Division of Plant Resources, Korea National Arboretum, Yangpyeong 12519, South Korea. ²Division of Horticulture & Medicinal Plant, Andong National University, Andong 36729, South Korea.

Received: 4 April 2019 Accepted: 20 May 2019

Published online: 21 June 2019

References

- Adams CA, Baskin JM, Baskin CC. Trait stasis versus adaptation in disjunct relict species: evolutionary changes in seed dormancy-breaking and germination requirements in a subclade of *Aristolochia* subgenus *Siphisia* (Piperales). *Seed Sci Res.* 2005;15:161–73.
- Baskin CC, Baskin JM. *Seeds: ecology, biogeography, and evolution of dormancy and germination*. San Diego: Academic; 1998.
- Baskin CC, Baskin JM. When breaking seed dormancy is a problem try a move-along experiment. *Nat Plants J.* 2003;4:17–21.

- Baskin CC, Baskin JM. Germination seeds of wildflowers, an ecological perspective. *HortTechnology*. 2004a;14:67–73.
- Baskin CC, Meyer SE, Baskin JM. Two types of morphophysiological dormancy in seeds of two genera (*Osmorhiza* and *Erythronium*) with an Arcto-Tertiary distribution pattern. *Am J Bot*. 1995;82:293–8.
- Baskin JM, Baskin CC. Germination ecophysiology of the woodland herb *Osmorhiza longistylis* (Umbelliferae). *Am J Bot*. 1984;71:687–92.
- Baskin JM, Baskin CC. Nondeep complex morphophysiological dormancy in seeds of *Osmorhiza claytonii* (Apiaceae). *Am J Bot*. 1991;78:588–93.
- Baskin JM, Baskin CC. A classification system for seed dormancy. *Seed Sci Res*. 2004b;14:1–16.
- Baskin JM, Davis BH, Baskin CC, Gleason SM, Cordell S. Physical dormancy in seeds of *Dodonaea viscosa* (Sapindales, Sapindaceae) from Hawaii. *Seed Sci Res*. 2004;14:81–90.
- De Souza TV, Voltolini CH, Santos M, Paulilo MTS. Water absorption and dormancy-breaking requirements of physically dormant seeds of *Schizolobium parahyba* (Fabaceae–Caesalpinioideae). *Seed Sci Res*. 2012;22:169–76.
- Doussi MA, Thanos CA. Ecophysiology of seed germination in Mediterranean geophytes. 1. *Muscari* spp. *Seed Sci Res*. 2002;12:193–201.
- Finch-Savage WE, Leubner-Metzger G. Seed dormancy and the control of germination. *New Phytol*. 2006;171:501–23.
- Geneve RL. Impact of temperature on seed dormancy. *HortScience*. 2003;38:336–40.
- Hidayati SN, Baskin JM, Baskin CC. Dormancy-breaking and germination requirements for seeds of *Diervilla lonicera* (Caprifoliaceae), a species with underdeveloped linear embryos. *Can J Bot*. 2000a;78:1199–205.
- Hidayati SN, Baskin JM, Baskin CC. Dormancy-breaking and germination requirements of seeds of four *Lonicera* species (Caprifoliaceae) with underdeveloped spatulate embryos. *Seed Sci Res*. 2000b;10:459–69.
- Jeong KS, Kim MS, Lee W, Park JH. Intraspecific variation and geographic study of *Lonicera insularis* (Caprifoliaceae) based on chloroplast DNA sequences. *Kor J Pl Taxon*. 2014;44:202–7.
- Kim SM, Won YH, Jeong K, Kim MJ, Chun WJ, Yang HJ, Kwon YS. Chemical constituents of *Lonicera maackii* leaves. *Kor J Pharmacogn*. 2016;47:117–21.
- Korea National Arboretum. 2019. <http://www.nature.go.kr>. Accessed 25 Jan 2019.
- Lee TB. Coloured flora of Korea. Hyangmunsa, Seoul; 2003.
- Martin AC. The comparative internal morphology of seeds. *Am Midl Naturalist*. 1946;36:513–660.
- Naugžemys D, Žilinskaitė S, Denkovskij J, Patamsytė J, Literiskis J, Žvingila D. RAPD based study of genetic variation and relationships among *Lonicera* germplasm accessions. *Biologija*. 2007;53:34–9.
- Nikolaeva MG. Factors controlling the seed dormancy pattern. In: *The physiology and biochemistry of seed dormancy and germination* (ed. A. A. Khan). Amsterdam: North-Holland Publishing. 1977. p. 51–74.
- Phartyal SS, Kondo T, Baskin JM, Baskin CC. Temperature requirements differ for the two stages of seed dormancy break in *Aegopodium podagraria* (Apiaceae), a species with deep complex morphophysiological dormancy. *Am J Bot*. 2009;96:1086–95.
- Ryuk JA, Lee HW, Ko BS. Discrimination of *Lonicera japonica* and *Lonicera confusa* using chemical analysis and genetic marker. *Kor J Herbology*. 2012;27:15–21.
- Santiago A, Herranz JM, Copete E, Ferrandis P. Species-specific environmental requirements to break seed dormancy: implications for selection of regeneration niches in three *Lonicera* (Caprifoliaceae) species. *Bot*. 2012;91:225–33.
- Theis N, Donoghue MJ, Li J. Phylogenetics of the Caprifoliaceae and Lonicera (Dipsacales) based on nuclear and chloroplast DNA sequences. *Syst Bot*. 2008;33:776–83.
- Walck JL, Hidayati SN, Okagami N. Seed germination ecophysiology of the Asian species *Osmorhiza aristata* (Apiaceae): comparison with its North American congeners and implications for evolution of types of dormancy. *Am J Bot*. 2002;89:829–35.
- Walck JL, Karlsson LM, Milberg P, Hidayati SN, Kondo T. Seed germination and seedling development ecology in world-wide populations of a circumboreal Tertiary relict. *AoB plants*. 2012;2012:pls007.
- Yuan Y, Song L, Li M, Liu G, Chu Y, Ma L, Zhou Y, Wang X, Gao W, Qin S, Yu J, Wang X, Huang L. Genetic variation and metabolic pathway intricacy govern the active compound content and quality of the Chinese medicinal plant *Lonicera japonica* thunb. *BMC Genomics*. 2012;13:195.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

