

Review

## Metabolic Engineering for Improved Fermentation of L-Arabinose

Suji Ye, Jeong-won Kim, and Soo Rin Kim\*

School of Food Science and Biotechnology, Kyungpook National University, Daegu 41566, Republic of Korea

Received: December 10, 2018  
Revised: January 20, 2019  
Accepted: January 29, 2019

First published online  
February 8, 2019

\*Corresponding author  
Phone: +53-950-7769;  
Fax: +53-950-7762;  
E-mail: soorinkim@knu.ac.kr

pISSN 1017-7825, eISSN 1738-8872

Copyright© 2019 by  
The Korean Society for Microbiology  
and Biotechnology

L-Arabinose, a five carbon sugar, has not been considered as an important bioresource because most studies have focused on D-xylose, another type of five-carbon sugar that is prevalent as a monomeric structure of hemicellulose. In fact, L-arabinose is also an important monomer of hemicellulose, but its content is much more significant in pectin (3–22%, g/g pectin), which is considered an alternative biomass due to its low lignin content and mass production as juice-processing waste. This review presents native and engineered microorganisms that can ferment L-arabinose. *Saccharomyces cerevisiae* is highlighted as the most preferred engineering host for expressing a heterologous arabinose pathway for producing ethanol. Because metabolic engineering efforts have been limited so far, with this review as momentum, more attention to research is needed on the fermentation of L-arabinose as well as the utilization of pectin-rich biomass.

**Keywords:** L-Arabinose, pectin, metabolic engineering, *Saccharomyces cerevisiae*, pentose

### Significance of L-Arabinose as a Bioresource

Bioconversion of lignocellulosic biomass such as agricultural residues and wood waste materials into fuels and value-added chemicals is technically challenging due to a variety of factors [1–4]. Lignocellulosic biomass is composed of 40–50% cellulose, 25–30% hemicellulose, and 15–20% lignin, and the high lignin content requires strong physical and chemical pretreatment for its decomposition [5]. Also, hemicellulose is hydrolyzed into a mixture of glucose, xylose, and other minor sugars such as L-arabinose, and their fermentation is not efficiently done by any native industrial microorganisms [6]. For these reasons, industrial bioprocesses utilizing lignocellulosic biomass have not been realized so far, and the search for other alternative renewable biomass continues [7–10].

Meanwhile, fruit processing wastes such as orange peels are becoming abundant with the growth of the fresh juice industry, but are not being efficiently utilized [11, 12]. Fruit processing wastes are high in pectin (12–35%, g/g dry weight) with much less lignin content (approximately 2%, g/g dry weight) than lignocellulosic biomass [13]. Therefore, this pectin-rich biomass can be easily broken down into monomers; however, pectin monomers, like

hemicellulose monomers, are not easily metabolized by common industrial hosts [14].

The primary chemical structure of pectin is methylated polygalacturonic acid in an alpha-(1-4) chain with branched oligosaccharides consisting of arabinose, galactose, xylose, and some minor sugars [15]. Among them, L-arabinose is one of the most abundant pentose sugars in pectin [16]. Arabinose content in various fruits and vegetables ranges from 3.3 to 21.6 g/l (summarized in Table 1) [14, 17, 18]. It is contradictory to lignocellulosic biomass which has limited arabinose content (approximately 0.2%, g/g dry weight) [19].

L-Arabinose is a five-carbon sugar like xylose. Unlike other sugars that naturally occur in the D-form, such as D-xylose, L-arabinose is a component of pectin and hemicellulose, and it is more common than D-arabinose in nature. Although studies have been conducted extensively for xylose metabolism to realize lignocellulosic bioprocesses [20], L-arabinose metabolism has not received much attention. In the present review, microbial strains that can natively metabolize L-arabinose are summarized. In some studies, the strains were engineered to produce useful products such as ethanol. Moreover, metabolic engineering efforts to develop efficient L-arabinose-fermenting strains

**Table 1.** Representative pectin-rich biomass and their arabinose content.

Source	Arabinose, % <sup>1)</sup>	Other major sugars <sup>2)</sup>	Refs
Sugar beet pulp	21.6 (0.28)	Glu	[17, 18]
Lime peel	8.5	Glu	[14]
Pear peel	6.0	Glu, Xyl, Fru	[14]
Orange peel	5.6 (0.20)	Glu, Fru	[14, 17]
Apple pomace	5.5	Glu, Fru, Suc	[14]
Mandarin peel	3.3	Glu, Fru, Suc	[14]

<sup>1)</sup>% Dry matter (g/g pectin).

<sup>2)</sup>Higher content than arabinose. Glu, Glucose; Xyl, xylose; Fuc, Fucose; Suc, Sucrose.

using non-native but industrial hosts are discussed, focusing on *Saccharomyces cerevisiae*.


### Native L-Arabinose-Fermenting Microorganisms

Arabinose catabolic pathways of native strains can be divided into the oxidoreductase (fungal) and isomerase (bacterial) pathways (Fig. 1). In both pathways, L-arabinose is converted into D-xylulose-5-phosphate, which is then canonically metabolized by the non-oxidative pentose phosphate pathway [21], or alternatively by the phospho-ketolase pathway such as that in *Clostridium acetobutylicum* [22].

In L-arabinose-fermenting native fungi, L-arabinose is converted into D-xylulose by two reduction and two oxidation reactions, which are composed of NAD(P)H-specific aldose reductase (AR or XR), NAD<sup>+</sup>-specific L-arabitol-4-dehydrogenase (LAD), NAD(P)H-specific L-xylulose reductase (LXR or ALX), and NAD<sup>+</sup>-specific xylitol dehydrogenase (XDH) [21]. Then, D-xylulose is phosphorylated into D-xylulose-5-phosphate by D-xylulokinase (XK). As highlighted in Fig. 1, fungal pathways of L-arabinose and D-xylose share three enzymes: AR (XR), XDH and XK. Although the cofactor preferences of AR and LXR (ALX) vary among fungal species, the first reductase enzyme usually prefers NADPH, while the two dehydrogenases strictly use NAD<sup>+</sup>. Therefore, redox balance of the pathway leading to efficient cell growth is achieved under aerobic conditions [23, 24]. Under oxygen-limited conditions, L-arabitol might be produced due to NAD<sup>+</sup> limitation, which is not found in the bacterial pathway of L-arabinose metabolism [25–27]. Because of the oxygen-dependent nature of the fungal pathway, ethanol production by these native strains is marginal [28, 29].

As early as 1990, a few native arabinose-fermenting fungi strains were identified, but the strains yielded a trace level of ethanol or even no ethanol production [26, 28]. In detail,

116 different yeast strains were screened for the ability to catabolize arabinose or xylose aerobically. As a result, four yeast strains (*Ambrosiozyma monospora*, and three *Candida spp.*) were found to ferment L-arabinose as a sole carbon source. Additionally, the ethanol yield was at most 0.18 (g/g



**Fig. 1.** Arabinose metabolic pathways in fungi (A) and bacteria (B).

AR(XR); Aldose reductase, LAD; L-arabitol-4-dehydrogenase, LXR(ALX); L-xylulose reductase, XDH; D-xylulose reductase, XK; Xylulokinase, AI; L-arabinose isomerase, RK; L-Ribulokinase, R5PE; L-Ribulose-5-P-4-epimerase.

**Table 2.** Native arabinose-assimilating microorganisms and their engineered strains producing ethanol.

Strain	Genotype <sup>1)</sup>	Fermentation conditions <sup>2)</sup>		Ethanol production		Refs
		Arabinose (g/l)	Aeration	Titer (g/l)	Yield (g/g)	
Yeast						
<i>Ambrosiozyma monospora</i> Y-1484	Wild type	80	OL	4.1	0.18	[28]
<i>Candida succiphila</i> Y-11998	Wild type	80	OL	3.9	0.05	[28]
Bacteria						
<i>Sarcina ventriculi</i>	Wild type	19	AN	4.7	0.31	[31]
<i>Klebsiella oxytoca</i> P2	<i>Zm_pdc, Zm_adhB</i>	80	OL	27.2	0.34	[32]
<i>Escherichia coli</i> FBR3	<i>Zm_pdc, Zm_adhB, ldhΔ, pflΔ</i>	100	OL	44.4	0.46	[33]

<sup>1)</sup>*Zm, Zymomonas mobilis*; *pdc*, pyruvate decarboxylase gene; *adhB*, alcohol dehydrogenase gene; *ldh*, lactate dehydrogenase gene; *pfl*, pyruvate formate lyase gene.

<sup>2)</sup>All fermentations were performed with complex media with an initial arabinose concentration as shown above. Arabinose was the only carbon source available. OL, oxygen-limited conditions; AN, anaerobic conditions.

consumed arabinose) with *A. monospora* and *C. succiphila* (Table 2).

Meanwhile, 15 xylose-fermenting microorganisms were screened to evaluate the ability to ferment L-arabinose to ethanol [26]. As a result, one bacterium (*Erwinia chrysanthemi*), six yeast strains (*C. tropicalis*, *C. shehatae*, *Pachysolen tannophilus* Y-2460, *P. tannophilus* Y-12891, *Scheffersomyces stipitis*, and *Torulopsis sonorensis*), and one mold strain (*Aspergillus oryzae*) were confirmed to assimilate arabinose with xylose and glucose as co-substrates. While *E. chrysanthemi* and *C. tropicalis* consumed xylose and arabinose simultaneously, all other fungal strains preferred xylose over arabinose. During arabinose metabolism, *S. stipitis* was the only fungal strain producing ethanol at a yield of 0.15 (g/g consumed sugar) and arabitol at a yield of 0.24 (g/g consumed sugar).

In comparison to the above-mentioned fungal pathways, bacterial pathways of L-arabinose metabolism are relatively simple; 1) only three enzymes are needed to convert to D-xylulose-5-phosphate, and 2) no cofactor is involved. The bacterial pathway consists of L-arabinose isomerase (AI), L-ribulose kinase (RK), and L-ribulose-5-phosphate-4-epimerase (R5PE) encoded by the *araA*, *araB*, and *araD* genes, respectively (Fig. 1B) [30]. In addition, native bacterial strains such as *Sarcina ventriculi* can ferment L-arabinose anaerobically and produce ethanol efficiently at a yield of 0.3 (g/g consumed arabinose) [31]. For some bacterial strains lacking the *pdc* and *adh* genes (encoding pyruvate decarboxylase and alcohol dehydrogenase, respectively) such as *Klebsiella oxytoca*, introducing the genes from *Zymomonas mobilis* enabled ethanol fermentation from L-arabinose [32]. For native L-arabinose-metabolizing *Escherichia coli*, in contrast, deletion of lactate dehydrogenase (*ldh*) and pyruvate formate lyase (*pfl*) genes was required to enable

ethanol production from L-arabinose [33]. As summarized in Table 2, native bacterial strains assimilating L-arabinose can be promising hosts for ethanol fermentation.

## Engineering *Saccharomyces cerevisiae* for L-Arabinose Fermentation

*S. cerevisiae*, the industrial host for bioethanol production, cannot utilize L-arabinose as efficiently as it can utilize xylose. For cellulosic ethanol production, a great amount of effort has been focused on the development of xylose-fermenting *S. cerevisiae* strains, while there have not been many examples for L-arabinose.

There are a limited number of studies on the development of *S. cerevisiae* expressing a heterologous fungal pathway of L-arabinose metabolism compared to that expressing a heterologous bacterial pathway [30, 34–38]. Usually, xylose-fermenting *S. cerevisiae* expressing heterologous AR, XDH, and XK is first engineered by expressing *Scheffersomyces stipitis* *XYL1*, *XYL2*, and *XYL3* genes, respectively, in most cases [20]. It has to be noted that an AR is NAD(P)H-specific aldose reductase with specificity for both xylose and L-arabinose [39] with a 50% higher rate for L-arabinose metabolism when using NADPH as a cofactor [40]. Next, the resulting strain is further engineered to express LAD and LXR, which are *T. reesei* *LAD1* and *A. monospora* *ALX1* genes, respectively [24, 34]. The *S. cerevisiae* 424A(LNH-ST)/pLXR<sub>NAD</sub>-LAD strain, which was developed as above, produced 10 g/l ethanol from 45 g/l L-arabinose [35] as summarized in Table 3. In another study, *T. reesei* *LXR1* gene was expressed instead of *A. monospora* *ALX1* gene, but the resulting strain only produced 0.1 g/l ethanol from 50 g/l L-arabinose [35, 41]. The low ethanol production can be explained by the fact that *T. reesei* *LXR1* gene is now

**Table 3.** Ethanol fermentation by engineered *Saccharomyces cerevisiae* using arabinose as the sole carbon source.

Strain names	Strain backgrounds	Arabinose pathways	Optimization strategies	Media	Arabinose (g/l)	Aeration	Titer (g/l)	Yield (g/g consumed)	Refs
Heterologous fungal pathways									
H2561	CEN.PK2	SsXYL1, SsXYL2, ScXKS1, TrLAD1, TrLXR1	-	Minimal	50	AN	0.1	-	[34]
424A(LNH-ST)/pLXR <sub>NAD</sub> -LAD	424A	SsXYL1, SsXYL2, SsXYL3, TrLAD1, AmALX1	-	Complex	45	OL	9.4	0.22	[35]
Heterologous bacterial pathways									
JBY25-4M	CEN.PK2-1C	BsaraA, EcaraB, EcaraD, ScGAL2	-	Minimal	20	OL	6	0.3	[36]
BWY1-4S	CEN.PK2-1C	BsaraA, EcaraB, EcaraD, ScGAL2	Codon optimization Adaptive evolution	Minimal	30	AN	9	0.39	[43]
IMS0002	CEN.PK2-1C	LparaA, LparaB, LparaD	PPP overexpression Adaptive evolution	Complex	20	AN	8.92	0.45	[30]
BSW3AP	CEN.PK102-3A	LparaA, LparaB, LparaD	PPP overexpression Adaptive evolution	Minimal	20	OL AN	6.9 -	0.43 0.42	[37]
BSW3AG	CEN.PK102-3A	LparaA, LparaB, LparaD, ScGAL2	PPP overexpression Adaptive evolution	Minimal	20	AN	-	0.43	[37]

OL, oxygen-limited conditions; AN, anaerobic conditions; Bs, *Bacillus subtilis*; Ec, *Escherichia coli*; Sc, *Saccharomyces cerevisiae*; Tr, *Trichoderma reesei*; Am, *Ambrosiozyma monospora*; Pi, *Piromyces sp.*; Lp, *Lactobacillus plantarum*; Nc, *Neurospora crassa*; Mt, *Myceliophthora thermophila*; Ss, *Scheffersomyces stipitis*; PPP, pentose phosphate pathway.

functionally identified as mannitol dehydrogenase [42]. Another factor determining L-arabinose fermentation efficiency is aeration. The fungal L-arabinose pathway is not redox-neutral because of the dual cofactor preference of AR and LXR (NADPH and NADH) while XDH and LAD are NAD<sup>+</sup>-specific. Therefore, the cofactor imbalance issue could be more severe than xylose fermentation requiring just AR and XDH.

Meanwhile, the bacterial L-arabinose pathway is redox-neutral; thus, more studies have been performed for heterologous expression in *S. cerevisiae* from genes of *Bacillus subtilis*, *Escherichia coli* and *Lactobacillus plantarum*. The bacterial L-arabinose pathway consisting of *araA*, *araB* and *araD* genes was tested with various combinations from different origins (Table 3). The *araA* gene from *Bacillus subtilis* [36], *Bacillus licheniformis* [43, 44] and *Lactobacillus plantarum* [30, 37] and the *araB* and *araD* genes from *Escherichia coli* [43, 44] or *L. plantarum* [30, 37] were tested. Ethanol production from the engineered strains varied between 6–9 g/l from 20 g/l L-arabinose. Regardless of the origin of the heterologous genes, various approaches to improve L-arabinose fermentation have been performed. In

general, the overexpression of the non-oxidative pentose phosphate pathway genes (*TAL1*, *TKL1*, *RPE1* and *RK11*) and adaptive evolution were required [30, 36, 43]. However, despite all optimizations, the fermentation productivity is limited by a bacterial pathway possibly because of the unfavorable thermodynamic properties of L-arabinose isomerase under ambient conditions [45].

It should be noted that adaptive evolution is proven to be an effective metabolic engineering strategy to improve xylose fermentation by engineered *S. cerevisiae* strains for both fungal and bacterial pathways [20, 46]. However, for L-arabinose fermentation, only the engineered strains with a bacterial pathway have been subjected to adaptive evolution (Table 2). It can be explained by the fact that L-arabinose fermentation has not been performed systematically and extensively compared to xylose fermentation. It is also possible that the heterologous expression of a fungal pathway in *S. cerevisiae* requires multiple strategies to be optimized to overcome the severe redox imbalance issue.

Arabinose fermentation can be improved by expressing L-arabinose-specific sugar transporters (Table 4). *S. cerevisiae*

**Table 4.** Characterization of putative arabinose transporters over-expressed in a hexose transporter null mutant of *Saccharomyces cerevisiae*.

Transporter genes	Arabinose uptake rate (mmol/h/g DCW)	Arabinose affinity (mM)	References
<i>Neurospora crassa</i> LAT-1	116.7	58.12	[50]
<i>Myceliophthora thermophila</i> LAT-1	10.29	29.39	[50]
<i>Penicillium chrysogenum</i> AraT	5.30	0.13	[51]
<i>Saccharomyces cerevisiae</i> GAL2	0.13	57.00	[48]
<i>Arabidopsis thaliana</i> Stp2	0.04	4.50	[48]
<i>Scheffersomyces stipites</i> AraT	0.02	3.80	[48]

could uptake arabinose through some hexose transporters such as Hxt5 and Hxt7 with low affinity [47]. *S. cerevisiae* Gal2 had the highest affinity to L-arabinose (57 mM) among other native hexose transporters [48, 49]. It is also reported that *S. cerevisiae* Gal2 contributed to anaerobic arabinose fermentation when arabinose is the sole carbon source [37]. Several heterologous arabinose transporters have been identified from *Neurospora crassa* [50], *Myceliophthora thermophila* [50], *Penicillium chrysogenum* [51], *Arabidopsis thaliana* [48], and *Scheffersomyces stipites* [48], and their ability to uptake L-arabinose varies significantly (0.02–116.7 mmol/h/g DCW). *N. crassa* LAT-1 was the most efficient L-arabinose transporter reported with a rate of 116.7 (mmol/h/g DCW), which is 2 orders of magnitude higher than that of *S. cerevisiae* GAL2 0.13 (mmol/h/g DCW) [50]. *P. chrysogenum* AraT was a high-affinity arabinose transporter with no activity with glucose and xylose, although it was still inhibited by the presence of glucose and xylose [51, 52]. At the present stage, no heterologous sugar transporter was reported to either improve L-arabinose fermentation or allow simultaneous uptake of arabinose and glucose. In arabinose metabolism, as in xylose's case [53, 54], it can be assumed that arabinose catabolism is currently more limiting than non-specific arabinose uptake in engineered *S. cerevisiae*.

### Engineering of Other Microorganisms for Arabinose Fermentation

Some non-native arabinose fermenting microorganisms have also been engineered to assimilate L-arabinose and produce ethanol or other products. *Z. mobilis* (pZB206), which natively carries *pdC* and *adhB* but lacked arabinose-assimilating enzymes, was constructed by introducing *E. coli* *araABD*, *talB*, *tktA* to metabolize arabinose to ethanol [55]. The resulting strain showed an ethanol yield of 0.49

by consuming 25 g/l L-arabinose. Unlike *S. cerevisiae*, *Corynebacterium glutamicum* expressing *E. coli* *araABD* was easily engineered to produce amino acids such as L-glutamate, L-lysine, L-ornithine and L-arginine with arabinose as the sole carbon source [56]. For example, one of the engineered *C. glutamicum* strains produced L-glutamate at a yield of 0.07 from 75 g/l L-arabinose. *Rhodococcus opacus* expressing *Streptomyces cattleya* *araABD* fermented 16 g/l L-arabinose as the sole carbon source and produced fatty acids at a yield of 0.13 (g/g consumed arabinose). The fatty acids were mostly palmitic acid with some cis-10-heptadecenoic acid, oleic acid, myristic acid, pentadecanoic acid, palmitoleic acid, heptadecanoic acid, and stearic acid. Although the examples are limited, non-*Saccharomyces cerevisiae* strains can also be engineered to ferment L-arabinose and produce various value-added products other than ethanol.

### Future Outlook

Both crop biomass and cellulosic biomass do not support sustainable bioprocesses due to their low contribution to greenhouse gas reduction [57] and limited technologies to overcome the recalcitrance [58], respectively. Alternatively, pectin-rich biomass such as fruit-processing wastes can be an attractive choice due to low lignin content and the growing demands for fresh juice. L-Arabinose is a primary sugar of pectin structure and its content is minimal in other biomass. As discussed in this review, research for L-arabinose fermentation is in an early stage. Thus, all options are open to either optimizing native arabinose-assimilating strains or engineering non-native strains such as *S. cerevisiae*, *Z. mobilis*, *C. glutamicum*, and *R. opacus* depending on the desired products. Engineered *S. cerevisiae* strains fermenting L-arabinose are still limited to ethanol production, however, various chemicals and value-added

products are expected to be studied as well. Additionally, the development of strains fermenting other pectin-derived monomers such as galacturonic acid and L-rhamnose needs to be considered.

## Acknowledgments

This work was carried out with the support of the “Cooperative Research Program for Agriculture Science & Technology Development (Project No. PJ01279801)” Rural Development Administration, Republic of Korea.

## Conflict of Interest

The authors have no financial conflicts of interest to declare.

## References

- Robertson GP, Hamilton SK, Barham BL, Dale BE, Izaurralde RC, Jackson RD, et al. 2017. Cellulosic biofuel contributions to a sustainable energy future: choices and outcomes. *Science* **356**(6345).
- Gírio FM, Fonseca C, Carvalheiro F, Duarte LC, Marques S, Bogel-Lukasik R. 2010. Hemicelluloses for fuel ethanol: a review. *Bioresour. Technol.* **101**: 4775-4800.
- Go AR, Ko JW, Lee SJ, Kim SW, Han SO, Lee J, et al. 2012. Process design and evaluation of value-added chemicals production from biomass. *Biotechnol. Bioprocess Eng.* **17**: 1055-1061.
- Trinh LTP, Lee Y-J, Lee J-W, Lee W-H. 2018. Optimization of ionic liquid pretreatment of mixed softwood by response surface methodology and reutilization of ionic liquid from hydrolysate. *Biotechnol. Bioprocess Eng.* **23**: 228-237.
- Alonso DM, Hakim SH, Zhou S, Won W, Hosseinaei O, Tao J, et al. 2017. Increasing the revenue from lignocellulosic biomass: maximizing feedstock utilization. *Science Adv.* **3**: e1603301.
- Liao JC, Mi L, Pontrelli S, Luo S. 2016. Fuelling the future: microbial engineering for the production of sustainable biofuels. *Nat. Rev. Microbiol.* **14**: 288-288.
- Wei N, Quarterman J, Jin YS. 2013. Marine macroalgae: an untapped resource for producing fuels and chemicals. *Trends Biotechnol.* **31**: 70-77.
- Park M-R, Kim S-K, Jeong G-T. 2018. Biosugar production from *Gracilaria verrucosa* with sulfamic acid pretreatment and subsequent enzymatic hydrolysis. *Biotechnol. Bioprocess Eng.* **23**: 302-310.
- Javier AG, Maria Cristina R, Oriana S, Maria Elena L. 2018. Saccharification of brown macroalgae using an arsenal of recombinant alginase lyases: potential application in the biorefinery process. *J. Microbiol. Biotechnol.* **28**: 1671-1682.
- Oh YR, Jung KA, Lee HJ, Jung GY, Park JM. 2018. A novel 3,6-anhydro-L-galactose dehydrogenase produced by a newly isolated *Raoultella ornithinolytica* B6-JMP12. *Biotechnol. Bioprocess Eng.* **23**: 64-71.
- Van Dyk JS, Gama R, Morrison D, Swart S, Pletschke BI. 2013. Food processing waste: problems, current management and prospects for utilisation of the lignocellulose component through enzyme synergistic degradation. *Renew. Sustain. Energy Rev.* **26**: 521-531.
- Lisandro GS, Raul NC, Maria TB, Miguel AI. 2018. Feasibility of bioethanol production from cider waste. *J. Microbiol. Biotechnol.* **28**: 1493-1501.
- Edwards MC, Doran-Peterson J. 2012. Pectin-rich biomass as feedstock for fuel ethanol production. *Appl. Microbiol. Biotechnol.* **95**: 565-575.
- Choi IS, Lee YG, Khanal SK, Park BJ, Bae HJ. 2015. A low-energy, cost-effective approach to fruit and citrus peel waste processing for bioethanol production. *Appl. Energy* **140**: 65-74.
- May CD. 1990. Industrial pectins: Sources, production and applications. *Carbohydr. Polym.* **12**: 79-99.
- Seiboth B, Metz B. 2011. Fungal arabinan and L-arabinose metabolism. *Appl. Microbiol. Biotechnol.* **89**: 1665-1673.
- Oosterveld A, Beldman G, Schols HA, Voragen AGJ. 1996. Arabinose and ferulic acid rich pectic polysaccharides extracted from sugar beet pulp. *Carbohydr. Res.* **288**: 143-153.
- Müller-Maatsch J, Bencivenni M, Caligiani A, Tedeschi T, Bruggeman G, Bosch M, et al. 2016. Pectin content and composition from different food waste streams in memory of Anna Surribas, scientist and friend. *Food Chem.* **201**: 37-45.
- van Maris AJA, Abbott DA, Bellissimi E, van den Brink J, Kuyper M, Luttik MAH, et al. 2006. Alcoholic fermentation of carbon sources in biomass hydrolysates by *Saccharomyces cerevisiae*: current status. *Antonie Van Leeuwenhoek Int. J. Gen. Mol. Microbiol.* **90**: 391-418.
- Kim SR, Park YC, Jin YS, Seo JH. 2013. Strain engineering of *Saccharomyces cerevisiae* for enhanced xylose metabolism. *Biotechnol. Adv.* **31**: 851-861.
- Hahn-Hägerdal B, Karhumaa K, Fonseca C, Spencer-Martins I, Gorwa-Grauslund MF. 2007. Towards industrial pentose-fermenting yeast strains. *Appl. Microbiol. Biotechnol.* **74**: 937-953.
- Servinsky MD, Germane KL, Liu S, Kiel JT, Clark AM, Shankar J, et al. 2012. Arabinose is metabolized via a phosphoketolase pathway in *Clostridium acetobutylicum* ATCC 824. *J. Ind. Microbiol. Biotechnol.* **39**: 1859-1867.
- Fonseca C, Romão R, Rodrigues De Sousa H, Hahn-Hägerdal B, Spencer-Martins I. 2007. L-Arabinose transport and catabolism in yeast. *FEBS J.* **274**: 3589-3600.
- Verho R, Putkonen M, Londesborough J, Penttilä M, Richard P. 2004. A novel NADH-linked L-xylulose reductase in the L-arabinose catabolic pathway of yeast. *J. Biol. Chem.* **279**: 14746-14751.
- Fonseca C, Spencer-Martins I, Hahn-Hägerdal B. 2007. L-Arabinose metabolism in *Candida arabinofementans* PYCC

- 5603T and *Pichia guilliermondii* PYCC 3012: influence of sugar and oxygen on product formation. *Appl. Microbiol. Biotechnol.* **75**: 303-310.
26. McMillan JD, Boynton BL. 1994. Arabinose utilization by xylose-fermenting yeasts and fungi. *Appl. Biochem. Biotechnol.* **45-46**: 569-584.
  27. Watanabe S, Kodak T, Makino K. 2006. Cloning, expression, and characterization of bacterial L-arabinose 1-dehydrogenase involved in an alternative pathway of L-arabinose metabolism. *J. Biol. Chem.* **281**: 2612-2623
  28. Dien BS, Kurtzman CP, Saha BC, Bothast RJ. 1996. Screening for L-arabinose fermenting yeasts. *Appl. Biochem. Biotechnol.* **57-58**: 233-242.
  29. Kurtzman CP, Dien BS. 1998. *Candida arabinofermentans*, a new L-arabinose fermenting yeast. *Antonie Van Leeuwenhoek Int. J. Gen. Mol. Microbiol.* **74**: 237-243.
  30. Wisselink HW, Toirkens MJ, Berriél MDRF, Winkler AA, Van Dijken JP, Pronk JT, et al. 2007. Engineering of *Saccharomyces cerevisiae* for efficient anaerobic alcoholic fermentation of L-arabinose. *Appl. Environ. Microbiol.* **15**: 488-491.
  31. Finn RK, Bringer S, Sahn H. 1984. Fermentation of arabinose to ethanol by *Sarcina ventriculi*. *Appl. Microbiol. Biotechnol.* **19**: 161-166.
  32. Bothast RJ, Saha BC, Flossenier V, Ingram LO. 1994. Fermentation of L-arabinose, D-xylose and D-glucose by ethanologenic recombinant *Klebsiella oxytoca* strain P2. *Biotechnol. Lett.* **16**: 401-406.
  33. Dien BS, Hespell RB, Wyckoff HA, Bothast RJ. 1998. Fermentation of hexoses and pentoses sugars using a novel ethanologenic *Escherichia coli* strain. *Enzyme Microb. Technol.* **23**: 366-371.
  34. Richard P, Verho R, Putkonen M, Londesborough J, Penttilä M. 2003. Production of ethanol from L-arabinose by *Saccharomyces cerevisiae* containing a fungal L-arabinose pathway. *FEMS Yeast Res.* **3**: 185-189.
  35. Bera AK, Sedlak M, Khan A, Ho NWY. 2010. Establishment of L-arabinose fermentation in glucose/xylose co-fermenting recombinant *Saccharomyces cerevisiae* 424A(LNH-ST) by genetic engineering. *Appl. Microbiol. Biotechnol.* **87**: 1803-1811.
  36. Becker J, Boles E. 2003. A modified *Saccharomyces cerevisiae* strain that consumes L-arabinose and produces ethanol. *Appl. Environ. Microbiol.* **69**: 4144-4150.
  37. Wang C, Shen Y, Zhang Y, Suo F, Hou J, Bao X. 2013. Improvement of L-arabinose fermentation by modifying the metabolic pathway and transport in *Saccharomyces cerevisiae*. *Biomed. Res. Int.* **2013**:461204.
  38. Wang C, Zhao J, Qiu C, Wang S, Shen Y, Du B, et al. 2017. Couitilization of D-glucose, D-xylose, and L-arabinose in *Saccharomyces cerevisiae* by coexpressing the metabolic pathways and evolutionary engineering. *Biomed. Res. Int.* **2017**: 5318232.
  39. Bettiga M, Bengtsson O, Hahn-Hägerdal B, Gorwa-Grauslund MF. 2009. Arabinose and xylose fermentation by recombinant *Saccharomyces cerevisiae* expressing a fungal pentose utilization pathway. *Microb. Cell Fact.* **8**: 1-12.
  40. Watanabe S, Utsumi Y, Sawayama S, Watanabe Y. 2016. Identification and characterization of D-arabinose reductase and D-arabinose transporters from *Pichia stipitis*. *Biosci. Biotechnol. Biochem.* **80**: 2151-2158.
  41. Richard P, Putkonen M, Väänänen R, Londesborough J, Penttilä M. 2002. The missing link in the fungal L-arabinose catabolic pathway, identification of the L-xylose reductase gene. *Biochem.* **41**: 6432-6437.
  42. Metz B, de Vries RP, Polak S, Seidl V, Seiboth B. 2009. The *Hypocrea jecorina* (syn. *Trichoderma reesei*) *lxr1* gene encodes a d-mannitol dehydrogenase and is not involved in L-arabinose catabolism. *FEBS Lett.* **583**: 1309-1313.
  43. Wiedemann B, Boles E. 2008. Codon-optimized bacterial genes improve L-arabinose fermentation in recombinant *Saccharomyces cerevisiae*. *Appl. Environ. Microbiol.* **74**: 2043-2050.
  44. Wang X, Yang J, Yang S, Jiang Y. 2019. Unraveling the genetic basis of fast l-arabinose consumption on top of recombinant xylose-fermenting *Saccharomyces cerevisiae*. *Biotechnol. Bioeng.* **116**: 283-293.
  45. Lee DW, Hong YH, Choe EA, Lee SJ, Kim SB, Lee HS, et al. 2005. A thermodynamic study of mesophilic, thermophilic, and hyperthermophilic L-arabinose isomerases: the effects of divalent metal ions on protein stability at elevated temperatures. *FEBS Lett.* **579**: 1261-1266.
  46. Jansen MLA, Bracher JM, Papapetridis I, Verhoeven MD, de Bruijn H, de Waal PP, et al. 2017. *Saccharomyces cerevisiae* strains for second-generation ethanol production: from academic exploration to industrial implementation. *FEMS Yeast Res.* **17**: fox044.
  47. Leandro MJ, Fonseca C, Gonçalves P. 2009. Hexose and pentose transport in ascomycetous yeasts: An overview. *FEMS Yeast Res.* **9**: 511-525.
  48. Subtil T, Boles E. 2011. Improving L-arabinose utilization of pentose fermenting *Saccharomyces cerevisiae* cells by heterologous expression of L-arabinose transporting sugar transporters. *Biotechnol. Biofuels* **4**: 1-10.
  49. Verhoeven MD, Bracher JM, Nijland JG, Bouwknegt J, Daran JG, Driessen AJM, et al. 2018. Laboratory evolution of a glucose-phosphorylation-deficient, arabinose-fermenting *S. cerevisiae* strain reveals mutations in GAL2 that enable glucose-insensitive l-arabinose uptake. *FEMS Yeast Res.* **18**: doi: 10.1093/femsyr/foy062.
  50. Li J, Xu J, Cai P, Wang B, Ma Y, Benz JP, et al. 2015. Functional analysis of two L-arabinose transporters from filamentous fungi reveals promising characteristics for improved pentose utilization in *Saccharomyces cerevisiae*. *Appl. Environ. Microbiol.* **81**: 4062-4070.
  51. Bracher JM, Verhoeven MD, Wisselink HW, Crimi B, Nijland JG, Driessen AJM, et al. 2018. The *Penicillium*

- chrysogenum* transporter Pc AraT enables high-affinity, glucose-insensitive L-arabinose transport in *Saccharomyces cerevisiae*. *Biotechnol. Biofuels* **11**:63.
52. Verhoeven MD, de Valk SC, Daran JG, van Maris AJA, Pronk JT. 2018. Fermentation of glucose-xylose-arabinose mixtures by a synthetic consortium of single-sugar-fermenting *Saccharomyces cerevisiae* strains. *FEMS Yeast Res.* **18**: doi: 10.1093/femsyr/foy075.
53. Kim SR, Ha S-J, Wei N, Oh EJ, Jin Y-S. 2012. Simultaneous co-fermentation of mixed sugars: a promising strategy for producing cellulosic ethanol. *Trends Biotechnol.* **30**: 274-282.
54. Lane S, Xu H, Oh EJ, Kim H, Lesmana A, Jeong D, et al. 2018. Glucose repression can be alleviated by reducing glucose phosphorylation rate in *Saccharomyces cerevisiae*. *Sci. Rep.* **8**: 2613.
55. Deanda K, Zhang MIN, Eddy C, Picataggio S. 1996. Development of an arabinose-fermenting *Zymomonas mobilis* strain by metabolic pathway engineering. *Microbiol.* **62**: 4465-4470.
56. Schneider J, Niermann K, Wendisch VF. 2011. Production of the amino acids L-glutamate, L-lysine, L-ornithine and L-arginine from arabinose by recombinant *Corynebacterium glutamicum*. *J. Biotechnol.* **154**: 191-198.
57. Searchinger T, Heimlich R, Houghton RA, Dong F, Elobeid A, Fabiosa J, et al. 2008. Use of U.S. croplands for biofuels increases greenhouse gases through emissions from land-use change. *Science* **423**: 1238-1241.
58. Lynd LR. 2017. The grand challenge of cellulosic biofuels. *Nat. Biotechnol.* **35**: 912-915.