

First Finding of a Bivalve-Inhabiting Hydrozoan (Cnidaria, Hydrozoa) from Korea

Shin Kubota^{1,*}, Jimin Lee², Cheon Young Chang³

¹Turritopsis Immortal Jellyfish Regenerative Biological Research/Experience Laboratory, Nishimuro 649-2211, Japan

²Marine Ecosystem Research Division, Korea Institute of Ocean Science and Technology, Busan 49111, Korea

³Department of Biological Science, College of Natural Sciences, Daegu University, Gyeongsan 38453, Korea

ABSTRACT

Eutima japonica Uchida, 1925, a bivalve-inhabiting hydrozoan was collected for the first time in Korea, associating with *Mytilus galloprovincialis*. The morphology of male medusae of this hydrozoan is clarified by culture and described as well as other developmental stages. As the present material from Korea is in good accord with that of the northern Japanese form of *E. japonica*, so the geographical distribution of the northern form of this species is widened, Japan, China and Korea. Green fluorescent protein distribution pattern of this medusa is also described and compared with that of the most related species *Eutima sapinhoa* Narchi and Hebling, 1975.

Keywords: culture, *Eutima japonica*, geographical distribution, green fluorescent protein, life cycle, medusan form, morphological variation, taxonomy

INTRODUCTION

The solitary, bivalve-inhabiting hydroid supposedly evolved from the free-living, colonial *Eutima* without drastic modification of the medusa stage, and appeared as a specialized, distinct member in the Eirenidae (see Kubota, 2000 for details of the evolution of bivalve-inhabiting hydrozoans and the study history of them; Kubota and Collins, 2017). In Japanese waters, records of *Eutima* refer only to the bivalve-inhabiting species, *E. japonica* Uchida, 1925 (Uchida, 1925; Yamazi, 1958; Kubota, 1992, 2000), which is divided into the four forms that are parapatrically distributed, changing their morphology of the earliest matured medusae step by step (Kubota, 1997, 1999, 2000). The medusa of *E. japonica* was also reported in the central part of the north Pacific (Kramp, 1965), Cochin Backwater, India (Santhakumari and Vannucci, 1971) and in the Jiulong River estuary near Amoy China (Zhenzu and Jiachi, 1983), while never found in the east coasts of the Pacific, from where no bivalve-inhabiting hydrozoans have been recorded (Kubota, 2000), except for a recent appearance of polyp by Tsunami transport

of the host *Mytilus galloprovincialis* that attached to small boat (Calder et al., 2014).

In Korea, 130 hydroid species, comprising 77 species or subspecies of the Thecata, 24 species of the Athecata and 29 species of the hydromedusae, are currently recorded in three monographical studies by the late Prof. Jung Hee Park (Park, 2010, 2011, 2012). However, notwithstanding her long-lasting efforts since her first paper published in 1979 (Rho and Park, 1979) and against our detailed survey at 30 sites along Korean coasts during the period between 1990 and 2006, no bivalve-inhabiting hydrozoans have so far been found (Kubota et al., 2006). Recently we conducted an additional survey in late April, 2018, and finally found a polyp colony associated with mussels of *Mytilus galloprovincialis* from Yeosu for the first time in Korea.

To determine the exact taxonomic status of a bivalve-associating hydroid as well as its morphological and geographical form, the morphology of mature medusa after culture is prerequisite and indispensable (Kubota, 1997). Furthermore, green fluorescent protein (GFP) distribution pattern of the medusa should be observed and compared with that

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

***To whom correspondence should be addressed**
Tel: 81-739-207402, Fax: 81-739-207402
E-mail: benikurage2018@gmail.com

of the most related species *Eutima sapinhoa* Narchi and Hebling, 1975 (Migotto et al., 2004). As a result of the studies, the mussel-inhabiting hydroid from Yeosu turned out to be *Eutima japonica* Uchida, 1925, which is a new record from Korea, and its form can be determined.

In this paper, we deal with the description of the male medusae of *E. japonica* as well as other developmental stages by culture, and with some brief comments on its morphological and geographical distinction according to the GFP distribution pattern of the medusa. We also provide herein a measurements table showing morphological variation of well-developed medusae reared in the laboratory after release from a mussel-inhabiting hydroid polyp.

MATERIALS AND METHODS

On April 30, 2018, 31 individuals of *Mytilus galloprovincialis*,

that were cultivated in nearby mussel farms and sold in Yeosu seafood market, were examined alive soon after purchase. The water temperature in Yeosu was about 15°C on these days, according to the “Korea Real-Time Database for NEAR-GOOS” by Korea Hydrographic and Oceanographic Agency. Polyps were found inside only in one individual of these mussels, then polyps with medusa buds were picked out from the host at the Kujyukusima Aquarium, Nagasaki Prefecture, Japan, on May 1, the next day after purchase. On the other hand, no such polyps were found from 30 mussels purchased at Jagalchi seafood market at Busan, Korea on April 28, 2018.

Laboratory culture of the polyps with medusa buds was done individually and bringing them back to the laboratory in Shirahama, Wakayama, Japan. For each polyp, an 80-mL polystyrene vessel of 60 mm in diameter and 30 mm high was filled with filtered seawater (32 psu) from Shirahama, Wakayama Prefecture (supplied from the Seto Marine Bio-




Fig. 1. A well-developed, but not yet fully matured, 1-month-old laboratory-reared medusa of *Eutima japonica* from Yeosu, Korea, 7.1 mm in diameter. A, Oral view; B, The manubrium with black line; C, Part of umbrella margin, showing tentacular bulbs, marginal warts, and statocysts; D, A tentacular bulb with one abaxial cirrus; E, Part of umbrella margin, showing marginal swellings with or without cirri and a statocyst containing many statoliths.

logical Laboratory, Kyoto University). In these vessels, material of every developmental stage was safely cultured and rarely died. Newly hatched *Artemia* nauplii were supplied in sufficient quantities to each medusa and polyp every day, then the seawater was changed daily. Water temperature was about 20–28°C (room temperature) during the culture. The last individual of the youngest medusa released from the polyp was obtained on May 19, 2018 in the laboratory in Shirahama, Wakayama Prefecture, Japan and cultured together with others released before.

For checking GFP patterns, a living individual was placed in a depression slide glass, and the GFP pattern was observed under OLYMPUS-SZX-RFL2 fluorescence microscope (Olympus, Tokyo, Japan) with blue light excitation, using the BP460-490 filter set, at Kinki University in Shirahama, Wakayama Prefecture.

SYSTEMATIC ACCOUNTS

Order Leptothecata Cornelius, 1992

Family Eirenidae Haeckel, 1879

¹*Genus *Eutima* McCrady, 1859

²**Eutima japonica* Uchida, 1925 (Figs. 1, 2)

Eutima japonica Uchida, 1925: 93, fig. 17; Yamazi, 1958: 136; Kramp, 1961: 197; 1965: 85; 1968: 96, fig. 260; Kubota, 1983: 296, figs. 1–25, Pl. 10; 1985: 144, figs. 1–2; 1990: 104.

Ostreohydra japonica Yamada, 1950: 117, fig. 1.

Eugymnanthea japonica: Crowell, 1957: 162; Yamada, 1959: 30; Rees, 1967: 221.

Eugymnanthea cirrhifera Kakinuma, 1964: 51, figs. 1–4; Uchida, 1964: 103, fig. 4; Rees, 1967: 221.

Eutima cirrhifera: Kubota, 1978: 125, figs. 1–11; 1979: 225,




Fig. 2. Green fluorescent protein distribution of the laboratory-reared medusa in a well-developed, 1-month-old medusa (A, B) and that of a newly liberated medusa 1 day old (C) of *Eutima japonica* from Yeosu, Korea.

Korean name: ¹*조개살이히드라속, ²*가랑잎조개살이히드라

Table 1. Measurements of laboratory-reared, well-developed medusa of *Eutima japonica* from Yeosu, Korea, about one month after release

No.	Diameter (mm)	Height (mm)	Manubrium length (mm)	Peduncle length (mm)	Thickness of mesoglea at bell apex (mm)	No. of tentacles, statocysts, marginal warts (no./quadrant), and statoliths (no./statocyst)	No. of lateral cirri, max. no./tentacular bulb or marginal wart, and max. no./quadrant
1	8.5	6.9	1.3	0.4	6.9	8, 8, 32 (4), 53 (6-8)	21, 2, 9
2	8	6	-	0.5	3.1	8, 8, 34 (4-5), 50 (5-7)	26, 2, 9
3	7.6	5.9	1.5	0.8	3.1	8, 8, 33 (3-5), 54 (6-8)	30, 2, 10
4	7.5	5.6	1.6	0.8	3.5	8, 8, 32 (4), 57 (6-8)	34, 3, 10
5	7.5	5.5	1.6	0.8	3.1	8, 8, 32 (4), 49 (5-7)	23, 3, 8
6	7.1	6.3	1.3	0.6	3.4	8, 8, 32 (4), 54 (5-8)	38, 2, 12
7	7	6.3	-	0.4	3.8	8, 8, 33 (4-5), 53 (6-9)	27, 2, 8
8	6.9	6.3	-	0.9	3	8, 8, 34 (4-5), 67 (8-9)	35, 2, 10
9	6.9	5.5	1.4	0.5	2.9	8, 8, 34 (4-5), 56 (6-8)	29, 2, 11
10	6.9	6	-	0.6	3.1	8, 8, 32 (2-5), 51 (6-7)	34, 2, 11
11	6.9	5.6	1.5	0.8	3.1	8, 8, 32 (4), 47 (5-7)	28, 2, 9
12	6.9	5.6	-	0.5	3.1	8, 8, 34 (4-5), 62 (6-13)	28, 2, 9
13	6.9	5.6	1.6	0.6	3.1	8, 8, 32 (4), 55 (6-9)	21, 2, 8
14	6.9	5	1.4	0.6	3.1	8, 8, 34 (4), 54 (6-7)	29, 2, 9
15	6.6	5	1.4	0.6	3	8, 8, 33 (4-5), 60 (6-9)	32, 2, 10
16	6.4	5	1.3	0.6	3.1	8, 8, 34 (4-5), 55 (6-8)	31, 2, 10
17	6.3	5	1.3	0.5	2.5	8, 8, 30 (3-4), 51 (5-7)	26, 2, 9
18	5.6	5	-	0.3	3.1	8, 8, 34 (4-5), 56 (6-8)	29, 2, 11

figs. 1-4.

Eucheilota intermedia Kubota, 1984: 454, figs. 1-3; 1985: 122, figs. 1-5, Pl. 1.

Material examined originally. Polyps, South Korea: Gyeongsangnam-do, Yeosu-si, Yeosu seafood market, 30 Apr 2018 (S. Kubota), from *Mytilus galloprovincialis*. After release from the polyps, 1-month-old, well-developed 18 medusae cultured in the laboratory, and among them 1.5-month-old 3 matured medusae.

Description.

Polyp: Among many zooids examined, a larger solitary polyp having 24 tentacles. Intertentacular web apparent. Medusa bud produced 1-4 in number per polyp, usually 1, on a lower part of hydrocaulus.

Newly released medusa: Medusae within one day after release from their polyps having 2-4 tentacles (N=41): 10 medusae with 2 tentacles; 3 medusae with 3 tentacles, and 28 medusae with 4 tentacles. Eight statocysts present. Each statocyst containing 1 statolith. Four marginal warts on interradial part. Mouth simple and cruciform. Many cirri present besides tentacular bulbs and marginal warts. GFP pattern observed and compared with grown-up medusa below.

Well-developed and mature medusae: A well-developed medusa about one month old, 5.6-8.5 mm in bell diameter, having 4 immature, oblong gonads, 8 tentacles, 8 statocysts, and a well-developed peduncle (Table 1, Fig. 1A). Manubrium not protruded from the umbrellar aperture (Fig. 1B). Each of 4 oral lips frilled only once (Fig. 1A, B). On

umbrellar margin, usually 4 marginal warts in each octant (Fig. 1C); 30-34 marginal warts per medusa. A total of 21-55 cirri per medusa, maximally 3 cirri on one or both sides of tentacular bulbs, marginal warts, and abaxial side of tentacular bulbs (Fig. 1D, E). A total of 47-67 statoliths, 5-13 statoliths per statocyst (Fig. 1E). Black pigmented regions on manubrium and tentacular bulbs (Fig. 1B).

GFP on manubrium, tentacular bulbs, tentacles, and umbrellar margin (Fig. 2A, B). Note that GFP on tentacular bulbs, tentacles, and interradial marginal warts of 1-day-old medusa unclear (Fig. 2C).

Three medusae reared for 1.5 months, and their morphology not undergo further noticeable changes (Table 1), except for three characters: (1) manubrium protruded from umbrellar aperture; (2) every cirrus disappeared; (3) they became male mature medusae. Largest one 9.3 mm wide, 6.4 mm high, 3.6 mm in mesoglea thickness, with a peduncle of 3.6 mm long, 82 marginal warts (10-11 warts per octant), and 115 statoliths (13-17 statoliths per statocyst).

Remarks. Although just one colony could be examined in the present study, the morphology of every developmental stage of the present material from Korea is in good accord with those of the northern Japanese form of *Eutima japonica* from Japan, that have such characteristics as a well-developed peduncle but no cirri on umbrellar margin (Kubota, 1978, 1983). Therefore, the geographical distribution of this form of the present species is widened, Japan, China and Korea. GFP pattern of medusa of the present species is clearly different from that of *Eutima sapinhoa* (Kubota, 1997;

Migotto et al., 2004; Kubota et al., 2010), and the mesoglea of the latter species is thin. It is noticeable that in Tshushima Island that is very near to the southern coasts of Korea, only intermediate form of *E. japonica* is distributed among four forms, and *Eugymnanthea japonica* is sympatrically distributed (Kubota, 1991, 2012).

ACKNOWLEDGMENTS

The author wishes to offer sincere thanks to Drs. Keitaro Kato and Yohei Washio, Kinki University, in their help to observe medusa GFP. We thank you to Mr. Hisashi Akiyama, Kujyukusima Aquarium, Nagasaki Prefecture, for occasions on examinations of polyps, and Dr. Mariko Kawamura, Seto Marine Biological Laboratory, Kyoto University, to supply the filtered, natural seawater from Shirahama for the present culture. Authors' sincere admiration is dedicated to the late Prof. Jung Hee Park who contributed to a great advance in the taxonomy of hydroids in Korea.

REFERENCES

- Calder DR, Choong HHC, Carlton JT, Chapman JW, Miller JA, Geller J, 2014. Hydroids (Cnidaria: Hydrozoa) from Japanese tsunami marine debris washing ashore in the northwestern United States. *Aquatic Invasions*, 9:425-440. <https://doi.org/10.3391/ai.2014.9.4.02>
- Crowell S, 1957. *Eugymnanthea*, a commensal hydroid living in pelecypods. *Pubblicazioni della Stazione Zoologica di Napoli*, 30:162-167.
- Kakinuma Y, 1964. A new commensal hydrozoan, *Eugymnanthea cirrhifera* n. sp. from Hachinohe. *Bulletin of the Marine Biological Station Asamushi*, 12:51-57.
- Kramp PL, 1961. Synopsis of the medusae of the world. *Journal of the Marine Biological Association of the United Kingdom*, 40:7-382. <https://doi.org/10.1017/S002531540007347>
- Kramp PL, 1965. The Hydromedusae of the Pacific and Indian Oceans. *Dana Reports*, 63:1-162.
- Kramp PL, 1968. The Hydromedusae of the Pacific and Indian Oceans. Sections II and III. *Dana-Reports*, 72:1-200.
- Kubota S, 1978. The polyp and medusae of a commensal eutimid hydroid associated with a mussel from Oshoro, Hokkaido. *Annotationes Zoologicae Japonenses*, 51:125-145.
- Kubota S, 1979. Occurrence of *Eutima cirrhifera* (Kakinuma) at Akkeshi, Hokkaido, associated with a cockle, *Clinocardium californiense*. *Annotationes Zoologicae Japonenses*, 52:225-234.
- Kubota S, 1983. Studies on life history and systematics of the Japanese commensal hydroids living in bivalves, with some references to their evolution. *Journal of Faculty of Science, Hokkaido University, Zoology*, 23:296-402.
- Kubota S, 1984. A new bivalve-inhabiting hydroid from central Japan, with reference to the evolution of the bivalve-inhabiting hydroids. *Journal of the Faculty of Science Hokkaido University, Series VI Zoology*, 23:454-467.
- Kubota S, 1985. Systematic study on a bivalve-inhabiting hydroid *Eucheilota intermedia* Kubota from central Japan. *Journal of Faculty of Science, Hokkaido University, Zoology*, 24:122-143.
- Kubota S, 1990. A new host of the southern form of *Eutima japonica* Uchida (Hydrozoa, Leptomedusae) in Japan. *Journal of Faculty of Science, Hokkaido University, Zoology*, 25:104-105.
- Kubota S, 1991. Crossing-experiments between Japanese populations of three hydrozoans symbiotic with bivalves. *Hydrobiologia*, 216/217:429-436. <https://doi.org/10.1007/BF00026496>
- Kubota S, 1992. *Eucheilota intermedia* Kubota is a distinct taxon and the third form of *Eutima japonica* Uchida (Hydrozoa: Leptomedusae). *Zoological Science*, 9:231-235.
- Kubota S, 1997. A new form of the bivalve-inhabiting hydrozoan *Eutima japonica* (Leptomedusae, Eirenidae) in Japan. *Publications of the Seto Marine Biological Laboratory*, 38:73-81. <https://doi.org/10.5134/176269>
- Kubota S, 1999. Biogeographical study on bivalve-inhabiting hydrozoans (Cnidaria, Hydrozoa) in Japan. In: *Updated progress in aquatic invertebrate zoology* (Eds., Okutani T, Ohta S, Ueshima R). Tokai University Press, Tokyo, pp. 35-39 (in Japanese).
- Kubota S, 2000. Parallel, pedomorphic evolutionary processes of the bivalve-inhabiting hydrozoans (Leptomedusae, Eirenidae) deduced from the morphology, life cycle and biogeography, with special reference to taxonomic treatment of *Eugymnanthea*. *Scientia Marina*, 64 (Supplement 1):241-247. <https://doi.org/10.3989/scimar.2000.64s1241>
- Kubota S, 2012. Distribution of two species of bivalve-inhabiting hydrozoans (Hydrozoa, Leptomedusae) at the northernmost place of Tsushima Island, Nagasaki Prefecture, Japan - GFP distribution pattern of intermediate form of *Eutima japonica*. *Bulletin of the Biogeographical Society of Japan*, 67:251-255 (in Japanese).
- Kubota S, Collins A, 2017. A single origin of bivalve-inhabiting hydrozoans (Cnidaria, Hydrozoa, Leptomedusae) in the family Eirenidae based on an analysis of 16S rRNA gene. *Biogeography*, 19:75-79.
- Kubota S, Lee J, Chang CY, 2006. Mysterious absence of bivalve-inhabiting hydrozoans (Hydrozoa, Leptomedusae) on the coasts of Korea. *Bulletin of the Biogeographical Society of Japan*, 61:71-74 (in Japanese).
- Kubota S, Nomaru E, Uchida H, Murakami A, 2010. Distribution pattern of GFP (green fluorescent protein) in a bivalve-inhabiting hydrozoan, *Eutima japonica* (Leptomedusae: Eirenidae). *Journal of Marine Biological Association of United Kingdom*, 90:1371-1374. <https://doi.org/10.1017/S0025315409991500>
- Migotto AE, Caobelli J, Kubota S, 2004. Redescription and life

- cycle of *Eutima sapinhoa* Narchi and Hebling, 1975 (Cnidaria, Hydrozoa, Leptothecata): a hydroid commensal with *Tivela mactroides* (Born) (Mollusca, Bivalvia, Veneridae). *Journal of Natural History*, 38:2533-2545. <https://doi.org/10.1080/00222930310001647316>
- Park JH, 2010. Invertebrate fauna of Korea. Vol. 4, No. 1. Cnidaria: Hydrozoa: Thecata. Thecates. Flora and Fauna of Korea. National Institute of Biological Resources, Incheon, pp. 1-183.
- Park JH, 2011. Invertebrate fauna of Korea. Vol. 4, No. 2. Cnidaria: Hydrozoa: Athecata. Athecates. Flora and fauna of Korea. National Institute of Biological Resources, Incheon, pp. 1-66.
- Park JH, 2012. Invertebrate fauna of Korea. Vol. 4, No. 3. Cnidaria: Hydrozoa: Hydromedusae. Thecates. Flora and fauna of Korea. National Institute of Biological Resources, Incheon, pp. 1-77.
- Rees WJ, 1967. A brief survey of the symbiotic associations of Cnidaria with Mollusca. *Proceedings of the Malacological Society of London*, 37:213-231. <https://doi.org/10.1093/oxfordjournals.mollus.a064991>
- Rho BJ, Park JH, 1979. A systematic study on the marine hydroids in Korea 5. Athecata hydroid. *Korean Journal of Zoology*, 22:165-174.
- Santhakumari V, Vannucci M, 1971. Monsoonal fluctuations in the distribution of the hydromedusae in the Cochin Backwater, 1968-1969. *Journal of the Marine Biological Association of India*, 13:211-219.
- Uchida T, 1925. Some Hydromedusae from northern Japan. *Japanese Journal of Zoology*, 1:77-100.
- Uchida T, 1964. Medusae of *Eugymnanthea*, an epizoic hydroid. *Publications of the Seto Marine Biological Laboratory*, 12:101-107. <https://doi.org/10.5134/175349>
- Yamada M, 1950. An epizoic athecate hydroid attached to the oyster body. *Annotationes Zoologicae Japonenses*, 23:117-118.
- Yamada M, 1959. Hydroid fauna of Japanese and its adjacent waters. *Publication of Akkeshi Marine Biological Station*, 9:1-101.
- Yamazi I, 1958. Preliminary check-list of plankton organisms found in Tanabe Bay and its environs. *Publications of the Seto Marine Biological Laboratory*, 7:111-163. <https://doi.org/10.5134/174597>
- Zhenzu X, Jiachi H, 1983. On the Hydromedusae, Siphonophora, Scyphomedusae and Ctenophora from the Jiulong River estuary of Fujian, China. *Taiwan Strait*, 2:99-110.

Received December 26, 2018
Revised April 7, 2019
Accepted April 7, 2019