



ARTICLE

## Effects of Heating Time and Temperature on Functional Properties of Proteins of Yellow Mealworm Larvae (*Tenebrio molitor* L.)

Ha-Jung Lee<sup>1</sup>, Ji-Han Kim<sup>2</sup>, Da-Som Ji<sup>1</sup>, and Chi-Ho Lee<sup>1,\*</sup>

<sup>1</sup>Department of Food Science and Biotechnology of Animal Resources, Konkuk University, Seoul 05029, Korea

<sup>2</sup>AgResearch Grasslands Research Centre, Palmerston North, 4442, New Zealand

OPEN ACCESS

Received March 21, 2019

Revised March 28, 2019

Accepted March 29, 2019

\*Corresponding author : Chi-ho Lee  
Department of Food Science and  
Biotechnology of Animal Resources,  
Konkuk University, Seoul 05029, Korea  
Tel: +82-2-450-3681  
Fax: +82-2-450-1948  
E-mail: [leech@konkuk.ac.kr](mailto:leech@konkuk.ac.kr)

\*ORCID  
Ha-jung Lee  
<https://orcid.org/0000-0001-5205-3727>  
Ji-han Kim  
<https://orcid.org/0000-0001-8197-0612>  
Da-som Ji  
<https://orcid.org/0000-0002-9212-7898>  
Chi-ho Lee  
<https://orcid.org/0000-0001-9406-8310>

**Abstract** Although the yellow mealworm (*Tenebrio molitor* L.) is a promising alternative protein source, the effects of processing conditions on functional properties are unclear. In this study, a protein extract of yellow mealworm larvae (PEYM) was subjected to different heat temperature (55°C, 75°C, and 95°C) with different time (20, 40, and 60 min) to evaluate the functional properties and protein oxidation. Different heat temperature treatment significantly affected the exposure of surface hydrophobicity of the proteins and protein molecule aggregation, which reached maximum levels at 95°C for 60 min. Protein oxidation was inversely proportional to the temperature. Both the highest carbonyl value (1.49 nmol/mg protein) and lowest thiol value (22.94 nmol/mg protein) were observed at 95°C for 60 min. The heating time-temperature interaction affected several functional properties, including solubility, emulsifying potential, and gel strength (GS). Solubility decreased near the isoelectric point (pH 5 to 6). As the temperature and heating time increased, emulsifying properties decreased and GS increased. The oil absorption capacity and foaming properties decreased and the water absorption capacity increased. These results confirmed that PEYM is a suitable source of proteins for processing and applications in the food industry.

**Keywords** yellow mealworm larvae, functional property, alternative protein, entomophagy, food processing

## Introduction

The increased demand for protein as a result of the rapid growth of the world population necessitates the development of sustainable protein production methods with minimal adverse environmental effects. Novel protein sources, such as single-cell proteins, fish protein concentrates, and edible insects, are crucial for resolving protein deficiency-mediated malnutrition (Ghaly and Aloaik, 2009; Zhao et al., 2016). In particular, interest in edible insects as protein sources is growing because they have several advantages over the livestock, including species diversity, efficient production,

and low environmental pollution with relatively high nutritional value (Van Huis, 2013). Accordingly, insect-based foods are consumed by over two billion people worldwide and are readily available in the US and European markets (Zielinska et al., 2018).

Entomophagy, the practice of eating insects, has been performed in various parts of the world since the emergence of humans (Ghaly and Aloaik, 2009). More than 2000 species of insects have been identified as edible. Insects are nutritious, with high protein and fat contents, biological value, and digestibility. Furthermore, insects are a source of micronutrients, including minerals and vitamins. Accordingly, insects are a potential supplement for various commercial foods (Barker et al., 1998; Rumpold and Schuler, 2013). Although whole insects are consumed in various regions, many consumers are still reluctant to accept this form. Therefore, processing into less recognizable forms may be required to increase consumer acceptability (Shelomi, 2015; Zielinska, et al., 2018).

Yellow mealworm (*Tenebrio molitor* L.), the larval form of the mealworm beetle, is gaining attention as an alternative protein source for various food applications. The Food and Agriculture Organization (FAO) has been estimating the potential of insects as human food and animal feed for convincing food security since 2010 (Van Huis, 2013). Many studies have reported genotoxicity, oral toxicity, nutrition composition, extraction methods, characteristics, and functional properties of proteins isolated from yellow mealworm (Finke, 2002; Han et al., 2014; Zhao et al., 2016). It is also important to understand the functional properties of proteins to improve food compositions. These functional properties, such as the water holding capacity, emulsion-forming ability, and gel formation, are dependent on physicochemical properties and pH (Zielinska et al., 2015). In addition, processing conditions, particularly heat treatment, induce conformational changes, which are associated with physicochemical properties of proteins (Lampart-Szczapa et al., 2006). Therefore, the functional properties of proteins and responses to heat treatment should be considered for practical applications.

Despite several studies of the functional properties of yellow mealworm-derived proteins, the effects of time and treatment duration are unclear. Therefore, the objective of this study was to investigate the effects of heat temperature and time on yellow mealworm-derived proteins, including physicochemical and functional properties, to provide a technical basis for their application in the food industry.

## Materials and Methods

### Materials and chemicals

Yellow mealworm (*Tenebrio molitor* L.) was purchased from a commercial market (Gyeongdong market, Seoul, Korea) and stored at  $-80^{\circ}\text{C}$  until use. Solvents were purchased from Samchun Pure Chemicals (Seoul, Korea). Other chemicals, including sodium dodecyl sulfate (SDS), 2-nitrobenzoic acid (DTNB), and 2,4-dinitrophenylhydrazine (DNPH), were obtained from Sigma-Aldrich (St. Louis, MO, USA).

### Preparation of samples

The protein extract of yellow mealworm larva (PEYM) was prepared according to the methods described by Zhao et al. (2016), with modifications. The yellow mealworm was defatted with a 5-fold volume of ether and evaporated under a vacuum at  $45^{\circ}\text{C}$  to remove the residual ether. Protein was extracted by mixing defatted yellow mealworm larvae and 0.25 M NaOH at a ratio of 1:15 (w/v) at  $45^{\circ}\text{C}$  for 60 min. The mixture was vortexed every 15 min during the extraction procedure. After centrifugation ( $3,500\times g$ , 20 min) of the slurry, the supernatant and gel fraction were separately harvested, and an

additional extraction procedure was performed using the pellet. Then, 2 mol/L HCl was added to the supernatant and gel fraction, to adjust the pH to 4.3. Following centrifugation (2,500×g, 15 min), precipitated pellets were washed twice with distilled water and lyophilized. These samples were heated in centrifuge tubes in a water bath, different heat temperature treatment (55°C, 75°C, and 95°C) and different time treatment (20, 40, and 60 min). After heating, the sample of PEYM was stored at -20°C until subsequent analyses.

### Protein solubility

The solubility of PEYM was assessed according to the methods described by Zielinska et al. (2018), with slight modifications. The PEYM was suspended in distilled water, and the pH was modified to values of 2 to 11 using 6 M HCl or NaOH. The volume of each suspension was adjusted to obtain a protein concentration of 10 mg/mL. Following solubilization in 0.5 M NaOH, the suspension was stirred for 90 min. After centrifugation (8,000×g, 15 min), the protein concentration of the supernatant was assessed by the Bradford method using BSA as a standard. Protein solubility was calculated as follows:

$$\text{Protein solubility (\%)} = (P_s/P_t) \times 100$$

where  $P_s$  is the protein content in the supernatant and  $P_t$  is the total protein content in PEYM.

### Determination of free thiol contents

The free thiol content of PEYM was determined according to Ellman's method, as described by Vossen and De Smet (2015), with minor modifications. Briefly, PEYM (2 g) was added to 50 mL of sodium dodecyl sulfate (SDS) buffer (5%, w/v) and incubated at 80°C for 1 h. After cooling and filtration, 0.5 mL of each sample was mixed with 2 mL of Tris buffer (0.1 M, pH 8.0) and 0.5 mL of 2-nitrobenzoic acid (DTNB) (10 mM in 0.1 M Tris buffer). The mixture without PEYM was used as a reagent blank, and the mixture without DTNB was used as the sample blank. All mixtures were reacted in the dark for 30 min, and absorbance was measured at 412 nm. The protein content of the mixture was determined by measuring the absorbance of the sample blank at 280 nm. The protein content was calculated using a BSA-standard curve. The thiol concentration was calculated referring to the Lambert-Beer equation ( $\epsilon_{412}=11,400 \text{ M}^{-1} \text{ cm}^{-1}$ ), and the results are presented in nM of thiol per mg protein.

### Determination of carbonyl contents

The carbonyl content of PEYM was measured as described by Vossen and De Smet (2015) using the 2,4-dinitrophenylhydrazine (DNPH) method. In particular, 1.5 g of PEYM was mixed with 15 mL of 20 mM phosphate buffer (pH 6.5) containing 0.6 mol/L NaCl. Aliquots (0.2 mL) of the mixture were mixed with 1 mL of ice-cold 10% (v/v) trichloroacetic acid (TCA) and placed in an ice bath for 15 min. Following centrifugation (2,000×g, 30 min), pellets were treated with 0.5 mL of 10 mmol/L DNPH, and the sample blank was treated with 0.5 mL of 2 M HCl. After reaction in the dark for 1 h, 0.5 mL of ice-cold 20% TCA was added. The pellets were collected by centrifugation (2,000×g, 20 min) and washed thrice with ethanol/ethyl acetate (1:1, v/v). The samples were dissolved in 1 mL of 6 M guanidine-HCl in 20 mM phosphate buffer (pH 6.5) and placed in the dark for 30 min. After centrifugation (9,500×g, 10 min), absorbance was measured at 280 and 370 nm. The relative protein concentration was calculated using BSA as standard. The carbonyl content was calculated using an absorption coefficient of  $0.021 \text{ nM}^{-1} \text{ cm}^{-1}$  at 370 nm, and results are presented as nM of carbonyl per mg of protein.

### Evaluation of aggregation

The aggregation ability of PEYM was determined according to the Nile red method described by Sante-Lhoutellier et al. (2008). Briefly, 1 mg of PEYM was suspended in 1 mL of 20 mM phosphate buffer (pH 6), and 10  $\mu$ L of Nile Red stock solution was added. The fluorescence intensity was measured at an excitation ( $\lambda_{ex}$ ) wavelength of 560 nm and an emission ( $\lambda_{em}$ ) wavelength of 620 nm using a microplate spectrofluorometer. Results are expressed as arbitrary units (au).

### Determination of surface hydrophobicity

Surface hydrophobicity was determined with the chromophore bromophenol blue (BPB) as described by Sante-Lhoutellier et al. (2008). Briefly, 2 mg of PEYM was suspended in 1 mL of 20 mM phosphate buffer (pH 6). Next, 1 mL of the sample was mixed with 40  $\mu$ L of BPB (1 mg/mL in distilled water) and reacted for 10 min. Following centrifugation (2,000 $\times$ g, 15 min), the absorbance of the supernatant was measured at 595 nm.

### Water absorption capacity

The water absorption capacity (WAC) was determined as described by Zielinska et al. (2018), with some modifications. Briefly, PEYM (0.5 g) was mixed with 20 mL of distilled water and stirred at 55 $\times$ g for 30 min. After centrifugation (8,000 $\times$ g, 15 min), the weight of the precipitate was compared with the initial weight. The results are expressed as g of absorbed water per g of sample.

### Oil absorption capacity

The oil absorption capacity (OAC) of PEYM was determined according to the methods described by Zielinska et al. (2018). The PEYM (0.5 g) was mixed with 10 mL of vegetable oil and stirred for 30 s. Following centrifugation (8,000 $\times$ g, 15 min), the sediment weight was compared with the initial weight. The results are expressed as g of oil absorbed per g of sample.

### Determination of emulsifying activity

The emulsifying properties were assessed according to the methods described by Zielinska et al. (2018), with some slight modifications. Briefly, 0.1 g of PEYM was suspended in 10 mL of distilled water and mixed with an equal volume of vegetable oil. Following homogenization (3,500 $\times$ g, 1 min), the mixture was centrifuged (3,000 $\times$ g, 5 min). The height of each layer was measured. The emulsion stability (ES) was determined by heating at 80°C for 30 min. The mixture was centrifuged (3,000 $\times$ g, 5 min). The emulsion activity (EA) and stability were calculated as follows:

$$\text{Emulsion activity (EA) (\%)} = (V_1/V) \times 100$$

$$\text{Emulsion stability (ES) (\%)} = (V_2/V_1) \times 100$$

where  $V$  is the total volume of the tube contents,  $V_1$  is the volume of the emulsified layer, and  $V_2$  is the volume of the emulsified layer after heating.

### Determination of foaming ability

The foaming properties of PEYM were determined by the methods described by Guo et al. (2015), with slight modifications. Briefly, 0.2 g of PEYM was suspended in 20 mL of distilled water and blended using a homogenizer (2,000×g, 2 min). The sample was transferred into a cylinder, and the total volume was determined. After standing for 30 min, the foaming stability (FS) was determined. The foaming capacity and stability were calculated as follows:

$$\text{Foaming capacity (FC) (\%)} = [(V_0 - V) / V] \times 100$$

$$\text{Foaming stability (FS) (\%)} = (V_{30} / V_0) \times 100$$

where  $V$  is the volume before whipping (mL),  $V_0$  is the volume after whipping (mL), and  $V_{30}$  is the volume after standing (mL).

### Evaluation of gel strength

The effect of PEYM on gel strength (GS) was evaluated following the methods described by Yi et al. (2013), with modifications. Gel strength was determined using a TX-XT2 instrument (Stable Micro Systems Ltd, Surrey, UK). A spherical stainless probe (0.25 inch) was used to penetrate 90% of the PEYM gel length (10×10 mm) and the peak penetration to reach the breaking point was determined. Gel strength (g×mm) was evaluated by the force-deformation curves based on the multiplication of force (g) by distance (mm).

### Statistical analysis

All experiments were conducted in triplicate, and the results are presented as means±SEM. Statistical analyses were performed using IBM SPSS 24.0 (SPSS, Chicago, IL, USA). Statistical differences were determined by one-way and two-way analyses of variance and *post hoc* Tukey HSD tests. A value of  $p < 0.05$  was considered statistically significant.

## Results and Discussion

### Protein solubility

The results of protein solubility are shown in Table 1, solubility was significantly lower at pH values of 5 and 6 ( $p < 0.05$ , respectively), than at alkaline (pH 11) and acidic (pH 2) values. Protein solubility gradually increased from pH 5 to 10 and showed a maximum of 88%. These results agree with those of Seena and Sridhar (2005), who reported that proteins have net positive and negative charges at alkaline and acidic pH values. Thus, the isoelectric point of PEYM was observed at pH 5 and 6, which was significantly affected by the heating time-temperature interaction ( $p < 0.01$  and  $p < 0.05$ , respectively). The solubility increased with increasing the heating time for 40 to 60 min at each different temperature while decreased with increasing the temperature at 75°C and 95°C for each different heating time. According to Timelisena et al. (2016) and Pelegrine and Gasparetto (2005), the solubility is influenced by protein-water interaction and unfolded protein molecule at under the protein denaturation temperature and affects the binding of the secondary and tertiary structures of the polypeptide chain, resulting in hydration. On the other hand, above the protein denaturation temperature, reducing the protein-water interactions and exposure of hydrophobicity groups, resulting in decreased solubility. These results correspond with the results of previous protein studies of whey proteins solubility (Pelegrine and Gasparetto, 2005). Thus, the protein solubility results indicated that pH as well as heating time and temperature interaction influence protein solubility.

**Table 1.** Effects of time and heat on the solubility of protein extracted from yellow mealworm larvae

Protein solubility	Protein extracted of yellow mealworm larvae										SEM
	Control	55°C (min)			75°C (min)			95°C (min)			
		20	40	60	20	40	60	20	40	60	
pH 2	69.81 <sup>aA</sup>	54.2 <sup>bB</sup>	54.82 <sup>bB</sup>	59.98 <sup>cB</sup>	52.97 <sup>bB</sup>	56.06 <sup>bB</sup>	52.63 <sup>cB</sup>	39.13 <sup>bB</sup>	42.81 <sup>bB</sup>	42.57 <sup>cB</sup>	±4.02
pH 3	50.88 <sup>aA</sup>	49.35 <sup>abB</sup>	40.67 <sup>abB</sup>	52.21 <sup>bB</sup>	38.11 <sup>abB</sup>	43.19 <sup>abB</sup>	47.67 <sup>bB</sup>	40.72 <sup>abAB</sup>	41.37 <sup>abAB</sup>	41.29 <sup>bAB</sup>	±3.23
pH 4	39.36 <sup>aAB</sup>	29.11 <sup>aB</sup>	35.96 <sup>aB</sup>	47.70 <sup>aB</sup>	33.29 <sup>aB</sup>	38.61 <sup>aB</sup>	48.81 <sup>aB</sup>	32.54 <sup>aA</sup>	27.07 <sup>aA</sup>	36.50 <sup>aA</sup>	±3.46
pH 5	38.48 <sup>aA</sup>	27.69 <sup>abB*</sup>	31.17 <sup>abB*</sup>	46.00 <sup>bB*</sup>	28.18 <sup>abB*</sup>	36.85 <sup>abB*</sup>	38.62 <sup>bB*</sup>	28.97 <sup>abA*</sup>	29.66 <sup>abA*</sup>	31.98 <sup>bA*</sup>	±2.36
pH 6	45.15 <sup>aA</sup>	29.28 <sup>bcC**</sup>	29.06 <sup>bcC**</sup>	44.15 <sup>cC**</sup>	28.43 <sup>bcC**</sup>	35.07 <sup>bcC**</sup>	37.78 <sup>cC**</sup>	27.24 <sup>bcB**</sup>	26.28 <sup>bB**</sup>	31.04 <sup>cB**</sup>	±2.30
pH 7	53.47 <sup>aA</sup>	29.52 <sup>bC</sup>	32.01 <sup>bC</sup>	42.82 <sup>bC</sup>	29.77 <sup>bBC</sup>	33.71 <sup>bBC</sup>	36.96 <sup>bBC</sup>	27.68 <sup>bB</sup>	31.11 <sup>bB</sup>	31.07 <sup>bB</sup>	±2.50
pH 8	56.30 <sup>aA</sup>	39.30 <sup>bB</sup>	41.40 <sup>bB</sup>	43.24 <sup>cB</sup>	38.76 <sup>bB</sup>	41.30 <sup>bB</sup>	40.10 <sup>cB</sup>	33.31 <sup>bB</sup>	29.84 <sup>bB</sup>	31.52 <sup>cB</sup>	±3.28
pH 9	63.68 <sup>aA</sup>	40.27 <sup>bB</sup>	48.70 <sup>bB</sup>	50.57 <sup>cB</sup>	48.82 <sup>bB</sup>	46.80 <sup>bB</sup>	43.02 <sup>cB</sup>	36.76 <sup>bB</sup>	35.58 <sup>bB</sup>	32.44 <sup>cB</sup>	±4.33
pH 10	85.42 <sup>aA</sup>	49.13 <sup>bB</sup>	63.57 <sup>bB</sup>	55.31 <sup>bB</sup>	60.24 <sup>bB</sup>	57.92 <sup>bB</sup>	51.16 <sup>bB</sup>	45.59 <sup>bB</sup>	66.60 <sup>bB</sup>	35.43 <sup>bB</sup>	±6.46
pH 11	97.88 <sup>aA</sup>	68.38 <sup>bB</sup>	82.92 <sup>bB</sup>	75.30 <sup>bB</sup>	75.27 <sup>bB</sup>	65.77 <sup>bB</sup>	88.82 <sup>bB</sup>	71.52 <sup>bB</sup>	66.45 <sup>bB</sup>	58.46 <sup>bB</sup>	±7.55

All values are expressed as means±SEM of triplicate experiments.

Indicate within a column significant difference between heating time and temperature interaction (\* p<0.05, \*\* p<0.01).


<sup>a-c</sup> denotes differences with respect to time (p<0.05) while <sup>A-C</sup> indicate significant effects of heat treatment (p<0.05).

### Aggregation

Aggregation results for PEYM are shown in Fig. 1A. The significant effects of time and the interaction on the aggregation was not detected (p>0.05). The aggregation tended to increase linearly with increasing the time, although there was not a significant difference. Aggregation increased significantly during heat temperature treatment (p<0.05) based on the increase in fluorescence as the temperature increased. Fig. 1A depicts the gradual increase in fluorescence with respect to heat and time. The most significant increase in fluorescence was observed at 95°C (64.25 au). According to Mahler et al. (2009), high temperatures affect the secondary, tertiary, and quaternary structures of polypeptide chains. During heating, aggregates form, releasing hydrophobic groups by protein-protein interactions. Our results indicated that protein aggregation increases with increasing temperatures.

### Surface hydrophobicity

Surface hydrophobicity influences the surface characteristics of proteins, such as protein-lipid interactions and protein-protein interactions (Timilsena et al., 2016). As shown in Fig. 1B, as the temperature increased, the surface hydrophobicity increased significantly (30.35 to 118.16 µg; p<0.01); however, heating time showed no significant difference (p>0.05). Chelh et al. (2006) suggested that oxidation increases surface hydrophobicity due to the cleavage of a specific peptide chain as the heating time and temperature increase. These results are in agreement with those of Sun et al. (2013), who reported that treatment at 40°C for 60 min significantly increased (p<0.05) hydrophobicity. Thus, the increase in surface hydrophobicity suggested that the structure of PEYM changed in response to heat treatment, and the exposure of hydrophobic groups was measured.




**Fig. 1. Aggregation (A) and surface hydrophobicity (B) of yellow mealworm larva protein, as measured by Nile Red fluorescence and BPB probes (Bromophenol blue).** All values are expressed as means $\pm$ SEM of triplicate experiments. Values with different letters are significantly different, \*  $p < 0.05$ , \*\*  $p < 0.01$  as compared to the control. 2M55, 20 min at 55°C; 2M75, 20 min at 75°C; 2M95, 20 min at 95°C; 4M55, 40 min, 55°C; 4M75, 40 min at 75°C; 4M95, 40 min at 95°C; 6M55, 60 min at 55°C; 6M75, 60 min at 75°C; 6M95, 60 min at 95°C.

### Protein oxidation

As shown in Fig. 2A, carbonyl contents increased significantly as the temperature increased (0.93 to 1.49 nmol/mg protein) ( $p < 0.01$ ). At 95°C, carbonyl contents showed a tendency to increase rapidly. Previous studies have reported increases in carbonyl contents by heat and time (Roldan et al., 2014). Heat and time induce deformation by increasing oxidative stress (Sante-Lhoutellier et al., 2008). According to Roldan et al. (2014), heat and time interaction increased carbonyl compounds and, compared to time treatment, heat treatment directly affects the formation of carbonyl compounds. However, the results of this experiment did not confirm the effect of time on the carbonyl content ( $p > 0.05$ ). Estevez (2011) found that the formation of carbonyl compounds from amino acid chains contributes to the denaturation of a protein and decreases its functionality, which affects protein quality. These results suggest that further studies are needed to understand the formation of carbonyl compounds with heat treatment under various time treatment.

The free thiol contents are reported to decrease due to disulfide conversion during initial protein oxidation, negatively affecting the digestibility and nutritional value of food (Soyer et al., 2010). As shown in Fig. 2B, the free thiol content



**Fig. 2.** Effects of time and temperature on carbonyl (A) and free thiol contents (B) of yellow mealworm larva protein. All values are expressed as means $\pm$ SEM of triplicate experiments. Values with different letters are significantly different, \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$  as compared to the control. 2M55, 20 min at 55°C; 2M75, 20 min at 75°C; 2M95, 20 min at 95°C; 4M55, 40 min, 55°C; 4M75, 40 min at 75°C; 4M95, 40 min at 95°C; 6M55, 60 min at 55°C; 6M75, 60 min at 75°C; 6M95, 60 min at 95°C.

decreased significantly ( $p < 0.001$ ) as the temperature increased (69.63 to 22.94 nmol/mg protein). Heat treatment enhances the exposure of free thiol groups and the exteriorization of hydrophobic residues, resulting in the destruction of hydrogen bonds, formation of disulfides, and induction of protein aggregation (Traore et al., 2012). The results of this experiment suggest that thiol contents in PEYM are reduced by heat treatment. The thiol content of PEYM was lower than those of egg white (58.5 to 9.64 mg of protein/mL) and meat (Lund et al., 2008; Van der Plancken et al., 2005), indicating that food processing involving heat treatment at 55°C, 75°C, and 95°C is possible.

### Water absorption capacity and oil absorption capacity

Water and protein interactions affect the functional properties and texture of food, as water absorption increases condensation and viscosity which has higher WAC, the higher the usefulness in the meat and bakery industries (Damodaran,



2017). As shown in Table 2, WAC did not differ significantly with respect to temperature and heating time (1.02 to 1.18 g/g), consistent with the results of Zhao et al. (2016) (1.29 g/g). Zielinska et al. (2018) reported that WAC could be enhanced depending on the treatment conditions for yellow mealworm protein (1.87 to 3.95 g/g). Thus, the WAC of PEYM can likely be improved by other factors, such as pH, concentration, and other extraction methods.

The result of OAC is presented in Table 2. OAC was significantly influenced by different temperatures (55°C, 75°C, and 95°C;  $p < 0.001$ ) and all the heating periods showed significantly lower values than those in the control group ( $p < 0.05$ ). OAC values decreased as the heating time increased at each different temperature treatments 55°C, 75°C, and 95°C (1.62 to 1.18, 1.66 to 1.32, and 1.74 to 1.33 g/g, respectively). Heat temperature treatment collapses the protein network and decreases OAC (Yin et al., 2008). Additionally, the formation of protein aggregates and the exposure of hydrophobic amino acid would decline the protein-oil absorption capacity (Zhao et al., 2016). These results indicated that the heating time and temperature affect the interactions of hydrophobic amino acids with oil, corresponding with the results of previous protein studies of soybean (1.1 g/g) and pea protein (1.2 g/g) (Naik et al., 2012; Shevkani et al., 2015). Furthermore, they support the use of PEYM as a food additive.

### Emulsion activity and emulsion stability

The results of EA and ES are shown in Table 2. EA and ES tended to decrease as the time increased at each different heat temperature treatments (55°C, 75°C, and 95°C). EA decreased significantly from 46.46% to 34.94% in response to the heating time-temperature interaction ( $p < 0.001$ ) and ES also decreased significantly from 87.59 to 69.39% in response to different heat temperature treatment ( $p < 0.001$ ). According to McClements (2004), the temperature exceeds the critical value and the protein unfolds exposing the internal non-polar amino groups, causing it to become insoluble or promoting protein-protein interactions. Heat denaturation also creates reactive thiol groups, promoting protein interactions and leading to changes in droplet flocculation and coalescence (McClements, 2004). Our results confirmed that EA and ES decrease by heat treatment and the observed decreases might be correlated with denaturation and droplet size. The EA results also agree with those of other studies of yellow mealworm protein (Gould and Wolf, 2018; Zielinska et al, 2018), suggesting that it is a new source of

**Table 2. Effects of time and heat treatment on functional properties of yellow mealworm larva protein**

Properties	Control	Protein extracted of yellow mealworm larvae									SEM
		55°C (min)			75°C (min)			95°C (min)			
		20	40	60	20	40	60	20	40	60	
WAC (g/g)	1.02 <sup>bb</sup>	1.05 <sup>abAB</sup>	1.08 <sup>abAB</sup>	1.10 <sup>aAB</sup>	1.07 <sup>abAB</sup>	1.08 <sup>abAB</sup>	1.12 <sup>aAB</sup>	1.07 <sup>abA</sup>	1.12 <sup>abA</sup>	1.18 <sup>aA</sup>	±0.04
OAC (g/g)	1.70 <sup>aA</sup>	1.62 <sup>bA</sup>	1.46 <sup>bA</sup>	1.18 <sup>bA</sup>	1.66 <sup>bB</sup>	1.44 <sup>bB</sup>	1.32 <sup>bB</sup>	1.74 <sup>bC</sup>	1.48 <sup>bC</sup>	1.33 <sup>bC</sup>	±0.05
EA (%)	49.21 <sup>aA</sup>	49.15 <sup>baA**</sup>	48.76 <sup>baA**</sup>	28.94 <sup>baA**</sup>	41.47 <sup>baA**</sup>	41.14 <sup>baA**</sup>	37.87 <sup>baA**</sup>	48.75 <sup>bbB**</sup>	42.37 <sup>bbB**</sup>	38.00 <sup>bbB**</sup>	±2.13
ES (%)	84.87 <sup>aA</sup>	94.09 <sup>aA</sup>	81.59 <sup>aA</sup>	64.97 <sup>aA</sup>	83.86 <sup>aAB</sup>	81.10 <sup>aAB</sup>	78.03 <sup>aAB</sup>	84.82 <sup>aB</sup>	87.89 <sup>aB</sup>	65.16 <sup>aB</sup>	±5.48
FC (%)	21.48 <sup>aA</sup>	9.70 <sup>bB</sup>	9.22 <sup>bB</sup>	6.50 <sup>cB</sup>	12.35 <sup>bB</sup>	10.97 <sup>bB</sup>	8.19 <sup>cB</sup>	8.18 <sup>bC</sup>	5.06 <sup>bC</sup>	3.58 <sup>cC</sup>	±0.85
FS (%)	88.26 <sup>cC</sup>	91.62 <sup>bbB</sup>	91.76 <sup>bbB</sup>	94.3 <sup>aB</sup>	89.98 <sup>baB</sup>	92.05 <sup>baB</sup>	93.82 <sup>aB</sup>	93.15 <sup>aA</sup>	96.80 <sup>aA</sup>	97.37 <sup>aA</sup>	±0.76
GS (g×mm)	312.47 <sup>cB*</sup>	234.21 <sup>bcB*</sup>	386.77 <sup>bbB*</sup>	1,080.6 <sup>abB*</sup>	369.07 <sup>bcB*</sup>	417.96 <sup>bbB*</sup>	1,796.68 <sup>abB*</sup>	446.92 <sup>bcA*</sup>	1,222.21 <sup>baA*</sup>	2,547.37 <sup>baA*</sup>	±175.85

Indicate within a column significant difference between heating time and temperature interaction (\*  $p < 0.01$ , \*\*  $p < 0.001$ ). All values are expressed as means±SEM of triplicate experiments.

Difference letters indicate significant differences; <sup>a-c</sup> indicate significant differences ( $p < 0.05$ ) and <sup>d-f</sup> indicate significant differences ( $p < 0.001$ ) with respect to time and <sup>A-C</sup> indicate significant effects of heat treatment ( $p < 0.05$ ).

WAC, water absorption capacity; OAC, oil absorption capacity; EA, emulsion activity; ES, emulsion stability; FC, foaming capacity; FS, foaming stability; GS, gel strength.

protein emulsifier for food formulations.

### **Foaming capacity and foaming stability**

The results of FC and FS are shown in Table 1, FC was significantly decreased as the temperature increased (12.3% to 3.58%,  $p < 0.001$ ) and FS was significantly increased as time and temperature increased (88.26% to 97.37%,  $p < 0.001$ ). The lowest value of FC was observed at 95°C (3.58%), while the highest value of FS was observed for 60 min (97.37%). Previous studies have shown that the high sugar content in mealworms induces passive protein-protein interactions, leading to a weak foaming property due to the formation of a weak interfacial membrane during foam formation (Yi et al., 2013; Zielinska et al., 2018). According to Bals and Kulozik (2003), protein denaturation is associated with structural changes, resulting in aggregation and reactive thiol groups. The aggregates do not readily form an interfacial film. The surface load becomes smaller, and foam rigidity becomes lower, negatively affecting FC and FS. Therefore, the FC of PEYM appeared to be low and was negatively influenced by time and heat.

### **Gel strength**

Gelation is induced by protein aggregation and network formation and contributes to the sensory properties and texture of food (Renkema and van Vliet, 2002). GS was significantly higher in the heating time-temperature interaction (234.21 to 2,547.37  $\text{g} \times \text{mm}$ ;  $p < 0.001$ ). In particular, the highest GS was obtained at 95°C for 60 min. A previous study has demonstrated that heating time during gel formation affects the unfolding of protein structures and protein formation (Totosaus et al., 2002). Increasing temperatures results in gel network formation via peptide aggregation (Totosaus et al., 2002). In this experiment, protein-protein bond formation increased by time and temperature treatment. These results confirmed that gels had stronger networks than those of the control group.

## **Conclusion**

In this study, we studied the effects of heating time and temperature on the functional properties and oxidation of PEYM. Notably, heat temperature treatment significantly influenced the surface hydrophobicity, aggregation, WAC, OAC, EA, ES, GS, carbonyl content, and thiol content of PEYM. High temperatures increased surface hydrophobicity and aggregation, induced protein oxidation, and altered the functional properties of proteins. This suggests that the heat treatment induces changes in protein properties; however, functional properties of PEYM can be maintained. Regarding the heat exposure, no significant change in functional properties and protein oxidation of PEYM were detected showing that PEYM is stable for a long time. In terms of protein denaturation (55°C to 95°C), the functional properties of PEYM were stable and appropriate for the parameters required for the food formulation and thermal processing. These results indicated that PEYM is a suitable alternative source of protein for thermal processing and the formulation of protein products. Further quantitative studies are needed to evaluate the practical applications of proteins derived from yellow mealworm larvae.

## **Conflict of Interest**

The authors declare no potential conflict of interest.

## Acknowledgements

This paper was supported by Konkuk University in 2018.

## Author Contributions

Conceptualization: Kim JH, Lee HJ. Data curation: Lee HJ. Formal analysis: Lee HJ. Methodology: Lee HJ, Kim JH, Ji DS. Software: Lee HJ, Kim JH. Validation: Lee HJ. Investigation: Lee HJ. Writing - original draft: Lee HJ, Kim JH. Writing - review & editing: Lee HJ, Kim JH, Ji DS, Lee CH.

## Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

## References

- Bals A, Kulozik U. 2003. Effect of pre-heating on the foaming properties of whey protein isolate using a membrane foaming apparatus. *Int Dairy J* 13:903-908.
- Barker D, Fitzpatrick MP, Dierenfeld ES. 1998. Nutrient composition of selected whole invertebrates. *Zoo Biol* 17:123-134.
- Chelh I, Gatellier P, Sante-Lhoutellier V. (2006). Technical note: A simplified procedure for myofibril hydrophobicity determination. *Meat Sci* 74:681-683.
- Damodaran S, Paraf A. 2017. *Food proteins and their applications*. Routledge, New York, NY, USA. p 694.
- Estevez M. 2011. Protein carbonyls in meat systems: A review. *Meat Sci* 89:259-279.
- Finke MD. 2002. Complete nutrient composition of commercially raised invertebrates used as food for insectivores. *Zoo Biol* 21:269-285.
- Ghaly AE, Alkoik FN. 2009. The yellow mealworm as a novel source of protein. *Am J Agric Biol Sci* 4:319-331.
- Gould J, Wolf B. 2018. Interfacial and emulsifying properties of mealworm protein at the oil/water interface. *Food Hydrocoll* 77:57-65.
- Guo F, Xiong YL, Qin F, Jian H, Huang X, Chen J. 2015. Surface properties of heat-induced soluble soy protein aggregates of different molecular masses. *J Food Sci* 80:C279-C287.
- Han SR, Yun EY, Kim JY, Hwang JS, Jeong EJ, Moon KS. 2014. Evaluation of genotoxicity and 28-day oral dose toxicity on freeze-dried powder of *Tenebrio molitor* larvae (Yellow Mealworm). *Toxicol Res* 30:121-130.
- Lampart-Szczapa E, Konieczny P, Nogala-Kalucka M, Walczak S, Kossowska I, Malinowska M. 2006. Some functional properties of lupin proteins modified by lactic fermentation and extrusion. *Food Chem* 96:290-296.
- Lund MN, Hviid MS, Claudi-Magnussen C, Skibsted LH. 2008. Effects of dietary soybean oil on lipid and protein oxidation in pork patties during chill storage. *Meat Sci* 79:727-733.
- Mahler HC, Friess W, Grauschopf U, Kiese S. 2009. Protein aggregation: Pathways, induction factors and analysis. *J Pharm Sci* 98:2909-2934.
- McClements DJ. 2004. Protein-stabilized emulsions. *Curr Opin Colloid Interface Sci* 9:305-313.
- Naik A, Raghavendra SN, Raghavarao KSMS. 2012. Production of coconut protein powder from coconut wet processing

- waste and its characterization. *Appl Biochem Biotechnol* 167:1290-1302.
- Pelegri DHG, Gasparetto CA. 2005. Whey proteins solubility as function of temperature and pH. *LWT-Food Sci Technol* 38:77-80.
- Renkema JMS, Van Vliet T. 2002. Heat-induced gel formation by soy proteins at neutral pH. *J Agric Food Chem* 50:1569-1573.
- Roldan M, Antequera T, Armenteros M, Ruiz J. 2014. Effect of different temperature-time combinations on lipid and protein oxidation of sous-vide cooked lamb loins. *Food Chem* 149:129-136.
- Rumpold BA, Schuler OK. 2013. Nutritional composition and safety aspects of edible insects. *Mol Nutri Food Res* 57:802-823.
- Sante-Lhoutellier V, Astruc T, Marinova P, Greve E, Gatellier P. 2008. Effect of meat cooking on physicochemical state and *in vitro* digestibility of myofibrillar proteins. *J Agric Food Chem* 56:1488-1494.
- Seena S, Sridhar KR. 2005. Physicochemical, functional and cooking properties of under explored legumes, *Canavalia* of the southwest coast of India. *Food Res Int* 38:803-814.
- Shelomi M. 2015. Why we still don't eat insects: Assessing entomophagy promotion through a diffusion of innovations framework. *Trends Food Sci Technol* 45:311-318.
- Shevkani K, Kaur A, Kumar S, Singh N. 2015. Cowpea protein isolates: Functional properties and application in gluten-free rice muffins. *LWT-Food Sci Technol* 63:927-933.
- Soyer A, Ozalp B, Dalmis U, Bilgin V. 2010. Effects of freezing temperature and duration of frozen storage on lipid and protein oxidation in chicken meat. *Food Chem* 120:1025-1030.
- Sun W, Zhou F, Sun DW, Zhao M. 2013. Effect of oxidation on the emulsifying properties of myofibrillar proteins. *Food Bioprocess Technol* 6:1703-1712.
- Timilsena YP, Adhikari R, Barrow CJ, Adhikari B. 2016. Physicochemical and functional properties of protein isolate produced from Australian chia seeds. *Food Chem* 212:648-656.
- Totosaus A, Montejano JG, Salazar, JA, Guerrero I. 2002. A review of physical and chemical protein-gel induction. *Int J Food Sci Technol* 37:589-601.
- Traore S, Aubry L, Gatellier P, Przybylski W, Jaworska D, Kajak-Siemaszko K, Sante-Lhoutellier V. 2012. Effect of heat treatment on protein oxidation in pig meat. *Meat Sci* 91:14-21.
- Van der Plancken I, Van Loey A, Hendrickx MEG. 2005. Changes in sulfhydryl content of egg white proteins due to heat and pressure treatment. *J Agric Food Chem* 53:5726-5733.
- Van Huis A. 2013. Potential of insects as food and feed in assuring food security. *Annu Rev Entomol* 58:563-583.
- Vossen E, De Smet S. 2015. Protein oxidation and protein nitration influenced by sodium nitrite in two different meat model systems. *J Agric Food Chem* 63:2550-2556.
- Yi L, Lakemond CMM, Sagis LMC, Eisner-Schadler V, Van Huis A, Van Boekel MAJS. 2013. Extraction and characterisation of protein fractions from five insect species. *Food Chem* 141:3341-3348.
- Yin SW, Tang CH, Cao JS, Hu EK, Wen QB, Yang XQ. 2008. Effects of limited enzymatic hydrolysis with trypsin on the functional properties of hemp (*Cannabis sativa* L.) protein isolate. *Food Chem* 106:1004-1013.
- Zhao X, Vazquez-Gutierrez JL, Johansson DP, Landberg, R, Langton M. 2016. Yellow mealworm protein for food purposes - Extraction and functional properties. *PLOS ONE* 11:e0147791.
- Zielinska E, Baraniak B, Karas M, Rybczynska, K, Jakubczyk A. 2015. Selected species of edible insects as a source of

nutrient composition. *Food Res Int* 77:460-466.

Zielinska E, Karas M, Baraniak B. 2018. Comparison of functional properties of edible insects and protein preparations thereof. *LWT-Food Sci Technol* 91:168-174.