

RESEARCH ARTICLE

Open Access



Protective effects of *Hizikia fusiforme* and *Chlorella* sp. extracts against lead acetate-induced hepatotoxicity in rats

Joo hyun Park¹, Jeong-Wook Choi², Min-Kyeong Lee², Youn Hee Choi³ and Taek-Jeong Nam^{1,2*} 

Abstract

In the present study, the protective effects of *Hizikia fusiforme* and *Chlorella* sp. extracts on lead acetate-induced hepatotoxicity were investigated. Hepatic damage was induced in rats by intraperitoneal (i.p.) injection of lead acetate and the protective effects of *H. fusiforme* (HZK) and *Chlorella* sp. (CHL) extracts on lead acetate-induced hepatic damage in rat liver were examined. The results revealed significantly increased glutamic oxaloacetate and glutamic pyruvic transaminase levels in the group treated with lead acetate only (Pb group); oral administration of HZK and CHL extracts tended to decrease the enzyme levels similar to those observed in the control group. Regarding antioxidant enzymes, superoxide dismutase activity was increased in the Pb group and decreased in a concentration-dependent manner in the HZK- and CHL-treated groups. Glutathione levels were increased in a concentration-dependent manner in the HZK- and CHL-treated groups. There was no significant difference in catalase activity. Western blot analysis showed inflammation-related protein expression in mitogen-activated protein kinase and Nrf2 pathways was affected in the HZK- and CHL-treated groups. Therefore, HZK and CHL extracts exerted antioxidant and anti-inflammatory effects against lead acetate-induced hepatotoxicity. Development of functional health foods containing HZK and CHL extracts, which have hepatoprotective effects against inhaled lead acetate, should be considered.

Keywords: Antioxidant, Anti-inflammation, *Hizikia fusiforme*, *Chlorella* sp.

Background

The development of industrial technology and the increased number of automobiles and factories cause various types of environmental pollution. For instance, air pollution is of serious concern, as it directly affects human health (Meo and Suraya 2015). Among air pollutants, fine dust causes various diseases. Since dust with small particle size has become a major health hazard, most countries have set environmental standards for particulate matter with a diameter of 10 μm (PM)₁₀ or 2.5 μm (PM)_{2.5} (Hsiao et al. 2000; Zhou et al. 2016). Dust is mainly emitted by automobile exhaust gases and factory chimneys. Heavy metals such as lead, cadmium, zinc, nickel, and chromium contained in fine dust are

absorbed into the human body and cause various respiratory and circulatory diseases through different mechanisms, such as the generation of active oxygen species (Donaldson and MacNee 2001). A previous study found that the lead content in fine dust was significantly higher than in soil (Hu et al. 2014).

Seaweed is consumed as food in large quantities in Korea, Japan, and China. The carbohydrates contained in seaweed are effective in promoting intestinal motility, heavy metal elimination, and improving hyperlipemia (Lee and Kim 2008; Kim et al. 2011). *Hizikia fusiforme* is not only a good source of dietary fiber but also fucoindans, in particular jaminaran, a neutral polysaccharide with functional properties such as blood coagulation and immunity enhancement (Wua et al. 2013). *Chlorella* is a single-celled green alga rich in vitamins and minerals, and is effective in improving overall health and well-being. *Chlorella* contains a physiologically active substance, chlorella growth factor, which reportedly

* Correspondence: namtj@pknu.ac.kr

¹Departments of Food Science and Nutrition, Pukyong National University, Yongso-ro, Nam-gu, Busan 48513, Republic of Korea

²Institute of Fisheries Science, Pukyong National University, Busan 46041, Republic of Korea

Full list of author information is available at the end of the article



possesses anti-cancer effects and is effective in child growth and development as well as immune enhancement (Hidalgo et al. 2014).

Given this background, we investigated the antioxidative and protective effects and related mechanisms of *H. fusiforme* and *Chlorella* sp. on lead-induced hepatotoxicity.

Methods

Sample preparation

H. fusiforme was washed three times with water to remove salt and impurities, cleaned, and dried in a freeze dryer (Eyela, Tokyo, Japan). Dried *H. fusiforme* was ground into a powder (below 50 mesh) using a food blender (Daesung, Gyeonggi-do, Korea). *Chlorella* sp. was purchased as 100% *Chlorella* powder (below 70 mesh) from Novarex Inc. (Seoul, Korea).

H. fusiforme and *Chlorella* sp. powders (40 g) were suspended in 1 L distilled water and stirred for 4 h at room temperature (RT). The suspension was centrifuged at 3000×g (4 °C, 10 min) and vacuum-filtered, followed by the addition of triple volumes (total quantity of filtrate: ×3) of 95% ethanol and remained for 24 h at RT. After that, the solution was filtered (Whatman no.4, GE Healthcare, Buckinghamshire, UK) and concentrated using rotary evaporation at 40 °C. The concentrated solution was distributed into 10 mL tubes and freeze-dried to produce *H. fusiforme* (HZK) and *Chlorella* sp. (CHL) extracts, which were stored at −70 °C until use (Fig. 1).

Sample protein and carbohydrate content

Extract protein contents were measured by protein quantification in a Bradford assay. To quantify protein concentration, 1 mg of extract powder was dissolve in 20 mL distilled water and 20 μL of dissolved extract were incubated with 180 μL of Bradford reagent at RT. The absorbance at 595 nm was read after 5 min. The


results showed that the protein content of *H. fusiforme* extract was 54.1% and the *Chlorella* sp. extract was 57.4%.

Extract carbohydrate contents were measured by phenol-sulfuric acid method. One milligram of extract powder was dissolve in 20 mL distilled water and 0.5 mL of dissolved extract was mixed with 0.5 mL of 5% phenol solution. After that, 2.5 mL sulfuric acid was added for 30 min. The absorbance was determined under the 490 nm. The results showed that the carbohydrate content of *H. fusiforme* extract was 19.1% and the *Chlorella* sp. extract was 22.6%.

Animals and experimental design

The animals used in the experiment were 6-week-old male Sprague-Dawley (S.D.) rats purchased from Samtako (Osan, Korea), pre-fed for 7 days, and adapted to the environment. The animals were kept in a stainless-steel cage at a temperature of 22 ± 2 °C and humidity of 45 ± 5% and subjected to a 12 h light/12 h dark cycle. The rats were fed a regular solid food diet and water ad libitum. Animals were randomly divided into six groups ($n = 7$, each group) and the animal experiment was designed to last 10 days.

The rats in the control (CON) group were administered an intraperitoneal (i.p.) injection of 0.5 mL saline solution (0.9% NaCl) daily for 10 days. The rats in the lead acetate (Pb) group were administered an i.p. injection of 0.5 mL lead acetate (25 mg/kg body weight) for the final 5 days of the experimental period. The rats in the HZK groups were orally administered 150 or 300 mg/kg body weight of HZK (HZK 150 and HZK 300 groups, respectively) for 10 days. Rats in the CHL groups were orally administered 150 or 300 mg/kg body weight of CHL (CHL 150 and CHL 300 groups, respectively) for 10 days. The rats in the HZK and CHL groups



were administered an i.p. injection of 0.5 mL lead acetate at a dose of 25 mg/kg body weight for the final 5 days of the experimental period.

Blood collection and tissue preparation

At the end of treatment, rats were fasted for 12 h, and then anesthetized with ether and sacrificed by decapitation. Blood samples were collected from the sacrificed animals and left on ice for 30 min before centrifugation. Livers were immediately washed using chilled saline solution, photographed, frozen in liquid nitrogen, and stored at -70°C .

Measurement of glutamic oxaloacetate and glutamic pyruvic transaminase activities in serum

Glutamic oxaloacetate (GOT) and glutamic pyruvic transaminase (GPT) activities were measured using the GOT and GPT kit according to the manufacturer's instructions (Asan Pharm, Seoul, Korea). The absorbance was measured at 505 nm using Synergy™ HTX and Synergy™ 2 Multi-Mode Readers (BioTek, Winooski, VT, USA).

Measurement of lipid peroxidation

The liver tissue was assessed for liver peroxidation using the thiobarbituric acid reactive substances (TBARS) assay kit according to the manufacturer's instructions (Oxford Biochemical Research, Auckland, New Zealand). The absorbance was measured at 532 nm using Synergy™ HTX and Synergy™ 2 Multi-Mode Readers (BioTek, Winooski, VT, USA).

Antioxidant enzyme activity in liver tissue

The superoxide dismutase (SOD) and catalase (CAT) activities, as well as glutathione (GSH) levels, were assessed in the liver homogenates.

SOD was measured using the SOD assay kit according to the manufacturer's instructions (Cayman, Ann Arbor, MI, USA). The absorbance was measured at 450 nm using Synergy™ HTX and Synergy™ 2 Multi-Mode Readers (BioTek, Winooski, VT, USA).

CAT was measured using the CAT assay kit according to the manufacturer's instructions (Cayman, Ann Arbor, MI, USA). The absorbance was measured at 540 nm using Synergy™ HTX and Synergy™ 2 Multi-Mode Readers (BioTek, Winooski, VT, USA).

GSH levels were measured using the GSH assay kit according to the manufacturer's instructions (Cayman, Ann Arbor, MI, USA). The absorbance was measured at 405 nm using Synergy™ HTX and Synergy™ 2 Multi-Mode Readers (BioTek, Winooski, VT, USA).

Analysis of protein expression in the liver tissue using Western blotting

Liver tissue (50 mg) was homogenized in RIPA lysis buffer and centrifuged ($12,000\times g$, 4°C , 20 min) to obtain the supernatant. Protein concentration was measured using a BCA protein assay kit (Pierce Biotechnology, Rockford, IL, USA). The same amount of protein was added at 50 $\mu\text{g}/\text{lane}$, mixed with 5 \times sample buffer, boiled in a water bath, and loaded on 7.5%, 10%, and 12.5% sodium dodecyl sulfate polyacrylamide gel electrophoresis gels. Standard molecular weights were measured using a dual color marker and transferred to a polyvinylidene difluoride membrane (Millipore, Billerica, MA, USA). The membranes were incubated with primary antibody (Santa Cruz Biotechnology, Inc., Santa Cruz, CA, USA) at room temperature for 2 h, washed twice with TBS-T [10 mM Tris-HCl (pH 7.5), 150 mM NaCl, and 0.1% Tween 20] (USB, Cleveland, OH, USA) for 15 min, and then diluted at a 1:10,000–5000 ratio for the second antibody (Santa Cruz Biotechnology, Inc., Santa Cruz, CA, USA). The membranes were washed twice with TBS-T for 15 min and then analyzed with a fluorescent gel image analyzer (Azure Biosystems, Dublin, CA, USA) using SuperSignal West Pico Luminol/Enhancer solution and SuperSignal West Pico Stable Peroxide solution (Pierce Biotechnology, Rockford, IL, USA). The protein expression level was confirmed by scanning the band.

Statistical analysis

The results were expressed as the mean and standard deviation (SD) for each group. All data were analyzed using SPSS software (IBM, Armonk, NY, USA) for Windows. Analysis of variance was used to identify differences between the experimental and control groups, and Duncan's multiple range test was used to test the significance of each group. All statistical significance levels were assessed at $P < 0.05$.

Results

Liver photographs

The rat livers in the CON group were red with a normal shape and shiny surface. In the Pb group, the livers showed a slight ashy coloration. Conversely, the color of rat livers in the HZK 150, HZK 300, CHL 150, and CHL 300 groups were similar to those in the CON group; no concentration-dependent change was observed (Fig. 2).

GOT and GPT activities in serum

GOT activity was significantly increased in the Pb group compared with the CON group and decreased in a concentration-dependent manner in the HZK 150, HZK 300, CHL 150, and CHL 300 groups compared with the Pb group.




Fig. 2 Photographs of liver after lead acetate or lead acetate with HZK and CHL (original magnification, × 100)

GPT activity was significantly increased in the Pb group, whereas the HZK 150, HZK 300, CHL 150, and CHL 300 groups showed less GPT activity than the Pb group (Fig. 3).

Effect on lipid peroxidation

The Pb, HZK 150, HZK 300, CHL 150, and CHL 300 groups had elevated TBARS levels compared with the CON group. The HZK 150, HZK 300, CHL 150, and CHL 300 groups showed decreased TBARS levels compared with the Pb group (Fig. 4). *H. fusiforme* and *Chlorella* sp. extracts significantly decreased TBARS levels and prevented lipid peroxidation caused by lead acetate (Newairy and Abdou 2009; Li et al. 2015a, b).

SOD activity in liver tissues

The Pb, HZK 150, HZK 300, CHL 150, and CHL 300 groups showed significantly increased SOD activity compared with the CON group. SOD activity in the HZK 150, HZK 300, CHL 150, and CHL 300 groups decreased in a concentration-dependent manner compared with the increased SOD activity in the Pb group (Fig. 5a).

CAT activity in liver tissues

The Pb group showed decreased CAT activity compared with the CON group and the CAT activity in the HZK 150, HZK 300, CHL 150, and CHL 300 groups was slightly, but there was no significant difference between each group (Fig. 5b).




Fig. 3 Effects of HZK and CHL on serum GOT and GPT activities in lead acetate-injected rats. The CON group consisted of control rats not administered lead acetate. The Pb group consisted of rats injected for the final 5 days of the experiment with distilled water (1.5 mL/rat) and lead acetate (25 mg/kg body weight). The HZK 150 and HZK 300 (HZK 150 or 300 mg/kg body weight, respectively) groups and the CHL 150 and CHL 300 (CHL 150 or 300 mg/kg body weight, respectively) groups consisted of rats orally administered HZK or CHL for 10 days and injected (i.p.) with 25 mg/kg of lead acetate for the final 5 days of the experimental period. Values are the mean ± SD based on Duncan's multiple range test ($P < 0.05$) of seven rats per group




Fig. 4 Effects of HZK and CHL on TBARS levels in liver tissues. The CON group consisted of rats not administered lead acetate. The Pb group consisted of rats injected for the final 5 days of the experiment with distilled water (1.5 mL/rat) and lead acetate (25 mg/kg body weight). The HZK 150 and HZK 300 (HZK 150 or 300 mg/kg body weight, respectively) groups and the CHL 150 and CHL 300 (CHL 150 or 300 mg/kg body weight, respectively) groups consisted of rats orally administered HZK or CHL for 10 days and injected (i.p.) with 25 mg/kg of lead acetate for the final 5 days of the experimental period. Values are the mean ± SD based on Duncan's multiple range test ($P < 0.05$) of seven rats per group




Fig. 5 Effects of HZK and CHL on antioxidant enzyme (SOD, CAT, and GSH) activity in liver tissues. The CON group consisted of rats not administered lead acetate. The Pb group consisted of rats injected for the final 5 days of five experiment with distilled water (1.5 mL/rat) and lead acetate (25 mg/kg body weight). The HZK 150 and HZK 300 (HZK 150 or 300 mg/kg body weight, respectively) groups and the CHL 150 and CHL 300 (CHL 150 or 300 mg/kg body weight, respectively) groups consisted of rats orally administered HZK or CHL for 10 days and injected (i.p.) with 25 mg/kg of lead acetate for the final 5 days of the experimental period. Values are the mean ± SD based on Duncan's multiple range test ($P < 0.05$) of seven rats per group

GSH activity in liver tissues

The Pb group showed significant decreased GSH activity compared with the CON group. GSH levels reduced due to lead acetate-induced liver damage which was significantly increased in a concentration-dependent manner in the HZK 150, HZK 300, CHL 150, and CHL 300 groups (Fig. 5c).

Analysis of mitogen-activated protein kinase pathway expression based on Western blot analysis

The mitogen-activated protein kinase (MAPK) pathway is responsible for a variety of cellular functions, such as cell signaling (Obata and Noguchi 2004). ERK is the first known MAPK pathway and is important for cell mitosis, differentiation, proliferation, and survival. p38 is engaged in cellular stress and inflammatory responses (Widmann et al. 1999). The phosphorylation of ERK, JNK, and p-38 in the MAPK pathway was confirmed and decreased in the HZK 150, HZK 300, CHL 150, and CHL 300 groups compared with the Pb group (Fig. 6).

Analysis of inflammation-related protein expression based on Western blot analysis

Phosphorylation of I-κB, an inflammatory response-related intracellular enzyme complex, inducible nitric oxide synthase (iNOS), and cyclooxygenase-2 (COX-2), an inflammatory protein, showed decreased expression in the HZK 150, HZK 300, CHL 150, and CHL 300 groups (Fig. 7).

Analysis of Nrf2 pathway expression based on Western blot analysis

Activation of the Nrf2 pathway contributes to inflammation control, antioxidant, drug resistance, and cancer-related functions. When activated by external stress [e.g., reactive oxygen species (ROS), carbonyls, hydroperoxides, and quinones], Nrf2 migrates into the nucleus and binds to the antioxidant response element (ARE) to express antioxidant and detoxifying enzymes such as NQO-1, HO-1, and glutathione S-transferase (GST) for cell protection (Li et al. 2015a, b). In this study, Nrf2 expression was increased by treatment with HZK and CHL. The expressions of NQO-1, HO-1, and




Fig. 6 Western blot analysis of MAPK phosphorylation in lead acetate-injected rats. The protein expression levels of ERK, p-ERK, JNK, p-JNK, p38, and p-p38 in the liver tissues were assessed by Western blot analysis. GAPDH was used as an internal standard




Fig. 7 Western blot analysis of inflammation-related protein expression in lead acetate-injected rats. The protein expression levels of IκB, p-IκB, iNOS and COX-2 in the liver tissues were assessed by Western blot analysis. GAPDH was used as an internal standard

GST were also increased (Fig. 8), and oral administration of HZK and CHL exerted protective effects against oxidative stress and inflammation in rats.

Discussion

Various nutrients and functional characteristics of seaweed have been demonstrated. Research on various properties such as antioxidant, anti-inflammation, anti-cancer, and diet is being conducted for functional health foods using bioactive substances of seaweed (Bocanegra et al. 2009). *H. fusiforme* contains calcium, iodine, iron, and is especially abundant in minerals. In addition, *H. fusiforme* has been used to study hepatic enzyme activity and recovery from oxidative stress (Kim and Kim 1998).

Chlorella sp. contains a significant amount of chlorophyll and various nutrients (Singh et al. 1998). Numerous recent studies have highlighted the biological functions of *Chlorella* sp., such as excretion of heavy metals in organisms (Pairat and Qiming 2001), anti-inflammation effects (Dao et al. 2001), and anti-cancer effects (Justo et al. 2001).

In this study, the effects of *H. fusiforme* and *Chlorella* sp. extracts on liver damage and oxidative stress in lead-induced hepatotoxicity were investigated. First, to confirm hepatotoxicity, analysis of GOT and GPT activities was performed, as blood biochemical indicators of lead-induced hepatotoxicity. In the HZK 150, HZK 300, CHL 150, and CHL 300 groups, the GOT and GPT enzyme activities showed recovery compared to the Pb

group, indicating that HZK and CHL inhibited cell destruction caused by lead.

Lipid peroxidation is caused by oxidative stress reactions in lead-induced hepatotoxicity (Johny et al. 2010). Lipid peroxidation (TBARS) was significantly higher in the Pb group than in the CON group, and decreased in a concentration-dependent manner in the HZK 150, HZK 300, CHL 150, and CHL 300 groups. This reduction in lipid peroxidation indicated that the administration of HZK and CHL prevented oxidative stress caused by lead-induced damage in liver tissue. SOD plays an important role as a superoxide radical scavenging enzyme (Fukai and Ushio-Fukai 2011). In this study, injection of lead acetate induced an abnormal increase in SOD activity in the Pb group compared with the CON group. The increase of SOD activity in the Pb group was considered a defense mechanism due to increased oxidative stress caused by the injection of lead (Carla et al. 2017). High SOD activities facilitate the conversion of superoxide anion (O_2^-) into hydrogen peroxide (H_2O_2) (Park et al. 2007). In the HZK 150, HZK 300, CHL 150, and CHL 300 groups, the decreased, concentration-dependent SOD activity was attributed to suppressed production of ROS by oral administration of HZK and CHL, resulting in suppressed oxidative stress (Kim and Lim 2005). CAT is an antioxidant enzyme that protects against cellular damage by reducing H_2O_2 into water and eliminating ROS free radicals. CAT also maintains cell homeostasis and prevents oxidative damage (Wang et al. 2013). In a recent




Fig. 8 Western blot analysis of Nrf2 pathway expression in lead acetate-injected rats. The protein expression levels of Nrf2, HO-1, NQO1, and GST in the liver tissues were assessed by Western blot analysis. GAPDH was used as an internal standard

study, significant differences in CAT activity were not observed. GSH acts as a substrate for the glutathione peroxidase and GST antioxidant enzymes, which removes highly reactive toxic metabolites and assists in cellular antioxidant defense (Yang et al. 2013). In the present study, GSH levels in rat liver were significantly decreased in the Pb group, but recovered with oral administration of HZK and CHL in a concentration-dependent manner. The GSH levels in the HZK- and CHL-treated groups were higher than in the CON group due to the reduction in H₂O₂ levels caused by lead administration.

MAPKs are a family of serine/threonine protein kinases that mediate fundamental biological processes and cellular responses to external stress signals (Cargnello and Roux 2011). Increased activity of MAPKs, in particular p38, and their involvement in the regulation of the synthesis of inflammation mediators at the translation and transcription levels, renders them potential targets for anti-inflammatory therapeutics (Bozena 2005). In this study, injection of lead acetate induced phosphorylation of ERK, JNK, and p38 proteins. However, ERK, JNK, and p38 phosphorylation were decreased when HZK and CHL were orally administered. Furthermore, NF-κB is a downstream signaling transcription factor in the MAPK pathway that produces cytokines and promotes development of cancer, inflammatory responses, and apoptosis (Didonato et al. 2012). iNOS is transcribed by NF-κB and promotes the production of NO, which is involved in inflammation and immune responses (Liu et al. 2014). In particular, iNOS expression

inhibits antioxidant enzymes (Aktan 2004). COX-2 is an enzyme induced by several types of stimuli and is involved in inflammatory diseases and tumorigenesis by producing prostaglandins, which cause inflammation and pain (Moon et al. 1998). I-κB phosphorylation was significantly increased after injection of lead, and confirmed the effects of *H. fusiforme* and *Chlorella* sp. extracts. Expression of iNOS and COX-2 was inhibited in the HZK 150, HZK 300, CHL 150, and CHL 300 groups. Therefore, oral administration of HZK and CHL regulated the expression of iNOS and COX-2 by inhibiting the NF-κB transcription factor, thereby confirming the inflammatory reaction caused by lead administration is regulated.

Activation of the Nrf2 pathway contributes to inflammation control, antioxidant, drug resistance, and cancer-related functions. When activated by external stress (e.g., ROS, carbonyls, hydroperoxides, and quinones), Nrf2 migrates into the nucleus and binds to AREs to express antioxidant and detoxifying enzymes such as NQO-1, HO-1, and GST for cell protection (Murakami and Motohashi 2015). The expression of Nrf2, HO-1, GST, and NQO-1 was increased in HZK 150, HZK 300, CHL 150, and CHL 300 groups.

Therefore, *H. fusiforme* and *Chlorella* sp. extracts exerted antioxidant and anti-inflammatory effects against lead acetate-induced hepatotoxicity. The development of functional health foods with hepatoprotective effects after lead intake using *H. fusiforme* and *Chlorella* sp. extract should be considered.

Funding

This study was supported by Pukyong National University grant no. C-D-2017-0360, Busan, Republic of Korea.

Availability of data and materials

The datasets supporting the conclusions of this article are included within the article. There are no additional data and materials to disclose.

Authors' contributions

JHP designed this study and drafted the manuscript. JWC, MKL, YHC, and TJN conceived and designed the study, and also revised the manuscript. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Animal experiments were performed according to the institutional guidelines for the care and the use of laboratory animals, and the protocol was approved by the ethics committee of Pukyong National University (Approval No. 2017-34).

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Departments of Food Science and Nutrition, Pukyong National University, Yongso-ro, Nam-gu, Busan 48513, Republic of Korea. ²Institute of Fisheries Science, Pukyong National University, Busan 46041, Republic of Korea. ³Departments of Marine Bio-Materials & Aquaculture, Pukyong National University, Busan 48513, Republic of Korea.

Received: 18 October 2018 Accepted: 10 January 2019

Published online: 25 January 2019

References

- Aktan F. iNOS-mediated nitric oxide production and its regulation. *Life Sci.* 2004; 75:639–53.
- Bocanegra A, Bastida S, Benedi J, Ródenas S, Sánchez-Muniz FJ. Characteristics and nutritional and cardiovascular-health properties of seaweeds. *J Med Food.* 2009;12:236–58.
- Bozena K. MAPK signalling pathways as molecular targets for anti-inflammatory therapy from molecular mechanisms to therapeutic benefits. *Biochim Biophys Acta.* 2005;1754:253–62.
- Cargnello M, Roux PP. Activation and function of the MAPKs and their substrates, the MAPK-activated protein kinases. *Microbiol Mol Biol Rev.* 2011;75:50–83.
- Carla SC, Gabriel TV, Janaina AS, Patrícia P, Sthefany TR, Carlos RT. Data on the effects of losartan on protein expression vascular reactivity and antioxidant capacity in the aorta of ethanol-treated rats. *Data Brief.* 2017;11:111–6.
- Dao VT, Floeren M, Kumpf S, Both C, Peter B, Balz V, Suvorava T, Kojda G. Catalase activity prevents exercise-induced up-regulation of vasoprotective proteins in venous tissue. *J Cell Mol Med.* 2001;15:2326–34.
- Didonato JA, Mercurio F, Karin M. NF- κ B and the link between inflammation and cancer. *Immunol Rev.* 2012;246:379–400.
- Donaldson K, MacNee W. Potential mechanisms of adverse pulmonary and cardiovascular effects of particulate air pollution (PM10). *Int J Hyg Environ Health.* 2001;203:411–5.
- Fukai T, Ushio-Fukai M. Superoxide dismutases: role in redox signaling, vascular function, and diseases. *Antioxid Redox Signal.* 2011;15:1583–606.
- Hidalgo S, Bisson JF, Duffaud A, Nejd A, Guerin L, Baert B, Saniez MH, Rozan P. Benefits of oral and topical administration of ROQUETTE *Chlorella* sp. on skin inflammation and wound healing in mice. *Antiinflamm Antiallergy Agents Med Chem.* 2014;13:93–102.
- Hsiao W, Mo ZY, Fang M, Shi XM, Wang F. Cytotoxicity of PM2.5 and PM2.5-10 ambient air pollutants assessed by the MTT and the comet assays. *Mutat Res.* 2000;471:45–55.
- Hu X, Sun Y, Ding Z, Zhang Y, Wu J, Lian H, Wang T. Lead contamination and transfer in urban environmental compartments analyzed by lead levels and isotopic compositions. *Environ Pollut.* 2014;187:42–8.
- Johny CP, Oscar PM, Rolando HM, Patricia VTD, Marco AJO. Protective effects of *Spirulina maxima* on hyperlipidemia and oxidative-stress induced by lead acetate in the liver and kidney. *Lipids Health Dis.* 2010;9:35.
- Justo GZ, Silva MR, Queiroz ML. Effects of the green algae *Chlorella vulgaris* on the response of the host hematopoietic system to intraperitoneal ehrlich ascites tumor transplantation in mice. *Immunopharmacol Immunotoxicol.* 2001;23:119–32.
- Kim HS, Choi EO, Park C, Choi YH, Hyun SK, Hwang HJ. Effect of *Hizikia fusiforme* extracts on antioxidant enzyme activity and vitamin E concentration in rats. *J Korean Soc Food Sci Nutr.* 2011;40:1556–61.
- Kim HS, Kim GJ. Effects of the feeding *Hizikia fusiforme* (Harvey) Okamura on lipid composition of serum in dietary hyperlipidemic rats. *J Korean Soc Food Sci Nutr.* 1998;27:718–23.
- Kim YC, Lim AK. The effect of UVB irradiation on the generating and scavenging enzymes of oxygen free radical in mouse skin. *J NERI.* 2005;10:49–55.
- Lee YA, Kim MH. The effect of sea tangle extract on serum lipid level in ovariectomized rats. *J Life Sci.* 2008;18:249–54.
- Li P, Su L, Li X, Di W, Zhang X, Zhang C, He T, Zhu X, Zhang Y, Li Y. Remote limb ischemic postconditioning protects mouse brain against cerebral ischemia/reperfusion injury via upregulating expression of Nrf2, HO-1 and NQO-1 in mice. *Int J Neurosci.* 2015a;17:1–8.
- Li XD, Sun GF, Zhu WB, Wang YH. Effects of high intensity exhaustive exercise on SOD, MDA, and NO levels in rats with knee osteoarthritis. *Genet Mol Res.* 2015b;14:12367–76.
- Liu CM, Ma JQ, Liu SS, Zheng GH, Feng ZJ, Sun JM. Proanthocyanidins improves lead-induced cognitive impairments by blocking endoplasmic reticulum stress and nuclear factor- κ B-mediated inflammatory pathways in rats. *Food Chem Toxicol.* 2014;72:295–302.
- Meo SA, Suraya F. Effect of environmental air pollution on cardiovascular diseases. *Eur Rev Med Pharmacol Sci.* 2015;19:4890–7.
- Moon TC, Murakami M, Ashraf MD, Kudo I, Chang HW. Regulation of cyclooxygenase-2 and endogenous cytokine expression by bacterial lipopolysaccharide that acts in synergy with c-kit ligand and Fc epsilon receptor 1 crosslinking in cultured mast cells. *Cell Immunol.* 1998;185:146–52.
- Murakami S, Motohashi H. Roles of Nrf2 in cell proliferation and differentiation. *Free Radic Biol Med.* 2015;88:168–78.
- Newairy AS, Abdou HM. Protective role of flax lignans against lead acetate induced oxidative damage and hyperlipidemia in rats. *Food Chem Toxicol.* 2009;47:813–8.
- Obata K, Noguchi K. MAPK activation in nociceptive neurons and pain hypersensitivity. *Life Sci.* 2004;74:2643–53.
- Pairat K, Qiming Y. Cadmium(II) removal from aqueous solutions by pre-treated biomass of marine alga *Padina* sp. *Environ Pollut.* 2001;112:209–13.
- Park EM, Ramnath N, Yang GY, Ahn JY, Park YR, Lee TY, Shin HS, Yu JH, Ip C, Park YM, High SOD. Low GPX activities in RBC predict susceptibility of lung cancer patients to radiation pneumonitis. *Free Radic Biol Med.* 2007;42:280–7.
- Singh A, Singh SP, Bamezai R. Perinatal influence of *Chlorella vulgaris* (E-25) on hepatic drug metabolizing enzymes and lipid peroxidation. *Anticancer Res.* 1998;18:1509–14.
- Wang C, Yue X, Lu X, Liu B. The role of catalase in the immune response to oxidative stress and pathogen challenge in the clam *Meretrix meretrix*. *Fish Shellfish Immunol.* 2013;34:91–9.
- Widmann C, Gibson S, Jarpe MB, Johnson GL. Mitogen-activated protein kinase: conservation of a three-kinase module from yeast to human. *Physiol.* 1999; 79:143–80.
- Wua M, Wua Y, Qub M, Li W, Yan X. Evaluation of antioxidant activities of water-soluble polysaccharides from brown alga *Hizikia fusiformis*. *Int J Biol Macromol.* 2013;56:28–33.
- Yang J, Wang XY, Xue J, Gu ZL, Xie ML. Protective effect of apigenin on mouse acute liver injury induced by acetaminophen is associated with increment of hepatic glutathione reductase activity. *Food Funct.* 2013;4:939–43.
- Zhou X, Cao Z, Ma Y, Wang L, Wu R, Wang W. Concentrations, correlations and chemical species of PM2.5/PM10 based on published data in China: potential implications for the revised particulate standard. *Chemosphere.* 2016;144:518–26.