

# Relationships between the diabetes awareness and clinical indices/nutrient intakes in Korean adults: Based on the 2012-2013 Korea National Health and Nutrition Examination Survey Data

Seul Lee<sup>1</sup>, Haeryun Park<sup>1</sup>, Youngmi Lee<sup>1</sup>, Onjeong Choi<sup>1</sup>, Jiwon Kim<sup>1</sup>, Heewon L Gray<sup>2</sup> and Kyunghee Song<sup>1§</sup>

<sup>1</sup>Department of Food and Nutrition, Myongji University, Yongin, Gyeonggi 17058, Korea

<sup>2</sup>Department of Community and Family Health, University of South Florida 13201, Florida, USA.

**BACKGROUND/OBJECTIVES:** Diabetes is diagnosed after an average of 10-12 years of diabetic development. Strict glycemic control in diabetic patients promotes the normalization of blood glucose and reduces cardiovascular diseases (CVDs) and diabetic complications. Therefore, early diagnosis in non-aware individuals is very important.

**SUBJECTS/METHODS:** Clinical indices and nutrient intakes in Korean diabetic adults aged 19-64 years were examined according to the awareness of diabetes, using 2012 and 2013 Korea National Health and Nutrition Examination Survey (KNHANES) data. The aware group was defined as individuals who were aware of having diabetes from diagnosis by physician before the survey and the non-aware group as individuals who were not aware of having diabetes.

**RESULTS:** The average age was higher in the aware group compared to the non-aware group in both men ( $P=0.002$ ) and women ( $P=0.004$ ). The prevalences of hypertension and dyslipidemia were not different between the two groups, but the diagnosis rate was significantly lower in the non-aware group. In the non-aware group, total and LDL-cholesterol were significantly higher ( $P < 0.001$ ), the risk for total cholesterol over 240 mg/dL was 3.4 times (95% CI: 1.58-7.52) higher ( $P=0.002$ ) and the risk for LDL-cholesterol over 160 mg/dL was 4.59 times (95% CI: 2.07-10.17) higher ( $P < 0.001$ ). The calorie intake of the female non-aware group was significantly higher compared to the female aware group ( $P=0.033$ ).

**CONCLUSION:** The results suggested that the recommendation of screening test is necessary even for young adults. Studies on the methodology for early diagnosis of diabetes are also needed.

Nutrition Research and Practice 2019;13(3):240-246; <https://doi.org/10.4162/nrp.2019.13.3.240>; pISSN 1976-1457 eISSN 2005-6168

**Keywords:** Diabetes, aware, clinical indices, nutrient intake

## INTRODUCTION

Diabetes is one of chronic diseases and the mortality rate caused by diabetes has increased every year worldwide [1]. Diabetes is a major risk factor for cardiocerebrovascular disease, which is a major cause of death in diabetic patients [2]. When HbA1c (6.5% and over) was included in the diagnostic criteria for diabetes, the prevalence of diabetes has increased in both men and women in Korea recently [3].

Only strict glycemic control can prevent complications of diabetes [4]. According to the guidelines for diabetes by the American Diabetes Association (ADA), controlling BMI, HbA1c, blood pressure and cholesterol is recommended [5]. Based on the Korean diabetes management guidelines, HbA1c  $< 6.5\%$ , blood pressure  $\leq 140/80$  mmHg, LDL-cholesterol  $\leq 100$  mg/dL, TG  $\leq 150$  mg/dL and HDL-cholesterol  $\geq 40$  mg/dL for men and  $\geq 50$  mg/dL for women are recommended [6].

Pharmacotherapy, dietary therapy and exercise therapy are commonly used to manage diabetes. Dietary therapy plays an

important role in delaying the development of diabetic complications during the course of treatment and progression of diabetes. Proper dietary therapy improves metabolic abnormalities and blood lipid profiles [7,8]. Studies on nutrient intakes in Korean diabetic patients showed that the intake of carbohydrates was more than 60% and even higher in women than in men [9,10]. The increased carbohydrate intake causes hyperglycemia, which can develop oxidative stress in the body and finally cause tissue damage [11]. In addition, excessive insulin secretion can increase body fat to induce obesity [12].

Among diabetic complications, microvascular complications are increased along with an increased duration of diabetes, while macrovascular complications are affected by the risk factors for cardiovascular diseases (CVDs) [8]. In addition, the prevalence of CVDs is more than 2-4 times higher in diabetic patients compared to non-diabetic patients [13,14]. The awareness of diabetes in Korea tends to be higher in women than in men. In particular, the awareness rate was the lowest in men in their 30 s and 40 s [3].

<sup>§</sup> Corresponding Author: Kyunghee Song, Tel. 82-31-330-6206, Fax. 82-31-330-6200, Email. [khsong@mju.ac.kr](mailto:khsong@mju.ac.kr)

Received: February 18, 2019, Revised: March 18, 2019, Accepted: May 20, 2019

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

According to a study by the American Medical Care System, the onset of diabetes, on average, is up to 10-12 years before clinical diagnosis [15]. Complications such as diabetic retinopathy or diabetic neuropathy are commonly found at the time of diagnosis [16] and CVDs such as peripheral arterial disease and coronary heart disease are co-diagnosed in many cases [17]. Therefore, early diagnosis is very important to prevent complications and comorbidities of diabetes, yet associations among patients' awareness of diabetes, its clinical indicators, and nutrient intakes have not been well investigated in the Korean population, especially using nationally representative data.

The purpose of this study was to examine clinical indices and nutrient intakes based on diabetes awareness among adults aged 19-64 years, using nationally representative data from the 2012-2013 Korea National Health and Nutrition Examination Survey (KNHANES).

## SUBJECTS AND METHODS

### *Study design and participants*

A cross-sectional secondary analysis was conducted using the 2012-2013 KNHANES data. Participants who were 19-64 years old and diagnosed with diabetes at the time of the measurement were included. For diabetes indicators, fasting blood glucose and HbA1c were used from the hematology section of the KNHANES. A total of 523 participants were divided into two groups: (1) the aware group (297; 148 men and 149 women), defined as individuals who were diagnosed by medical doctors; and (2) the non-aware group (226; 103 men and 123 women), defined as individuals who were not clinically diagnosed at the time of the measurement but had 8-h fasting plasma glucose over 126 mg/dL or HbA1c over 6.5%. The Institutional Review Board (IRB) of the Korea Center for Disease Control and Prevention approved the KNHANES (IRB. 2012-01EXP-01-2C and 2013-07CON-03-4C).

### *General characteristics*

Information on gender, age, income level, education level, exercise, drinking and smoking patterns was obtained from the KNHANES data. The exercise patterns were classified into two levels: moderate/walking was classified as more than 10 minutes per session and more than 20 minutes of daily exercise for at least 3 days a week and vigorous exercise was classified as more than 10 minutes per session and more than 30 minutes of daily exercise for at least 5 days a week. Drinking pattern was classified into a binary variable: either equal to greater vs. fewer than once a month during the past one year. Height, weight and waist circumference data were extracted from the anthropometric measurement section of the KNHANES.

### *Prevalence and diagnosis of hypertension and dyslipidemia*

The prevalence of hypertension was defined as systolic pressure  $\geq 140$  mmHg or diastolic pressure  $\geq 90$  mmHg. The prevalence of dyslipidemia was defined as total cholesterol  $\geq 240$  mg/dL or triglyceride  $\geq 200$  mg/dL or HDL-cholesterol  $< 40$  mg/dL or LDL-cholesterol  $\geq 160$  mg/dL. The diagnosis rate for each of hypertension and dyslipidemia was defined as individuals who were diagnosed by a medical doctor. LDL-

cholesterol was calculated with the Friedewald equation for those with serum triglyceride level  $\leq 400$  mg/dL [18].

### *Clinical indices*

The following data were obtained from the health examination survey: systolic pressure, diastolic pressure, total-cholesterol, HDL-cholesterol, triglyceride, hemoglobin, hematocrit, blood urea nitrogen (BUN) and blood creatinine.

### *Nutrient intakes*

The intake of nutrients was analyzed through the type and amount of food consumed the day before the survey using the 24h recall method. The ratio of intakes of protein, fat and carbohydrates was calculated using the presented nutrient values. For the analysis of the intake ratio by nutrients, the Dietary Reference Intakes for Koreans (2010) was used for different age groups.

### *Statistical analysis*

Statistical analysis was performed after adjusting for the KNHANES sample weights, strata and clusters. Clinical indices and nutrient intakes were analyzed after controlling for age. Group comparisons were performed based on the aware vs. non-aware groups. Independent samples t-tests were performed for continuous data and  $\chi^2$ -tests were performed for nominal data. General linear models were used to examine differences in biochemical indices, energy and nutrient intakes by group, after controlling for age. To analyze the proper ratio of protein, fat and carbohydrates intake by group, logistic regressions were performed. Odds ratio (OR) and 95% confidence interval (CI) are presented. All data in the study were analyzed using SPSS 19.0 (SPSS Inc., Chicago, USA) and the statistical significances were tested at  $P < 0.05$ .

## RESULTS

### *General characteristics*

The general characteristics of the participants by gender and diabetes awareness are shown in Table 1. The average ages of men and women were significantly higher at 53.2 and 53.6 years in the aware group compared to 48.8 and 49.3 years in the non-aware group ( $P=0.002$  and  $P=0.004$ ), respectively. There were significant differences in the distribution of age categories between the aware and the non-aware groups. A large percentage of men and women in the non-aware group was in their 50 s (41.6%) and 60 s or older (23.5%), while 43.1% of the aware group was in their 60 s or older ( $P=0.001$ ). The average BMI in men was significantly higher at 25.8 kg/m<sup>2</sup> in the non-aware group compared to 24.7 kg/m<sup>2</sup> in the aware group ( $P=0.031$ ). The average BMI in women was not significantly different between two groups, but the distribution of BMI was significantly different ( $P=0.026$ ). The obesity rate was 50.3% in the female aware group and 64.2% in the female non-aware group.

### *Prevalence and diagnosis of hypertension and dyslipidemia*

The comorbidity status of hypertension and dyslipidemia is shown in Table 2. The prevalence of hypertension in women

**Table 1.** General characteristics of subjects by sex and diabetes awareness.

	Male (n = 251)			Female (n = 272)			Total (n = 523)		
	AG <sup>1)</sup> (n = 148)	NAG <sup>2)</sup> (n = 103)	P-value	AG (n = 149)	NAG (n = 123)	P-value	AG (n = 297)	NAG (n = 226)	P-value
Age (yrs)	53.2 ± 1.0 <sup>3)</sup>	48.8 ± 1.0	0.002**	53.6 ± 0.9	49.3 ± 1.3	0.004**	53.4 ± 0.7	49.0 ± 0.8	0.001***
19-29	2 (1.4) <sup>4)</sup>	3 (2.9)	0.028*	2 (1.3)	4 (3.3)	0.009**	4 (1.3)	7 (3.1)	0.001***
30-39	6 (4.1)	11 (10.7)		4 (2.7)	11 (8.9)		10 (3.4)	22 (9.7)	
40-49	21 (14.2)	23 (22.3)		21 (14.1)	27 (22.0)		42 (14.1)	50 (22.1)	
50-59	54 (36.5)	41 (39.8)		59 (39.6)	53 (43.1)		113 (38.0)	94 (41.6)	
≥ 60	65 (43.9)	25 (24.3)		63 (42.3)	28 (22.8)		128 (43.1)	53 (23.5)	
Height (cm)	169.3 ± 0.7	171.3 ± 0.7	0.024*	155.1 ± 0.5	158.0 ± 0.7	0.001**	163.0 ± 0.7	165.8 ± 0.6	0.003**
Weight (cm)	71.0 ± 1.2	76.0 ± 1.5	0.009**	63.5 ± 1.8	69.3 ± 2.0	0.034*	67.7 ± 1.0	73.2 ± 1.2	0.001***
Waist circumference (cm)	86.9 ± 0.8	88.8 ± 1.2	0.159	86.8 ± 1.5	88.3 ± 1.4	0.468	86.8 ± 0.8	88.6 ± 0.9	0.148
BMI (kg/m <sup>2</sup> ) <sup>5)</sup>	24.7 ± 0.3	25.8 ± 0.4	0.031*	26.4 ± 0.7	27.6 ± 0.7	0.217	25.5 ± 0.4	26.6 ± 0.4	0.047*
Underweight	1 (0.7)	1 (1.0)	0.076	3 (2.0)	1 (0.8)	0.026*	4 (1.35)	2 (0.9)	0.006**
Normal	41 (27.7)	21 (20.4)		40 (26.8)	16 (13.0)		81 (27.3)	37 (16.4)	
Overweight	46 (31.1)	19 (18.4)		31 (20.8)	27 (22.0)		77 (25.9)	46 (20.4)	
Obesity	60 (40.5)	62 (60.2)		75 (50.3)	79 (64.2)		135 (45.5)	141 (62.4)	
House income									
Low	19 (12.8)	11 (10.7)	0.463	34 (22.8)	18 (14.6)	0.491	53 (17.8)	29 (12.8)	0.317
Middle-low	34 (23.0)	24 (23.3)		56 (37.6)	36 (29.3)		90 (30.3)	60 (26.5)	
Middle-high	44 (29.7)	30 (29.1)		34 (22.8)	41 (33.3)		78 (26.3)	71 (31.4)	
High	51 (34.5)	38 (36.9)		25 (16.8)	28 (22.8)		76 (25.6)	66 (29.2)	
Education									
≤ Elementary school	36 (24.3)	13 (12.6)	0.016*	58 (38.9)	33 (26.8)	0.267	94 (31.6)	46 (20.4)	0.019*
Middle school	26 (17.6)	12 (11.7)		25 (16.8)	30 (24.4)		51 (17.2)	42 (18.6)	
High school	41 (27.7)	39 (37.9)		51 (34.2)	38 (30.9)		92 (31.0)	77 (34.1)	
≥ College	45 (30.4)	39 (37.9)		15 (10.1)	22 (17.9)		60 (20.2)	61 (27.0)	
Exercise									
Yes	72 (48.6)	39 (37.9)	0.126	65 (43.6)	53 (43.1)	0.763	160 (53.9)	134 (59.3)	0.293
No	76 (51.4)	64 (62.1)		84 (56.4)	70 (57.0)		137 (46.1)	92 (40.7)	
Smoking									
Current	54 (36.5)	48 (46.6)	0.085	6 (4.0)	6 (4.9)	0.589	157 (52.9)	131 (58.0)	0.052
Ex-smoker	75 (50.7)	40 (38.8)		5 (3.4)	1 (0.8)		80 (26.9)	41 (18.1)	
None	19 (12.8)	15 (14.6)		138 (92.6)	116 (94.3)		60 (20.2)	54 (23.9)	
Drinking									
Yes	44 (29.7)	24 (23.3)	0.375	38 (25.5)	44 (35.8)	0.187	155 (52.2)	103 (45.6)	0.103
No	104 (70.3)	79 (76.7)		111 (74.5)	79 (64.2)		142 (47.8)	123 (54.4)	
Fasting blood glucose (mg)	139.6 ± 3.8	136.7 ± 3.6	0.526	151.9 ± 5.6	138.4 ± 5.2	0.094	145.3 ± 3.4	137.3 ± 3.1	0.081
HbA1C (%)	7.4 ± 0.2	6.9 ± 0.1	0.017*	7.9 ± 0.2	7.4 ± 0.2	0.038*	7.6 ± 0.1	7.1 ± 0.1	0.002*

<sup>1)</sup> Aware group<sup>2)</sup> Non-aware group<sup>3)</sup> Mean ± SD<sup>4)</sup> N (%)<sup>5)</sup> Underweight: < 18.5 kg/m<sup>2</sup>, Normal: 18.5-22.9 kg/m<sup>2</sup>, Overweight: 23.0-24.9 kg/m<sup>2</sup>, Obesity: ≥ 25.0 kg/m<sup>2</sup>

\*P &lt; 0.05, \*\*P &lt; 0.01, \*\*\*P &lt; 0.001

**Table 2.** Prevalence and diagnosis of hypertension and dyslipidemia of subjects by sex and diabetes awareness.

	Male (n = 251)			Female (n = 272)			Total (n = 523)		
	AG <sup>1)</sup> (n = 148)	NAG <sup>2)</sup> (n = 103)	P-value	AG (n = 149)	NAG (n = 123)	P-value	AG (n = 297)	NAG (n = 226)	P-value
Prevalence of hypertension <sup>3)</sup>	72 (48.6) <sup>4)</sup>	50 (48.5)	0.425	82 (55.0)	44 (35.8)	0.013*	154 (51.9)	94 (41.6)	0.417
Diagnosed of hypertension	66 (44.6)	26 (25.2)	0.024*	79 (53.0)	37 (30.1)	0.008**	145 (48.8)	63 (27.9)	0.001**
Prevalence of dyslipidemia <sup>5)</sup>	93 (62.8)	70 (68.0)	0.210	98 (65.8)	80 (65.0)	0.755	191 (64.3)	150 (66.4)	0.218
Diagnosed of dyslipidemia	46 (31.1)	12 (11.7)	0.001***	70 (47.0)	21 (17.1)	0.001***	116 (39.1)	33 (14.6)	0.001***

<sup>1)</sup> Aware group<sup>2)</sup> Non-aware group<sup>3)</sup> Prevalence of hypertension: systolic pressure ≥ 140 mmHg or diastolic pressure ≥ 90 mmHg<sup>4)</sup> N (%)<sup>5)</sup> Prevalence of dyslipidemia: Total Cholesterol ≥ 240 mg/dL or HDL-Cholesterol < 40 mg/dL or LDL-Cholesterol ≥ 60 mg/dL or Triglyceride ≥ 200 mg/dL

\*P &lt; 0.05, \*\*P &lt; 0.01, \*\*\*P &lt; 0.001

**Table 3.** The biochemical indices of subjects by sex and diabetes awareness.

	Male (n = 251)			Female (n = 272)			Total (n = 523)		
	AG <sup>1)</sup> (n = 148)	NAG <sup>2)</sup> (n = 103)	P-value	AG (n = 149)	NAG (n = 123)	P-value	AG (n = 297)	NAG (n = 226)	P-value
Systolic pressure	124.3 ± 1.5 <sup>3)</sup>	126.2 ± 1.8	0.424	121.4 ± 1.6	122.3 ± 1.6	0.698	123.0 ± 1.2	124.6 ± 1.3	0.368
Diastolic pressure	78.7 ± 1.0	83.3 ± 1.4	0.010*	74.9 ± 1.0	78.9 ± 0.9	0.004**	77.0 ± 0.8	81.5 ± 0.9	0.001***
Total cholesterol (mg/dL)	177.4 ± 3.7	201.1 ± 3.8	0.001***	182.0 ± 3.8	214.2 ± 4.4	0.001***	179.4 ± 2.6	206.6 ± 2.9	0.001***
HDL-cholesterol (mg/dL)	44.2 ± 1.0	42.7 ± 1.1	0.333	46.3 ± 1.1	46.9 ± 1.1	0.686	45.1 ± 0.7	44.5 ± 0.8	0.547
LDL-cholesterol (mg/dL)	102.5 ± 3.2	118.6 ± 3.3	0.001***	106.1 ± 3.5	136.3 ± 4.2	0.001***	104.0 ± 2.3	126.0 ± 2.7	0.001***
Triglyceride (mg/dL)	153.8 ± 7.6	199.3 ± 10.1	0.001***	148.0 ± 7.1	155.0 ± 7.3	0.481	151.3 ± 5.3	180.8 ± 7.0	0.001***
Hemoglobin (g/dL)	14.9 ± 0.1	15.5 ± 0.1	0.002**	13.0 ± 0.2	13.6 ± 0.1	0.019*	14.0 ± 0.1	14.7 ± 0.1	0.002**
Hematocrit (%)	43.6 ± 0.3	45.2 ± 0.4	0.002**	39.1 ± 0.5	40.8 ± 0.3	0.006**	41.6 ± 0.4	43.4 ± 0.3	0.001***
BUN (mg/dL)	16.3 ± 0.4	15.4 ± 0.4	0.157	15.5 ± 1.3	13.9 ± 0.4	0.292	15.9 ± 0.6	14.8 ± 0.3	0.125
Creatinine (mg/dL)	1.2 ± 0.0	1.0 ± 0.0	0.001***	0.9 ± 0.1	0.7 ± 0.0	0.200	1.0 ± 0.1	0.9 ± 0.0	0.008**

Adjusted for age

<sup>1)</sup> Aware group<sup>2)</sup> Non-aware group<sup>3)</sup> Mean ± SD\*  $P < 0.05$ , \*\*  $P < 0.01$ , \*\*\*  $P < 0.001$ **Table 4.** Comparison of nutrient intake of subjects by sex and diabetes awareness.

	Male (n = 251)			Female (n = 272)			Total (n = 523)		
	AG <sup>1)</sup> (n = 148)	NAG <sup>2)</sup> (n = 103)	P-value	AG (n = 149)	NAG (n = 123)	P-value	AG (n = 297)	NAG (n = 226)	P-value
Energy (kcal)	2251 ± 77 <sup>3)</sup>	2452 ± 132	0.172	1665 ± 74	1891 ± 69	0.033*	1990 ± 60	2218 ± 84	0.025*
Protein (g)	78.5 ± 3.3	90.6 ± 6.3	0.096	61.3 ± 3.7	68.8 ± 4.3	0.196	70.9 ± 2.4	81.5 ± 4.2	0.038*
Fat (g)	43.2 ± 2.5	44.7 ± 3.8	0.739	33.1 ± 2.7	37.4 ± 2.4	0.267	38.7 ± 1.9	41.6 ± 2.5	0.342
Carbohydrate (g)	341.4 ± 11.5	362.7 ± 18.6	0.284	280.7 ± 13.0	319.2 ± 13.3	0.048*	314.3 ± 9.6	344.6 ± 12.4	0.044*
Fiber (g)	9.3 ± 0.5	8.5 ± 0.6	0.391	7.7 ± 0.6	7.9 ± 0.5	0.717	8.5 ± 0.4	8.3 ± 0.4	0.669
Vitamin A (mg)	919.2 ± 103.5	744.4 ± 61.1	0.160	754.1 ± 116.3	954.0 ± 102.3	0.167	844.5 ± 77.1	833.0 ± 57.3	0.902
Vitamin B <sub>1</sub> (mg)	1.8 ± 0.1	2.3 ± 0.2	0.054	1.5 ± 0.1	1.7 ± 0.1	0.130	1.7 ± 0.1	2.0 ± 0.1	0.020*
Vitamin B <sub>2</sub> (mg)	1.4 ± 0.1	1.5 ± 0.1	0.441	1.1 ± 0.1	1.3 ± 0.1	0.042*	1.3 ± 0.0	1.4 ± 0.1	0.065
Niacin (mg)	18.7 ± 0.9	19.9 ± 1.2	0.446	14.8 ± 1.0	15.8 ± 0.8	0.456	17.0 ± 0.7	18.2 ± 0.8	0.261
Vitamin C (mg)	112.7 ± 9.1	127.8 ± 15.9	0.400	104.4 ± 10.3	126.5 ± 13.0	0.170	109.0 ± 7.0	127.3 ± 10.9	0.150
Calcium (mg)	566.9 ± 26.7	533.5 ± 33.7	0.427	442.0 ± 26.9	499.9 ± 38.3	0.203	510.9 ± 20.5	519.9 ± 25.6	0.777
Phosphorus (mg)	1,321.0 ± 46.5	1,336.1 ± 65.8	0.851	1,035.7 ± 53.0	1,071.0 ± 46.5	0.632	1,193.7 ± 36.8	1,226.0 ± 44.8	0.590
Sodium (mg)	4,940.3 ± 288.0	5,132.2 ± 317.2	0.654	3,962.5 ± 389.8	4,576.0 ± 350.2	0.234	4,506.5 ± 236.1	4,899.0 ± 235.5	0.232
Potassium (mg)	3,481.0 ± 151.4	3,456.4 ± 171.6	0.912	2,798.8 ± 166.6	3,140.6 ± 230.5	0.220	3,177.2 ± 115.4	3,325.1 ± 139.6	0.409
Iron (mg)	25.9 ± 7.1	20.0 ± 2.3	0.448	14.5 ± 0.8	17.3 ± 1.1	0.050	20.8 ± 4.0	18.9 ± 1.5	0.654
Distribution									
% Protein	14.0 ± 0.3	14.7 ± 0.6	0.296	14.9 ± 0.5	14.4 ± 0.6	0.516	14.4 ± 0.3	14.6 ± 0.4	0.761
% Fat	16.4 ± 0.8	16.0 ± 0.9	0.709	17.0 ± 0.8	17.6 ± 0.9	0.628	16.7 ± 0.6	16.7 ± 0.7	0.977
% Carbohydrate	63.3 ± 1.5	62.2 ± 1.8	0.626	68.4 ± 1.2	67.7 ± 1.2	0.667	65.6 ± 1.0	64.5 ± 1.2	0.460

Adjusted for age

<sup>1)</sup> Aware group<sup>2)</sup> Non-aware group<sup>3)</sup> Mean ± SD\*  $P < 0.05$ 

was significantly higher in the aware group (55.0%) compared to 35.8% in the non-aware group ( $P = 0.013$ ) and also more women in the aware group were clinically diagnosed with hypertension (53.0%) compared to the non-aware group (30.1%;  $P = 0.008$ ). For men, there was no significant difference in prevalence but a significantly higher percentage of men in the aware group (44.6%) had been clinically diagnosed with hypertension compared to the non-aware group (25.2%;  $P = 0.024$ ). The prevalence of dyslipidemia was not significantly different

between the two groups. However, there were significant differences in the rate of clinical diagnosis of dyslipidemia in both men and women: in men, it was significantly higher in the aware group (31.1%) compared to the non-aware group (11.7%;  $P = 0.001$ ); and in women, it was 47.0% in the aware group and 17.1% in the non-aware group ( $P < 0.001$ ).

#### Clinical indices

The clinical indices of the subjects are shown in Table 3. Total

**Table 5.** Percentage of nutrient intake of Korean Reference Nutrient Intake (RNI) by sex and diabetes awareness.

	Male (n = 251)			Female (n = 272)			Total (n = 523)		
	AG <sup>1)</sup> (n = 148)	NAG <sup>2)</sup> (n = 103)	P-value	AG (n = 149)	NAG (n = 123)	P-value	AG (n = 297)	NAG (n = 226)	P-value
Energy	97.2 ± 3.1 <sup>3)</sup>	109.0 ± 6.4	0.103	90.1 ± 3.9	102.7 ± 3.8	0.025*	94.0 ± 2.6	106.3 ± 4.1	0.013*
Protein	148.8 ± 6.0	177.8 ± 12.8	0.043*	134.0 ± 7.4	154.1 ± 9.9	0.103	142.2 ± 4.6	167.9 ± 8.7	0.012*
Vitamin A	125.4 ± 14.0	106.4 ± 8.7	0.244	126.1 ± 20.6	150.1 ± 16.7	0.356	125.7 ± 11.7	124.6 ± 8.7	0.936
Vitamin B <sub>1</sub>	149.6 ± 6.4	192.0 ± 17.5	0.024*	135.6 ± 7.1	154.1 ± 10.3	0.093	143.4 ± 5.1	176.3 ± 11.1	0.007**
Vitamin B <sub>2</sub>	91.5 ± 4.0	104.3 ± 7.3	0.118	88.6 ± 6.0	111.2 ± 7.5	0.014*	90.2 ± 3.5	107.1 ± 5.5	0.008**
Niacin	114.0 ± 4.6	127.8 ± 8.4	0.158	104.8 ± 6.8	113.9 ± 5.9	0.306	109.9 ± 4.0	122.0 ± 5.6	0.085
Vitamin C	113.6 ± 9.2	126.7 ± 15.8	0.467	106.4 ± 10.6	124.0 ± 13.3	0.300	110.4 ± 7.1	125.6 ± 10.9	0.239
Calcium	77.6 ± 3.7	75.7 ± 5.0	0.752	62.9 ± 3.9	68.2 ± 5.4	0.423	71.1 ± 2.7	72.6 ± 3.7	0.734
Phosphorus	183.4 ± 6.4	196.7 ± 10.8	0.287	147.8 ± 7.0	153.2 ± 6.9	0.594	165.8 ± 4.9	178.6 ± 7.1	0.218
Sodium	343.9 ± 19.8	358.1 ± 23.6	0.652	275.9 ± 26.2	318.2 ± 24.7	0.231	313.6 ± 16.1	341.5 ± 17.1	0.237
Potassium	98.7 ± 4.4	99.6 ± 5.3	0.888	80.6 ± 4.6	88.9 ± 7.0	0.321	90.6 ± 3.3	95.2 ± 4.2	0.407
Iron	271.8 ± 75.6	221.2 ± 28.8	0.539	163.7 ± 8.8	173.5 ± 11.4	0.500	223.7 ± 42.6	201.3 ± 17.6	0.633

<sup>1)</sup> Aware group

<sup>2)</sup> Non-aware group

<sup>3)</sup> Mean ± SD

\*  $P < 0.05$ , \*\*  $P < 0.01$

**Table 6.** Odds ratio (OR) and 95% confidence interval (CI) of biochemical indices and nutrients intake distribution by diabetes awareness.

		Crude OR (95% CI)	P-value	Adjusted OR <sup>1)</sup> (95% CI)	P-value	Adjusted OR <sup>2)</sup> (95% CI)	P-value
Waist circumference M ≥ 90 cm, F ≥ 85 cm	Aware	1		1		1	
	Non-aware	1.93 (1.25-2.98)	0.003**	2.25 (1.30-3.90)	0.004**	2.64 (1.18-5.88)	0.018*
Systolic pressure ≥ 140 mmHg	Aware	1		1		1	
	Non-aware	0.89 (0.48-1.64)	0.710	1.26 (0.59-2.68)	0.548	1.25 (0.60-2.60)	0.545
Diastolic pressure ≥ 90 mmHg	Aware	1		1		1	
	Non-aware	2.10 (1.16-3.8)	0.015*	1.61 (0.76-3.41)	0.212	1.61 (0.78-3.36)	0.200
Total cholesterol ≥ 240 mg/dL	Aware	1		1		1	
	Non-aware	4.08 (2.04-8.14)	0.001***	3.68 (1.74-7.82)	0.001***	3.44 (1.58-7.52)	0.002**
HDL-cholesterol < 40 mg/dL	Aware	1		1		1	
	Non-aware	1.46 (0.95-2.26)	0.088	1.67 (0.96-2.92)	0.071	1.62 (0.93-2.81)	0.088
LDL-cholesterol ≥ 160 mg/dL	Aware	1		1		1	
	Non-aware	5.37 (2.68-10.74)	0.001***	4.46 (2.05-9.07)	0.001***	4.59 (2.07-10.17)	0.001***
Triglyceride ≥ 200 mg/dL	Aware	1		1		1	
	Non-aware	1.98 (1.23-3.17)	0.005**	2.14 (1.17-3.09)	0.013*	1.80 (0.94-3.42)	0.074
Protein intake > 20%	Aware	1		1		1	
	Non-aware	0.89 (0.37-2.12)	0.791	0.73 (0.26-20.4)	0.540	0.77 (0.24-2.82)	0.666
Fat intake > 25%	Aware	1		1		1	
	Non-aware	1.55 (0.87-2.78)	0.137	1.51 (0.73-2.13)	0.261	1.42 (0.68-0.99)	0.348
Carbohydrate intake > 60%	Aware	1		1		1	
	Non-aware	0.76 (0.47-1.22)	0.250	0.79 (0.43-1.46)	0.448	0.94 (0.48-1.85)	0.860

<sup>1)</sup> Adjusted for sex, age, income, education and diabetes duration

<sup>2)</sup> Adjusted for sex, age, income, education, diabetes duration, smoking, drinking, physical activity and BMI

cholesterol and LDL-cholesterol were significantly higher in the non-aware group compared to the aware group in both men and women ( $P < 0.001$ ). In men, triglyceride was significantly higher at 199.3 mg/dL in the non-aware group compared to 153.8 mg/dL in the aware group ( $P < 0.001$ ). Hemoglobin was 14.9 g/dL in the aware group and 15.5 g/dL in the non-aware group ( $P = 0.002$ ). In women, hemoglobin was significantly lower in the aware group ( $P = 0.019$ ). Hematocrit was significantly lower in the aware group compared to the non-aware group in both men ( $P = 0.002$ ) and women ( $P = 0.006$ ).

#### Nutrient intakes

Daily calorie and nutrient intakes are shown in Table 4. There was no significant difference between the aware and the non-aware groups among men. In women, calorie intake was 1,665 kcal in the aware group and 1,891 kcal in the non-aware group ( $P = 0.033$ ). Carbohydrate intake was significantly higher in the non-aware group (319.2 g) compared to the aware group (280.7 g;  $P = 0.048$ ). Vitamin B<sub>2</sub> intake was significantly higher in the non-aware group (1.3 mg) compared to the aware group (1.1 mg;  $P = 0.042$ ). In total, calorie intake was 1,990 kcal in the

aware group and 2,218 kcal in the non-aware group ( $P = 0.025$ ). Protein intake was significantly higher in the non-aware group (81.5 g) compared to the aware group (70.9 g;  $P = 0.038$ ). Carbohydrate intake was significantly higher in the non-aware group (344.6 g) compared to the aware group (314.3 g;  $P = 0.044$ ).

The results of nutrient intake of Korean Reference Nutrient Intake (RNI) of the subjects are shown in Table 5. The ratio of calorie intake to RNI was significantly higher at 106.3% in the non-aware group compared to 94.0% in the aware group ( $P = 0.013$ ), and both men and women in the non-aware group had an intake higher than Korean RNI. In men, protein intake was significantly higher at 177.8% in the non-aware group than 148.8% in the aware group ( $P = 0.043$ ). In men, vitamin B<sub>1</sub> intake was significantly higher in the non-aware group compared to the aware group ( $P = 0.024$ ). In women, vitamin B<sub>2</sub> intake was significantly higher in the non-aware group compared to the aware group ( $P = 0.014$ ).

The degree of risk exerted by diabetes awareness on clinical indices and nutrient intakes is shown in Table 6. The crude risk for abnormal values was significantly higher for waist circumference, diastolic blood pressure, total cholesterol, LDL-cholesterol and triglycerides in the non-aware group. After controlling for gender, age, income, education level, and the duration of diabetes, the risk for abnormal values was significantly higher for waist circumference, total cholesterol, LDL-cholesterol and triglycerides in the non-aware group. After additional correction for smoking, drinking, exercise and BMI, the risk for waist circumference over 90cm in men and over 85cm in women was 2.64 times (95% CI: 1.18-5.88) higher in the non-aware group ( $P = 0.018$ ); and the risk for total cholesterol over 240 mg/dL was 3.44 times (95% CI: 1.58-7.52) higher in the non-aware group ( $P = 0.002$ ). The risk for LDL-cholesterol over 160 mg/dL was 4.59 times (95% CI: 2.07-10.17) higher in the non-aware group ( $P < 0.001$ ). Among nutrient intakes, abnormal intake ratios of protein, fat and carbohydrates were not significantly different.

## DISCUSSION

This is the first study analyzing the health-related habits, clinical indices and nutrient intakes based on the diabetes awareness in the Korean adult population using nationally representative data (2012-2013 KNHANES) [3]. Early diagnosis and treatment of diabetes significantly reduces the risks for CVDs [19]. Therefore, providing men in their 30s with diabetes education would benefit the population health in Korea by reducing the prevalence and enhancing early treatment of diabetes in young men.

BMI was significantly higher in the non-aware group compared to the aware group and the ratio of BMI over 25 kg/m<sup>2</sup> was 45.5% and 62.4% in the aware and non-aware groups, respectively. Obesity is a major preceding factor of diabetes and a factor for preventing and delaying diabetes in individuals with the possibility of progressing diabetes. Weight control for obesity is needed in both the aware and non-aware groups.

Hypertension is a common comorbidity in diabetic patients and a major risk factor for CVDs [20,21]. Strict blood pressure control can reduce the mortality and delay complications such

as diabetic retinopathy and reduced vision [22]. The study results indicated that the non-aware group was unaware of not only diabetes but also accompanying diseases such as hypertension and dyslipidemia. The diagnosis rate for hypertension and dyslipidemia was lower than the prevalence rate, demonstrating uncontrolled blood pressure and blood lipid profiles in the aware group. Screening and strict management of blood pressure, LDL-cholesterol, triglycerides and HDL-cholesterol are recommended for diabetic individuals. The results of this study were consistent with those in the studies on diabetic patients by Kwon *et al.* [23] and on diabetes awareness by Kim *et al.* [5]. Cardiovascular diseases rank as the 2<sup>nd</sup> and 3<sup>rd</sup> causes of death in Korea and individuals with diabetes have higher risks for CVDs [24]. Because the prevalence of CVD is 2-4 times higher in diabetic patients than in non-diabetic patients, early diagnosis in the non-aware group can promote the regulation of risk factors for CVDs through glycemic control and blood lipid control [25,26].

Park *et al.* [27] showed a significant relationship between carbohydrate intake and waist circumference after controlling for total energy intake and BMI. The ADA recommends that the total amount rather than the type of carbohydrates is important in the intake of carbohydrates. Thus, education on carbohydrate control is highly recommended.

Limitations of the study include that the definition of the non-aware group was based on either fasting blood glucose or HbA<sub>1c</sub> data from the KNHANES. Because it relied on one measurement for each individual, it is possible that some participants might have been miss-categorized as diabetic.

Although the currently recommended target for diabetes screening is individuals over 40 years or over 30 years in case of having diabetic risk factors, this study suggested that diabetes screening before 40 years of age through regular health check-up would be beneficial. In the aware group, continuous and systematic management and education are necessary to prevent diabetic complications. Careful self-monitoring for glycemic control is also recommended for diabetic patients. To increase overall diabetes awareness, not only individual-level education but also social marketing or other community-level education strategies will be beneficial. Further studies determining effective strategies to increase diabetes awareness and screening as well as rigorous diabetes management programs are needed.

## CONFLICT OF INTEREST

The authors declare no potential conflicts of interests.

## ORCID

Seul Lee: <https://orcid.org/0000-0003-4145-9865>

Haeryun Park: <https://orcid.org/0000-0003-2627-6518>

Youngmi Lee: <https://orcid.org/0000-0001-9965-0748>

Onjeong Choi: <https://orcid.org/0000-0001-5093-8476>

Jiwon Kim: <https://orcid.org/0000-0002-3307-1441>

Heewon L Gray: <https://orcid.org/0000-0002-5257-5012>

Kyunghee Song: <https://orcid.org/0000-0001-9549-0716>

## REFERENCES

1. World Health Organization. The-top-10-causes-of-death[Internet]. Geneva: World Health Organization; 2018 [cited 2019 May 27]. Available from: <https://www.who.int/news-room/fact-sheets/detail/the-top-10-causes-of-death>.
2. Bloomgarden ZT. Diabetes and cardiovascular disease. *Diabetes Care* 2011;34:e24-30.
3. Korean Diabetes Association. Diabetes Fact Sheet in Korea [Internet]. Seoul: Korean Diabetes Association; 2018 [cited 2019 May 27]. Available from: <http://www.diabetes.or.kr/pro/news/admin.php?category=A&code=admin&number=1428&mode=view>.
4. Gregg EW, Cadwell BL, Cheng YJ, Cowie CC, Williams DE, Geiss L, Engelgau MM, Vinicor F. Trends in the prevalence and ratio of diagnosed to undiagnosed diabetes according to obesity levels in the U.S. *Diabetes Care* 2004;27:2806-12.
5. Kim SM, Lee JS, Lee J, Na JK, Han JH, Yoon DK, Baik SH, Choi DS, Choi KM. Prevalence of diabetes and impaired fasting glucose in Korea: Korean National Health and Nutrition Survey 2001. *Diabetes Care* 2006;29:226-31.
6. Korean Diabetes Association. Treatment Guideline for Diabetes, 5th ed [Internet]. Seoul: Korean Diabetes Association; 2015 [cited 2019 May 27]. Available from: [http://www.diabetes.or.kr/pro/publish/guide.php?code=guide&mode=list&year\\_v=2015](http://www.diabetes.or.kr/pro/publish/guide.php?code=guide&mode=list&year_v=2015).
7. Franz MJ, Bantle JP, Beebe CA, Brunzell JD, Chiasson JL, Garg A, Holzmeister LA, Hoogwerf B, Mayer-Davis E, Mooradian AD, Purnell JQ, Wheeler M. Evidence-based nutrition principles and recommendations for the treatment and prevention of diabetes and related complications. *Diabetes Care* 2002;25:148-98.
8. Lim JH, Lee YS, Chang HC, Moon MK, Song Y. Association between dietary patterns and blood lipid profiles in Korean adults with type 2 diabetes. *J Korean Med Sci* 2011;26:1201-8.
9. Park YM, Sohn CM, Jang HC. Correlation of carbohydrate intake with obesity in type 2 diabetes mellitus patients. *J Korean Diet Assoc* 2006;12:254-63.
10. Kim EK, Lee JS, Hong HO, Yu CH. Association between glycemic index, glycemic load, dietary carbohydrates and diabetes from Korean National Health and Nutrition Examination Survey 2005. *Korean J Nutr* 2009;42:622-30.
11. Ceriello A, Testa R. Antioxidant anti-inflammatory treatment in type 2 diabetes. *Diabetes Care* 2009;32 Suppl 2:S232-6.
12. Weiss R, Dufour S, Taksali SE, Tamborlane WV, Petersen KF, Bonadonna RC, Boselli L, Barbetta G, Allen K, Rife F, Savoye M, Dziura J, Sherwin R, Shulman GI, Caprio S. Prediabetes in obese youth: a syndrome of impaired glucose tolerance, severe insulin resistance, and altered myocellular and abdominal fat partitioning. *Lancet* 2003;362:951-7.
13. Girach A, Manner D, Porta M. Diabetic microvascular complications: can patients at risk be identified? A review. *Int J Clin Pract* 2006;60:1471-83.
14. Harris MI, Klein R, Welborn TA, Knudman MW. Onset of NIDDM occurs at least 4-7 yr before clinical diagnosis. *Diabetes Care* 1992;15:815-9.
15. UK Prospective Diabetes Study Group. UK Prospective Diabetes Study 6. Complications in newly diagnosed type 2 diabetic patients and their association with different clinical and biochemical risk factors. *Diabetes Res* 1990;13:1-11.
16. Siitonen O, Uusitupa M, Pyörälä K, Voutilainen E, Länsimies E. Peripheral arterial disease and its relationship to cardiovascular risk factors and coronary heart disease in newly diagnosed non-insulin-dependent diabetics. *Acta Med Scand* 1986;220:205-12.
17. Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* 1972;18:499-502.
18. Harris MI. Undiagnosed NIDDM: clinical and public health issues. *Diabetes Care* 1993;16:642-52.
19. Harris MI, Eastman RC. Early detection of undiagnosed diabetes mellitus: a US perspective. *Diabetes Metab Res Rev* 2000;16:230-6.
20. Wingard DL, Barrett-Connor E. Heart disease and diabetes. In: Harris MI, Cowie CC, Stern MS, editors. *Diabetes in America*. Washington, D.C.: National Institutes of Health; 1995. p.429-48.
21. UK Prospective Diabetes Study Group. Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38. *BMJ* 1998;317:703-13.
22. Arauz-Pacheco C, Parrott MA, Raskin P. The treatment of hypertension in adult patients with diabetes. *Diabetes Care* 2002;25:134-47.
23. Kwon JY, Chung HY. Study on the correlation between the nutrient intakes and clinical indices of type 2 diabetes patients. *Korean J Food Nutr* 2013;26:909-18.
24. Committee for Guidelines for Management of Dyslipidemia. 2015 Korean guidelines for management of dyslipidemia. *J Lipid Atheroscler* 2015;4:61-92.
25. Jee SH, Jang Y, Oh DJ, Oh BH, Lee SH, Park SW, Seung KB, Mok Y, Jung KJ, Kimm H, Yun YD, Baek SJ, Lee DC, Choi SH, Kim MJ, Sung J, Cho B, Kim ES, Yu BY, Lee TY, Kim JS, Lee YJ, Oh JK, Kim SH, Park JK, Koh SB, Park SB, Lee SY, Yoo CI, Kim MC, Kim HK, Park JS, Kim HC, Lee GJ, Woodward M. A coronary heart disease prediction model: the Korean Heart Study. *BMJ Open* 2014;4:e005025.
26. Collins R, Armitage J, Parish S, Sleight P, Peto R; Heart Protection Study Collaborative Group. MRC/BHF Heart Protection Study of cholesterol-lowering with simvastatin in 5963 people with diabetes: a randomised placebo-controlled trial. *Lancet* 2003;361:2005-16.
27. Park S, Park MS, Ko JA. The association between carbohydrate intake and waist circumference. *Korean J Obes* 2008;17:175-81.