



Chemical Constituents of *Impatiens balsamina* Stems and Their Biological Activities

Dong Hyun Kim¹, Tae Hyun Lee¹, Lalita Subedi^{2,3}, Sun Yeou Kim^{2,3}, and Kang Ro Lee^{1,*}

¹Natural Products Laboratory, School of Pharmacy, Sungkyunkwan University, Suwon 16419, Korea

²Gachon Institute of Pharmaceutical Science, Gachon University, 191 Hambakmoero, Yeonsu-gu, Incheon 21936, Republic of Korea

³College of pharmacy, Gachon University, 191 Hambakmoero, Yeonsu-gu, Incheon 21936, Republic of Korea

Abstract – The purification of the MeOH extract from *Impatiens balsamina* by repeated column chromatography led to the isolation of one new tetrahydronaphthalene (**1**), together with eleven known compounds (**2** - **12**). The structure of the new compound (**1**) was determined by spectral data analysis (¹H and ¹³C-NMR, ¹H-¹H COSY, HSQC, HMBC, NOESY, and HR-ESI-MS). Isolated compounds (**1** - **12**) were evaluated for their inhibitory effects on NO production in LPS-activated murine microglial BV-2 cells and their effects on NGF secretion from C6 glioma cells. Compounds **3**, **7**, and **10** reduced NO levels in LPS-activated murine microglial cells with IC₅₀ values of 26.89, 25.59, and 44.21 μM, respectively. Compounds **1**, **5**, and **9** upregulated NGF secretion to 153.09 ± 4.66, 156.88 ± 8.86, and 157.34 ± 3.30%, respectively.

Keywords – *Impatiens balsamina*, balsaminaceae, tetrahydronaphthalene, neuroprotective effect

Introduction

Impatiens balsamina (Balsaminaceae), also known as Garden balsam, is widely distributed in Korea, China and India. *I. balsamina* has been used as Chinese medicine to treat anti-cancer and anti-inflammatory herb, also has been used as a Korean traditional medicine for the treatment of scrofulosis, carbuncle, and dysentery.¹⁻³ Previous phytochemical investigations on *I. balsamina* reported the isolation of triterpenoid saponins, quinones, coumarin, flavonoids, and phenolic compounds, and there were several reports on various biological activities.⁴⁻¹⁰ Our earlier phytochemical investigation on white petal of *I. balsamina* resulted in the isolation of anti-neurodegenerative biflavonoid glycosides and anti-inflammatory phenolic compounds.^{9,10} In a continuing search for bioactive constituents from Korean medical plant sources, we investigated the MeOH extract of stems of *I. balsamina*. The purification of hexane and EtOAc fractions by repeated column chromatography led to the isolation of one new tetrahydronaphthalene (**1**), together with eleven known compounds (**2** - **12**). The structure of **1** was established on spectral data analysis (¹H and ¹³C-NMR, ¹H-¹H COSY, HSQC, HMBC, and NOESY). Isolated com-

pounds (**1** - **12**) were evaluated for their inhibitory effects on nitric oxide (NO) production in lipopolysaccharide (LPS)-activated murine microglial cells and their effects on NGF secretion from C6 glioma cells.

Experimental

General experimental procedures – Optical rotations were measured on a Jasco P-2000 polarimeter using methanol solvent. Ultraviolet (UV) spectra were recorded with a Shimadzu UV-1601 UV-vis spectrometer. High Resolution ESI Mass Spectrometer data were obtained with a Waters SYNAPT G2 mass spectrometer. Infrared (IR) spectra were recorded on a Bruker IFS-66/S Fourier-transform IR spectrometer. NMR spectra were recorded on Varian UNITY INOVA 500 NMR spectrometer at 500 MHz (¹H) and 125 MHz (¹³C), and Bruker AVANCE III 700 NMR spectrometer at 700 MHz (¹H) and 175 MHz (¹³C). Preparative HPLC was performed using a Gilson 321 pump with a Shodex Refractive Index Detector, YMC-Triart C₁₈ 5 μm column (250 × 10 mm) and HAISIL 100 silica 5 μm column (250 × 10 mm). Silica gel 60 (Merck, 70 - 230 mesh and 230 - 400 mesh) and RP-C₁₈ silica gel (Merck, 230 - 400 mesh) was used for column chromatography. LPLC was performed over a LiChrorep Lobar-A silica 60 column (Merck, 240 mm × 10 mm i.d.) equipped with a FMI QSY-0 pump. Merck precoated

*Author for correspondence

Kang Ro Lee, Natural Products Laboratory, School of Pharmacy, Sungkyunkwan University, Suwon 16419, Korea
Tel: +82-31-290-7710; E-mail: krlee@skku.ac.kr

silica gel F₂₅₄ plates and reversed-phase (RP)-18 F_{254s} plates (Merck) were used for thin-layer chromatography (TLC). Spots were detected on TLC under UV light or by heating after spraying with anisaldehyde-sulfuric acid

Plant materials – The dried stems of *I. balsamina* (2 kg) were collected at Asan in Chungcheongnam-Do, Korea, in August 2014. The plants were identified by one of the authors (K.R.L.). A voucher specimen (SKKU-NPL-1425) of the plant was deposited in the herbarium of the School of Pharmacy at Sungkyunkwan University, Suwon, Korea.

Extraction and isolation – The dried stems of *I. balsamina* (2 kg) were extracted with 80% MeOH at room temperature. The filtrate was evaporated *in vacuo* to yield MeOH extract (294 g), which was suspended in distilled H₂O (2.4 L) and then successively partitioned with hexane, CHCl₃, EtOAc and *n*-BuOH, yielding 20, 11, 6, and 28 g, respectively. The EtOAc fraction (5 g) was separated over a silica gel column with a solvent system of CHCl₃/MeOH/water (7:1:0.1) to yielded five sub-fractions (E1-E5). Sub-fraction E1 (1.8 g) was chromatographed on a RP-C₁₈ silica gel column with 30% aqueous MeOH to yield nine sub-fractions (E11-E19). Fraction E12 (430 mg) was purified by RP-C₁₈ semi-prep. HPLC (30% MeOH) to yield **2** (9 mg, *t*_R = 17.1 min). Fraction E2 (1.1 g) was separated by RP-C₁₈ silica gel column with 30% aqueous MeOH to give nine sub-fractions (E21-E29). Sub-fraction E23 (161 mg) was purified using RP-C₁₈ semi-prep. HPLC (30% MeOH) to yield **1** (20 mg, *t*_R = 15.1 min), **6** (24 mg, *t*_R = 21.1 min), and **3** (7 mg, *t*_R = 22.4 min). E3 (557 mg) was subjected to RP-C₁₈ silica gel column with 45% aqueous MeOH to give five sub-fractions (E31-E35). Fraction E31 (148 mg) was purified by RP-C₁₈ semi-prep. HPLC (30% CH₃CN) to

yield **4** (100 mg, *t*_R = 11.1 min). Fraction E5 (235 mg) was purified using RP-C₁₈ semi-prep. HPLC (20% CH₃CN) to yield **5** (32 mg, *t*_R = 21.2 min). The hexane soluble fraction was chromatographed on a silica gel column with hexane/EtOAc (20:1 to 1:1) to give eight sub-fractions (H1-H8). Fraction H2 (546 mg) was subjected to using a RP-C₁₈ silica gel column with 95% aqueous MeOH to yield **11** (9 mg, *t*_R = 34.4 min). Fraction H3 (918 mg) was separated to RP-C₁₈ silica gel column with 90% aqueous MeOH to yield eight sub-fractions (H31-H38). Subfraction H32 (85 mg) was purified using RP-C₁₈ semi-prep. HPLC (90% MeOH) to yield **7** (9 mg, *t*_R = 36.9 min). Fraction H36 (117 mg) was purified by semi-prep. HPLC (CHCl₃/MeOH 120:1) to give **12** (88 mg, *t*_R = 11.0 min). Fraction H5 (2.4 g) was chromatographed on a RP-C₁₈ silica gel column (95% aqueous MeOH) to give twelve sub-fractions (H51-H512). Subfraction H56 (89 mg) was purified using RP-C₁₈ semi-prep. HPLC (95% MeOH) to yield **8** (4 mg, *t*_R = 17.8 min). Fraction H57 (169 mg) was purified by semi-prep. HPLC (hexane/EtOAc 3:1) to yield **9** (3 mg, *t*_R = 15.3 min). Fraction H512 (112 mg) was purified using semi-prep. HPLC (hexane/EtOAc 3:1) to give **10** (7 mg, *t*_R = 8.8 min).

1 β ,2 α ,4 β -Triol-1,2,3,4-tetrahydronaphthalene (1) – Colorless gum. [α]_D²⁵: -35.8 (*c* 0.10, MeOH); IR (KBr) ν_{\max} cm⁻¹: 3348, 2938, 2836, 1448, 1412, 1055, 1025; UV (MeOH) λ_{\max} (log ϵ) 201 (0.75), 213 (0.45), 217 (0.38), 262 (0.02) nm; ¹H and ¹³C NMR : see Table 1; HRESIMS *m/z* 181.0865 [M+H]⁺; (calcd. for C₁₀H₁₃O₃, 181.0865).

1 β ,2 β ,4 β -Triol-1,2,3,4-tetrahydronaphthalene (2) – Colorless gum. [α]_D²⁵: -57.8 (*c* 0.11, MeOH); IR (KBr) ν_{\max} cm⁻¹: 3340, 2944, 2832, 1460, 1408, 1054, 1021; UV (MeOH) λ_{\max} (log ϵ) 203 (1.28), 211 (1.11), 215 (0.92), 256 (0.04) nm; ¹H and ¹³C NMR : see Table 1.

Table 1. ¹H and ¹³C NMR data of **1** and **2** in CD₃OD. (δ in ppm, 500 MHz for ¹H and 125 MHz for ¹³C)^a

Position	1		2	
	δ_{H}	δ_{C}	δ_{H}	δ_{C}
1	4.41 d (7.3)	73.6	4.64 d (3.5)	70.1
2	4.09 ddd (10.1, 7.3, 3.4)	68.9	3.89 dt (10.9, 3.5)	67.7
3	2.12 ddd (13.5, 10.1, 4.7)	37.1	2.15 ddd (12.2, 10.9, 9.5)	34.9
			2.21 dddd (12.2, 6.1, 3.5, 1.0)	
4	4.89 m	66.6	4.73 dd (9.5, 6.1)	67.4
4a		137.5		139.4
5	7.41 dd (7.3, 1.5)	128.1	7.57 d (7.5)	126.7
6	7.34 td (7.3, 1.5)	127.1	7.30 td (7.5, 1.6)	127.9
7	7.29 td (7.3, 1.5)	127.6	7.34 td (7.5, 1.6)	127.3
8	7.56 d (7.3)	127.4	7.41 dd (7.5, 1.6)	129.6
8a		137.6		135.9

^aJ values are in parentheses and reported in Hz; the assignments were based on ¹H-¹H COSY, HSQC, and HMBC experiments.




Fig. 1. Structures of compounds 1 - 12 isolated from *I. balsamina*.

(7*R*,8*S*)-Dihydrodehydrodiconiferyl alcohol-9- β -D-glucopyranoside (3) – White gum; ^1H NMR (CD_3OD , 500 MHz): δ 6.99 (1H, d, $J = 1.9$ Hz, H-2), 6.85 (1H, dd, $J = 8.1, 1.9$ Hz, H-6), 6.79 (1H, s, H-6'), 6.75 (1H, d, $J = 8.1$ Hz, H-5), 6.72 (1H, s, H-2'), 5.58 (1H, d, $J = 6.4$ Hz, H-7), 4.35 (1H, d, $J = 7.8$ Hz, H-1''), 3.85 (3H, s, 3'-OCH₃), 3.82 (3H, s, 3-OCH₃), 3.56 (2H, t, $J = 6.5$ Hz, H-9'), 2.62 (2H, m, H-7'), 1.81 (2H, tt, $J = 13.1, 6.5$ Hz, H-8'); ^{13}C NMR (CD_3OD , 125 MHz): δ 147.6 (C-3), 146.1 (C-4), 146.0 (C-4'), 143.8 (C-3'), 135.6 (C-1'), 133.2 (C-1), 128.3 (C-5'), 118.4 (C-6), 116.8 (C-6'), 114.6 (C-5), 112.8 (C-2'), 109.4 (C-2), 102.8 (C-1''), 87.8 (C-7), 76.8 (C-5''), 76.7 (C-3''), 73.7 (C-2''), 70.9 (C-9), 70.2 (C-4''), 61.3 (C-6''), 60.8 (C-9'), 55.4 (3'-OCH₃), 55.1 (3-OCH₃), 51.5 (C-8), 34.4 (C-8'), 31.5 (C-7').

Kaempferol-3- O - β -D-glucopyranoside (4) – Yellow needles; ^1H NMR (CD_3OD , 500 MHz): δ 8.08 (2H, d, $J = 8.8$ Hz, H-2', 6'), 6.91 (2H, d, $J = 8.8$ Hz, H-3', 5'), 6.43 (1H, s, H-8), 6.23 (1H, s, H-6), 5.28 (1H, d, $J = 7.4$ Hz, H-1''); ^{13}C NMR (CD_3OD , 125 MHz): δ 178.1 (C-4), 164.7 (C-7), 161.7 (C-5), 160.1 (C-4'), 157.7 (C-9), 157.1 (C-2), 134.0 (C-3), 130.8 (C-2', 6'), 121.4 (C-1'), 114.7 (C-3', 5'), 104.3 (C-10), 102.7 (C-1''), 98.5 (C-6), 93.6 (C-8), 77.2 (C-5''), 76.7 (C-3''), 74.3 (C-2''), 70.0 (C-4''), 61.2 (C-6'').

Nicotiflorin (5) – Yellow needles; ^1H NMR (CD_3OD , 500 MHz): δ 8.09 (2H, d, $J = 8.8$ Hz, H-2', 6'), 6.91 (2H, d, $J = 8.8$ Hz, H-3', 5'), 6.42 (1H, s, H-8), 6.23 (1H, d, $J = 1.8$ Hz, H-6), 5.15 (1H, d, $J = 7.4$ Hz, H-1''), 4.54 (1H, d, $J = 0.9$ Hz, H-1'''), 3.66 (1H, m, H-2'''), 3.55 (1H, dd, $J = 9.5, 3.3$ Hz, H-3'''), 1.14 (3H, d, $J = 6.2$ Hz, H-6'''); ^{13}C

Table 2. Inhibitory effects of compounds **1** - **12** on NO production in LPS-activated BV-2 cells

Compound	IC ₅₀ (μM) ^a	Cell Viability (% of LPS) ^b
1	50.29	89.42 ± 6.36
2	>500	93.08 ± 5.10
3	26.89	109.90 ± 3.47
4	60.27	107.24 ± 2.72
5	>500	92.35 ± 7.51
6	108.55	112.35 ± 4.80
7	25.59	111.09 ± 4.11
8	>500	82.11 ± 2.89
9	>500	91.41 ± 7.19
10	44.21	101.07 ± 2.86
11	94.38	97.45 ± 11.50
12	>500	86.83 ± 4.21
L-NMMA ^c	21.40	100.14 ± 2.54

^aIC₅₀ value of each compound was defined as the concentration (μM) that caused

50% inhibition of NO production in LPS-activated BV-2 cells.

^bCell viability after treatment with 20 μM of each compound was determined by

MTT assay and is expressed in percentage (%). The results are averages of three

independent experiments, and the data are expressed as mean ± SD.

^cL-NMMA as positive control.

NMR (CD₃OD, 125 MHz): δ 178.0 (C-4), 164.8 (C-7), 161.6 (C-5), 160.1 (C-4'), 158.0 (C-9), 157.2 (C-2), 134.1 (C-3), 130.1 (C-2', 6'), 121.4 (C-1'), 114.7 (C-3', 5'), 104.2 (C-10), 103.2 (C-1''), 101.0 (C-1'''), 98.6 (C-6), 93.5 (C-8), 76.8 (C-3''), 75.8 (C-5''), 74.4 (C-2''), 72.5 (C-4''), 70.9 (C-2'''), 70.7 (C-3'''), 70.0 (C-4'''), 68.3 (C-5'''), 67.2 (C-6''), 16.5 (C-6''').

p-Hydroxybenzoic acid (6) – White gum; ¹H NMR (CD₃OD, 500 MHz): δ 7.90 (2H, d, *J* = 8.8 Hz, H-2', 6'), 6.84 (2H, d, *J* = 8.8 Hz, H-3', 5'); ¹³C NMR (CD₃OD, 125 MHz): δ 170.1 (C-7), 163.3 (C-4), 133.0 (C-2, 6), 122.8 (C-1), 116.0 (C-3, 5).

β-Amyrin (7) – Colorless gum; ¹H NMR (CDCl₃, 500 MHz): δ 5.19 (1H, t, *J* = 3.6 Hz, H-12), 3.22 (1H, dd, *J* = 11.0, 4.6 Hz, H-3), 1.14 (3H, s, H-27), 1.01 (3H, s, H-26), 0.97 (3H, s, H-23), 0.94 (3H, s, H-25), 0.88 (6H, s, H-29, 30), 0.84 (3H, s, H-28), 0.80 (3H, s, H-24); ¹³C NMR (CDCl₃, 175 MHz): δ 145.4 (C-13), 121.9 (C-12), 79.3 (C-3), 55.4 (C-5), 47.8 (C-9), 47.4 (C-18), 47.0 (C-19), 41.9 (C-14), 40.0 (C-8), 39.0 (C-4), 38.8 (C-1), 37.3 (C-10), 37.2 (C-22), 34.9 (C-21), 33.5 (C-29), 32.8 (C-7), 32.7 (C-17), 31.3 (C-20), 28.6 (C-28), 28.3 (C-23), 27.3 (C-2), 27.1 (C-16), 26.3 (C-15), 26.2 (C-27), 23.9 (C-30), 23.7 (C-11), 18.5 (C-6), 17.0 (C-26), 15.8 (C-25), 15.7 (C-24).

Table 3. Effects of compounds **1** - **12** on NGF secretion in C6 cells

Compound	NGF secretion (%) ^a	Cell Viability (%) ^b
1	153.09 ± 4.66	98.69 ± 3.69
2	102.90 ± 2.03	98.46 ± 11.25
3	105.83 ± 1.44	97.02 ± 8.02
4	137.97 ± 1.34	96.55 ± 7.38
5	156.88 ± 8.86	96.81 ± 17.29
6	149.21 ± 0.70	86.72 ± 0.02
7	129.76 ± 0.91	91.60 ± 0.70
8	94.75 ± 8.47	91.65 ± 1.20
9	157.34 ± 3.30	95.52 ± 0.38
10	130.55 ± 11.41	98.64 ± 4.12
11	151.05 ± 2.56	103.14 ± 0.62
12	104.93 ± 4.75	83.43 ± 2.48
6-Shogaol ^c	164.74 ± 1.95	100.34 ± 7.91

^aC6 cells were treated with 20 μM of the compounds. After 24 h, the content of

NGF secreted into C6-conditioned media was measured by ELISA. The level of

secreted NGF is expressed as a percentage of the untreated control (set at 100%).

Data are the mean ± SD of three independent experiments performed in triplicate.

^bCell viability after treatment with 20 μM of each compound was determined by

the MTT assay and is expressed as a percentage (%). Results are the mean of three

independent experiments, and the data are expressed as the mean ± SD.

^c6-shogaol was used as positive control.

Erythrodiol (8) – Colorless gum; ¹H NMR (CDCl₃, 500 MHz): δ 5.20 (1H, t, *J* = 3.5 Hz, H-12), 3.55 (1H, d, *J* = 10.9 Hz, H-28a), 3.22 (1H, d, *J* = 10.9 Hz, H-28b), 3.22 (1H, d, *J* = 10.9 Hz, H-3), 1.17 (3H, s, H-27), 1.00 (3H, s, H-23), 0.95 (3H, s, H-26), 0.94 (3H, s, H-25), 0.89 (3H, s, H-29), 0.88 (3H, s, H-30), 0.79 (3H, s, H-24); ¹³C NMR (CDCl₃, 175 MHz): δ 144.2 (C-13), 122.4 (C-12), 79.0 (C-3), 69.7 (C-28), 55.2 (C-5), 47.6 (C-9), 46.5 (C-19), 42.4 (C-18), 41.7 (C-14), 39.8 (C-8), 38.9 (C-4), 38.6 (C-1), 37.0 (C-10, 17), 34.1 (C-21), 33.2 (C-29), 32.6 (C-7), 31.0 (C-20), 30.9 (C-22), 28.0 (C-23), 27.2 (C-2), 25.9 (C-27), 25.6 (C-15), 23.6 (C-11), 23.5 (C-30), 22.0 (C-16), 18.3 (C-6), 16.7 (C-26), 15.6 (C-24), 15.5 (C-25).

α-Spinasterol (9) – White needles; ¹H NMR (CDCl₃, 500 MHz): δ 5.16 (1H, dd, *J* = 15.7, 8.7 Hz, H-22), 5.03 (1H, dd, *J* = 15.2, 8.6 Hz, H-23), 3.60 (1H, tt, *J* = 10.9, 4.5 Hz, H-3), 1.03 (3H, d, *J* = 6.6 Hz, H-3), 0.85 (3H, d, *J* = 6.5 Hz, H-26), 0.56 (3H, s, H-18); ¹³C NMR (CDCl₃, 175 MHz): δ 139.8 (C-8), 139.4 (C-22), 129.7 (C-23), 117.7 (C-7), 71.3 (C-3), 56.1 (C-17), 55.4 (C-14), 51.5 (C-24), 49.7 (C-9), 43.5 (C-13), 41.0 (C-20), 40.5 (C-5), 39.7 (C-12), 38.2 (C-4), 37.4 (C-1), 34.4 (C-10), 32.1 (C-

25), 31.7 (C-2), 29.9 (C-6), 28.7 (C-16), 25.6 (C-28), 23.2 (C-15), 21.8 (C-11), 21.6 (C-26), 21.3 (C-21), 19.2 (C-27), 13.2 (C-19), 12.5 (C-29), 12.3 (C-18).

29-Nor-20-oxolupeol (10) – White gum; ^1H NMR (CDCl_3 , 500 MHz): δ 3.19 (1H, dd, $J = 11.4, 4.8$ Hz, H-3), 2.58 (1H, td, $J = 11.3, 6.0$ Hz, H-19), 2.15 (3H, s, H-30), 1.02 (3H, s, H-26), 0.97 (6H, s, H-23, 27), 0.83 (3H, s, H-25), 0.78 (3H, s, H-28), 0.76 (3H, s, H-24); ^{13}C NMR (CDCl_3 , 175 MHz): δ 213.1 (C-20), 79.2 (C-3), 55.5 (C-5), 52.9 (C-19), 50.5 (C-9), 49.9 (C-18), 43.3 (C-14), 42.9 (C-17), 41.0 (C-8), 40.1 (C-22), 39.1 (C-4), 38.9 (C-1), 37.4 (C-10), 37.3 (C-13), 35.2 (C-16), 34.4 (C-7), 29.4 (C-30), 28.2 (C-23), 27.9 (C-21), 27.6 (C-15), 27.5 (C-2), 27.4 (C-12), 21.1 (C-11), 18.5 (C-6), 18.2 (C-28), 16.3 (C-25), 16.1 (C-26), 15.6 (C-24), 14.7 (C-27).

Lupenone (11) – Colorless gum; ^1H NMR (CDCl_3 , 500 MHz): δ 4.69 (1H, d, $J = 2.4$ Hz, H-29a), 4.57 (1H, dd, $J = 2.4, 1.4$ Hz, H-29b), 1.69 (3H, brs, H-30), 1.07 (6H, s, H-24, 26), 1.03 (3H, s, H-23), 0.96 (3H, s, H-27), 0.93 (3H, s, H-25), 0.80 (3H, s, H-28); ^{13}C NMR (CDCl_3 , 175 MHz): δ 218.2 (C-3), 150.9 (C-20), 109.4 (C-29), 54.9 (C-5), 49.8 (C-9), 48.2 (C-18), 47.9 (C-19), 47.3 (C-4), 43.0 (C-14), 42.9 (C-17), 40.8 (C-8), 40.0 (C-22), 39.6 (C-2), 38.2 (C-13), 36.9 (C-10), 35.5 (C-16), 34.1 (C-1), 33.6 (C-7), 29.8 (C-21), 27.4 (C-15), 26.6 (C-23), 25.1 (C-12), 21.5 (C-11), 21.0 (C-24), 19.7 (C-6), 19.3 (C-30), 18.0 (C-28), 16.0 (C-25), 15.8 (C-26), 14.5 (C-27).

Lupeol (12) – White gum; ^1H NMR (CDCl_3 , 500 MHz): δ 4.69 (1H, d, $J = 2.3$ Hz, H-29a), 4.57 (1H, dd, $J = 2.4, 1.4$ Hz, H-29b), 3.19 (1H, dd, $J = 11.4, 4.9$ Hz, H-3), 2.37 (1H, td, $J = 11.1, 5.8$ Hz, H-19), 1.68 (3H, brs, H-30), 1.03 (3H, s, H-26), 0.97 (3H, s, H-27), 0.95 (3H, s, H-23), 0.83 (3H, s, H-25), 0.79 (3H, s, H-24), 0.76 (3H, s, H-28); ^{13}C NMR (CDCl_3 , 175 MHz): δ 151.2 (C-20), 109.4 (C-29), 79.1 (C-3), 55.6 (C-5), 50.5 (C-9), 48.7 (C-19), 48.3 (C-18), 43.2 (C-17), 43.1 (C-14), 41.1 (C-8), 40.3 (C-22), 39.0 (C-1, 4), 38.3 (C-13), 37.2 (C-10), 35.6 (C-16), 34.4 (C-7), 30.1 (C-21), 28.4 (C-23), 27.9 (C-15), 27.7 (C-2), 25.4 (C-12), 21.3 (C-11), 19.7 (C-30), 18.6 (C-6), 18.3 (C-28), 16.5 (C-25), 16.3 (C-26), 15.7 (C-24), 14.7 (C-27).

Measurement of NO Production and Cell Viability – Murine microglial BV-2 cells were plated into a 96-well plate (3×10^4 cells/well). After 24 h, cells were pretreated with samples for 30 min and then stimulated with 100 ng/mL of LPS for another 24 h. Nitrite, a soluble oxidation product of NO, was measured in culture media using the Griess reaction. The supernatant (50 μL) was harvested and mixed with an equal volume of Griess reagent (1% sulfanilamide, 0.1% *N*-1-naphthylethylenediamine dihydrochloride in 5% phosphoric acid). After 10 min, the

absorbance at 540 nm was measured using a microplate reader. Sodium nitrite was used as a standard to calculate the NO_2^- concentration. Cell viability was tested by the 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl-tetrazolium bromide (MTT) assay. N^G -Mono-methyl-L-arginine was evaluated as a positive control.

NGF and cell viability assays – C6 glioma cells were used to measure NGF release into the medium. C6 cells were purchased from the Korean Cell Line Bank and maintained in DMEM supplemented with 10% FBS and 1% penicillin-streptomycin in a humidified incubator with 5% CO_2 . C6 cells were seeded into 24-well plates (1×10^5 cells/well). After 24 h, the cells were treated with DMEM containing 2% FBS and 1% penicillin-streptomycin with 20 μM of each sample for one day. The cell viability of the C6 cells was tested via an MTT assay. 6-shogaol was evaluated as a positive control.

Result and Discussion

The structures of compounds **2** - **12** were determined by comparison of their spectroscopic data with those in the literatures to be $1\beta,2\beta,4\beta$ -triol-1,2,3,4-tetrahydronaphthalene (**2**),¹¹ (7*R*,8*S*)-dihydrodehydrodiconiferyl alcohol-9-*O*- β -D-glucopyranoside (**3**),¹² kaempferol-3-*O*- β -D-glucopyranoside (**4**),¹³ nicotiflorin (**5**),¹⁴ *p*-hydroxybenzoic acid (**6**),¹⁵ β -amyrin (**7**),¹⁶ erythrodiol (**8**),¹⁷ α -spinasterol (**9**),¹⁸ 29-nor-20-oxolupeol (**10**),¹⁹ lupenone (**11**),²⁰ and lupeol (**12**).²¹

Compound **1** was obtained as a colorless gum. The molecular formula was determined to be $\text{C}_{10}\text{H}_{12}\text{O}_3$, based on molecular ion peak $[\text{M}+\text{H}]^+$ at m/z 181.0865 (calcd for $\text{C}_{10}\text{H}_{13}\text{O}_3$, 118.0865) in the positive-ion HR-ESI-MS. The IR spectrum of **1** indicated the presence of hydroxy group (3348 cm^{-1}). The ^1H NMR spectrum of **1** showed the presence of a 1,2-disubstituted aromatic ring protons at δ_{H} 7.56 (1H, d, $J = 7.3$ Hz), 7.41 (1H, dd, $J = 7.3, 1.5$ Hz), 7.34 (1H, td, $J = 7.3, 1.5$ Hz), and 7.29 (1H, td, $J = 7.3, 1.5$ Hz), three oxygenated proton signals at δ_{H} 4.89 (1H, m), 4.41 (1H, d, $J = 7.3$ Hz), and 4.09 (1H, ddd, $J = 10.1, 7.3, 3.4$ Hz) and one methylene proton signal at δ_{H} 2.21 (1H, ddd, $J = 13.5, 4.7, 3.4$ Hz), and 2.12 (1H, ddd, $J = 13.5, 10.1, 4.7$ Hz). The ^{13}C NMR spectrum displayed 10 carbon signals, including aromatic carbon signals at δ_{C} 137.6, 137.5, 128.1, 127.6, 127.4, and 127.1, three oxygenated carbon signals at δ_{C} 73.6, 68.9, and 66.6, one methylene carbon signal at δ_{C} 37.1. These NMR data were similar to $1\alpha,2\alpha,4\beta$ -triol-1,2,3,4-tetrahydronaphthalene, except for chemical shifts and coupling constants of methine protons.²² The location of hydroxyl groups was confirmed by ^1H - ^1H COSY correlations and HMBC data




Fig. 2. Key ^1H - ^1H COSY (—), HMBC (→) and NOESY (←-----→) correlations of **1**.

(Fig. 2). The relative configuration of **1** established based on coupling constants ($^3J_{1,2} = 7.3$ Hz, $^3J_{2,3} = 10.1$, 3.4 Hz, and $^3J_{3,4} = 4.7$ Hz) of ^1H -NMR spectra, leading oxygenated protons at C-1, C-2, and C-4 to be as axial, axial, and equatorial positions, respectively (Fig. 2).^{22,23} In the NOESY spectrum, correlations of H-3_{ax} with H-1 and H-4 indicated that OH groups at C-1 and C-4 were to be in the same orientation, and cross peak H-2/H-3_{eq} showed that OH groups at C-1 and C-2 were to be in the opposite orientation (Fig. 2). Thus, the structure of **1** was established as $1\beta,2\alpha,4\beta$ -triol-1,2,3,4-tetrahydronaphthalene.

The isolated compounds (**1** - **12**) were evaluated for their inhibitory effects on NO production in LPS-activated murine microglial cells and their effects on NGF secretion from C6 glioma cells. Among them, compounds **3**, **7**, and **10** reduced NO levels in LPS-activated murine microglial BV-2 cells with IC_{50} values of 26.89, 25.59, and 44.21 μM , respectively. Compounds **1**, **5**, and **9** upregulated NGF secretion to 153.09 ± 4.66 , 156.88 ± 8.86 , and $157.34 \pm 3.30\%$, respectively.

Acknowledgments

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2012R1A5A2A 28671860). We are thankful to the Korea Basic Science Institute (KBSI) for the NMR and mass spectrum data.

References

- (1) Meenu, B.; Neeraja, E. D.; Rejimon, G.; Varghese, A. *J. Chem. Pharm. Res.* **2015**, *7*, 16-21.
- (2) Ding, Z. S.; Jiang, F. S.; Chen, N. P.; Lv, G. Y.; Zhu, C. G. *molecules* **2008**, *13*, 220-229.
- (3) Kang, S. N.; Goo, Y. M.; Yang, M. R.; Ibrahim, R. I. H.; Cho, J. H.; Kim, I. S.; Lee, O. H. *molecules* **2013**, *18*, 6356-6365.
- (4) Shoji, N.; Umeyama, A.; Yosikawa, K.; Nagai, M. Arihara, S. *Phytochemistry*, **1994**, *37*, 1437-1441.
- (5) Panichayupakaranant, P.; Noguchi, H.; De-Eknamkul, W.; Sankawa, U. *Phytochemistry*, **1995**, *40*, 1141-1143.
- (6) Li, Q.; Zhang, X.; Cao, J.; Guo, Z.; Lou, Y.; Ding, M.; Zhao, Y. *Fitoterapia* **2015**, *105*, 234-239.
- (7) Ishiguro, K.; Ohira, Y.; Oku, H. *J. Nat. Prod.* **1998**, *61*, 1126-1129.
- (8) Imam, M. Z.; Nahar, N.; Akter, S.; Rana, M. S. *J. Ethnopharmacol.* **2012**, *142*, 804-810.
- (9) Kim, C. S.; Subedi, L.; Kim, S. Y.; Choi, S. U.; Choi, S. J.; Son, M. W.; Kim, K. H.; Lee, K. R. *Phytochem. lett.* **2015**, *14*, 215-220.
- (10) Kim, C. S.; Bae, M.; O, J.; Subedi, L.; Suh, W. S.; Choi, S. Z.; Son, M. W.; Kim, S. Y.; Choi, S. U.; Oh, D. C.; Lee, K. R. *J. Nat. Prod.* **2017**, *80*, 471-478.
- (11) Rashid, A.; Read, G. *J. Chem. Soc.* **1969**, *15*, 2053-2058.
- (12) Zhao, X. C.; Du, J. L.; Xie, Y. G.; Zhang, Y.; Jin, H. Z. *Chem. Nat. Compd.* **2018**, *54*, 556-558.
- (13) Jayasinghe, U. L. B.; Balasooriya, B. A. I. S.; Bandara, A. G. D.; Fujimoto, Y. *Nat. Prod. Res.* **2004**, *18*, 499-502.
- (14) Park, S. Y.; Kim, S. J.; Lee, S. Y.; Bae, K. H.; Kang, S. S. *Nat. Prod. Sci.* **2008**, *14*, 281-288.
- (15) Rho, T. W.; Yoon, K. D. *Nat. Prod. Sci.* **2017**, *23*, 253-257.
- (16) Wang, H.; Tian, X.; Chen, Y. Z. *J. Chin. Chem. Soc.* **2002**, *49*, 433-436.
- (17) Ge, Y. C.; Zhang, H. J.; Lei, J. X.; Wang, K. W. *Chem. Nat. Compd.* **2018**, *54*, 600-602.
- (18) Choi, S. Z.; Lee, S. O.; Choi, S. U.; Lee, K. R. *Arch. Pharm. Res.* **2003**, *26*, 521-525.
- (19) Kotowicz, C.; Catalan, C. A. N.; Griffin, C. L.; Herz, W. *Biochem. Syst. Ecol.* **2005**, *33*, 737-742.
- (20) Puapairoj, P.; Naengchomnong, W.; Kijjoa, A.; Pinto, M. M.; Pedro, M.; Nascimento, M. S.; Silva, A. M.; Herz, W. *Planta Med.* **2005**, *71*, 208-213.
- (21) Fotie, J.; Bohle, D. S.; Leimanis, M. L.; Georges, E.; Rukunga, G.; Nkengfack, A. E. *J. Nat. Prod.* **2006**, *69*, 62-67.
- (22) Chen, X. M.; Qian, S. H.; Feng, F. *Chine. Chem. Lett.* **2010**, *21*, 440-442.
- (23) Naemura, K.; Wakebe, T.; Hirose, K.; Tobe, Y. *Tetrahedron: Asymmetry* **1997**, *15*, 2585-2595.

Received November 27, 2018

Revised December 12, 2018

Accepted December 12, 2018