

RESEARCH ARTICLE

Open Access



Stock identification of minor carp, *Cirrhinus reba*, Hamilton 1822 through landmark-based morphometric and meristic variations

Rokhsana Ethin^{1†}, Md Shakhawate Hossain^{1,2†}, Animesh Roy¹ and Marcellin Rutegwa^{3*}

Abstract

Background: Wild fish populations stock is continuously diminishing in the Indo-Ganges river basin, and the population status of most fishes is unidentified. The identification of the population status and the conservation of commercially important and endemic wild fish populations in this region are crucial for the management. The aim of this paper was to identify the population status of *Cirrhinus reba*, a promising aquaculture but vulnerable species in the Indo-Ganges river basin in Bangladesh.

Methods: *C. reba* samples were collected from four isolated populations of the Brahmaputra ($n = 30$), the Padma (33), the Karatoya (31), and the Jamuna Rivers (30) in Bangladesh, and the population status was evaluated using morphometric and landmark comparisons. Data were analyzed with the Kruskal-Wallis test, univariate analysis, discriminant function analysis, and the formation of a dendrogram.

Results: Three meristic characters (Pectoral fin rays, caudal fin rays, scale in lateral lines), four morphometric characters (head length, pre-orbital length, post-orbital length, maximum body depth), and truss measurement (4–7) were significantly different among the stocks. The step-wise discriminant function analysis retained 15 variables from morphometric and landmark measurements that significantly differentiated the populations based on the constructed DFI and DFII. Discriminate function analysis also showed that 91.2% of the original groups were classified into their correct samples. The cluster analysis of Euclidean distances placed the Jamuna population in one cluster and the Brahmaputra, the Padma, and the Karatoya populations in the second one.

Conclusion : Morphological differences among the stock were probably due to different ancestral origin. This is the first report about population status of *C. reba* in their natural habitat of the Indian subcontinent. Further genetic studies and the evaluation of environmental impact on *C. reba* populations in Bangladesh are suggested to support our findings.

Keywords: Conservation, *Cirrhinus reba*, Population status, Threatened, Reba carp

* Correspondence: idaruteg@gmail.com

[†]Rokhsana Ethin and Md Shakhawate Hossain contributed equally to this work.

³Faculty of Education, University of Rwanda-College of Education, P.O. Box 55, Rwamagana, Eastern Province, Rwanda

Full list of author information is available at the end of the article



Background

The management and conservation strategy for different species, subspecies, stocks, and races depends on the information of biology and population structure of that particular group (Turan et al. 2006). Intraspecific variability in life history attributes in terms of growth, maturation, and development induce morphometric differences between species (Cadrin 2000). Morphometric variance among stocks of a species is considered an important marker for assessing the population dynamics and evaluation of the stocks for their management and sustainable harvest (Khan et al. 2013; Randall and Pyle 2008; Vishalakshi and Singh 2008). Intra- and interspecific biometric variation, especially in geometric terms, has long been investigated for overall patterns of racial, geographic variations (Gunawickrama 2007; Hajje et al. 2011). Equally, meristic characters have also been found to be valid in stock identification for fishery purposes (Turan 2004). In addition, landmark or truss network systems are recent powerful tools for stock separation based on morphometric and meristic characters (Cadrin 2000), which are low cost compared to molecular techniques. In a small country like Bangladesh, where there are probably very small environmental changes from place to place, the resulting morphological differences in fish may be so small that they might be impossible to discern with gross morphometric characters.

Cirrhinus reba is a commercially important indigenous minor carp species, distributed over the Indian subcontinent (Rahman 2005; Talwar and Jhingran 1991) and locally known as reba carp, bhagna bata, raik, or tatkini. In Bangladesh, the natural distributions of *C. reba* included most of the rivers, small creeks, natural depressions, and floodplains (Akhteruzzaman et al. 1998; Hussain and Mazid 2001). It is highly popular to consumers due to its oily flesh and taste qualities as well as having a reasonable amount of protein, minerals, and fatty acid (Afroz and Begum 2014; Rahman 2005). The *C. reba* are omnivores column feeder in nature with a maximum length of 40–50 cm and can attain the sexual maturity at the age of 1+ year (Akhteruzzaman et al. 1998; Ali 1997; Rahman 2005). The spawning season of *C. reba* begins in April and lasts until August and fecundity is between 200,000 and 250,000 egg kg^{-1} body weight of females (Hussain and Mazid 2001). Though induced breeding and nursing of fry are mentioned by some authors (Akhteruzzaman et al. 1998; Rahman et al. 2009; Sarkar et al. 2004) the species is not yet subjected to commercial production (Mohanta et al. 2008). During the last two decades, the populations of this species from natural water bodies have declined in the Indian subcontinent due to overfishing (Sarkar et al. 2004) and anthropogenic activities leading to siltation, aquatic pollution, and loss of natural habitat for spawning and

growth (Ahmad et al. 2013; Sarkar et al. 2004). Currently, *C. reba* is considered as a vulnerable species in Bangladesh (IUCN 2000). Therefore, it is necessary to identify the distinct wild stocks of *C. reba* for the development of good hatchery stocks through effective breeding technology, which is an important measure for mass seed production and sustainable aquaculture as well as conservation (Sarkar et al. 2004). Nevertheless, there is no known information available on the stocks of *C. reba* populations inhabiting natural waters of Bangladesh due to the scarcity of scientific works on the structural variations and no attempts have been made to assess the stock variability of different populations. The present study was therefore carried out to identify the stock structure of *C. reba* based on morphometric characters and truss network measurements in four rivers, the Padma, Jamuna, Brahmaputra, and Karatoya of Bangladesh.

Materials and methods


Fish sampling and preservation

In total, 124 individuals of *C. reba* were collected from the Brahmaputra River ($n = 30$), the Padma River ($n = 33$), the Karatoya River ($n = 31$), and the Jamuna River ($n = 30$) (Fig. 1, Table 1) with the help of local fishermen. The sampling sites were selected randomly to represent the major river systems of Bangladesh. Captured fish were wrapped into individual plastic bags and preserved and transported in an ice box to the laboratory for morphometric, meristic, and landmark measurements. Morphological identification of *C. reba* was done based on Talwar and Jhingran (1991).

Meristic and morphometric data and landmark distances

Eight meristic (Additional file 1: Table S1) and nine morphometric (Additional file 2: Figure S1) characters were recorded from the undamaged fish only immediately after arriving at the laboratory. The principle fin rays were counted manually. A magnifying glass was used to ease the counting of the fin rays. Morphometric characters were measured to an accuracy of 0.1 mm with digital calipers following the methods described by Hubbs and Lagler (2004).

For measurement of landmark distances, the truss network system described for fish body morphometric was used to construct a network on the fish body. Ten landmarks were measured (Bookstein 1997) by first placing an individual fish on a water-resistant paper and then by marking the points corresponding to the anatomical landmarks on the paper using needles (Additional file 3: Figure S2). Afterwards, the paper was removed from the fish, and the distances between perforated places on the graph paper were measured using digital calipers. Each landmark point was marked with colored pointers to enable accurate and consistent measurements.



Data analysis

The Kruskal-Wallis non-parametric test (Hoaglin et al. 2009) was used to test fish from the four localities had significantly different meristic counts. The dorsal fin rays (DFR) and the scales below the lateral line (SBLL) were

not analyzed using the Kruskal-Wallis test due to identical values in all samples. A multivariate discriminant analysis was used to identify the combination of morphometric variables that best separates the *C. reba* populations from the four rivers. Prior to the analysis, the

Table 1 Sampling details of *C. reba* collected from different sources in Bangladesh

Sample sources	Location	Number of individuals	Total length (mean ± SD)	Month of capture
Brahmaputra River	Mymensingh	30	184.01 ± 16.74	February 2016
Karatoya River	Rangpur	31	158.83 ± 26.44	April 2016
Padma River	Faridpur	33	224.11 ± 12.36	May 2016
Jamuna River	Sirajgonj	30	194.08 ± 9.23	June 2016

size effects from the dataset were eliminated through the allometric formula of Elliott et al. (1995).

$$M_{\text{adj}} = M(\text{Ls}/\text{Lo})^b$$

Where M is the original measurement, M_{adj} the size-adjusted measurement, Lo the standard length of fish, Ls the overall mean of standard length for all fish from all samples in each analysis. The parameter b was estimated for each character from the observed data as the slope of the regression of $\log M$ on $\log \text{Lo}$ using the fish from all groups. The total length was excluded from the final analyses and transformed data were checked by testing the significance of the correlation between the transformed variables and the total length.

A univariate analysis of variance (ANOVA) was carried out to test if there was any significant difference in morphological parameters among individuals of different location ($P < 0.05$) on the basis of size-adjusted morphological and landmark distance data. In addition, all size-adjusted morphometric and landmark distance data were standardized and submitted to a discriminant function analysis (DFA). A dendrogram of the populations based on the morphometric and landmark distances data were drawn by the squared Euclidean dissimilarity distance method. All statistical analyses were done using SPSS Version.16 (SPSS, Chicago, IL, USA).

Results

Meristic characters

Among the counted meristic parameters, only the pectoral fin rays (PtFR), and the scales in lateral line (SLL) were varied significantly (Kruskal-Wallis, $P < 0.05$) among the stocks (Additional file 4: Table S2).

Morphometric and landmark distances

The allometric transformation of the morphometric (8) and truss measurements (22) did not show a significant correlation with the standard length. The univariate statistics (ANOVA) showed that among the eight size-adjusted general morphometric measurements, four measurements (head length, HL; pre-orbital length, pre-OL; post-orbital length, post-OL; and maximum body depth, MBD) and only one truss measurement (4–7) were significantly ($P < 0.05$) different among the stocks (Additional file 5: Table S3).

Pooled within groups correlation between discriminant variables and DFs revealed that among the morphometric measurements, the post-OL and the HL dominantly contributed to the first DF while, the pre-OL and the pre-dorsal length (Pre-DL) contributed to the second DF and the rest contributed to the third DF (Additional file 6: Table S4). Landmark measurements, 4 to 5, and 3 to 9 contributed to the first DF, whereas 5 to 7, 5 to 8, 5 to 6, 2 to 10, 1 to 10, 4 to 8, 3 to 10, 1 to 2, and 6 to 7 contributed to the second DF and the remaining 11


landmark measurements contributed to the third DF (Additional file 7: Table S5).

Discriminant function analysis produced three types of discriminant functions (DF1, DF2, and DF3) for both morphometric and landmark measurements. For morphometric measurements, the first DF accounted for 62.7%, the second DF 34.7%, and the third DF 2.6% of the total variability among groups respectively. In sum, the first DF and second DF explained 97.45% of the total variability among groups. For landmark measurements, the first DF accounted for 78.8%, the second DF 17.9%, and the third DF 3.3% of the total variability among the groups respectively. In sum, the first DF and second DF explained 96.33% of the total variability among groups. In canonical discriminant function, the Jamuna population sample formed a distinct group while the samples from the Padma, Brahmaputra, and Korotuya were clustered around their centroid value while overlapping to a certain degree with each other (Fig. 2). The discriminant function analysis also showed that the overall allotment of individuals into their original populations was high (91.2%). All of the fishes (100%) of Jamuna River were correctly classified into their own group. While 95, 92.9, and 80% of fishes were correctly classified for the Brahmaputra, Padma, and Karatoya River samples, respectively. On the other hand, the overlapping of samples from the Karatoya (20%) and the Padma (7.1%) rivers implied some misclassification (Table 2).

The dendrogram derived from the cluster analysis of Euclidean distances among groups of centroids showed that the four populations of *C. reba* segregated from each other into two distinct clusters (Fig. 3). The Padma populations formed one cluster with the Brahmaputra and the Karatoya populations. However, the Brahmaputra and the Karatoya populations formed another sub-cluster. Finally, the Jamuna River population formed a completely separate third group.

Discussion

This study demonstrates that meristic attributes of *C. reba* including the number of pectoral fin rays (PtFR), caudal fin rays (CFR), and scales in the lateral line (SLL) varied significantly within the studied populations. This indicates that these populations underwent some extent morphological changes over the course of the time due to geographical isolation or might be of from different of ancestors' origin. They may also be resulting from different environmental and habitat conditions variations. Since, fish shows higher plasticity in morphological traits to environmental changes (Wimberger 1992; Allendorf 1988; Stearns 1983; Swain et al. 1991). The variability of meristic counts of *C. reba* have also been reported in the findings of Hamilton (1822), Rahman (2005), Talwar and Jhingran (1991) and Galib (2008), while, Shafi and



Quddus (2001) found no difference in meristic counts among different populations. The later researchers may have sampled populations that were experiencing similar environmental conditions or had a common ancestor and therefore were not undergo differentiation despite being isolated geographically. Similarly, variable meristic counts have also been observed in other fish species of isolated stocks, such as *Labeo calbasu*, Mesopotamian spiny eel, (Çakmak and Alp 2010); *Heteropneustes fossilis*, (Rahman et al. 2014) and *Rhinomugil corsula*, (Lipták et al. 2019).

In addition to meristic characters, a truss network system is another important criterion to discern species, subspecies, races, and strains and have been employed by different authors (Khan et al. 2013; Mir et al. 2013; Sajina et al. 2011; Siddik et al. 2016; Ujjania and Kohli 2011). These characters can be also modified by environmental fluctuations during early life-history stages of fish (Cheverud 1988; Reyment 1985). In the present study,

we did not find any significant difference among the measured landmarked distance except one (4–7). Size-adjusted morphometric and landmark data showed phenotypic divergence among *C. reba* samples. The discriminant function analysis and dendrogram revealed the existence of two distinct morphologically differentiated stocks. The Jamuna River population itself is a unique stock, highly distinct from the other three stocks, and formed the first group due its geographical location. The Padma, the Brahmaputra, and the Karatoya River populations formed the second group. The stock from the Brahmaputra and Karatoya River sites had similarity in all examined parameters in addition to overlapping between the two populations. This confirmed their phenotypic similarity. Both the rivers are the distributary and tributary of the Jamuna River and point of origin not geographically isolated (Fig. 1). Fish could migrate from Jamuna River to these two rivers. This similarity indicates that the *C. reba* populations in these two rivers might be are of the same ancestral origin. On the other hand, the Jamuna stock highly deviated from the Brahmaputra and the Karatoya as well as from the Padma River samples. The sampling sites of Jamuna and Padma River are geographically far from the sampling sites of Karatoya and Brahmaputra River (see Fig. 1). These might make the Jamuna population unique from others. In accordance to geographical distance and sampling site position (downstream from the Jamuna sampling site) the Padma population could be connected with the Jamuna population. But, the Padma population was

Table 2 Percentage of individuals correctly classified into their original populations using discriminate function analysis

Population	Predicted group membership ¹				Total
	Brahmaputra	Karatoya	Padma	Jamuna	
% Brahmaputra	95.0	0.0	5.0	0.0	100.0
Karatoya	0.0	80.0	20.0	0.0	100.0
Padma	0.0	7.1	92.9	0.0	100.0
Jamuna	0.0	0.0	0.0	100.0	100.0

¹91.2% of original grouped cases correctly classified

Additional file 7: Table S5. Pooled within-groups correlations between discriminating variables and standardized canonical discriminant functions based on truss measurement (variables ordered by absolute size of correlation within function, *denotes the largest absolute correlation between each variable and any discriminant functions). (DOCX 16 kb)

Acknowledgements

Authors would like to thank the funding agencies that supported financially this study and the local fishermen who helped in sample collection.

Consent of publication

Not applicable

Authors' contributions

MSH designed the experiment, RE and MSH conducted the experiment, RE and AR collected the samples. RE and MSH drafted the manuscript. MR helped in data analysis and final revision. All authors read and approved the final manuscript.

Funding

The Ministry of Education, Youth, financially supported the study and Sports of the Czech Republic-project "CENAKVA" (LM2018099) and by the Grant Agency of the University of South Bohemia (no. 017/2016/Z).

Availability of data and materials

The datasets and/or analyzed during current study available from the corresponding author on reasonable request.

Ethics approval and consent to participate

No specific permissions were required for the locations and activities involved in this study. All experimental manipulations (capture and measurements) were conducted according to the principles of the Ethical Committee of Bangabandhu Sheikh Mujibur Rahman Agricultural University, Bangladesh, and National Research Ethics Committee (NREC) of Bangladesh. This study also follows the rules for the Protection of Animals in Research of the University of South Bohemia, Faculty of Fisheries and Protection of Waters, Research Institute of Fish Culture and Hydrobiology, Vodňany, based on the EU harmonized animal welfare act of Czech Republic. The principles of laboratory animal care and the national laws 246/1992 and regulations on animal welfare were followed (ref. number 22761/2009-17210).

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Fisheries Biology and Aquatic Environment, Bangabandhu Sheikh Mujibur Rahman Agricultural University, Gazipur 1706, Bangladesh.

²University of South Bohemia in České Budejovice, Faculty of Fisheries and Protection of Waters, South Bohemian Research Center of Aquaculture and Biodiversity of Hydrocenoses, Zátěží 728/II, 389 25, Vodňany, Czech Republic.

³Faculty of Education, University of Rwanda-College of Education, P.O. Box 55, Rwamagana, Eastern Province, Rwanda.

Received: 25 March 2019 Accepted: 13 June 2019

Published online: 21 June 2019

References

- Afroz H, Begum M. Analysis of nutritional value and mineral contents of three different parts of body of *Cirrhinus reba*. *Int J Sci Eng Res*. 2014;5:2301–6.
- Ahmad S, Muralidharan M, Venkateshwarlu M, Arunachalam M. Distribution pattern, endemism, threat status and conservation measures of fishes in the Tunga and Bhadra rivers of Western Ghats, India. *Environ Biol Fish*. 2013; 96(10-11):1245–56.
- Akhteruzzaman M, Kohinoor AHM, Islam MS, Modak PC. Observations on the induced breeding of indigenous small fish Bhagna (*Cirrhinus reba* Ham) in Bangladesh. *Progress Agric*. 1998;9(1-2):281–4.
- Ali MY. Small indigenous fish species culture in Bangladesh. Dhaka: Paper presented at the Proceedings of national workshop on small indigenous fish culture in Bangladesh, IFADEP-SP2; 1997.
- Allendorf FW. Conservation biology of fishes. *Conserv Biol*. 1988;2(2):145–8.
- Bookstein FL. *Morphometric tools for landmark data: geometry and biology*. Cambridge: Cambridge University Press; 1997.
- Cadrin SX. Advances in morphometric identification of fishery stocks. *Rev Fish Biol Fish*. 2000;10(1):91–112.
- Çakmak, Esen, & Alp, Ahmet. Morphological Differences Among the Mesopotamian Spiny Eel, *Mastacembelus mastacembelus* (Banks & Solander 1794), Populations. *Turkish J Fish Aquatic Sci*. 2010;10:87–92.
- Cheverud JM. A comparison of genetic and phenotypic correlations. *Evolution*. 1988;42(5):958–68.
- Elliott NG, Haskard K, Koslow JA. Morphometric analysis of orange roughy (*Hoplostethus atlanticus*) off the continental slope of southern Australia. *J Fish Biol*. 1995;46(2):202–20.
- Galib SM. A study on fish diversity and fishing gears of Chalan beel with reference to preservation of catches. Bangladesh: Honors dissertation submitted to the Department of Fisheries, University of Rajshahi; 2008.
- Gunawickrama KBS. Morphological heterogeneity and population differentiation in the green chromid *Ectopplus suratensis* (Pisces: Cichlidae) in Sri Lanka. *Ruhuna J Sci*. 2007;2:70–81.
- Hajjej G, Hattour A, Hajjej A, Allaya H, Jarboui A, Bouain A. Biometry, length-length and length-weight relationships of little tuna *Euthynnus alletteratus* in the Tunisian waters. *J Fish Aquatic Sci*. 2011;6(3):256.
- Hamilton F. An account of the fishes found in the river Ganges and its branches: Printed for A. Constable and company; 1822.
- Haque MA, Jewel MAS, Sultana MP. Assessment of physicochemical and bacteriological parameters in surface water of Padma River, Bangladesh. *Appl Water Sci*. 2019;9(1):10.
- Hoaglin DC, Mosteller F, Tukey JW. *Fundamentals of exploratory analysis of variance*, vol. 367: Wiley; 2009.
- Hubbs CL, Lagler KF. *Fishes of the Great Lakes region* (revised ed.): The University of Michigan Press; 2004.
- Hussain MG, Mazid MA. Genetic improvement and conservation of carp species in Bangladesh: Research Institute and International Center for Living Aquatic Resources Management, Bangladesh; 2001.
- Islam MS, Ferdous J, Nasmi S, Kabir MH. Seasonal variations in water quality and its effects on fish production in the Brahmaputra River at Mymensingh of Bangladesh. *Bang J Environ Sci*. 2015;29:18–25.
- Islam MS, Kabir MH, Sifat SA, Meghla NT, Tushar TR. Status of water quality from the Padma river at Bheramara point of Kushtia in Bangladesh. *Bang J Environ Sci*. 2014;27:110–5.
- IUCN, Bangladesh. *Red Book of Threatened Fishes of Bangladesh: IUCN-The world conservation union*; 2000. p. 116.
- Keivany Y, Mousavi SMA, Dorafshan S, Zamani-Faradonbe M. Morphological variations of *Alburnus mossulensis* Heckel, 1843 populations in the Tigris tributaries of the Persian Gulf basin in Iran (Teleostei: Cyprinidae). *Iran J Ichthyol*. 2016;3(3):190–202.
- Khan MA, Miyan K, Khan S. Morphometric variation of snakehead fish, *Channa punctatus*, populations from three Indian rivers. *J Appl Ichthyol*. 2013;29(3):637–42.
- Lipták B, Veselý L, Ercoli F, Bláha M, Buřič M, Ruokonen TJ, Kouba A. Trophic role of marbled crayfish in a lentic freshwater ecosystem. *Aquat Invas*. 2019; Accepted.
- Mir JI, Sarkar UK, Dwivedi AK, Gusain OP, Jena JK. Stock structure analysis of *Labeo rohita* (Hamilton, 1822) across the Ganga basin (India) using a truss network system. *J Appl Ichthyol*. 2013;29(5):1097–103.
- Mohanta KN, Subramanian S, Komarpant N, Saurabh S. Alternate carp species for diversification in freshwater aquaculture in India. *Aquaculture Asia*. 2008; 13(1):11.
- Moore GA. The cutaneous sense organs of barbeled minnows adapted to life in the muddy waters of the Great Plains region. *Transac Am Microscop Soc*. 1950;69(1):69–95.
- Murta AG. Morphological variation of horse mackerel (*Trachurus trachurus*) in the Iberian and North African Atlantic: implications for stock identification. *ICES J Marine Sci*. 2000;57(4):1240–8.
- Rahman AKA. *Freshwater fishes of Bangladesh* (second ed.). Dacca: Zoological Society of Bangladesh; 2005.
- Rahman MA, Zaher M, Azimuddin KM. Development of fingerling production techniques in nursery ponds for the critically endangered reba carp, *Cirrhinus ariza* (Hamilton, 1807). *Turkish J Fish Aquatic Sci*. 2009;9(2):165–72.
- Rahman MM, Sharker MR, Sumi KR, Alam MA, Hossen MS. Landmark-based morphometric and meristic variations of stinging catfish, *Heteropneustes fossilis* (Bloch) among three isolated stocks, the Old Brahmaputra river and the Tangaur haor and a hatchery. *Int J Fish Aquatic Stud*. 2014;1(3):163–70.

- Randall JE, Pyle RL. *Synodus orientalis*, a new lizardfish (Aulopiformes: Synodontidae) from Taiwan and Japan, with correction of the Asian record of *S. lobeli*. *Zool Stud.* 2008;47(5):657–62.
- Rao RJ. Biological resources of the Ganga river, India. *Hydrobiologia.* 2001;458(1): 159–68.
- Reyment RA. Multivariate morphometrics and analysis of shape. *J Int Assoc Math Geol.* 1985;17(6):591–609.
- Sajina AM, Chakraborty SK, Jaiswar AK, Pazhayamadam DG, Sudheesan D. Stock structure analysis of *Megalaspis cordyla* (Linnaeus, 1758) along the Indian coast based on truss network analysis. *Fish Res.* 2011;108(1):100–5.
- Sarkar UK, Negi RS, Deepak PK, Singh SP, Srivastava SM, Roy D. Captive breeding of vulnerable Indian carp *Cirrhinus reba* with Ovaprim for conservation of wild populations. *Aquaculture Asia.* 2004;9:5–7.
- Shafi M, Quddus MMA. *Bangladesher Matsho Shampad* (Fisheries of Bangladesh)(in Bengali), Kabir publication. Dhaka; 2001. p. 153–4.
- Siddik M, Chaklader M, Hanif M, Islam M, Sharker M, Rahman M. Stock identification of critically endangered olive barb, *Puntius sarana* (Hamilton, 1822) with emphasis on management implications. *J Aquacult Res Dev.* 2016;7(2):411.
- Stearns SC. A Natural Experiment in Life-History Evolution: Field Data on the Introduction of Mosquitofish (*Gambusia affinis*) to Hawaii. *Evolution.* 1983: 601–17.
- Sun P, Hawkins W, Overstreet R, Brown-Peterson N. Morphological deformities as biomarkers in fish from contaminated rivers in Taiwan. *Int J Environ Res Public Health.* 2009;6(8):2307–31.
- Swain DP, Riddell BE, Murray CB. Morphological differences between hatchery and wild populations of coho salmon (*Oncorhynchus kisutch*): environmental versus genetic origin. *Can J Fish Aquatic Sci.* 1991;48(9):1783–91.
- Talwar PK, Jhingran AG. *Inland fisheries of India and adjacent countries* (Vol. 2). New Delhi: Oxford and IBH Publishing; 1991.
- Turan C. Stock identification of Mediterranean horse mackerel (*Trachurus mediterraneus*) using morphometric and meristic characters. *ICES J Marine Sci.* 2004;61(5):774–81.
- Turan C, Oral M, Öztürk B, Düzgüneş E. Morphometric and meristic variation between stocks of Bluefish (*Pomatomus saltatrix*) in the Black, Marmara, Aegean and northeastern Mediterranean Seas. *Fish Res.* 2006;79(1):139–47.
- Uddin MN, Alam MS, Mobin MN, Miah MA. An Assessment of the River water quality parameters: A case of Jamuna River. *J Environ Sci Nat Res.* 2014;7(1): 249–56.
- Ujjania NC, Kohli MPS. Landmark-based morphometric analysis for selected species of Indian major carp (*Catla catla*, Ham. 1822). *Int J Food Agric Vet Sci.* 2011;1:64–74.
- Vishalakshi C, Singh BN. Differences in Morphological Traits between Two Sibling Species, *Drosophila ananassae* and *D. pallidosa*. *Zool Stud.* 2008;47(3):352–9.
- Wimberger PH. Plasticity of fish body shape. The effects of diet, development, family and age in two species of *Geophagus* (Pisces: Cichlidae). *Biol J Linnean Soc.* 1992;45(3):197–218.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

