



## Corrigendum: Dental alloplastic bone substitutes currently available in Korea

Jeong-Kui Ku<sup>1,2</sup>, Inseok Hong<sup>3</sup>, Bu-Kyu Lee<sup>1</sup>, Pil-Young Yun<sup>4</sup>, Jeong Keun Lee<sup>5</sup>

<sup>1</sup>*Department of Oral and Maxillofacial Surgery, Asan Medical Center, Seoul,*

<sup>2</sup>*Department of Oral and Maxillofacial Surgery, Section of Dentistry, Armed Forces Capital Hospital, Seongnam,*

<sup>3</sup>*Department of Oral and Maxillofacial Surgery, School of Dentistry and Institute of Oral Bioscience,  
Research Institute of Clinical Medicine of Chonbuk National University-Biomedical Research Institute of  
Chonbuk National University Hospital, Chonbuk National University, Jeonju,*

<sup>4</sup>*Department of Oral and Maxillofacial Surgery, Section of Dentistry, Seoul National University Bundang Hospital, Seongnam,*

<sup>5</sup>*Department of Oral and Maxillofacial Surgery, Institute of Oral Health Science, Ajou University School of Medicine, Suwon, Korea*

J Korean Assoc Oral Maxillofac Surg 2019;45:51-67.

<https://doi.org/10.5125/jkaoms.2019.45.2.51>

This correction is being published to correct the 4th author's affiliation in above article.  
The authors apologize for this error.

Before correction

Jeong-Kui Ku<sup>1,2</sup>, Inseok Hong<sup>3</sup>, Bu-Kyu Lee<sup>1</sup>, Pil-Young Yun<sup>2</sup>, Jeong Keun Lee<sup>4</sup>

<sup>1</sup>*Department of Oral and Maxillofacial Surgery, Asan Medical Center, Seoul,*

<sup>2</sup>*Department of Oral and Maxillofacial Surgery, Section of Dentistry, Armed Forces Capital Hospital, Seongnam,*

<sup>3</sup>*Department of Oral and Maxillofacial Surgery, School of Dentistry and Institute of Oral Bioscience, Research Institute of Clinical Medicine of  
Chonbuk National University-Biomedical Research Institute of Chonbuk National University Hospital, Chonbuk National University, Jeonju,*

<sup>4</sup>*Department of Oral and Maxillofacial Surgery, Institute of Oral Health Science, Ajou University School of Medicine, Suwon, Korea*

After correction

Jeong-Kui Ku<sup>1,2</sup>, Inseok Hong<sup>3</sup>, Bu-Kyu Lee<sup>1</sup>, Pil-Young Yun<sup>4</sup>, Jeong Keun Lee<sup>5</sup>

<sup>1</sup>*Department of Oral and Maxillofacial Surgery, Asan Medical Center, Seoul,*

<sup>2</sup>*Department of Oral and Maxillofacial Surgery, Section of Dentistry, Armed Forces Capital Hospital, Seongnam,*

<sup>3</sup>*Department of Oral and Maxillofacial Surgery, School of Dentistry and Institute of Oral Bioscience, Research Institute of Clinical Medicine of  
Chonbuk National University-Biomedical Research Institute of Chonbuk National University Hospital, Chonbuk National University, Jeonju,*

<sup>4</sup>*Department of Oral and Maxillofacial Surgery, Section of Dentistry, Seoul National University Bundang Hospital, Seongnam.*

<sup>5</sup>*Department of Oral and Maxillofacial Surgery, Institute of Oral Health Science, Ajou University School of Medicine, Suwon, Korea*