



Simple universal cutoff point of waist-height ratio for metabolic risk in Korean children and adolescents

Hae Soon Kim, MD, PhD

Department of Pediatrics, Ewha Womans University College of Medicine, Seoul, Korea

The prevalence of obesity among children and adolescents has been increasing over 30 years, worldwide.¹⁾ Pediatric obesity is closely related to metabolic syndrome during childhood and adolescence and adult obesity and cardiovascular risks. Recently, many studies from developed countries have reported that childhood obesity prevalence rates stabilized around the late 1990s and early 2000s. A similar trend was noted in Korea where the prevalence of childhood overweight and obesity stabilized from 2001 to 2012 for both boys and girls.²⁾ Similarly, Kim et al.³⁾ reported no significant secular change in waist circumference (WC) and the waist-height ratio (WHtR) over 10 years (2005–2015) in Korean children and adolescents. Nevertheless, obesity-related metabolic disorders, such as hyperlipidemia, type 2 diabetes mellitus, hypertension, and nonalcoholic fatty liver disease are recently emerging medical problems among children and adolescents.

It is somewhat difficult for parents to identify their children's obesity, because normal weight children have less subcutaneous fat and lower normal body mass index (BMI) than do adults. However, early detection of obesity is the most important to prevent adult obesity and obesity-related cardiovascular risks in adulthood, because managing childhood obesity is challenging.

Several anthropometric measurements have been developed to define obesity. BMI is mainly used as obesity index because it is simply calculated using body weight and height; however, it reflects not only adiposity but also muscle mass. WC is widely used as an abdominal obesity index for adult and children.⁴⁾ Both are considered useful obesity indices to identify obesity-related cardiometabolic risks. However, the limitations of these 2 indices are ethnicity and sex-specificity in adults. For example, compared with men in western countries, a lower cutoff of BMI and WC are used for women in the Asian population. Furthermore, age and sex-specific percentiles should assist in diagnosis of obesity and abdominal obesity for children and adolescents. In a pediatric clinical setting, the pediatrician should use the BMI and WC percentiles for age and sex-specific tables for diagnosing pediatric obesity so that the patient's indices can be compared to age and sex related cutoff points. Another limitation of WC as an abdominal obesity index is its association with height. Taller subjects have higher WC regardless of their lower cardiovascular risk.

Studies on alternative measurements for abdominal obesity have suggested that compared with BMI and WC, the WHtR calculated by dividing the WC by the height, has advantages in identifying cardiometabolic risk, thus downscaling the unfavorable effect of WC on cardiovascular risk.⁴⁻⁶⁾

According to systemic reviews among adults, a universal WHtR cutoff point of 0.5 is widely accepted as a screening tool for metabolic syndrome risks independent of sex and ethnicity in adults and even in children and adolescents.⁷⁾

Contrary to results from adults, inconsistent results on cutoff points for the associations between the WHtR with cardiovascular disease risk factors for children have been obtained.

Corresponding author: Hae Soon Kim, MD, PhD
 Department of Pediatrics, Ewha Womans University Mokdong Hospital, 1071 Anyangcheon-ro, Yangcheon-gu, Seoul 07985, Korea
 Tel: +82-2-2650-5569
 Fax: +82-2-2653-3718
 E-mail: hyesk@ewha.ac.kr
<http://orcid.org/0000-0002-6976-6878>

Received: 10 January, 2019
 Revised: 4 March, 2019
 Accepted: 28 March, 2019

Copyright © 2019 by The Korean Pediatric Society

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.




Fig. 1. Waist-height ratio percentile (5th, 50th, 90th, and 95th) by sex and age. The closed line denotes boys and the dotted line denotes girls.³⁾

Studies have suggested that a WHtR cutoff point of 0.5 can be applied independent of sex and age in children. Studies from many countries validated that a universal WHtR \geq 0.5 is a better predictor of metabolic disorder risk in adolescents, independent of sex or age.⁸⁾ Another study suggested showed that compared with BMI z score and WC, the WHtR was the worst at predicting metabolic risk in adolescents.⁹⁾ Kim et al.³⁾ reported a WHtR cutoff point of 0.48 for abdominal obesity with metabolic disorder in 13- to 18-year-old Korean adolescents based on the Korea National Health and Nutrition Examination Survey (2005–2015) data. They showed that the WHtR is a universal indicator independent of age and sex in children and adolescents and a cutoff point of 0.48 is near the 90th percentile of the WHtR for both sexes (Fig. 1). According to these results, the WHtR may be a universal, simple, and sensitive indicator to assess obesity-related metabolic risks and is a practical and cost-effective option for children and adolescents, even though ethnic differences in the WHtR have been noted.

In conclusion, the WHtR is a simple and practical indicator with a cutoff point that can be used as a screening tool for predicting metabolic disorder risk independent of sex, age, height, and ethnicity in children and adolescents. A universal cutoff point suggests the following proposal for a practical primary screening tool in children

and adolescents:¹⁰⁾ ‘Maintain your WC at less than half of your height.’

Conflicts of interest

No potential conflict of interest relevant to this article was reported.

References

1. Swinburn BA, Sacks G, Hall KD, McPherson K, Finegood DT, Moodie ML, et al. The global obesity pandemic: shaped by global drivers and local environments. *Lancet* 2011;378:804–14.
2. Bahk J, Khang YH. Trends in measures of childhood obesity in Korea from 1998 to 2012. *J Epidemiol* 2016;26:199–207.
3. Kim MS, Kim SY, Kim JH. Secular change in waist circumference and waist–height ratio and optimal cutoff of waist–height ratio for abdominal obesity among Korean children and adolescents. *Korean J Pediatr* 2018 Dec 3 [Epub]. <https://doi.org/10.3345/kjp.2018.07038>.
4. Lee CM, Huxley RR, Wildman RP, Woodward M. Indices of abdominal obesity are better discriminators of cardiovascular risk factors than BMI: a meta-analysis. *J Clin Epidemiol* 2008;61:646–53.
5. Ashwell M, Hsieh SD. Six reasons why the waist-to-height ratio is a rapid and effective global indicator for health risks of obesity and how its use could simplify the international public health message on obesity. *Int J Food Sci Nutr* 2005;56:303–7.
6. Freedman DS, Kahn HS, Mei Z, Grummer-Strawn LM, Dietz WH, Srinivasan SR, et al. Relation of body mass index and waist-to-height ratio to cardiovascular disease risk factors in children and adolescents: the Bogalusa Heart Study. *Am J Clin Nutr* 2007;86:33–40.
7. Browning LM, Hsieh SD, Ashwell M. A systematic review of waist-to-height ratio as a screening tool for the prediction of cardiovascular disease and diabetes: 0.5 could be a suitable global boundary value. *Nutr Res Rev* 2010;23:247–69.
8. Chung IH, Park S, Park MJ, Yoo EG. Waist-to-height ratio as an index for cardiometabolic risk in adolescents: results from the 1998–2008 KNHANES. *Yonsei Med J* 2016;57:658–63.
9. Gomes TN, Nevill A, Katzmarzyk PT, Pereira S, Dos Santos MM, Buranarugsa R, et al. Identifying the best body-weight-status index associated with metabolic risk in youth. *Scand J Med Sci Sports* 2018; 28:2375–83.
10. Ashwell M, Gibson S. A proposal for a primary screening tool: ‘Keep your waist circumference to less than half your height’. *BMC Med* 2014;12:207.