

¹H-NMR Analysis of Metabolic Changes Induced by Snf1/AMP-Activated Protein Kinase During Environmental Stress Responses

Jiyoung Kim^{a*} , Junsang Oh^{b*} , Deok-Hyo Yoon^b  and Gi-Ho Sung^{b,c} 

^aJeonju AgroBio-Materials Institute, Jeonju-si, Korea; ^bTranslational Research Division, Biomedical Institute of Mycological Resource, International St. Mary's Hospital and College of Medicine, Catholic Kwandong University, Incheon, Korea; ^cDepartment of Microbiology, College of Medicine, Catholic Kwandong University, Gangneung-si, Korea

ABSTRACT

AMP-activated protein kinase sucrose non-fermenting 1 (Snf1) is a representative regulator of energy status that maintains cellular energy homeostasis. In addition, Snf1 is involved in the mediation of environmental stress such as salt stress. Snf1 regulates metabolic enzymes such as acetyl-CoA carboxylase, indicating a possible role for Snf1 in metabolic regulation. In this article, we performed nuclear magnetic resonance (NMR) spectroscopy to profile the metabolic changes induced by Snf1 under environmental stress. According to our NMR data, we suggest that Snf1 plays a role in regulating cellular concentrations of a variety of metabolites during environmental stress responses.

ARTICLE HISTORY

Received 10 September 2018
Revised 15 May 2019
Accepted 28 May 2019

KEYWORDS

Snf1/AMPK; stress signals; nuclear magnetic resonance; metabolic changes; yeast

Since yeast mainly uses glucose as a carbon source, sensing of glucose levels is important for mediating yeast energy metabolism. Sucrose non-fermenting 1 (Snf1)/AMP-activated protein kinase is a representative controller of energy status that maintains cellular energy homeostasis [1–3]. Snf1 is a regulatory kinase that is highly conserved in eukaryotic cells and plays a critical role in regulating a variety of activators and repressors required for energy balance mechanism [1–3]. In addition, Snf1 is involved in the mediation of environmental stimuli such as salt stress [4–11].

Snf1 phosphorylates and regulates metabolic enzymes such as acetyl-CoA carboxylase, suggesting a crucial role for Snf1 in metabolic control [12]. To date, there have not been any reports describing the metabolic changes induced by Snf1 during environmental stress responses. Nuclear magnetic resonance (NMR) spectroscopy is a useful technique for structure elucidation due to its various two-dimensional measurements, which makes NMR an ideal tool for metabolic analysis [13,14]. Here, we report that Snf1 regulates metabolic changes in response to environmental stresses in *Saccharomyces cerevisiae*.


We investigated whether Snf1 is involved in responses to multiple stresses such as salt, heat, and energy (potassium cyanide (KCN), a specific inhibitor of cytochrome c oxidase for ATP synthesis).

Consistent with previous reports [9,11], the $\Delta snf1$ mutant showed increased sensitivity to salt and heat stresses (Supplementary Figure S1). In addition, our result revealed that Snf1 is required for a proper response against KCN stress (Supplementary Figure S1). Snf1 functions as a key sensor of energy status for the maintenance of cellular energy homeostasis [2,3,9]. Therefore, our work indicates that Snf1 might play a role in energy regulation as a protective mechanism under various stress conditions.

To explore changes in yeast stress responses from a metabolic perspective, we profiled the metabolites in yeast cells using ¹H-NMR spectroscopy (Figure 1 and Supplementary Table S1). ¹H-NMR spectroscopy and raw data processing were essentially performed as described previously [15]. Chemical shifts of signals were assigned to the metabolites in the area of amino acids, organic acids, carbohydrates, and nucleotide derivatives. On the basis of the ¹H-NMR spectra, we identified 36 metabolites in the whole cell extracts with chemical shifts and coupling patterns (Figure 1 and Supplementary Table S1). The main variations are summarized in the form of a heatmap shown in Figure 2(A) and Supplementary Table S2. ¹H-NMR spectra of wild-type and $\Delta snf1$ exhibited differentiated resonance spectra under various stress conditions. Partial least squares-discriminant analysis (PLS-DA), a supervised

CONTACT Gi-Ho Sung  sung97330@gmail.com

*These authors contributed equally to this work.

 Supplemental data for this article can be accessed [here](#).

© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group on behalf of the Korean Society of Mycology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. $^1\text{H-NMR}$ spectra of the metabolites of *S. cerevisiae*. $^1\text{H-NMR}$ spectroscopy and raw data processing were essentially performed as described previously [15].




Figure 2. Heatmap of main metabolite variations (A) and PLS-DA score plot (B) in wild-type and $\Delta snf1$ under various stress conditions. The *S. cerevisiae* wild-type (W303-1A) and $\Delta snf1$ strains were grown on YPD medium. Yeast cells were treated with 0.8 M NaCl, 5 mM KCN, or 40 °C for 1 h, and then subjected to $^1\text{H-NMR}$. Similar results were obtained from three independent experiments. $^1\text{H-NMR}$ spectroscopy and raw data processing were essentially performed as described previously [15].




Figure 3. Snf1 regulates cellular concentrations of trehalose and glucose under salt stress. Yeast cells were treated with 0.8 M NaCl for 1 h, and then subjected to $^1\text{H-NMR}$. Cellular concentrations of glucose and trehalose are differentially regulated in wild-type and $\Delta snf1$ under salt stress. Experimental values are the means of three independent experiments with standard deviation.

multivariate data analysis method, was used to investigate intrinsic variation in $^1\text{H-NMR}$ data. In the wild-type PLS-DA score plots, two principal components, PC1 and PC2, were calculated with the R^2Y and Q^2Y parameters of 0.95 and 0.77, and the $\Delta snf1$ mutant PLS-DA score plots were calculated with the R^2Y and Q^2Y parameters of 0.93 and 0.63, respectively (Figure 2(B)). As shown in Figure 2(B), the PLS-DA score plot of $^1\text{H-NMR}$ spectra showed a clear separation between wild-type and $\Delta snf1$ under various stress conditions.

Our $^1\text{H-NMR}$ data suggest that Snf1 plays a role in the regulation of metabolic changes induced by stressful environments in yeast (Figure 2). Especially, cellular concentrations of glucose and trehalose are increased in wild-type under salt stress, but not in $\Delta snf1$ (Figure 3). Trehalose functions as a representative osmolyte in order to cope with changes in osmotic pressure [16]. Meanwhile, maintenance of ATP balance is vital for all cells and the hydrolysis of ATP is the main energy source. In response to salt stress, a rapid increase in cellular ATP metabolism may reflect the higher energy demands required for salt stress tolerance [17,18]. Under normal condition, glucose production was slightly reduced in $\Delta snf1$ compared to wild-type (WT). In response to salt stress, glucose production was increased in WT, however it was not increased in $\Delta snf1$ at all (Figure 3). At present, we do not know whether Snf1 is involved in the mechanism of directly increasing glucose production in response to salt stress. Nevertheless, it can be assumed that Snf1 is related to glucose production under salt stress response. Our $^1\text{H-NMR}$ data will make a

contribution to our understanding of metabolic changes induced by Snf1 during environmental stress in yeast.

Acknowledgments

We thank Dr. Enrique Herrero for providing us with the $\Delta snf1$ yeast mutants.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

This research was supported by Bio-industry Technology Development Program [316025-05] of IPET (Korea Institute of Planning and Evaluation for Technology in Food, Agriculture, Forestry, and Fisheries) and the National Research Foundation (NRF) grant funded by the Korea government (MSIT) [No. 2019R1A2C2005157].

ORCID

Jiyoung Kim <http://orcid.org/0000-0002-5258-6889>
 Deok-Hyo Yoon <http://orcid.org/0000-0003-4422-3532>
 Gi-Ho Sung <http://orcid.org/0000-0002-1861-5543>

References

- [1] Hardie DG, Carling D, Carlson M. The AMP-activated/SNF1 protein kinase subfamily: metabolic sensors of the eukaryotic cell? *Annu Rev Biochem.* 1998;67:821–855.
- [2] Hardie DG. AMP-activated/SNF1 protein kinases: conserved guardians of cellular energy. *Nat Rev Mol Cell Biol.* 2007;8:774–785.
- [3] Kayikci Ö, Nielsen J. Glucose repression in *Saccharomyces cerevisiae*. *FEMS Yeast Res.* 2015; 15:fov068.
- [4] Thompson-Jaeger S, Francois J, Gaughran JP, et al. Deletion of *SNF1* affects the nutrient response of yeast and resembles mutations which activate the adenylate cyclase pathway. *Genetics.* 1991;129: 697–706.
- [5] Alepuz PM, Cunningham KW, Estruch F. Glucose repression affects ion homeostasis in yeast through the regulation of the stress-activated *ENA1* gene. *Mol Microbiol.* 1997;26:91–98.
- [6] McCartney RR, Schmidt MC. Regulation of Snf1 kinase. Activation requires phosphorylation of threonine 210 by an upstream kinase as well as a distinct step mediated by the Snf4 subunit. *J Biol Chem.* 2001;276:36460–36466.
- [7] Dubacq C, Chevalier A, Mann C. The protein kinase Snf1 is required for tolerance to the ribonucleotide reductase inhibitor hydroxyurea. *Mol Cell Biol.* 2004;24:2560–2572.
- [8] Portillo F, Mulet JM, Serrano R. A role for the non-phosphorylated form of yeast Snf1: tolerance

- to toxic cations and activation of potassium transport. FEBS Lett. 2005;579:512–516.
- [9] Hong SP, Carlson M. Regulation of *snf1* protein kinase in response to environmental stress. J Biol Chem. 2007;282:16838–16845.
- [10] Hedbacker K, Carlson M. SNF1/AMPK pathways in yeast. Front Biosci. 2008;13:2408–2420.
- [11] Pastor MM, Proft M, Pascual-Ahuir A. Mitochondrial function is an inducible determinant of osmotic stress adaptation in yeast. J Biol Chem. 2009;284:30307–30317.
- [12] Woods A, Munday MR, Scott J, et al. Yeast SNF1 is functionally related to mammalian AMP-activated protein kinase and regulates acetyl-CoA carboxylase *in vivo*. J Biol Chem. 1994;269:19509–19515.
- [13] Beckonert O, Keun HC, Ebbels TM, et al. Metabolic profiling, metabolomic and metabolomic procedures for NMR spectroscopy of urine, plasma, serum and tissue extracts. Nat Protoc. 2007;2:2692–2703.
- [14] Kim HK, Choi YH, Verpoorte R. NMR-based metabolomic analysis of plants. Nat Protoc. 2010;5:536–549.
- [15] Kim J, Oh J, Sung GH. MAP kinase Hog1 regulates metabolic changes induced by hyperosmotic stress. Front Microbiol. 2016;7:732.
- [16] Hounsa CG, Brandt EV, Thevelein J, et al. Role of trehalose in survival of *Saccharomyces cerevisiae* under osmotic stress. Microbiology. 1998;144:671–680.
- [17] Olz R, Larsson K, Adler L, et al. Energy flux and osmoregulation of *Saccharomyces cerevisiae* grown in chemostats under NaCl stress. J Bacteriol. 1993;175:2205–2213.
- [18] Oren A. Bioenergetic aspects of halophilism. Microbiol Mol Biol Rev. 1999;63:334–348.