

Set-theoretic Kripke-style Semantics for Weakly Associative Substructural Fuzzy Logics*

Eunsuk Yang

【Abstract】 This paper deals with Kripke-style semantics, which will be called set-theoretic Kripke-style semantics, for weakly associative substructural fuzzy logics. We first recall three weakly associative substructural fuzzy logic systems and then introduce their corresponding Kripke-style semantics. Next, we provide set-theoretic completeness results for them.

【Key Words】 (Set-theoretic) Kripke-style semantics, Relational semantics, Fuzzy logic, Substructural logic, Weakly associative logic.

1. Introduction

The aim of this paper is to introduce *set-theoretic* Kripke-style semantics for weakly associative substructural fuzzy logic. For this, we first note that Yang introduced two kinds of (binary) Kripke-style semantics, i.e., algebraic and non-algebraic Kripke-style semantics, for logics with pseudo-Boolean (briefly, pB) and de Morgan (briefly, dM) negations in Yang (2015b), logics with weak-Boolean (briefly, wB) negations, which can be regarded as paraconsistent logics in Yang (2014b, 2015a), and weakening-free non-commutative substructural fuzzy logics in Yang (2016a, 2018b). Recently, Yang (2018a) further introduced *algebraic* Kripke-style semantics for some weakly associative substructural fuzzy logics. However, he did not consider *set-theoretical* semantics for them. Thus, it is not clear whether this semantics works for weakly associative substructural fuzzy logic systems. (Note that as Kripke's works show in Kripke (1963, 1965a, 1965b), this semantics is interesting in the sense that it can treat logics to which algebraic semantics cannot be applied.)

This is a tough question because Kripke-style semantics for well-known substructural core fuzzy logic systems are algebraic, but not set-theoretical. As Yang mentioned in Yang (2014a), after algebraic semantics for t-norm¹⁾ (based) logics were introduced, their corresponding algebraic Kripke-style semantics have been introduced. For instance, after algebraic semantics for monoidal

¹⁾ T-norms are commutative, associative, increasing, binary functions with identity 1 on the real unit interval $[0,1]$.

t-norm (based) logics were introduced by Esteva and Godo in Esteva & Godo (2001), their corresponding algebraic Kripke-style semantics were introduced in Montagna & Ono (2002), Montagna & Sacchetti (2003; 2004), and Diaconescu & Georgescu (2007).²⁾

These facts give rise to the following question:

- Can we introduce set-theoretical Kripke-style semantics for weakly associative substructural fuzzy logics?

The answer to the question is positive in the sense that we can provide such Kripke-style semantics for the weakly associative substructural fuzzy logics introduced in Yang (2016b, 2018b). For this, first, in Section 2 we recall the wta-monoidal uninorm logic **WAMUL** and its two axiomatic extensions and then introduce their corresponding Kripke-style semantics. In Section 3, using set-theoretic method, we provide soundness and completeness results for them..

For convenience, we shall adopt the notations and terminology similar to those in Cintula (2006), Metcalfe & Montagna (2007), Montagna & Sacchetti (2003; 2004), and Yang (2012; 2014a; 2016a), and we assume reader familiarity with them (along with results found therein).

²⁾ See Yang (2018b) for more examples.

2. Weakly associative substructural fuzzy logics and Kripke-style semantics

Here we first briefly recall the systems introduced in Yang (2016b) as preliminaries. Weakly associative substructural fuzzy logics are based on a countable propositional language with formulas Fm built inductively as usual from a set of propositional variables VAR , binary connectives \rightarrow , $\&$, \wedge , \vee , and constants \mathbf{T} , \mathbf{F} , \mathbf{f} , \mathbf{t} , with defined connectives:

$$\text{df1. } \neg\phi := \phi \rightarrow \mathbf{f},$$

$$\text{df2. } \phi \leftrightarrow \psi := (\phi \rightarrow \psi) \wedge (\psi \rightarrow \phi).$$

We may define \mathbf{t} as $\mathbf{f} \rightarrow \mathbf{f}$. We moreover define ϕ_t^n as $\phi_t \& \dots \& \phi_t$, n factors, where $\phi_t := \phi \wedge \mathbf{t}$. For the rest of this paper, we use the customary notations and terminology, and the axiom systems to provide a consequence relation.

We start with the following axiom schemes and rules for the weak \mathbf{t} -associative monoidal uninorm logic $\mathbf{WA}_t\mathbf{MUL}$ and its two axiomatic extensions.

Definition 2.1 (Yang (2016b))

(i) $\mathbf{WA}_t\mathbf{MUL}$ consists of the following axiom schemes and rules:

$$\text{A1. } \phi \rightarrow \phi \quad (\text{self-implication, SI})$$

$$\text{A2. } (\phi \wedge \psi) \rightarrow \phi, (\phi \wedge \psi) \rightarrow \psi \quad (\wedge\text{-elimination, } \wedge\text{-E})$$

$$\text{A3. } ((\phi \rightarrow \psi) \wedge (\phi \rightarrow \chi)) \rightarrow (\phi \rightarrow (\psi \wedge \chi)) \quad (\wedge\text{-introduction, } \wedge\text{-I})$$

- A4. $\phi \rightarrow (\phi \vee \psi), \psi \rightarrow (\phi \vee \psi)$ (\vee -introduction, \vee -I)
 A5. $((\phi \rightarrow \chi) \wedge (\psi \rightarrow \chi)) \rightarrow ((\phi \vee \psi) \rightarrow \chi)$ (\vee -elimination, \vee -E)
 A6. $\mathbf{F} \rightarrow \phi$ (ex falsum quodlibet, EF)
 A7. $(\phi \& \psi) \rightarrow (\psi \& \phi)$ ($\&$ -commutativity, $\&$ -C)
 A8. $\phi \leftrightarrow (\mathbf{t} \rightarrow \phi)$ (push and pop, PP)
 A9. $\phi \rightarrow (\psi \rightarrow (\psi \& \phi))$ ($\&$ -adjunction, $\&$ -Adj)
 A10. $(\phi_t \& \psi_t) \rightarrow (\phi \wedge \psi)$ ($\&$ \wedge)
 A11. $(\psi \& (\phi \& (\phi \rightarrow (\psi \rightarrow \chi)))) \rightarrow \chi$ (residuation, Res')
 A12. $(\phi \rightarrow ((\phi \& (\phi \rightarrow \psi)) \& (\psi \rightarrow \chi))) \rightarrow (\phi \rightarrow \chi)$ (T')
 A13. $((\delta \& \varepsilon) \rightarrow (\delta \& (\varepsilon \& (\phi \rightarrow \psi)_t))) \vee (\delta' \rightarrow (\varepsilon' \rightarrow ((\varepsilon' \& \delta') \& (\psi \rightarrow \phi)_t)))$

(PL)

- A14. $(\phi_t \& (\psi_t \& \chi_t)) \leftrightarrow ((\phi_t \& \psi_t) \& \chi_t)$ (weak \mathbf{t} -associativity, \mathbf{wAS}_t)
 $\phi \rightarrow \psi, \phi \vdash \psi$ (modus ponens, mp)
 $\phi \vdash \phi_t$ (adj_u)
 $\phi \vdash (\delta \& \varepsilon) \rightarrow (\delta \& (\varepsilon \& \phi))$ (α)
 $\phi \vdash \delta \rightarrow (\varepsilon \rightarrow ((\varepsilon \& \delta) \& \phi))$ (β).

(ii) The following are weakly associative substructural fuzzy logics that axiomatically extend $\mathbf{WA}_t\mathbf{MUL}$:

- \mathbf{t} -associative (ta-) monoidal uninorm logic $\mathbf{A}_t\mathbf{MUL}$ is $\mathbf{WA}_t\mathbf{MUL}$ plus

- (\mathbf{AS}_t) $(\phi \& (\psi \& \chi))_t \leftrightarrow ((\phi \& \psi) \& \chi)_t$;
 (\mathbf{RE}_t) $(\phi \rightarrow (\psi \rightarrow \chi))_t \leftrightarrow ((\phi \& \psi) \rightarrow \chi)_t$;
 (\mathbf{SF}_t) $(\phi \rightarrow \psi)_t \rightarrow ((\psi \rightarrow \chi) \rightarrow (\phi \rightarrow \chi))$;
 (\mathbf{PF}_t) $(\psi \rightarrow \chi)_t \rightarrow ((\phi \rightarrow \psi) \rightarrow (\phi \rightarrow \chi))$; and
 (\mathbf{MT}_t) $(\phi \rightarrow \psi)_t \rightarrow ((\phi \& \chi) \rightarrow (\psi \& \chi))$.

- Strong ta-monoidal uninorm logic $\mathbf{SA}_t\mathbf{MUL}$ is $\mathbf{A}_t\mathbf{MUL}$ plus
 (\mathbf{sAS}_t) $(\phi_t \& (\psi \& \chi)) \leftrightarrow ((\phi_t \& \psi) \& \chi)$.

For easy reference, we let L_s be a set of weakly associative substructural fuzzy logics defined previously.

Definition 2.2 $L_s = \{\mathbf{WA}_t\mathbf{MUL}, \mathbf{A}_t\mathbf{MUL}, \mathbf{SA}_t\mathbf{MUL}\}$.

In $L \in L_s$, \mathbf{f} can be defined as $\neg \mathbf{t}$ and vice versa.

A *theory* over $L \in L_s$ is a set T of formulas. A *proof* in a sequence of formulas whose each member is either an axiom of L or a member of T or follows from some preceding members of the sequence using a rule of L . $T \vdash \phi$, more exactly $T \vdash_L \phi$, means that ϕ is *provable* in T w.r.t. L , i.e., there is an L -proof of ϕ in T . A theory T is *inconsistent* if $T \vdash \mathbf{F}$; otherwise it is *consistent*.

The deduction theorem for L is as follows:

Proposition 2.3 Let T be a theory, and ϕ, ψ formulas.

(i) (Cintula, Horčík, & Noguera (2013, 2015)) $T \cup \{\phi\} \vdash_L \psi$ iff $T \vdash_L \forall(\phi) \rightarrow \psi$ for some $\forall \in \Pi(\mathbf{bDT}^*)$.³⁾

(ii) (Yang (2009)) For $L \in \{\mathbf{A}_t\mathbf{MUL}, \mathbf{SA}_t\mathbf{MUL}\}$, $T \cup \{\phi\} \vdash_L \psi$ iff there is n such that $T \vdash_L \phi_t^n \rightarrow \psi$.

For convenience, “ \neg ,” “ \wedge ,” “ \vee ,” and “ \rightarrow ” are used ambiguously as propositional connectives and as frame operators, but context should clarify their meanings.

Next we provide Kripke-style semantics for L_s . First, Kripke

³⁾ For \forall and $\Pi(\mathbf{bDT}^*)$, see Cintula, Horčík, & Noguera (2013; 2015) and Yang (2015a).

frames are defined as follows.

Definition 2.4 (i) (Kripke frame) A *Kripke frame* is a structure $\mathbf{X} = (X, \top, \perp, t, f, \leq, *, \rightarrow)$ such that $(X, \top, \perp, t, f, \leq, *, \rightarrow)$ is a linearly ordered pointed bounded commutative rlu-groupoid.⁴⁾ The elements of \mathbf{X} are called *nodes*.

(ii) (MICAL frame) An *MICAL frame* is a Kripke frame, where $*$ is conjunctive (i.e., $\perp * \top = \perp$) and left-continuous (i.e., whenever $\sup\{x_i : i \in I\}$ exists, $x * \sup\{x_i : i \in I\} = \sup\{x * x_i : i \in I\}$), and so its residuum \rightarrow is defined as $x \rightarrow y := \sup\{z: x * z \leq y\}$ for all $x, y \in X$.

Definition 2.5 (L frame) A *WA_tMUL frame* is an MICAL frame satisfying $(wAS_t^{A'}) x_t * (y_t * z_t) = (x_t * y_t) * z_t$, for all $x, y, z \in A$; an *A_tMUL frame* is an MICAL frame satisfying $(AS_t^{A'}) (x * (y * z))_t = ((x * y) * z)_t$, for all $x, y, z \in A$; and an *SA_tMUL frame* is an A_tMUL frame satisfying $(sAS_t^{A'}) x_t * (y * z) = (x_t * y) * z$, for all $x, y, z \in A$. We call all these frames *L frames*.

Definition 2.4 (ii) ensures that an MICAL frame has a supremum w.r.t. $*$, i.e., for every $x, y \in X$, the set $\{z: x * z \leq y\}$ has the supremum. \mathbf{X} is said to be *complete* if \leq is a complete order.

An *evaluation* or *forcing* on a set-theoretic Kripke frame is a

⁴⁾ For more detailed interpretation of the notations for rlu-groupoids, see Galatos et al. (2007).

relation \Vdash between nodes and propositional variables, and arbitrary formulas subject to the conditions below: for every propositional variable p ,

(AHC) if $x \Vdash p$ and $y \leq x$, then $y \Vdash p$;

(min) $\perp \Vdash p$; and

for arbitrary formulas,

(t) $x \Vdash \mathbf{t}$ iff $x \leq \mathbf{t}$;

(f) $x \Vdash \mathbf{f}$ iff $x \leq \mathbf{f}$;

(\perp) $x \Vdash \mathbf{F}$ iff $x = \perp$;

(\wedge) $x \Vdash \phi \wedge \psi$ iff $x \Vdash \phi$ and $x \Vdash \psi$;

(\vee) $x \Vdash \phi \vee \psi$ iff $x \Vdash \phi$ or $x \Vdash \psi$;

($\&$) $x \Vdash \phi \& \psi$ iff there are $y, z \in X$ such that $y \Vdash \phi$, $z \Vdash \psi$, and $x \leq y * z$;

(\rightarrow) $x \Vdash \phi \rightarrow \psi$ iff for all $y \in X$, if $y \Vdash \phi$, then $x * y \Vdash \psi$.

An evaluation or forcing on an L frame is an evaluation or forcing further satisfying that (max) for every atomic sentence p , $\{x : x \Vdash p\}$ has a maximum.

Definition 2.6 (i) (Kripke model) A *Kripke model* is a pair (\mathbf{X}, \Vdash) , where \mathbf{X} is a Kripke frame and \Vdash is a forcing on \mathbf{X} .

(ii) (L model) An *L model* is a pair (\mathbf{X}, \Vdash) , where \mathbf{X} is an L frame and \Vdash is a forcing on \mathbf{X} . an L model (\mathbf{X}, \Vdash) is said to

be *complete* if \mathbf{X} is a complete frame and \Vdash is a forcing on \mathbf{X} .

Definition 2.7 Given a Kripke model (\mathbf{X}, \Vdash) , a node x of \mathbf{X} and a formula ϕ , we say that x *forces* ϕ to express $x \Vdash \phi$. We say that ϕ is *true* in (\mathbf{X}, \Vdash) if $t \Vdash \phi$, and that ϕ is *valid* in the frame \mathbf{X} (expressed by $\mathbf{X} \models \phi$) if ϕ is true in (\mathbf{X}, \Vdash) for every forcing \Vdash on \mathbf{X} .

3. Soundness and completeness for Ls

We first introduce two lemmas, which can be easily proved:

Lemma 3.1 (Cf, Yang (2016b)) (Hereditary Lemma, HL) Let \mathbf{X} be a Kripke frame. For any sentence ϕ and for all nodes $x, y \in \mathbf{X}$, if $x \Vdash \phi$ and $y \leq x$, then $y \Vdash \phi$.

Lemma 3.2 $t \Vdash \phi \rightarrow \psi$ iff for all $x \in \mathbf{X}$, if $x \Vdash \phi$, then $x \Vdash \psi$.

We then provide soundness and completeness results for Ls.

Proposition 3.3 (Soundness, Yang (2018b)) If $\vdash_{\mathbf{L}} \phi$, then ϕ is valid in every L frame.

Now we provide completeness results for Ls using set-theoretical Kripke-style semantics. A theory T is said to be *linear* if, for each pair ϕ, ψ of formulas, we have $T \vdash \phi \rightarrow \psi$

or $T \vdash \psi \rightarrow \phi$. By an L-theory, we mean a theory T closed under rules of L . As in relevance logic, by a regular L-theory, we mean an L-theory containing all of the theorems of L . Since we have no use of irregular theories, by an L-theory, we henceforth mean an L-theory containing all of the theorems of L .

Moreover, where T is a linear L-theory, we define the *canonical L frame* determined by T to be a structure $\mathbf{X} = (X_{\text{can}}, \top_{\text{can}}, \perp_{\text{can}}, t_{\text{can}}, f_{\text{can}}, \leq_{\text{can}}, *_{\text{can}})$, where $\top_{\text{can}} = \{\phi : T \vdash_L \mathbf{T} \rightarrow \phi\}$, $\perp_{\text{can}} = \{\phi : T \vdash_L \mathbf{F} \rightarrow \phi\}$, $t_{\text{can}} = T$, $f_{\text{can}} = \{\phi : T \vdash_L \mathbf{f} \rightarrow \phi\}$, X_{can} is the set of linear L-theories extending t_{can} , \leq_{can} is \supseteq restricted to X_{can} , i.e, $x \leq_{\text{can}} y$ iff $\{\phi : x \vdash_L \phi\} \supseteq \{\phi : y \vdash_L \phi\}$, and $*_{\text{can}}$ is defined as $x *_{\text{can}} y := \{\phi \ \& \ \psi : \text{for some } \phi \in x, \psi \in y\}$ satisfying groupoid properties corresponding to L frames on $(X_{\text{can}}, t_{\text{can}}, \leq_{\text{can}})$. Note that the base t_{can} is constructed as the linear L-theory that excludes nontheorems of L , i.e., excludes ϕ such that $\not\vdash_L \phi$. The partial orderedness and the linear orderedness of the canonical L frame depend on \leq_{can} restricted on X_{can} . Then, first, the following is obvious.

Proposition 3.4 A canonical L frame is linearly ordered.

Proof: Since \leq_{can} is an order reversed subset relation, it is obvious that a canonical L frame is partially ordered. For linearly orderedness, suppose for contradiction that neither $x \leq_{\text{can}} y$ nor $y \leq_{\text{can}} x$. Then, there exist ϕ, ψ such that $\phi \in y$, $\phi \notin x$, $\psi \in x$, and $\psi \notin y$. Since t_{can} is a linear theory, we have that $\phi \rightarrow \psi \in t_{\text{can}}$ or $\psi \rightarrow \phi \in t_{\text{can}}$. Let $\phi \rightarrow \psi \in t_{\text{can}}$. Then, since ϕ

$\rightarrow \psi \in y$, by (mp), we have $\psi \in y$, a contradiction. The case, where $\psi \rightarrow \phi \in t_{\text{can}}$, is analogous. \square

Next, we define a canonical evaluation as follows:

$$(a) \ x \Vdash_{\text{can}} \phi \text{ iff } \phi \in x.$$

We then consider the following two lemmas.

Lemma 3.5 $t_{\text{can}} \Vdash_{\text{can}} \phi \rightarrow \psi$ iff for all $x \in X_{\text{can}}$, if $x \Vdash_{\text{can}} \phi$, then $x \Vdash_{\text{can}} \psi$.

Proof: By (a), we instead show that $\phi \rightarrow \psi \in t_{\text{can}}$ iff for all $x \in X_{\text{can}}$, if $\phi \in x$, then $\psi \in x$. For the left-to-right direction, we assume $\phi \rightarrow \psi \in t_{\text{can}}$ and $\phi \in x$, and show $\psi \in x$. By the suppositions and the definition of $*_{\text{can}}$, we have that $\phi \& (\phi \rightarrow \psi) \in x *_{\text{can}} t_{\text{can}} = x$. Then, since $(\phi \& (\phi \rightarrow \psi)) \rightarrow \psi \in t_{\text{can}}$ and thus $(\phi \& (\phi \rightarrow \psi)) \rightarrow \psi \in x$, we also obtain that $\psi \in x$ by (mp). For the right-to-left direction, suppose contrapositively that $\phi \rightarrow \psi \notin t_{\text{can}}$. Set $x_0 = \{Z : \text{there exists } X \in t_{\text{can}} \text{ and } t_{\text{can}} \vdash X \rightarrow (\phi \rightarrow Z)\}$. Clearly, $x_0 \supseteq t_{\text{can}}$, $\phi \in x_0$, and $\psi \notin x_0$. (Otherwise, $t_{\text{can}} \vdash X \rightarrow (\phi \rightarrow \psi)$; therefore, $t_{\text{can}} \vdash \phi \rightarrow \psi$, a contradiction, by (mp), since $t_{\text{can}} \vdash X$.)

Then, by the Linear Extension Property of Theorem 12.9 in Cintula, Horčík, & Noguera (2015), we have a linear theory $x \supseteq x_0$ with $\psi \notin x$; therefore $\phi \in x$ but $\psi \notin x$. \square

Lemma 3.6 (Canonical Evaluation Lemma) \Vdash_{can} is an evaluation.

Proof: First, consider the conditions for propositional variables.

For (AHC), we need to show that: for every propositional variable p ,

$$\text{if } x \Vdash_{\text{can}} p \text{ and } y \leq_{\text{can}} x, \text{ then } y \Vdash_{\text{can}} p.$$

Assume that $x \Vdash_{\text{can}} p$ and $y \leq_{\text{can}} x$. By (a), we have that $p \in x$ and $x \subseteq y$, and thus $p \in y$; therefore, $y \Vdash_{\text{can}} p$.

For (min), we need to show that: for every propositional variable p ,

$$\perp_{\text{can}} \Vdash_{\text{can}} p.$$

By (a), we need to show that $p \in \perp_{\text{can}}$. Since $\perp_{\text{can}} = \{\phi : T \vdash_{\perp} \mathbf{F} \rightarrow \phi\}$, we have that $p \in \perp_{\text{can}}$; therefore, $\perp_{\text{can}} \Vdash_{\text{can}} p$ by (a).

We next consider the conditions for propositional constants \mathbf{t} , \mathbf{f} , and \mathbf{F} .

For (t), we need to show that:

$$x \Vdash_{\text{can}} \mathbf{t} \text{ iff } x \leq_{\text{can}} t_{\text{can}}.$$

Since $t_{\text{can}} = T$ and x is a theory extending T , we can ensure that $\mathbf{t} \in x$ iff $x \supseteq t_{\text{can}}$; therefore, $x \Vdash_{\text{can}} \mathbf{t}$ iff $x \leq_{\text{can}} t_{\text{can}}$ by

(a).

The proof for (f) and (\perp) is analogous to that of (t).

Now we consider the conditions for arbitrary formulas.

For (\wedge), we need to show

$$x \Vdash_{\text{can}} \phi \wedge \psi \text{ iff } x \Vdash_{\text{can}} \phi \text{ and } x \Vdash_{\text{can}} \psi.$$

By (a), we instead show that $\phi \wedge \psi \in x$ iff $\phi \in x$ and $\psi \in x$. The left-to-right direction follows from (\wedge -E) and (mp). The right-to-left direction follows from (adj).

For (\vee), we must show

$$x \Vdash_{\text{can}} \phi \vee \psi \text{ iff } x \Vdash_{\text{can}} \phi \text{ or } x \Vdash_{\text{can}} \psi.$$

By (a), we instead show that $\phi \vee \psi \in x$ iff $\phi \in x$ or $\psi \in x$. The left-to-right direction follows from the fact that linear theories are also prime theories in L (see Cintula & Noguera (2011)). The right-to-left direction follows from (\vee -I) and (mp).

For ($\&$), we need to show

$$x \Vdash_{\text{can}} \phi \& \psi \text{ iff there are } y, z \in X \text{ such that } y \Vdash_{\text{can}} \phi, z \Vdash_{\text{can}} \psi, \text{ and } x \leq_{\text{can}} y *_{\text{can}} z.$$

The definition of $*_{\text{can}}$ ensures that $\phi \& \psi \in x$ iff there are $y, z \in X$ such that $\phi \in y$, $\psi \in z$, and $x \leq_{\text{can}} y *_{\text{can}} z$. Then, by (a), we obtain the claim.

For (\rightarrow), we need to show

$x \Vdash_{\text{can}} \phi \rightarrow \psi$ iff for all $y \in X$, if $y \Vdash_{\text{can}} \phi$, then $x \ast_{\text{can}} y \Vdash_{\text{can}} \psi$.

By (a), we instead show that $\phi \rightarrow \psi \in x$ iff for all $y \in X$, if $\phi \in y$, then $\psi \in x \ast_{\text{can}} y$. For the left-to-right direction, we assume $\phi \rightarrow \psi \in x$ and $\phi \in y$, and show $\psi \in x \ast_{\text{can}} y$. The definition of \ast_{can} ensures $(\phi \rightarrow \psi) \& \phi \in x \ast_{\text{can}} y$. Since $((\phi \rightarrow \psi) \& \phi) \rightarrow \psi \in t_{\text{can}}$ and thus $((\phi \rightarrow \psi) \& \phi) \rightarrow \psi \in x \ast_{\text{can}} y$, by (mp), we have that $\psi \in x \ast_{\text{can}} y$. For the right-to-left direction, suppose contrapositively that $\phi \rightarrow \psi \notin x$. As in Lemma 3.5, we can construct a linear theory y such that $\phi \in y$ and $\psi \notin x \ast_{\text{can}} y$. \square

Let us call a model $\mathbf{M}, = (\mathbf{X}, \Vdash_{\text{can}})$ (i.e., $(X_{\text{can}}, \top_{\text{can}}, \perp_{\text{can}}, t_{\text{can}}, f_{\text{can}}, \leq_{\text{can}}, \ast_{\text{can}}, \Vdash_{\text{can}})$), for L , an L model. Then, by Lemma 3.6, the canonically defined $(\mathbf{X}, \Vdash_{\text{can}})$ is an L model. Thus, since, by construction, t_{can} excludes our chosen nontheorem ϕ , and the canonical definition of models agrees with membership, we can state that, for each nontheorem ϕ of L , there is an L model in which ϕ is $t_{\text{can}} \not\vdash_L \phi$. It gives us the weak completeness of L as follows.

Theorem 3.7 (Weak completeness) If $\models_L \phi$, then $\vdash_L \phi$.

Furthermore, using Lemma 3.6 and the Linear Extension Property, we can show the strong completeness of L as follows.

Theorem 3.8 (Strong completeness) L is strongly complete w.r.t. the class of all L -frames.

4. Concluding remark

Here we investigated set-theoretic Kripke-style semantics for some weakly associative substructural fuzzy logics. Note that, while Yang provided algebraic semantics for other non-associative substructural fuzzy logics in Yang (2015a, 2016c, 2017a, 2017b), he did not consider set-theoretic Kripke-style semantics for those systems. To provide such semantics for these logics remains a problem to be solved.

References

- Cintula, P. (2006), “Weakly Implicative (Fuzzy) Logics I: Basic properties”, *Archive for Mathematical Logic* 45: pp. 673-704.
- Cintula, P., Horčík, R., and Noguera, C. (2013), “Non-associative substructural logics and their semilinear extensions: axiomatization and completeness properties”, *Review of Symbol. Logic*, 12, pp. 394-423.
- Cintula, P., Horčík, R., and Noguera, C. (2015), “The quest for the basic fuzzy logic”, *Mathematical Fuzzy Logic*, P. Hájek (Ed.), Springer.
- Cintula, P. and Noguera, C. (2011), A general framework for mathematical fuzzy logic, *Handbook of Mathematical Fuzzy Logic*, vol 1, P. Cintula, P. Hájek, and C. Noguera (Eds.), London, College publications, pp. 103-207.
- Diaconescu, D. and Georgescu, G. (2007), “On the forcing semantics for monoidal t-norm based logic”, *Journal of Universal Computer Science* 13: pp. 1550-1572.
- Esteva, F. and Godo, L. (2001), “Monoidal t-norm based logic: towards a logic for left-continuous t-norms”, *Fuzzy Sets and Systems* 124: pp. 271-288.
- Galatos, N., Jipsen, P., Kowalski, T., and Ono, H., (2007). *Residuated lattices: an algebraic glimpse at substructural logics*, Amsterdam, Elsevier.
- Kripke, S. (1963). “Semantic analysis of modal logic I: normal modal propositional calculi”, *Zeitschrift für mathematische Logik und Grundlagen der Mathematik* 9, pp. 67-96.
- Kripke, S. (1965a). “Semantic analysis of intuitionistic logic I”,

- Formal systems and Recursive Functions*, J. Crossley and M. Dummett (Eds.), Amsterdam, North-Holland Publ. Co., pp. 92-129.
- Kripke, S. (1965b). "Semantic analysis of modal logic II", *The theory of models*, J. Addison, L. Henkin and A. Tarski (Eds.), Amsterdam, North-Holland Publ. Co., pp. 206-220
- Metcalf, G., and Montagna, F. (2007), "Substructural Fuzzy Logics", *Journal of Symbolic Logic*, 72, pp. 834-864.
- Montagna, F. and Ono, H. (2002) "Kripke semantics, undecidability and standard completeness for Esteva and Godo's Logic MTL_{\forall} ", *Studia Logica*, 71, pp. 227-245.
- Montagna, F. and Sacchetti, L. (2003) "Kripke-style semantics for many-valued logics", *Mathematical Logic Quarterly*, 49, pp. 629-641.
- Montagna, F. and Sacchetti, L. (2004) "Corrigendum to "Kripke-style semantics for many-valued logics", *Mathematical Logic Quarterly*, 50, pp. 104-107.
- Yang, E. (2009) "Non-associative fuzzy-relevance logics: strong t-associative monoidal uninorm logics", *Korean Journal of Logic*, 12(1), pp. 89-110.
- Yang, E. (2012) "Kripke-style semantics for UL", *Korean Journal of Logic*, 15 (1), pp. 1-15.
- Yang, E. (2014a) "Algebraic Kripke-style semantics for weakening-free fuzzy logics", *Korean Journal of Logic*, 17 (1), pp. 181-195.
- Yang, E. (2014b) "Algebraic Kripke-style semantics for relevance logics", *Journal of Philosophical Logic*, 43, pp. 803-826.
- Yang, E. (2015a) "Weakening-free, non-associative fuzzy logics: micanorm-based logics", *Fuzzy Sets and Systems*, 276, pp.

43-58.

- Yang, E. (2015b) “Some an axiomatic extension of the involutive micanorm logic IMICAL”, *Korean Journal of Logic*, 18(2), pp. 197-215.
- Yang, E. (2016a) “Algebraic Kripke-style semantics for substructural fuzzy logics”, *Korean Journal of Logic* 19(2), pp. 295-322.
- Yang, E. (2016b) “Weakly associative fuzzy logics”, *Korean Journal of Logic* 19(3), pp. 437-461.
- Yang, E. (2016c) “Basic substructural core fuzzy logics and their extensions: Mianorm-based logics”, *Fuzzy Sets and Systems* 276, pp. 43-58.
- Yang, E. (2017a) “Involutive basic substructural core fuzzy logics: Involutive mianorm-based logics”, *Fuzzy Sets and Systems*, 320, pp. 1-16.
- Yang, E. (2017b) “A non-associative generalization of continuous t-norm-based logics”, *Journal of Intelligent & Fuzzy Systems* 33, pp. 3743-3752.
- Yang, E. (2018a) “Set-theoretical Kripke-style semantics for an extension of HpsUL, CnHpsUL*”, *Korean Journal of Logic* 21(1), pp. 39-57.
- Yang, E. (2018b) “Algebraic Kripke-style semantics for weakly associative fuzzy logics”, *Korean Journal of Logic* 21(2), pp. 155-173.

전북대학교 철학과, 비판적사고와논술연구소

Department of Philosophy & Institute of Critical Thinking and Writing, Chonbuk National University

eunsyang@jbnu.ac.kr

약한 결합 원리를 갖는 준구조 퍼지 논리를 위한 집합 이론적 크립키형 의미론

양 은 석

이 글에서 우리는 (곱 연언 &의) 약한 형식의 결합 원리를 갖는 준구조 퍼지 논리를 위한 집합 이론적 크립키형 의미론을 연구한다. 이를 위하여 먼저 약한 결합 원리를 갖는 세 준구조 퍼지 논리 체계들을 상기한 후 이 체계들에 상응하는 크립키형 의미론을 소개한다. 다음으로 집합 이론적 방식을 이용하여 이 체계들이 완전하다는 것을 보인다.

주요어: (집합 이론적) 크립키형 의미론, 관계 의미론, 퍼지 논리, 약한 결합 원리, 준구조 논리