

RESEARCH NOTE

Dermea piceina (Dermateaceae): An Unrecorded Endophytic Fungus of Isolated from *Abies koreana*


Ju-Kyeong Eo^{1*}, Eunsu Park¹, and Han-Na Choe²¹Division of Climate and Ecology, Bureau of Conservation & Assessment Research, National Institute of Ecology, Seocheon 33657, Korea²Biological Resource Center, Korea Research Institute of Bioscience and Biotechnology, Jeongseup 56212, Korea

*Corresponding author: abiesendo@gmail.com

ABSTRACT

We found an unrecorded endophytic fungus, *Dermea piceina* J.W. Groves, isolated from alpine conifer *Abies koreana*. Until now only one *Dermea* species, *D. cerasi*, has been reported in Korea. In this study, we compared morphological characteristics and DNA sequences, including internal transcribed spacer and 28S ribosomal DNA, of *D. piceina* isolated from *A. koreana* with those of related species. Here, we present morphological and molecular characters of this fungus for the first time in Korea.

Key word: *Abies koreana*, *Dermea piceina*, Endophytic fungi, Korea



OPEN ACCESS

pISSN : 0253-651X
eISSN : 2383-5249Kor. J. Mycol. 2020 December, 48(4): 485-489
<https://doi.org/10.4489/KJM.20200046>**Received:** October 30, 2020**Revised:** October 30, 2020**Accepted:** December 15, 2020

© 2020 THE KOREAN SOCIETY OF MYCOLOGY.



This is an Open Access article distributed under the terms of the Creative Commons Attribution-Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

The genus *Dermea* Fr. contains 24 species worldwide [1]. Until now only one species, *D. cerasi*, had been discovered in Korea on fallen branches in the national park of Byeonsanbando [2]. Since then, there was no additional record of *Dermea* species therein. The apothecium of *Dermea* is very hard, leathery, dark brown to black in color and has a clavate ascus with eight ascospores. In the asexual stage, the macroconidium is sickle-shaped or filiform with 0-3 septa, and the microconidium is rod or filiform without septa [3].

Alpine conifers are vulnerable to climate change [4]. National Institute of Ecology (NIE) has selected seven vulnerable species - *Abies koreana* E. H. Wilson, *Abies nephrolepis* (Trautv.) Maxim., *Juniperus chinensis* var. *sargentii* A. Henry, *Picea jezoensis* (Siebold et Zucc.) Carriere, *Pinus pumila* (Pall.) Regel., *Taxus cuspidata* Siebold et Zucc. *Thuja koraiensis* Nakai - and is studying their basic ecology to help conserve these species [5]. In this context, study of the biodiversity of endophytic fungi on firs is important. Through this, we have found an unrecorded endophytic fungus, *Dermea piceina*, on *A. koreana* in Korea and reported it here.

Needle leaves of *A. koreana* were harvested from Mt. Halla (33° 20' N, 126° 31' E, 1,952 m) in the Jeju

Special Self-Governing Province of Korea in 2018. The samples were transported in a zipper bags, and fungi were isolated within 24 h. All samples were washed with tap water and cut into 1-cm pieces. Surface sterilization was performed; samples were immersed in 96% ethyl alcohol for 1 min, sodium hypochlorite for 3 min, and then 96% ethyl alcohol for 30 s; finally, they were washed twice with sterilized water. Each sample was placed on potato dextrose agar (PDA; MBcell, Seoul, Korea) and incubated in the dark for 4 weeks at 25°C to isolate endophytic fungi [6]. PDA and maltose extract agar (MEA; MBcell, Seoul, Korea) media were used for the pure culture of endophytic fungi. The macroscopic and microscopic of the fungus were measured by light microscope (DM2500, Leica Microsystems, Wetzlar, Germany). The fungus was deposited in the Korean Collection for Type Cultures (KCTC).

Genomic DNA was extracted from the fungus using a plant tissues genomic DNA extraction kit (Xi'an Tianlong Science & Technology, Shaanxi, Taiwan) following the manufacturer's instructions. Polymerase chain reaction (PCR) was performed using primers ITS1 and LR3, which can selectively amplify from the internal transcribed spacer 1 (ITS1) region 1 to the D2 region of 28S ribosomal DNA [7,8]. The PCR conditions were as follows: Pre-denaturing for 5 min at 94°C with one cycle, denaturing for 30 s at 94°C, annealing for 30 s at 50°C, extending for 1 min at 72°C in 30 cycles, and then finally stabilizing for 10 min at 72°C for one cycle. The PCR product was confirmed by electrophoresis using 1.5% agarose gel.

DNA sequencing was supplied to Macrogen (Seoul, Korea), and the resulting sequence was then identified based on similarity with the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>) using the basic local alignment search tool. A maximum-likelihood tree was generated by MEGA 10.0.5 based on the Kimura-2 parameter distance model with the 1,000-times bootstrap method [9].

***Dermea piccina* J.W. Groves**

[MB#286055]

This fungus grew slowly on the chosen media. Colony diameter after 14 days was <1.0-2.0 mm on PDA and MEA. However, the colony diameter after 180 days was 15.5-19.6 mm on PDA and 21.5-30.4 mm on MEA. The mycelium was denser in PDA than in MEA. The surface color was light amberish gray (MEA) (Munsell color notation: 5Y 8/2) to light vermilionish gray (PDA) (Munsell color notation: 7.5YR 8/2) with an irregular margin, velvety texture at the center to scant at the margin, and no exudates. The reverse was grayish gamboge (MEA) (Munsell color notation: 2.5Y 7/4) to light brownish gray (PDA) (Munsell color notation: 2.5Y 9/2) [10]. Macroconidia measured $97.6\text{-}109.6 \times 4.0\text{-}6.6 \mu\text{m}$ ($n=20$), were strongly curved to sickle-shaped with pointed ends, hyaline, dyed well with lactophenol cotton blue, and displays 0-3 septa inside. Microconidia were not observed. (Fig. 1, Table 1).

Specimen examined: Mt. Halla, Jeju Special Self-Governing Province, Korea, 16 October 2018., isolated




Fig. 1. Cultures of *Dermea piceina* strain NIE7236 isolated from *Abies koreana*. A, B, front and reversed sides of the colonies on potato dextrose agar after 180 days; C, D, front and reversed sides of the colonies on maltose extract agar after 180 days; E, macroconidia. Scale bars: E=50 µm.

Table 1. Morphological characteristics of *Dermea piceina* NIE7236 isolated from needle leaves of *Abies koreana*.

Strain	<i>D. piceina</i> NIE7236	<i>D. piceina</i> [3]	<i>D. cerasi</i> [3]
Colony	on MEA & PDA, 25°C, 180 days	No observation	No observation
color	on MEA, light amberish gray; reverse grayish gamboge on PDA, light vermilionish gray; reverse light brownish gray	No observation	No observation
Size	on MEA, 21.5-30.4 mm; on PDA, 15.5-19.6 mm in diameter after 180 days	No observation	No observation
Shape	Velvety texture at the center to scanty pale yellow mycelium, margin irregular	No observation	No observation
Conidia	Macroconidia hyaline, 0-3 septa, strongly curved to sickle-shaped with pointed ends, 97.6-109.6×4.0-6.6 µm in diameter No observation	Macroconidia hyaline, 0-3 septa, elongate-fusiform, strongly curved to nearly straight, pointed at the ends 22.0-40.0×3.0-5.0 µm in diameter Microconidia hyaline, aseptate, filiform, strongly curved, ends rounded, 9.0-15.0×1.0-1.5 µm in diameter	Macroconidia hyaline to faintly greenish, 0-1 septa, elongate-fusiform to subfiliform, sickle-shaped or sigmoid to almost straight, pointed at the ends 35.0-65.0×2.5-4.5 µm in diameter Microconidia hyaline, aseptate, filiform, almost straight or curved, 12.0-23.0×1.0-1.5 µm in diameter

PDA, potato dextrose agar; MEA, maltose extract agar.

from leaves of *Abies koreana*, strain NIE7236, KCTC no. 56711, GenBank no. MW186178 for ITS rDNA

Note: *Dermea* species were identified by morphological characters (ascospore number, conidia size etc.) and host plants [3]. However, at present, host specificity is ambiguous because some studies reported their multiple host plants [11,12]. *D. piceina* was found on *Picea* sp. in its first report by Groves [3], whereas in our study, it was discovered on *Abies* sp.. Recently, *Dermea chinensis* C. M. Tian & N. Jiang, isolated from Chinese red birch (*Betula albosinensis* Burkill), was reported as a new species in China [1]. Most

studies of *Dermea* spp. have been carried out in North America, Europe, and India [3] so host specificity of genus *Dermea* was thought to show bias due to insufficient research in other regions. The DNA sequence of *D. piceina* in the present study was analyzed from ITS region to the D2 region of 28S ribosomal DNA and revealed 96.8% similarity to *D. piceina* (MH856143.1) (Figure 2). The genus *Dermea* is associated with nineteen host plant genera: *Abies*, *Acer*, *Amelanchier*, *Betula*, *Capparis*, *Chionanthus*, *Fraxinus*, *Hamamelis*, *Ilex*, *Libocedrus*, *Nemopanthus*, *Picea*, *Pinus*, *Prunus*, *Pseudotsuga*, *Rhamnus*, *Sorbus*, *Tsuga*, *Viburnum* [1,3,12]. Through this study, we present the first report of *D. piceina* in Korea, and thus two species of *Dermea* - *D. cerasi* and *D. piceina* - are distributed in Korea. Many preceding studies of *Dermea* spp. focused on apothecium (ascospore) in the bark or leaf, but lack information about the morphology of colony in the media. Therefore, further studies are needed for exact measurements of their morphological characters from cultures maintained under controlled laboratory conditions.




Fig. 2. Phylogenetic tree of *Dermea piceina* strain NIE7236 isolated from *Abies koreana*. The internal transcribed spacer region, including 5.8S ribosomal DNA, and 28S ribosomal DNA including D1 and D2 region, were used for the sequence analysis to confirm the topological appropriation of the fungal isolates. *Diploccarpon earlianum* was used as an out-group and bootstrap values are shown at the branches (1,000 replicates).

ACKNOWLEDGEMENTS

This study was supported with funds from the National Institute of Ecology under project No. NIE-C-2018-19 and the Ministry of Environment of Korea as a part of basic ecological research

REFERENCE

1. Jiang N, Tian CM. Re-collection of *Dermea prunus* in China, with a description of *D. chinensis* sp. nov. MycoKeys 2019;50:79-91.
2. Cho DH. The mycoflora of higher fungi in Pyonsan Peninsula National Park. The Report of the Korean Association for Conservation of Nature 1995;34:167-93.
3. Groves JW. North American species of *Dermea*. Mycologia 1946;38:351-431.
4. Kong WS, Kim K, Lee S, Park H, Cho SH. Distribution of high mountain plants and species vulnerability against climate change. J Environ Impact Assess 2014;23:119-36.
5. Lee JH, Shin HS, Cho HJ, Yun CW. Subalpine conifer forest communities. Seocheon: National Institute of Ecology; 2014.
6. Eo JK, Kim CK, Lee HB, Eom AH. Diversity of endophytic fungi isolated from *Pinus densiflora* and *Larix kaempferi* in Mt. Oser, Korea. Kor J Mycol 2013;41:137-41.
7. White TJ, Bruns T, Lee S, Taylor J. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. PCR protocols: A guide to methods and applications. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ, editors. San Diego: Academic Press, Inc.; 1990. p. 315-22.
8. Hopple Jr JS, Vilgalys R. Phylogenetic relationships among coprinoid taxa and allies based on data from restriction site mapping of nuclear rDNA. Mycologia 1994;86:96-107.
9. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular evolutionary genetics analysis across computing platforms. Mol Biol Evol 2018;35:1547-9.
10. Munsell Color. Munsell book of color glossy edition. Grand Rapids: Munsell Color; 2012.
11. Funk A. The genus *Dermea* and related conidial states on Douglas fir. Can J Bot 1976;54:2852-6.
12. Mehrabi M, Asgari B, Wijayawardene NN, Hyde KD. Description of *Dermea persica* (Dermateaceae, Helotiales), a new asexual Ascomycete from Iran, and an updated key to *Dermea* species. Phytotaxa 2018;367:25-37.