

Optimization of Ascorbic Acid Extraction from Rugosa Rose (*Rosa rugosa* Thunb.) Fruit Using Response Surface Methodology and Validation of the Analytical Method¹

Min UM² · Ji-Woo KIM² · Jae-Won LEE^{2,†}

ABSTRACT

In this study, ultrasound-assisted extraction was performed to extract ascorbic acid from rugosa rose (*Rosa rugosa* Thunb.) fruit. The optimal conditions were investigated by response surface methodology, using two variable including reaction time (16-44 min) and temperature (16-44°C). The ascorbic acid extraction was sensitive to the reaction time rather than the reaction temperature, and the optimal conditions for ascorbic acid extraction were 25°C and 30 min. Ascorbic acid and gallic acid in the rugosa rose fruit extract were completely separated by HPLC, with a resolution factor of over 1.5 between the two. The correlation coefficient of the ascorbic acid was 0.999 in a linearity test for 50-150 µg/mL concentration of extract. The limit of detection and limit of quantification values were 0.16 µg/mL and 29.89 µg/mL, respectively. The relative standard deviations (RSD) for repeatability and reproducibility were determined, and each RSD showed good precision at less than 5% (N=6).

Keywords: rugosa rose, ultrasound-assisted extraction, response surface methodology, ascorbic acid, validation

1. INTRODUCTION

Antioxidant compounds in plants have attracted attention due to their potential value as food preservation agents or nutraceuticals (Kim *et al.*, 2010; Kim *et al.*, 2014). Rose species have been used for food and medicinal purposes in many countries. Rose leaves and flowers have been used mainly in cosmetics and perfumes. Rose fruits are used in many food products and drinks including teas, jellies, jams, and alcoholic beverages (Grochowski, 1990; Zhang *et al.*, 2008). The rose fruits

have a high antioxidant content and are enriched in vitamin C (ascorbic acid) (Czyzowska *et al.*, 2015; Um *et al.*, 2018). Vitamin C is essential for humans but is not synthesized in the body. It is acquired by ingestion. Although the rose fruit extracts in food have been confirmed to act as health supplements, the research is insufficient.

The rugosa rose (*Rosa rugosa* Thunb.) is a halophyte originating from East Asia. It has traditionally been used as food to prevent intractable diseases, such as diabetes (Cho *et al.*, 2004). It is naturally grow well

¹ Date Received February 25, 2020, Date Accepted April 28, 2020

² Department of Wood Science and Engineering, College of Agriculture and Life Sciences, Chonnam National University, Gwangju 61186, Republic of Korea

† Corresponding author: Jae-Won LEE (e-mail: ljw43376@chonnam.ac.kr, ORCID: 0000-0003-2528-1905)

in various environments, from coastal to inland habitats, and is resistant to harsh climatic conditions and salinity. Thus, this plant is expected to have various bioactive compounds. Several studies on the antioxidant activity of the rugosa rose have been reported, which have used different extraction methods and investigated different parts of the plant (Demir *et al.*, 2014; Kim *et al.*, 2018; Um *et al.*, 2018).

Extraction is the initial step in the process of obtaining bioactive compounds from plants (Jung *et al.*, 2017; Manurung *et al.*, 2019; Min *et al.*, 2019). The extraction conditions differ depending on the target bioactive compound in the same plant. Various extraction methods have been developed to separate bioactive components from plants. Soxhlet extraction, heating reflux extraction, and maceration have been used as traditional extraction methods (Kim *et al.*, 2017). However, these methods require high temperatures and are time-consuming, and thus their extraction efficiency is low in terms of the energy consumed during extraction (Yang *et al.*, 2013). In order to overcome these disadvantages, ultrasound-assisted extraction (UAE) was suggested. This method improves the extraction efficiency by increasing the contact area between the solvent and the target compounds present in the cell wall of the plant (Rodríguez-Rojo *et al.*, 2012). The method has the advantage of rapidly extracting target compounds using a small amount of solvent. The extract yield differs depending on several variables that include the ultrasonic power, extraction temperature and time, and solvent type. It is important that the optimal extraction conditions for each target compound be determined, taking into account the several variables on which UAE depends (Vilkhu *et al.*, 2008; Wang *et al.*, 2008).

UAE has higher extraction efficiency than conventional methods at the laboratory scale. However, as the extraction scale increases, the energy required for UAE increases rapidly (Preece *et al.*, 2017). Therefore, it is important to determine the scale at which UAE

has maximum extraction efficiency at the most feasible absorbed power density and absorbed energy density in the extraction efficiency.

In this study, we performed UAE to extract ascorbic acid from the rugosa rose fruit. The conditions of solvent concentration (50% ethanol) and liquid/solid ratio (1:20) for UAE were the same as described previously (Um *et al.*, 2018). Based on these parameters, the optimal UAE conditions were investigated by response surface methodology, using the two independent variables of reaction time and temperature. In addition, we validated the ascorbic acid extract to ensure the reliability of our results. The specificity, linearity, limit of detection, limit of quantification, precision, and robustness were assessed according to the guidelines of the International Conference on Harmonization (ICH) were conducted for method validation to determine ascorbic acid extraction. Finally, we analyzed the extraction efficiency by taking into account the absorbed power density (W/mL) and absorbed energy density (J/mL) for UAE at the laboratory scale.

2. MATERIALS and METHODS

2.1. Materials and ultrasound-assisted extraction (UAE) processing

The rugosa rose was grown at Chonnam National University (35°10'25.6"N, 126°54'00.8"E, South Korea) and harvested in November 2016. After harvesting, the fruit was washed in water and freeze-dried for 2 days. The dried fruit was then ground to pass through a 40 mesh sieve and stored at 4°C in a sealed state.

The extraction from the rugosa rose fruit was carried out in an ultrasonic water bath (NXP-1505P, KODO Technical Research Co., Ltd., Hwaseong, Korea) operating at an ultrasonic power and frequency of 220 W and 40 kHz, respectively.

2.2. Analysis of ascorbic acid using high performance liquid chromatography (HPLC)

Ascorbic acid (No 23024, 99.7-100.5%) and gallic acid (No G7384, 97.5-100.5%) were purchased from Sigma Aldrich (St. Louis, MO, USA). Methanol for HPLC analysis was purchased from Duksan Pure Chemicals Co. (Ansan, Korea). The ascorbic acid was analyzed by HPLC (e2695, Separations module, Waters Co., Milford, MA, USA), equipped with a SunFire® C18 column (SKU: 186002560 Waters Co.) and a UV/Vis Detector (2489, Waters Co.). The analysis was performed at 254 nm and a column temperature of 30°C for 10 min (Nováková *et al.*, 2008). In addition, 0.1% phosphoric acid (w/v) in 30% methanol was used as the mobile phase, and the flow rate was 0.6-1.0 mL/min.

2.3. Design and analysis of extraction condition using response surface methodology (RSM)

A central composite design was used to optimize the extraction of ascorbic acid from rugosa rose fruit (Nováková *et al.*, 2008). The temperature (X1, 16-44°C), reaction time (X2, 16-44 min) and five replicates at the central point were chosen as the independent variables by central composite design. The ascorbic acid concentration (Y1, mg/g) was used as the dependent output variable. Further, the other extraction conditions selected were the ethanol concentration (50%) and liquid/solid ratio (1:20), which were in accordance with a previous study (Um *et al.*, 2018). After each extraction process, the extract was separated from solid fraction by vacuum filtration and subsequently concentrated using a rotary evaporator. The concentrated extract was then freeze-dried for further analysis. Design Expert 10 (State Ease Inc., Minneapolis, MN, USA) was used for model building and data analysis.

2.4. Method validation

The specificity, linearity, limit of detection (LOD), limit of quantification (LOQ), precision, and robustness were assessed according to the guidelines of the International Conference on Harmonization (ICH) for method validation in order to determine the presence of ascorbic acid in the extract obtained by RSM at optimal conditions (Shabir, 2003; Sahu *et al.*, 2018).

The specificity was evaluated by HPLC as described above. Specificity implies the property of accurately distinguishing the target component (ascorbic acid) from other components by HPLC. Methanol was used as solvent. Ascorbic acid and gallic acid were used as the standard components to determine the chromatogram resolution. The standard component concentration was 100 µg/mL.

The linearity was evaluated for the extracts diluted at five different concentrations (50, 75, 100, 125, and 150 µg/mL). The ascorbic acid concentration in each diluted sample was measured by HPLC as described above. The LOD and LOQ were calculated for the analyses in six repeats.

The precision was evaluated to ensure repeatability of the extraction process. The precision for ascorbic acid was determined by HPLC for three different extract concentration (50 µg/mL, 100 µg/mL, and 150 µg/mL), by different analysts (analyst 1 and analyst 2), and at different dates (day 1 and day 2). The precision experiments were carried out both intra-day (within a day) and inter-day (between days).

The robustness was evaluated to assess the change in the relative standard deviation (RSD) by modification factors such as the column temperature (30°C and 35°C), solvent type (different manufacturers), and flow rate (0.6 mL/min and 1.0 mL/min). The ascorbic acid was detected by HPLC in six repeats for each condition.

2.5. Statistical analysis

All experiments in this study were performed in more than triplicate (RSM $N = 3$, Validation $N = 6$). Two way ANOVA was performed to determine the significant differences between mean values. The Pearson correlation coefficient was used to determine the correlation between the variables used, and the differences were considered significant at a p -value < 0.05 .

2.6. Energy consumption for extraction

The absorbed power density (APD, W/mL) and absorbed energy density (AED, J/mL) were investigated according to the UAE scale (solid/solvent: 2 g/40 mL, 5 g/100 mL, 10 g/200 mL, and 20 g/400 mL) (Chan *et al.*, 2013), using the equations (1)-(3).

$$APD = Q/(V \times t_H) \quad (1)$$

$$Q = m_L C_p \Delta T \quad (2)$$

$$AED = APD \times t \quad (3)$$

Q: total heat absorbed in the solvent during the UAE heating (J) (Incropera *et al.*, 2007)

V: solvent volume (mL)

t_H : UAE heating time (min)

m_L : initial mass of the extraction solvent

C_p : heat capacity of the extraction solvent

ΔT : temperature differential after UAE

t: extraction time (min)

3. RESULTS and DISCUSSION

3.1. Optimization for ascorbic acid extraction by RSM design and analysis

The independent variables (solid/solvent ratio, temperature, time, and extraction method) that define the extraction conditions can have either a concerted or an independently effect on target compound extraction

Table 1. Ascorbic acid extraction from rugosa rose fruit by central composite design

Run	Temperature (°C), X_1	Time (min), X_2	Response (Y)
			Ascorbic acid (mg/g)
1	30	30	6.48
2	40	40	5.67
3	30	30	6.33
4	30	30	6.58
5	20	40	5.45
6	30	44	5.63
7	30	30	6.39
8	20	20	4.58
9	40	20	4.80
10	30	30	6.82
11	16	30	6.68
12	30	16	4.02
13	44	30	5.93

(Vaez *et al.*, 2012). In previous study, the optimum solvent concentration for ascorbic acid extraction from rugosa rose fruit was suggested to be 50% ethanol (Noordin *et al.*, 2004; Rodríguez-Rojo *et al.*, 2012). In this study, ascorbic acid was extracted from rugosa rose fruit using a central composite design that considered two other independent variables: temperature and time. The experimental results according to the extraction conditions are shown in Table 1. The highest ascorbic acid concentration was 6.82 mg/g when the rugosa rose fruit underwent extraction at 30°C for 30 min. On the other hand, the extract concentration was lowest, 4.02 mg/g at 30°C and an extraction time of 16 min. These results suggest that ascorbic acid extraction is sensitive to the reaction time rather than the reaction temperature. This is due to the ascorbic acid extraction increase as the time increases at the same reaction temperature.

The ANOVA analysis of the experiment results is shown in Table 2. The p -value was used to confirm the significance of each factor ($p < 0.05$ was considered statistically significant). The ascorbic acid concentration was affected by the independent variable

Table 2. ANOVA analysis of central composite design for optimizing ascorbic acid extraction from rugosa rose fruit

Source	Sum of squares	Degree of freedom	Mean square	<i>f</i> -value	<i>p</i> -value	
Model	8.50	5	1.70	13.96	0.0016	Significant
A ^a	0.05	1	0.05	0.40	0.5495	
B ^b	2.02	1	2.02	16.55	0.0048	
AB	0.00	1	0.00	0.00	1.0000	
A ²	0.33	1	0.33	2.70	0.1443	
B ²	6.38	1	6.38	52.34	0.0002	
Residual	0.85	7	0.12			
Lack of Fit	0.70	3	0.23	6.34	0.0532	Not significant
Pure Error	0.15	4	0.04			
Cor. Total ^c	9.36	12				
SD ^d	0.35		R ²	0.908841		
Mean	5.80		Adj R ²	0.843727		


^a Temperature^b Time^c Total all information corrected for the mean^d Standard deviation

of time (p -value < 0.05). However, the interaction between the two variables (temperature and time) showed no significant effects, with the p -value being higher than 0.05 (p -value = 1.00).

For this model the ‘‘Lack of Fit F-test’’ was 6.34, which implied that the lack of fit was not significant relative to the pure error. To fit the model, the lack of fit needs to be non-significant. The model calculated for the response (Y : ascorbic acid concentration, X_1 : temperature, X_2 : time) is represented in equation (4).

$$Y = 6.52 - 0.07758X_1 + 0.50211X_2 + X_1X_2 - 0.2175X_1^2 - 0.9575X_2^2 \quad (4)$$

The equation (4) represents the response surface of the result in Fig. 1. The temperature had no significant influence on the experimental design compared to the time, indicating that a high temperature is not necessary to obtain a high concentration of ascorbic acid. These results are similar to those of

**Fig. 1.** Response surface plot presenting effect of temperature and time for ascorbic acid extraction from rugosa rose fruit.

a previous study (Um *et al.*, 2018). However, the ascorbic acid extraction was sensitive to the reaction time, because the extract concentration decreased significantly for a relatively short or long reaction time.

The extraction conditions corresponding to a max-

imum concentration of ascorbic acid were incorporated in the experimental design. In general, as the temperature increases, the movement of molecules accelerates, and the diffusivity of the bioactive compounds improves, leading to higher extract yields (Yang *et al.*, 2010). However, ascorbic acid is sensitive to temperature changes, as it decomposes at high temperatures (Van den Broeck *et al.*, 1998). Therefore, high temperatures are not suited for the extraction of ascorbic acid. Based on the experimental design, the optimal temperature and time for ascorbic acid extraction from rugosa rose fruit were 25°C and 30 min (6.50 mg/g), respectively. The experimental result was 6.55 mg/g, indicating that the models can mathe-

matically present the ascorbic acid extraction from rugosa rose fruit.

3.2 Method validation

The specificity, linearity, limit of detection (LOD), limit of quantification (LOQ), precision, and robustness were assessed in order to confirm the detection of ascorbic acid according to the guidelines of the International Conference on Harmonization (ICH) (Sahu *et al.*, 2018).

The specificity for ascorbic acid is shown in Fig. 2. Ascorbic acid and gallic acid were used as standard chemicals to confirm the detected peaks from rugosa




Fig. 2. Specificity for ascorbic acid using HPLC at 100 µg/mL extract; (A) methanol, (B) rugosa rose fruit extract, (C) ascorbic acid and gallic acid standard, (D) ascorbic acid, (E) gallic acid.

rose fruit extract. Ascorbic acid and gallic acid from rugosa rose fruit extract were detected in an HPLC chromatogram, and the major peak was that for ascorbic acid (Fig. 2B). The retention times of the two compounds were in agreement with those of the standard chemicals (Fig. 2C). It was confirmed that the peaks were detected without interaction between the two components. The resolution factor and resolution limit value between the two chemicals were 1.85 and 1.56, respectively. The ascorbic acid and gallic acid in the rugosa rose fruit extract were completely separated by HPLC analysis with a resolution factor of over 1.5 between the two, confirming HPLC to be suitable for assessing the ascorbic acid in the extract (Song and Wang, 2003).

Ascorbic acid concentration in the extract ranged from 50-150 µg/mL. The correlation between the ascorbic acid concentration and the peak areas was confirmed. The ascorbic acid concentration and the peak areas showed a linear proportional relationship. The correlation coefficient (R^2) for the extracted ascorbic acid was 0.999 (Table 3). The LOD and LOQ values were calculated based on the standard deviation

and the slope of linearity curve which are shown in Table 3.

The precision of the analytical method in detecting ascorbic acid was assessed by analyzing repeatability relative standard deviation (RSD, intra-day) and a reproducibility relative standard deviation (RSD, inter-day) (Table 4) (Zhao *et al.*, 2016). Each RSD showed good precision at less than 5% (N=6). The ascorbic acid extracted from rugosa rose fruit can be considered to provide accurate quantitative analysis data at the proposed concentrations (50-150 µL) (Singh *et al.*, 2006).

Robustness refers to the reliability of the analysis. This property is determined by intentionally changing the variables, such as flow rate, column temperature, and solvent manufacturer, during analysis (Vander Heyden *et al.*, 2001). In this study, experimental robustness determinations were carried out for different solvent manufacturers (A and B), flow rates (1.0 and 0.6 mL/min), and column temperatures (30 and 35°C) (Table 5). The flow rate had the strongest influence on the ascorbic acid analysis. The peak shape was relatively wide at a flow rate 0.6 mL/min of flow rate

Table 3. Linearity range, limit of detection and limit of quantification for ascorbic acid from rugosa rose fruit

Linearity matrix	Ascorbic acid
Linear range (µg/mL)	50-150
Correlation coefficient, R	0.999
Coefficient of determination, R^2	0.999
y-axis intercept	2370
Slope	793
Retention time (min)	2.623
N ^a	6
Equation (Y)	7.93e+002X + 2.37e+003
LOD ^b (µg/mL)	0.1598
LOQ ^c (µg/mL)	29.8865

^a Number of repeated experiment

^b Limit of detection

^c Limit of quantification

Optimization of Ascorbic Acid Extraction from Rugosa Rose (*Rosa rugosa* Thunb.) Fruit Using Response Surface Methodology and Validation of the Analytical Method

Table 4. Repeatability relative standard deviation and reproducibility relative standard deviation for ascorbic acid from rugosa rose fruit

Rugosa rose extract		Retention time (min)	Area (μ AU)	Ascorbic acid concentration (μ g/mL)	Ascorbic acid (%)
Intra-day (Day 1, analyst 1)					
50 μ g/mL	Average	2.62	51594	0.27	73.46
	SD ^a	0.00	1523	0.01	1.80
	RSD ^b	0.02	2.95	3.70	2.45
100 μ g/mL	Average	2.625	105604	0.54	77.94
	SD	0.00	308	0.00	1.95
	RSD	0.07	0.29	0.81	2.50
150 μ g/mL	Average	2.62	157362	2.68	79.52
	SD	0.00	1442	0.06	2.09
	RSD	0.06	0.97	2.22	2.63
Intra-day (Day 2, analyst 2)					
50 μ g/mL	Average	2.62	50322	0.28	77.97
	SD	0.01	255	0.00	1.99
	RSD	0.25	0.51	0.54	2.56
100 μ g/mL	Average	2.63	106289	0.57	76.05
	SD	0.01	658	0.03	1.29
	RSD	0.20	0.62	4.76	1.70
150 μ g/mL	Average	2.63	156271	2.63	76.27
	SD	0.00	2301	0.10	2.75
	RSD	0.17	1.47	3.61	3.61
Inter-day (RSD)					
50 μ g/mL		0.16	2.35	3.23	3.96
100 μ g/mL		0.13	0.56	4.56	2.34
150 μ g/mL		0.12	1.16	2.83	3.61

^a Standard deviation

^b Relative standard deviation (%)

Table 5. Validation of robustness depending on solvent manufacturer, flow rate, and column temperature

		Retention time (min)	Area (μ AU)	Ascorbic acid concentration (μ g/mL)	Ascorbic acid (%)	
Solvent manufacturer	A	2.62	106010	0.56	74.14	
		(0.00)	(450)	(0.02)	(3.02)	
	B	2.61	105796	0.55	73.86	
		(0.00)	(287)	(0.01)	(1.46)	
	Average		2.62	105903	0.55	74.00
	SD ^a		0.01	357	0.02	2.13
RSD ^b		0.29	0.34	2.79	2.87	

Table 5. (Continued)

		Retention time (min)	Area (μ AU)	Ascorbic acid concentration (μ g/mL)	Ascorbic acid (%)
Flow rate (mL/min)	0.6	4.32 (0.00)	190584 (5261)	0.51 (0.01)	53.57 (5.57)
	1.0	2.62 (0.00)	105983 (322)	0.56 (0.01)	75.44 (0.28)
	Average	3.47	148284	0.53	64.51
	SD	0.93	46458	0.03	12.49
	RSD	26.82	31.33	4.90	19.36
Column temperature ($^{\circ}$ C)	30	2.62 (0.00)	106270 (685)	0.57 (0.03)	76.27 (0.89)
	35	2.60 (0.00)	106028 (895)	0.55 (0.04)	74.50 (3.82)
	Average	2.61	106149	0.56	75.38
	SD	0.01	725	0.02	2.66
	RSD	0.46	0.68	3.47	3.53

^a Standard deviation

^b Relative standard deviation (%)

The parenthesis contains the standard deviation.

when a C18 column was used for HPLC. The peak exhibited tailing, and the detected ascorbic acid concentration decreased slightly, from 0.56 to 0.51 μ g/mL. On the other hand, the ascorbic acid analysis showed reproducibility for solvent manufacturer and column temperature under the tested conditions. The RSD values were 2.79 and 3.47%, respectively. The RSD values were less than 5% for all conditions. This implies that ascorbic acid can be analyzed stably under the given conditions.

3.3. Extraction efficiency depending on scale based on energy consumption

For UAE carried out at laboratory scale, the absorbed power density (APD) and absorbed energy density (AED) were investigated. APD represents the power (W) used per volume of solvent. The AED conveys the energy (J) when consuming 1 W power

for a certain period of time. The AED and APD for the extraction of ascorbic acid were proportional according to the extraction scale. The amount of energy consumption decreased as the amount of solvent increased. At an extraction scale of 2:40, the AED was 7002 J/mL, and the energy consumption decreased as the extraction scale increased (616 J/mL, 20:400). Similarly, the APD was 4.67 W/mL at 2:40 scale, and its value decreased to 0.41 W/mL when the extraction scale reached 20:400, which corresponds to a 91.22% reduction. However, the ascorbic acid extraction yield decreased as the extraction scale increased (Fig. 3). The ascorbic acid extraction yield was 6.4 mg/g at an extraction scale of 2:40, and 7 mg/g at 20:400, indicating a 9.7% decrease in the extraction yield. At the 20:400 extraction scale, the AED and APD decreased 91.20% (J/mL) and 91.22% (W/mL), respectively, but the ascorbic acid extraction yield decreased by 9.69% compared to that at the 2:40 extraction scale.

Optimization of Ascorbic Acid Extraction from Rugosa Rose (*Rosa rugosa* Thunb.) Fruit Using Response Surface Methodology and Validation of the Analytical Method




Fig. 3. Absorbed power density (APD) and extract yield of ascorbic acid depending on ultrasound-assisted extraction scale.

4. CONCLUSION

ANOVA for the ascorbic acid extraction from rugosa rose fruit using RSM revealed that the optimum conditions for ascorbic acid extraction by UAE were 25°C for 30 min. The validation of the extracts was performed by HPLC using an extract concentration in the range of 50-150 µg/mL. In the extracts, ascorbic acid was detected along with gallic acid, and the resolution factor between the two compounds was over 1.5. The correlation coefficient of ascorbic acid in a linearity test was 0.999. The repeatability and reproducibility RSDs were less than 5%. These results suggest that the method is highly repeatable and reproducible. These results further indicate that the validation method is appropriate for the extraction of ascorbic acid and determination of its concentration. The energy consumption (AED and APD) decreased by more than 90% when the extraction scale was increased 10 times, from 2:40 to 20:400. At the same conditions, the ascorbic acid extraction yield decreased by 9.69%. These results can be used as data to reduce the time and energy required to analyze ascorbic acid from such extracts, and can provide basic data for the development of cosmetics development and functional food processing using preparations from rugosa rose fruit.

REFERENCES

- Chan, C.H., Yusoff, R., Ngoh, G.C. 2013. Modeling and prediction of extraction profile for microwave-assisted extraction based on absorbed microwave energy. *Food Chemistry* 140(1-2): 147-153.
- Cho, E.J., Yokozawa, T., Kim, H.Y., Shibahara, N., Park, J.C. 2004. *Rosa rugosa* attenuates diabetic oxidative stress in rats with streptozotocin-induced diabetes. *The American Journal of Chinese Medicine* 32(4): 487-496.
- Czyzowska, A., Klewicka, E., Pogorzelski, A., Nowak, A. 2015. Polyphenols, vitamin C and antioxidant activity in wines from *Rosa canina* L. and *Rosa rugosa* Thunb. *Journal of Food Composition and Analysis* 39: 62-68.
- Demir, N., Yildiz, O., Alpaslan, M., Hayaloglu, A.A. 2014. Evaluation of volatiles, phenolic compounds and antioxidant activities of rose hip (*Rosa* L.) fruits in Turkey. *LWT - Food Science and Technology* 57(1): 126-133.
- Grochowski, W. 1990. *Ubozna produkcja leśna*. PWN. Warszawa, Poland, pp 379-383.
- Jung, J.Y., Ha, S.Y., Yang, J.K. 2017. Response surface optimization of phenolic compounds extraction from steam exploded oak wood (*Quercus mongolica*). *Journal of the Korean Wood Science and Technology* 45(6): 809-827.
- Kim, J.K., Kwon, D.J., Lim, S.S., Bae, Y.S. 2010. Antioxidant and anti-inflammatory activity of stem bark extracts from *Ulmus davidiana* var. *japonica*. *Journal of the Korean Wood Science and Technology* 38(5): 444-449.
- Kim, J.S., Lee, H.C., Jo, J.S., Jung, J.Y., Ha, Y.L., Yang, J.K. 2014. Evaluation of antioxidant and anticancer activity of steam extract from the bamboo species. *Journal of the Korean Wood Science and Technology* 42(5): 543-554.
- Kim, J.W., Um, M., Lee, J.W. 2018. Antioxidant

- activities of hot water extracts from different parts of Rugosa rose (*Rosa rugosa* Thunb.). Journal of the Korean Wood Science and Technology 46(1): 38-47.
- Kim, S.H., Lee, S.Y., Cho, S.M., Hong, C.Y., Park, S.Y., Park, M.J., Choi, I.G. 2017. Antioxidant activities of *Cryptomeria japonica* leaves extracts by extraction methods. Journal of the Korean Wood Science and Technology 45(5): 495-510.
- Manurung, H., Sari, R.K., Syafii, W., Cahyaningsih, U., Ekasari, W. 2019. Antimalarial activity and phytochemical profile of ethanolic and aqueous extracts of bidara laut (*Strychnos ligustrina* Blum) Wood. Journal of the Korean Wood Science and Technology 47(5): 587-596.
- Min, H.J., Kim, E.J., Shin, S.W., Bae, Y.S. 2019. Antidiabetic activities of Korean red pine (*Pinus densiflora*) inner bark extracts. Journal of the Korean Wood Science and Technology 47(4): 498-508.
- Noordin, M.Y., Venkatesh, V.C., Sharif, S., Elting, S., Abdullah, A. 2004. Application of response surface methodology in describing the performance of coated carbide tools when turning AISI 1045 steel. Journal of Materials Processing Technology 145(1): 46-58.
- Nováková, L., Solich, P., Solichová, D. 2008. HPLC methods for simultaneous determination of ascorbic and dehydroascorbic acids. Trends in Analytical Chemistry 27(10): 942-958.
- Preece, K.E., Hooshyar, N., Krijgsman, A.J., Fryer, P.J., Zuidam, N.J. 2017. Intensification of protein extraction from soybean processing materials using hydrodynamic cavitation. Innovative Food Science and Emerging Technologies 41: 47-55.
- Rodríguez-Rojo, S., Visentin, A., Maestri, D., Cocero, M.J. 2012. Assisted extraction of rosemary antioxidants with green solvents. Journal of Food Engineering 109(1): 98-103.
- Sahu, P.K., Ramiseti, N.R., Cecchi, T., Swain, S., Patro, C.S., Panda, J. 2018. An overview of experimental designs in HPLC method development and validation. Journal of Pharmaceutical and Biomedical Analysis 147: 590-611.
- Shabir, G.A. 2003. Validation of high-performance liquid chromatography methods for pharmaceutical analysis: Understanding the differences and similarities between validation requirements of the US Food and Drug Administration, the US Pharmacopeia and the International Conference on Harmonization. Journal of Chromatography A 987(1-2): 57-66.
- Singh, S., Singh, B., Bahuguna, R., Wadhwa, L., Saxena, R. 2006. Stress degradation studies on ezetimibe and development of a validated stability-indicating HPLC assay. Journal of Pharmaceutical and Biomedical Analysis 41(3): 1037-1040.
- Song, D., Wang, J. 2003. Modified resolution factor for asymmetrical peaks in chromatographic separation. Journal of Pharmaceutical and Biomedical Analysis 32(6): 1105-1112.
- Um, M., Han, T.H., Lee, J.W. 2018. Ultrasound-assisted extraction and antioxidant activity of phenolic and flavonoid compounds and ascorbic acid from rugosa rose (*Rosa rugosa* Thunb.) fruit. Food Science and Biotechnology 27(2): 375-382.
- Vaez, M., Moghaddam, A.Z., Alijani, S. 2012. Optimization and modeling of photocatalytic degradation of azo dye using a response surface methodology (RSM) based on the central composite design with immobilized *Titania Nanoparticles*. Industrial and Engineering Chemistry Research 51(11): 4199-4207.
- Van den Broeck, I., Ludikhuyze, L., Weemaes, C., Van Loey, A., Hendrickx, M. 1998. Kinetics for isobaric – isothermal degradation of L-ascorbic acid. Journal of Agricultural and Food Chemistry 46(5): 2001-2006.

Optimization of Ascorbic Acid Extraction from Rugosa Rose (*Rosa rugosa* Thunb.) Fruit Using Response Surface Methodology and Validation of the Analytical Method

- Vander Heyden, Y., Nijhuis, A., Smeyers-Verbeke, J., Vandeginste, B.G.M., Massart, D.L. 2001. Guidance for robustness/ruggedness tests in method validation. *Journal of Pharmaceutical and Biomedical Analysis* 24(5-6): 723-753.
- Vilkhu, K., Mawson, R., Simons, L., Bates, D. 2008. Applications and opportunities for ultrasound assisted extraction in the food industry-A review. *Innovative Food Science and Emerging Technologies* 9(2): 161-169.
- Wang, J., Sun, B., Cao, Y., Tian, Y., Li, X. 2008. Optimization of ultrasound-assisted extraction of phenolic compounds from wheat bran. *Food Chemistry* 106(2): 804-810.
- Yang, L., Cao, Y.L., Jiang, J.G., Lin, Q.S., Chen, J., Zhu, L. 2010. Response surface optimization of ultrasound-assisted flavonoids extraction from the flower of *Citrus aurantium* L. var. *amara* Engl. *Journal of Separation Science* 33(9): 1349-1355.
- Yang, Y.C., Wei, M.C., Huang, T.C., Lee, S.Z., Lin, S.S. 2013. Comparison of modified ultrasound-assisted and traditional extraction methods for the extraction of baicalin and baicalein from *Radix Scutellariae*. *Industrial Crops and Products* 45: 182-190.
- Zhang, G.Q., Huang, X.D., Wang, H., Leung, A.K.N., Chan, C.L., Fong, E.W.F., Yub, Z.L. 2008. Anti-inflammatory and analgesic effects of the ethanol extract of *Rosa multiflora* Thunb. hips. *Journal of Ethnopharmacology* 118(2): 290-294.
- Zhao, M., Araújo, M.M., Dal, S., Sigrist, S., Bergaentzlé, M., Ramanitrahasimbola, D., Andrianjara, C., Marchioni, E. 2016. Development and validation of a selective and effective pressurized liquid extraction followed by liquid chromatography-mass spectrometry method for the determination of fructosazine analogues in the ammonia treated extract of *Eugenia jambolana* Lamarck seeds. *Journal of Chromatography A* 1473(18): 66-75.