

# The Effects of Clinical Practice Stress and Resilience on Nursing Students' Academic Burnout

Sujin Shin<sup>1</sup>, Eunhee Hwang<sup>2</sup>

<sup>1</sup>College of Nursing, Ewha Womans University, Seoul; <sup>2</sup>Department of Nursing, Wonkwang University, Iksan, Korea

The high degree of academic burnout experienced during academic life indicates that job skill levels during the first year following graduation are low, and the correlation with turnover intention is high. We investigated the effects of clinical practice stress and resilience on nursing students' burnout, and searched for factors that can prevent or control burnout. We recruited a convenience sample of 202 nursing students. Academic burnout, general characteristics, clinical practice stress, and resilience were assessed via self-reported questionnaires. The mean total score of academic burnout was 44.0 points; exhaustion was the highest at 18.5 points, inefficacy was 15.9 points, and cynicism was 9.6 points. High levels of clinical practice stress affected academic burnout ( $\beta=0.194$ ,  $p=0.003$ ), while high resilience was a factor that lowered the degree of academic burnout ( $\beta=-0.449$ ,  $p<0.001$ ). Based on our results, factors affecting students' experiences of academic burnout were clinical practice stress and resilience. We therefore propose the implementation of a new curriculum aimed at increasing satisfaction with the major, reducing clinical practice stress, and increasing resilience, including an efficient peer mentoring program for clinical practice.

**Keywords:** Burnout, Exhaustion, Nursing education

## Corresponding author

Eunhee Hwang  
Department of Nursing, Wonkwang University, 460 Iksan-daero, Iksan 54538, Korea  
Tel: +82-63-850-6071  
Fax: +82-63-850-6060  
E-mail: ehh@wku.ac.kr  
<https://orcid.org/0000-0002-6229-5946>

Received: December 24, 2019

1st revised: February 26, 2020

2nd revised: March 16, 2020

Accepted: March 17, 2020

## INTRODUCTION

Nursing and medical schools require emerging healthcare professionals to practice in the clinical field. Reportedly, 80% of the causes of stress experienced by medical students are academic in nature due to the competitive atmosphere [1], and academic burnout of junior students is highest among nursing students [2]. If this stress persists, it can lead to depression, anxiety, anger, and exhaustion [1]. As is the case with medical students, nursing students who have to complete both theoretical education and clinical practice are also likely to experience a high degree of academic burnout [3].

Academic burnout is defined as emotional exhaustion, cynicism, and inefficacy due to excessive and persistent academic stress [4]. The academic burnout experienced by nursing students often leads to a loss of confidence in nursing and acts as a negative factor leading to frustration with studying within their nursing major [5]. In addition, the high degree of exhaustion experienced during academic life indicates that job skill levels during the first year following graduation are low, and the correlation with turnover intention is high [6]. Therefore, it is necessary to find a support plan that can reduce exhaustion to help nursing students

adapt successfully to university life and transfer this adaptation to their roles as nurses.

Nursing students experience a lot of stress due to the unique characteristics of the nursing department, in addition to the general stress experienced by most university students as they adapt to college life [7]. Due to difficulties with studying vast amounts of information pertaining to their academic major at the time of enrollment, having a less flexible curriculum and a more competitive atmosphere, and undertaking the necessary preparations for the national nursing licensing qualifications [8], high levels of stress lead to academic burnout. Among the various stressors experienced by nursing students, clinical practice stress is a major factor that causes exhaustion in clinical practice [9].

Nursing students must devote more time to their clinical studies because they are required to spend 3 hours per credit to complete one clinical practice course related to their major, as opposed to students of other majors who are required to spend only 1 hour to complete each clinical practice course [10]. In other words, they spend more time studying than students of other majors. Within 2 years, they will undergo 1,000 hours of clinical training and complete shift work in the practice field as early as dawn. This can be a major stress factor. In addition, clinical

practice education exposes students to unfamiliar environments and this often leads to serious anxiety and tension [11]. Most nursing colleges operate a block system in order to meet practice hours. Therefore, nursing students have a relatively narrow curriculum from which to choose compared to students of other majors. The stress related to clinical practice can be a factor in exacerbating the exhaustion of nursing students, so it is necessary to ascertain the current situation of exhaustion and preemptive effort.

Among the variables that can alleviate negative influences such as stress, resilience is considered to help nursing students prepare to face challenges and adversity, and not only survive but also thrive in the face of additional life events and challenges with hope and optimism for future successful outcomes, improved well-being, and career longevity [7,12]. The self-resilient person is one who has the necessary patience and ability to appropriately adjust to the stress level and react flexibly in unfamiliar situations, thus enabling successful adaptation [13]. Ego resilience can reduce the negative impact of psychological exhaustion, and high resilience scores have been reported to have negative correlations with the three dimensions of burnout [14,15]. Resilience is accepted as an important concept in nursing education. Ego resilience promotion leads to positive effects such as a successful university life, the ability to cope with crises and challenges, and the improvement of nursing college students' and faculty's satisfaction [12]. Ego resilience can be considered as a measure to reduce the burnout of nursing students.

The results of a study of nursing students who experienced clinical practice showed that a large portion (49.2%) wanted to change their major, suggesting that there are problems with current nursing education such as excessive academic demands and major maladjustment [16]. Academic burnout affects life satisfaction beyond the negative effects of school life [17]. In order to increase the retention of nursing students, it is needed to reduce academic burnout and increase adaptability to the clinical field. Therefore, it is important to make efforts to minimize academic burnout and to identify the factors that affect it. This proactive approach will help nursing students grow academically, teach nursing students how to increase their own resilience, and ultimately increase nursing student retention.

The purpose of this study was to investigate the effects of clinical practice stress and ego resilience on nursing students' burnout, and to search for factors that can prevent or control

academic burnout. The specific objectives were as follows: (1) understand the subjects' clinical practice stress, resilience, and academic burnout; (2) identify differences in academic burnout according to general characteristics; and (3) analyze the effects of clinical practice stress and resilience on nursing students' academic burnout.

## METHODS

### 1. Study design

This was a descriptive research study conducted to investigate the effects of clinical practice stress and ego resilience on academic burnout among nursing students.

### 2. Study sample

The study sample consisted of 202 nursing students from two universities in different cities. The inclusion criteria were: (1) being third- or fourth-year students and (2) having more than 6 months of clinical practice experience. The exclusion criteria were: (1) having a psychiatric history and (2) having a native language other than Korean.

For the multiple regression analysis, the significance level was 0.05, the power was 0.95, the effect size was 0.15, the number of predictors was 10, and the required number of samples was 172. Considering an inadequate response rate of about 20%, 206 nursing students were chosen based on convenience sampling, and the data from 202 of the students were included in the final analysis.

### 3. Measures

The questionnaire collected data on general characteristics such as age, sex, academic year, mentor status (have/don't have), major satisfaction, intention to transfer, clinical practice stress, ego resilience, and academic burnout.

#### 1) Academic burnout

To measure academic burnout, we used the Maslach Burnout Inventory-Student Survey instrument, which was developed by Schaufeli et al. [4] and validated for Korean students by Shin et al. [18]. This tool consists of 15 items within three sub-factors (i.e., exhaustion, cynicism, and inefficacy). The score is rated on a 5-point Likert scale, with 1 meaning "not at all," and 5 meaning "strongly agree." The inefficacy score is decoded after

measuring competence, and the total score is between 15 and 75 points. The higher the score, the higher the tendency of academic burnout. A study by Schaufeli et al. [4] reported the reliability of Cronbach's  $\alpha$  to be 0.62-0.68, and a study by Shin et al. [18] reported Cronbach's  $\alpha$  to be 0.82-0.86. The reliability of this study was indicated by a Cronbach's  $\alpha$  of 0.80-0.84.

### 2) Clinical practice stress

Clinical practice stress was measured by Beck and Srivastava [19], and was modified by Lee and Kim [20]. This tool is comprised of 24 questions in total. The questionnaire consists of 5 items on the practical training environment, 6 items on undesirable role models, 4 items on the burden of practical work, 4 items on interpersonal conflict, and 5 items on conflict with patients. For each item, the 5-point Likert scale responses ranged from 1 ("not at all") to 5 ("very much agree"); the higher the score, the higher the clinical practice stress level. A study by Lee and Kim [20] reported Cronbach's  $\alpha$  to be 0.91. The reliability of this study was indicated by a Cronbach's  $\alpha$  of 0.85.

### 3) Resilience

To measure the degree of ego resilience, we used Yu and Chae's [21] modification of the scale developed by Block and Kremen [22]. This tool measures responses on a 4-point Likert scale, with a total of 14 possible points, from 1 ("not at all") to 4 ("very much agree"); the higher the average score of all items, the higher the ego resilience level. A study by Block and Kremen [22] reported the reliability of Cronbach's  $\alpha$  to be 0.76, and a study by Yu and Chae [21] reported Cronbach's  $\alpha$  to be 0.67. The reliability of this study was indicated by a Cronbach's  $\alpha$  of 0.80.

### 4. Data collection

This study was conducted by a research assistant trained for ethical consideration of subjects in the classroom who explained to participants the purpose of the study, their guaranteed anonymity and confidentiality before data collection, and the fact that the questionnaire did not affect their grade point averages (GPAs). The questionnaire was distributed to only those students who agreed. A questionnaire collection box was placed at the exit of the classroom and the questionnaires of participants who completed them were voluntarily placed in the collection box. Data collection was conducted from June to August 2018. The

questionnaire completion response time was about 10-15 minutes.

### 5. Data analysis

The collected data were analyzed using the IBM SPSS for Windows ver. 24.0 (IBM Corp., Armonk, NY, USA) program. The differences in the degree of academic burnout according to general characteristics were analyzed using t-tests and analysis of variance, followed by an least significant difference test for post-hoc comparison. Hierarchical multiple regression analysis was used to analyze the factors affecting academic burnout.

### 6. Ethical considerations

This study was conducted in compliance with the Helsinki Declaration and the Good Clinical Practice Guidelines. The study subjects were informed that there were no potential disadvantages if they refused to participate in the research. The purpose of the study, the participation process, and the potential risks and benefits of participating in the research were all explained. After the explanations were complete, written consent was voluntarily submitted to the researchers. In addition, research assistants with no connection to either the professor or the participants completed the data collection process. The respondents were given a small gift as compensation for their time.

## RESULTS

### 1. General characteristics of the subjects

Ninety-six percent of the subjects were female; 46.0% were in their third year, and 54.0% were in their fourth. A majority, 72.3%, of the respondents said that they did not have mentors, and 49.0% of them were satisfied with their majors, while 10.9% of them were unsatisfied. About half, 45.5%, of the respondents had intentions to transfer (Table 1).

### 2. Mean scores for academic burnout, clinical practice stress, and resilience

The mean total score of academic burnout was 44.0 points (possible range, 15-75 points); exhaustion was the highest at 18.5 points (possible range, 5-25 points), inefficacy was 15.9 points (possible range, 5-25 points), and cynicism was 9.6 points (possible range, 5-25 points). Clinical practice stress was 71.8 points (possible range, 24-100 points), and resilience was 38.0 points (possible range, 14-64 points) (Table 2).

### 3. Differences in academic burnout by general characteristics

There was a statistically significant difference in the degree of burnout among general characteristics according to GPA, mentor presence, major satisfaction, and intention to transfer. Specifically, the degree of academic burnout of the third-year students was higher than that of the fourth-year students ( $t=2.56$ ,  $p=0.011$ ), and the degree of academic burnout with a mentor was lower ( $t=-3.54$ ,  $p<0.001$ ). Nursing students who were

dissatisfied with their majors had a high degree of academic burnout ( $F=40.32$ ,  $p<0.001$ ). Moreover, if subjects expressed a desire to transfer, their degree of academic burnout was high ( $t=3.85$ ,  $p<0.001$ ). For the subscales, the third-year students showed higher disability scores than did the fourth-year students ( $t=2.89$ ,  $p=0.004$ ). Additionally, the scores for exhaustion, cynicism, and inefficacy were low with the presence of a mentor and high satisfaction with the major. Students with previous experiences of attempting to transfer or transferring had higher scores on exhaustion and cynicism ( $t=3.05$ ,  $p=0.003$ ;  $t=4.25$ ,  $p=0.001$ ) (Table 3).

**Table 1.** General characteristics of participants (N = 202)

Characteristic	Category	Value
Age (yr)		23.01 ± 5.52
Sex	Male	6 (3.0)
	Female	195 (96.5)
	Missing	1
Academic year	3rd	93 (46.0)
	4th	109 (54.0)
Mentor	Yes	55 (27.2)
	No	146 (72.3)
	Missing	1
Major satisfaction	Satisfied	99 (49.0)
	Moderate	81 (40.1)
	Unsatisfied	22 (10.9)
Intention to transfer to other major or other university	Yes	92 (45.5)
	No	110 (54.5)

Values are presented as mean ± standard deviation or number (%).

### 4. Effect of clinical practice stress and resilience on academic burnout

In order to identify the effects of dependent variables on academic burnout, clinical practice stress, and ego resilience were used as independent variables and regression factors (Table 4). As a result of the regression analysis assumption for the independent variables, the variance inflation factor of the variables included in all models was less than 10, and there was no problem of multi-collinearity. The F statistic value of the regression model was 31.458 ( $p<0.001$ ), and the coefficient of determination  $R^2$  was 0.235, accounting for 23.5% of the academic burnout on

**Table 2.** Descriptive statistics of academic burnout, stress of clinical practicum, and resilience (N = 202)

Variable	Mean ± standard deviation	Minimum	Maximum
Total academic burnout	44.0 ± 7.86	24.0	67.0
Exhaustion	18.5 ± 3.60	8.0	25.0
Cynicism	9.6 ± 3.16	4.0	20.0
Inefficacy	15.9 ± 3.59	7.0	26.0
Stress of clinical practice	71.8 ± 10.4	42.0	97.0
Resilience	38.0 ± 5.68	25.0	56.0

**Table 3.** Differences of burnout according to general characteristics (N = 202)

Variable	Category	Total academic burnout				Exhaustion				Cynicism				Inefficacy			
		Mean ± SD	t or F	p-value	LSD	Mean ± SD	t or F	p-value	LSD	Mean ± SD	t or F	p-value	LSD	Mean ± SD	t or F	p-value	LSD
Academic year	3rd	45.6 ± 7.82	2.56			18.8 ± 3.60	1.08			10.0 ± 3.42	1.92			16.7 ± 3.50	2.89		
	4th	42.7 ± 7.68		0.011		18.2 ± 3.59		0.281		9.2 ± 2.88		0.056		15.3 ± 3.55		0.004	
Mentor	Yes	40.8 ± 8.85	-3.54			17.5 ± 4.03	-2.44			8.7 ± 3.31	-2.38	0.018		14.6 ± 3.81	-3.27		
	No	45.2 ± 7.17		<0.001		18.9 ± 3.37		0.016		9.9 ± 3.04				16.5 ± 3.37		0.001	
Major satisfaction	Satisfied	40.1 ± 6.81	40.32	<0.001	a<b<c	17.4 ± 3.60	10.31	<0.001	a<b<c	8.0 ± 2.36	43.63	<0.001	a<b<c	14.6 ± 3.40	18.79	<0.001	a<b<c
	Moderate	46.7 ± 5.72				19.4 ± 2.66				10.5 ± 2.83				16.8 ± 3.04			
	Unsatisfied	51.1 ± 8.76				20.0 ± 4.96				13.3 ± 3.13				18.8 ± 3.84			
Intention to transfer to other major or other university	Yes	46.3 ± 7.78	3.85	<0.001		19.3 ± 3.56	3.05	0.003		10.6 ± 3.26	4.25	<0.001		16.4 ± 3.51	1.70	0.090	
	No	42.2 ± 7.33				17.8 ± 3.49				8.7 ± 2.83				15.6 ± 3.63			

SD, standard deviation; LSD, least significant difference.

**Table 4.** Effect of clinical practice stress and resilience on academic burnout (N=202)

Variable	$\beta$	t-value	p-value	Variance inflation factor
(Constant)		12.048	<0.001	
Stress of clinical practicum	0.194	3.128	0.002	1.000
Resilience	-0.449	-7.230	<0.001	1.000

Adjusted  $R^2=0.235$ ,  $F=31.458$ ,  $p<0.001$ , Durbin-Watson = 1.787.

variance. High levels of clinical practice stress affected academic burnout ( $\beta=0.194$ ,  $p=0.002$ ), while high resilience was a factor that lowered the degree of academic burnout ( $\beta=-0.449$ ,  $p<0.001$ ).

### DISCUSSION

This study uncovered the factors influencing academic burnout and provided basic data for the establishment of educational strategies aimed at reducing nursing students' burnout.

In a previous study that measured nursing students' academic burnout in Korea using the same tools, the average level of burnout was 2.93 to -3.24 [8,23]. From a comparison with the result of 2.86 points reported by medical students [24], it can be seen that nursing and medical students experience similar levels of academic burnout. On the other hand, according to the results of this study, exhaustion was the highest at 3.70 points, inefficacy was at 3.18, and cynicism was at 1.92; therefore, exhaustion and inefficacy were rather high compared with cynicism. In other previous studies, exhaustion, inefficacy, and cynicism scores were found to be 3.18, 2.77, 2.48, 3.25, 3.39, and 2.13 points, respectively [3,8]. This means that the differences between sub-dimensions were relatively small. The difference between exhaustion and inefficacy scores among the subjects in this study is considered to be due to differences in the subjects. This study was aimed at nursing students in the third and fourth grade, but the subjects of the preceding study were all nursing students. The degree of burnout in the third year was higher than that of the fourth year. In addition, exhaustion and inefficacy were higher in the fourth grades than in the third grades students. This finding in the present study corresponds to previous results.

As a result of analyzing the general characteristics related to burnout, respondents who answered that they were dissatisfied with their major showed higher degrees of burnout, which is consistent with previous research results [3,8]. On the other hand, respondents who answered that they had a mentor had a low degree of burnout, as well as significantly lower exhaustion,

cynicism, and inefficacy. This can be interpreted as a mentor helping the mentee get relief, helping them overcome difficulties, and helping to reduce their anxiety [25]. In addition, degree of burnout was high in the case of intention to transfer. In particular, exhaustion and cynicism scores were high, suggesting that intention to transfer is related to academic burnout. One of the important strategies for nursing students' retention and success is peer mentoring-tutoring [26]. In particular, peer mentoring programs for third- and fourth-year nursing students participating in hands-on training at the university can be considered effective because the learners can benefit from the hidden extracurricular activities [27]. In addition, the higher the clinical practice stress, the higher the degrees of burnout. The results of this study can be interpreted within the same context as those of the previous study, in that the degree of burnout was low when the satisfaction level with the practice was high [3]; the higher the clinical practice stress, the higher the clinical practice exhaustion [9]. Further, nursing students with high academic resilience had low levels of clinical practice stress [16].

In regression analysis, the lower the clinical practice stress and the higher the resilience, the lower the degree of burnout. This is in agreement with the findings of a study finding academic burnout affected by resilience [28]. Nursing students with high levels of academic resilience were more satisfied with their major and experienced less stress related to clinical practice stress [16]. These results identified the importance of the protective role of resilience. Especially, students with high levels of burnout but high levels of resilience were able to maintain their competency and reflect deeply on personal and professional issues despite stress. Reflection has facilitated self-awareness, which is essential to resilience [29]. Teaching strategies using reflection are recommended to prepare students to actively adapt using effective mechanisms for learning in clinical placement [30] and to develop self-awareness and resilience among nursing students at the individual and educational levels.

Therefore, in order to reduce the degree of burnout, it is necessary to lower clinical practice stress, and improve resilience.

Nursing students who participate in effective peer mentoring are reported to experience benefits such as increased competence levels, autonomy and work independence, personal and professional growth, and self-esteem and confidence [28]. In order to induce such a positive experience, it is essential to strengthen resilience through effective mentoring programs, rather than simply recommending or placing students with mentors. Self-efficacy, aggressive stress coping strategy, and social support were found to affect the resilience of nursing students [31]. It is believed that providing training programs that can accumulate successful experiences in solving stressful problems recognized by nursing students can enhance adaptation by enhancing self-efficacy and promoting resilience. The support of professors, clinical educators, and peers will play an important mediator role.

In addition, it is necessary to review the academic achievement evaluation system during curriculum operation in order to alleviate the excessive amounts of the study that cause exhaustion among nursing students, as well as the psychological pressure to maintain high achievement and high GPAs. Assessment should be used to motivate learners and, more importantly, assessments should be made to allow students to experience what they are learning and knowing as they grow academically [32]. Therefore, it is suggested that mentor support and instructor intervention and formation evaluation should be implemented together.

This study has a limitation in that it does not include the educational environment in schools and clinical settings, and thus does not provide a comprehensive analysis of the academic burnout. Therefore, we propose a repeated study that considers a number of factors that affect academic burnout in the future. Despite these limitations, this study is meaningful in that it has identified the factors affecting nursing students' academic burnout and suggests strategies for improving resilience to alleviate nursing students' academic burnout.

### Authors' Contributions

The first author Shin Sujin conceived the study and analyzed the data; the corresponding author Hwang Eunhee originated the basic concept, designed the research, and wrote the manuscript.

## REFERENCES

1. Lee SJ, Choi YJ, Chae H. The effects of personality traits on academic burnout in Korean medical students. *Integr Med Res.* 2017;6(2):207-13.
2. An M, Kang AY, Kim YA, Kim MJ, Kim YL, Kim HW, et al. Comparison of academic engagement, academic burnout, stress, and social support by grade among undergraduate nursing students. *J Korean Soc Sch Health.* 2017;30(2):113-23.
3. Cho HH, Kang JM. Effect of resilience, coping, and mental health on burnout of student nurses. *Child Health Nurs Res.* 2018;24(2):199-207.
4. Schaufeli WB, Martinez IM, Pinto AM, Salanova M, Bakker AB. Burnout and engagement in university students: a cross-national study. *J Cross Cult Psychol.* 2002;33(5):464-81.
5. Ko CM. Mediating effect of stress on relationship between emotional intelligence and burnout among nursing college students. *J Korean Soc Sch Health.* 2015;28(3):239-47.
6. Rudman A, Gustavsson JP. Burnout during nursing education predicts lower occupational preparedness and future clinical performance: a longitudinal study. *Int J Nurs Stud.* 2012;49(8):988-1001.
7. Lee E. Effect of resilience on academic burnout of nursing students. *J Korea Acad Ind Coop Soc.* 2019;20(6):178-87.
8. Lee MK, Lim SH. Relationships among academic burnout, stress coping and college adjustment in nursing students. *Asia Pac J Multimed Serv Converg Art Humanit Sociol.* 2017;7(6):521-34.
9. Lee DY, Park JK, Choi MY. The relation between stress of clinical practice and burnout among nursing students: the mediation effect of spiritual well-being. *J Korean Acad Soc Nurs Educ.* 2017;23(3):300-8.
10. Shin S, Hwang E. Effects of academic resilience and interpersonal competence on the happiness of nursing students. *Korean Med Educ Rev.* 2018;20(3):156-63.
11. Yang YK, Han KS, Bae MH, Yang SH. Social support, academic stress, clinical practice stress in college student of nursing. *Korean J Stress Res.* 2014;22(1):23-34.
12. Stephens TM. Nursing student resilience: a concept clarification. *Nurs Forum.* 2013;48(2):125-33.
13. Block J, Block JH. The role of ego-control and ego-resiliency in the organization of behavior. In: Collins WA, editor. *Development of cognition, affect, and social relations: the Minnesota symposia on child psychology.* Hillsdale (NJ): Erlbaum; 1980. p. 39-101.
14. Lee YM. A moderating effect of academic resilience on the relationship between academic stress and academic burnout. *Think Dev.* 2014;10(4):79-101.
15. Rios-Risquez MI, Garcia-Izquierdo M, Sabuco-Tebar EL, Carrillo-Garcia C, Martinez-Roche ME. An exploratory study of the relationship between resilience, academic burnout and psychological health in nursing students. *Contemp Nurse.* 2016;52(4):430-9.
16. Hwang E, Shin S. Characteristics of nursing students with high levels of academic resilience: a cross-sectional study. *Nurse Educ Today.* 2018;71:54-9.
17. Jung ES, Ha JH. Mediating effects and moderating effects of anticipated risks, anticipated benefits in the relationships between academic burnout and life satisfaction. *J Korea Acad Ind Coop Soc.* 2015;16(9):6009-18.
18. Shin H, Puig A, Lee J, Lee JH, Lee SM. Cultural validation of the



- Maslach Burnout Inventory for Korean students. *Asia Pac Educ Rev.* 2011;12(4):633-9.
19. Beck DL, Srivastava R. Perceived level and sources of stress in baccalaureate nursing students. *J Nurs Educ.* 1991;30(3):127-33.
  20. Lee JE, Kim SL. Relationship among stress, coping strategies, and self-esteem in nursing students taking clinical experience. *J Korean Acad Soc Nurs Educ.* 2005;11(1):98-106.
  21. Yu JH, Chae SJ. Differences in academic burnout and achievement goal orientation by perfectionism of medical students. *J Health Inf Stat.* 2016;41(1):88-95.
  22. Block J, Kremen AM. IQ and ego-resiliency: conceptual and empirical connections and separateness. *J Pers Soc Psychol.* 1996;70(2):349-61.
  23. Yang S, Jung MR. The influences of academic-burnout, self-efficacy, major satisfaction on nursing professionalism in nursing students. *J Learn Cent Curric Instr.* 2016;16(11):613-29.
  24. Yoo SK, Hong SH, Choi BY. Mediating effects of ego-resilience, attachment, and existential spirituality between family risks and adaptation. *Korean J Educ Psychol.* 2004;18(1):393-408.
  25. Szabo S, Lloyd B, McKellar D, Myles H, Newton H, Schutz J, et al. 'Having a mentor helped me with difficult times': a trainee-run mentoring project. *Australas Psychiatry.* 2019;27(3):230-3.
  26. Jeffreys MR. Jeffreys's Nursing Universal Retention and Success model: overview and action ideas for optimizing outcomes A-Z. *Nurse Educ Today.* 2015;35(3):425-31.
  27. Yen-Ju Lin B, Liu PC, Ku KT, Lee CC. Adaptation of medical students during clinical training: effects of holistic preclinical education on clerkship performance. *Teach Learn Med.* 2019;31(1):65-75.
  28. Sibiya MN, Ngxongo TS, Beepat SY. The influence of peer mentoring on critical care nursing students' learning outcomes. *Int J Workplace Health Manag.* 2018;11(3):130-42.
  29. Thomas LJ, Asselin M. Promoting resilience among nursing students in clinical education. *Nurse Educ Pract.* 2018;28:231-4.
  30. Amsrud KE, Lyberg A, Severinsson E. Development of resilience in nursing students: a systematic qualitative review and thematic synthesis. *Nurse Educ Pract.* 2019;41:102621.
  31. Park MH. Factors influencing resilience in nursing students. *J Korea Acad Ind Coop Soc.* 2018;19(11):614-25.
  32. Lee SH, Jeon WT. The relationship between academic self-efficacy and academic burnout in medical students. *Korean J Med Educ.* 2015; 27(1):27-35.